Coexisting relapsing polychondritis and sarcoidosis: an unusual association

Florian Pasquet, Vincent Cottin, Nadia Sivova, Julie Scanff, Christiane Broussolle, Pascal Sève

► To cite this version:

Florian Pasquet, Vincent Cottin, Nadia Sivova, Julie Scanff, Christiane Broussolle, et al.. Coexisting relapsing polychondritis and sarcoidosis: an unusual association. *Rheumatology International*, 2009, 30 (11), pp.1507-1509. 10.1007/s00296-009-1102-5 . hal-00562258

HAL Id: hal-00562258

<https://hal.science/hal-00562258>

Submitted on 3 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Coexisting relapsing polychondritis and sarcoidosis: an unusual association.

Florian Pasquet¹, Vincent Cottin², Nadia Sivova², Julie Le Scanff¹, Christiane Broussolle¹, and Pascal Sève ¹

¹Department of Internal Medicine, Hospices Civils de Lyon, Université Claude Bernard, Lyon, Hôtel-Dieu, 1 place de l'Hôpital, 69288 Lyon Cedex 02, France

²Department of Pneumology, Hospices civils de Lyon, Université de Lyon, Centre de référence des maladies pulmonaires rares, hôpital Louis-Pradel, 69677 Bron Cedex, France.

Address for correspondence and reprint requests: Pascal Sève, MD, PhD. Department of Internal Medicine, Hôtel Dieu, 1 place de l'Hôpital, 69288 Lyon, France.

Tel: (33).4.72.41.33.00. Fax: (33).4.72.41.31.21

e-mail: pascal.seve@chu-lyon.fr.

Abstract

Relapsing polychondritis is an episodic and progressive systemic inflammatory disease characterized by auricular chondritis, polyarthrititis, nasal and respiratory tract chondritis. About 30% of the patients have additional autoimmune and or hematological diseases, most frequently systemic vasculitis, rheumatoid arthritis, myelodysplastic syndromes or systemic lupus erythematosus. So far, only one case of coexisting relapsing polychondritis and sarcoidosis in a patient with AIDS has been reported. We describe here a case of sarcoidosis and relapsing polychondritis in an immunocompetent patient. Physicians should be aware of this possible association.

KEY WORDS: Relapsing polychondritis, sarcoidosis, autoimmunity

Introduction

Relapsing polychondritis (RP) is an episodic and progressive inflammatory disease of the cartilaginous structures, including elastic cartilage of the ear and nose, hyaline cartilage of peripheral joints, fibrocartilage at axial sites, and cartilage of the tracheobronchial tree (1; 2). Inflammation of other proteoglycan-rich structures, such as the eyes, heart, blood vessels, and kidneys may also be present. Diagnostic criteria for RP are listed in Table 1 (1; 2). Although the etiology of RP is unclear, an autoimmune mechanism is considered to be most likely. RP is associated with autoimmune disorders in about one third of the patients, most frequently with connective tissue diseases and vasculitides (3; 4).

Sarcoidosis is a systemic granulomatous disease that most frequently involves the lungs, lymph nodes, skin and eyes (5). An increase in polyclonal gamma-globulin serum level is frequent, with rarely autoantibodies

We report here an unusual observation of coexisting RP and sarcoidosis. To our knowledge, this association in an immunocompetent patient has not been previously reported in the literature.

Case report

A 42-year-old woman was admitted to our hospital with complaints of swelling and pain of the left ear and arthralgia of the hands for 3 weeks. Her past medical history was significant for pulmonary sarcoidosis of 2 years. The diagnosis of sarcoidosis was based on the clinical picture of chronic cough, mediastinal lymphadenopathies, diffuse micronodular interstitial lung disease at chest imaging, and demonstration of typical granulomas at bronchial biopsy. Infectious causes were ruled out. As pulmonary function was normal and chest CT scan was stable, oral corticosteroids were not considered.

Upon examination the significant findings were: left pinna swelling, tenderness and warmth, sparing the non cartilaginous lobe (Fig. 1) and bilateral symmetrical polyarthritis involving wrists and metacarpophalangeal joints. The patient was afebrile. Heart and lung examinations were normal, and there was no evidence of peripheral lymphadenopathy, hepatosplenomegaly or eye involvement. The anamnesis revealed a first episode of chondritis of the right ear 3 years previously, and an episode of nasal chondritis 1 year before, both of which recovered spontaneously. Laboratory results revealed elevated serum level of C-reactive protein of 25 mg/L, and fibrinogen of 4.6g/L, with inflammatory profile of blood proteins electrophoresis. Blood count, liver enzymes, kidney function tests results and serum angiotensin-converting enzyme level were normal. Repeated blood cultures were sterile. Serologic investigations for HIV, hepatitis C and B viruses, *Parvovirus* B19, syphilis and *Borrelia burgdorferi* were negative. Rheumatoid factor, anti anticyclic citrullinated cyclic peptides, antinuclear, anti-double-stranded DNA and type-II anticollagen antibodies were not detected. Serum complement levels were normal. The HLA class II DR4 was not present. Bone radiographs of hands showed no erosions. Spirometry and chest CT results remained unchanged. Fiberoptic bronchoscopy showed absence of laryngeal or tracheal abnormality.

Based on the clinical presentation, the diagnosis of RP was established and treatment with nonsteroid anti-inflammatory agents and colchicine was initiated, with clinical improvement within one week. At 6 months follow-up, she was completely asymptomatic, and physical examination was normal. Fibrinogen and C-reactive protein levels were normal. The CT scan of the chest was unchanged.

Discussion:

According to Mc Adam et al.(1), 25% to 35% of patients with RP have a concurrent disease (Table 2). Coexisting RP and sarcoidosis is very unusual since it has been described only once. Zandman-Goddard et al. reported a case of coexisting RP and sarcoidosis in a patient with AIDS whose disease improved with antiretroviral drug therapy (6). We report here the first observation in an immunocompetent patient. Sarcoidosis can lead to sclerouveitis, middle ear involvement, polyarthritis and saddle nose deformity that may mimic RP but does not cause chondritis (7). Moreover, hilar lymphadenopathies, interstitial lung disease, and tissue granulomas are very rare in RP. In our patient, the clinical findings described above were conclusive with RP.

Although the etiopathogenesis of RP remains unknown, there are many arguments to support the role of the immune system. First, as seen above, RP is frequently associated with autoimmune diseases, and responds to corticosteroids. Second, RP is linked to the HLA class II DR4 isotype, which is frequently associated with other autoimmune diseases (8). Third, autoantibodies to type II, IX and XI collagen and to matrilin-I, a cartilage protein, can be found in the sera of patients with RP (3; 9; 10). Moreover immunofluorescence studies of involved cartilages have shown deposits of Immunoglobulins G, M, A and C3 as well as the presence of CD4⁺ T lymphocytes and plasma cells (11). Mice experimentally immunised with type 2 collagen or matrilin-1 develop auricular chondritis and symptoms resembling RP in humans (12).

Sarcoidosis is a systemic granulomatous disease that preferentially involves the lungs and lymph nodes. The etiopathogenesis also implicates the immune system. Sarcoidosis is the consequence of an exaggerated immune reaction mediated by CD4⁺ T lymphocytes in response to unknown antigens (13). T-cells present in alveoli and pulmonary interstitium secrete chemotactic factors to monocytes and macrophages, in particular monocyte chemotactic protein-1, which is also secreted by CD4⁺ T lymphocytes in patients with RP (5, 14).

Sarcoidosis and RP seem to share some etiopathogenic mechanisms: CD4⁺ T lymphocytes are implicated by an immune reaction through recruitment of the monocytes that contribute to inflammatory lesions. This could explain why these 2 diseases can coexist in the same patient.

In conclusion, sarcoidosis should be added to the list of diseases potentially coexisting with RP. Given the multitude of associated diseases, it has been proposed that RP be considered as a syndrome which can be primary or secondary.

Acknowledgments: We thank Bonnie Possest for her administrative assistance

REFERENCES

1. McAdam LP, O'Hanlan MA, Bluestone R, Pearson CM (1976) Relapsing polychondritis: prospective study of 23 patients and a review of the literature. *Medicine (Baltimore)* 55:193-215
2. Michet CJ, Jr., McKenna CH, Luthra HS, O'Fallon WM (1986) Relapsing polychondritis. Survival and predictive role of early disease manifestations. *Ann Intern Med* 104:74-78
3. Gergely P, Jr., Poor G (2004) Relapsing polychondritis. *Best Pract Res Clin Rheumatol* 18:723-738
4. Kent PD, Michet CJ, Jr., Luthra HS (2004) Relapsing polychondritis. *Curr Opin Rheumatol* 16:56-61
5. Baughman RP, Lower EE, du Bois RM (2003) Sarcoidosis. *Lancet* 361:1111-1118
6. Zandman-Goddard G, Peeva E, Barland P (2002) Combined autoimmune disease in a patient with AIDS. *Clin Rheumatol* 21:70-72
7. Letko E, Zafirakis P, Baltatzis S, Voudouri A, Livir-Rallatos C, Foster CS (2002) Relapsing polychondritis: a clinical review. *Semin Arthritis Rheum* 31:384-395
8. Lang B, Rothenfusser A, Lanchbury JS, Rauh G, Breedveld FC, Urlacher A, Albert ED, Peter HH, Melchers I (1993) Susceptibility to relapsing polychondritis is associated with HLA-DR4. *Arthritis Rheum* 36:660-664
9. Foidart JM, Abe S, Martin GR, Zizic TM, Barnett EV, Lawley TJ, Katz SI (1978) Antibodies to type II collagen in relapsing polychondritis. *N Engl J Med* 299:1203-1207
10. Alsalameh S, Mollenhauer J, Scheuplein F, Stoss H, Kalden JR, Burkhardt H, Burmester GR (1993) Preferential cellular and humoral immune reactivities to native and denatured collagen types IX and XI in a patient with fatal relapsing polychondritis. *J Rheumatol* 20:1419-1424
11. Dolan DL, Lemmon GB, Jr., Teitelbaum SL (1966) Relapsing polychondritis. Analytical literature review and studies on pathogenesis. *Am J Med* 41:285-299
12. McCune WJ, Schiller AL, Dynesius-Trentham RA, Trentham DE (1982) Type II collagen-induced auricular chondritis. *Arthritis Rheum* 25:266-273
13. Iannuzzi MC, Rybicki BA, Teirstein AS (2007) Sarcoidosis. *N Engl J Med* 357:2153-2165
14. Stabler T, Piette JC, Chevalier X, Marini-Portugal A, Kraus VB (2004) Serum cytokine profiles in relapsing polychondritis suggest monocyte/macrophage activation. *Arthritis Rheum* 50:3663-7
15. Trentham DE, Le CH (1998) Relapsing polychondritis. *Ann Intern Med* 129:114-122
16. Asadi AK (2003) Relapsing polychondritis. *Dermatol Online J* 9:3
17. Rapini RP, Warner NB (2006) Relapsing polychondritis. *Clin Dermatol* 24:482-485
18. Karaca NE, Aksu G, Yildiz B, Gulez N, Turk B, Dereli T, Kutukculer N (2009) Relapsing polychondritis in a child with common variable immunodeficiency. *Int J Dermatol* 48:525-528
19. Frances C, el Rassi R, Laporte JL, Rybojad M, Papo T, Piette JC (2001) Dermatologic manifestations of relapsing polychondritis. A study of 200 cases at a single center. *Medicine (Baltimore)* 80:173-179
20. Cohen PR (2006) Granuloma annulare, relapsing polychondritis, sarcoidosis, and systemic lupus erythematosus: conditions whose dermatologic manifestations may occur as hematologic malignancy-associated mucocutaneous paraneoplastic syndromes. *Int J Dermatol* 45:70-80
21. Wright MJ, Bel'eed K, Sellars L, Richmond I (1997) Relapsing polychondritis and myelodysplasia. *Nephrol Dial Transplant* 12:1704-1707
22. Cohen PR (2007) Paraneoplastic relapsing polychondritis. *Arch Dermatol* 143:949-950
23. Yanagi T, Matsumura T, Kamekura R, Sasaki N, Hashino S (2007) Relapsing polychondritis and malignant lymphoma: is polychondritis paraneoplastic? *Arch Dermatol* 143:89-90

Table 1 Major diagnostic criteria for relapsing polychondritis (1; 2)

McAdam criteria (three of six conditions recommended to be present)
Michet et al. criteria (one of two conditions recommended to be present)
<ol style="list-style-type: none"> 1. Recurrent chondritis of both auricles 2. Inflammatory polyarthritis 3. Chondritis of nasal cartilage 4. Inflammation of ocular structures including conjunctivitis, keratitis, scleritis/episcleritis and uveitis 5. Chondritis of the respiratory tract involving laryngeal and tracheal cartilages 6. Cochlear or vestibular damage manifested by neurosensory hearing loss, tinnitus and

vertigo
Michet et al. criteria (one of two conditions recommended to be present) <ol style="list-style-type: none"> 1. Proven inflammation in two of three of the auricular, nasal or laryngotracheal cartilages 2. Proven inflammation in one of three of the auricular, nasal or laryngotracheal cartilages plus 2 other signs including ocular inflammation, vestibular dysfunction, seronegative arthritis and hearing loss

Table 2 Diseases associated with relapsing polychondritis

Diseases	Références
Vasculitis	(15)
Wegener's granulomatosis	(15; 16)
Polyarteritis nodosa	(15; 16)
MAGIC syndrome (mouth and genital ulcers with inflamed cartilage syndrome)	(15; 16)
Behçet's disease	(15; 16)
Hypothyroidism	(17)
Systemic lupus erythematosus	(15)
Rheumatoid arthritis	(15)
Sjögren's syndrome	(15)
Mixed connective tissue disease	(15)
Pernicious anemia	(17)
Diabetes mellitus	(17)
Myasthenia gravis	(17)
Ulcerative colitis	(17)
Common variable immunodeficiency	(18)
Myelodysplastic syndrome	(19; 20)
Hematologic malignancies	(21-23)
Solid tumor malignancies	(21-23)

Figure 1. Auricular chondritis of the patient