Lipid biomarkers in suspended particles from a subtropical estuary: Assessment of seasonal changes in sources and transport of organic matter

Yunping Xu, Rudolf Jaffé

► To cite this version:

Yunping Xu, Rudolf Jaffé. Lipid biomarkers in suspended particles from a subtropical estuary: Assessment of seasonal changes in sources and transport of organic matter. *Marine Environmental Research*, 2007, 64 (5), pp.666. 10.1016/j.marenvres.2007.07.004 . hal-00562988

HAL Id: hal-00562988

<https://hal.science/hal-00562988>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Lipid biomarkers in suspended particles from a subtropical estuary: Assessment of seasonal changes in sources and transport of organic matter

Yunping Xu, Rudolf Jaffé

PII: S0141-1136(07)00090-6

DOI: [10.1016/j.marenvres.2007.07.004](https://doi.org/10.1016/j.marenvres.2007.07.004)

Reference: MERE 3135

To appear in: *Marine Environmental Research*

Received Date: 1 June 2006

Revised Date: 2 February 2007

Accepted Date: 8 July 2007

Please cite this article as: Xu, Y., Jaffé, R., Lipid biomarkers in suspended particles from a subtropical estuary: Assessment of seasonal changes in sources and transport of organic matter, *Marine Environmental Research* (2007), doi: [10.1016/j.marenvres.2007.07.004](https://doi.org/10.1016/j.marenvres.2007.07.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Lipid biomarkers in suspended particles from a subtropical estuary:
Assessment of seasonal changes in sources and transport of organic matter.

Yunping Xu¹ and Rudolf Jaffé*

Environmental Geochemistry Group, Southeast Environmental Research Center and Department of
Chemistry & Biochemistry, Florida International University, Miami, 33199, USA

*Corresponding author: Tel: +1-305-348-3095; fax: +1-305-348-4096

E-mail address: jaffer@fiu.edu (R. Jaffé)

¹Present address: Department of Physical & Environmental Sciences, S433, University of
Toronto, Toronto, ON M1C 1A4, Canada.

Abstract

Temporal and spatial variations in the composition of particulate organic matter (POM) from Florida Bay, USA were examined. The predominance of short chain homologues for *n*-alkanes, *n*-alcohols and *n*-fatty acids as well as relatively high abundance of C₂₇ and C₂₈ sterols suggested that an autochthonous/marine source of OM was dominant bay-wide. Several biomarker proxies such as *Paq* [(C₂₃+C₂₅)/(C₂₃+C₂₅+C₂₉+C₃₁) *n*-alkanes], short/long chain *n*-alkanes, (C₂₉+C₃₁) *n*-alkanes and taraxerol indicated a spatial shift in OM sources, where terrestrial OM rapidly decreased while seagrass and microbial OM markedly increased along a northeastern to southwestern transect. Regarding seasonal variations, POM collected during the dry season was enriched in terrestrial constituents relative to the wet season, likely as a result of reduced primary productivity of planktonic species and seagrasses during the dry season. Principal component analysis (PCA) classified the sample set into sub-groups based on PC1 which seemed to be spatially

controlled by OM origin (terrestrial-mangrove vs. marine-planktonic/seagrass). The PC2 seemed to be more seasonally controlled suggesting that hydrological fluctuations and seasonal primary productivity are the drivers controlling the POM composition in Florida Bay.

Key words: suspended particulate organic matter, Florida Bay, lipid biomarkers, seagrass, mangroves

1. Introduction

Relative to open oceans, coastal areas are highly vulnerable to environmental change due to their small size and location close to urban areas (Canuel, Cloern, Ringelberg, Guckert, & Rau, 1995). Coastal areas serve as sites of organic material exchange between land and oceans and are also active zones where OM is produced by higher plants, phytoplankton, benthic algae and bacteria (e.g. Jaffé, Mead, Hernandez, Peralba, & DiGuida, 2001; Jaffé, Boyer, Lu, Maie, Yang, & Scully et al., 2004; McCallister, Bauer, Ducklow, & Canuel, 2006). Additionally, aquatic vegetation is often abundant in coastal wetlands and estuarine areas and also serves as a significant source of OM (e.g. Campbell, & Miller, 2002; Jaffé et al., 2001; Paula, Fidalgo Ecosta, Martins, & Gove, 2001). This is the case of Florida Bay, where dense seagrass prairies dominate the benthic communities (Zieman, Fourqurean, & Iverson, 1989). The strength of each OM source can vary spatially as well as temporally on seasonal to tidal cycle time scales.

Various methods have been employed to characterize OM in aquatic environments. Among them, lipid tracers or biomarkers are particularly useful because they can reveal valuable information on OM sources at the molecular level (Meyers, 2003). Previous

studies have successfully used this approach to assess sedimentary OM sources in coastal areas, estuaries, rivers and lakes (e.g. Bianchi, Mitra, & McKee, 2002; Jaffé et al., 2001; Mead, Xu, Chong, & Jaffé, 2005; Prartono, & Wolff, 1998; Volkman, Rohjans, Rullkotter, Scholz-Bottcher, & Liebezeit, 2000). However, OM preserved in sediments may not always accurately reflect original inputs. In fact over 90% of organic material produced by the source organisms is degraded or consumed before reaching the sediments (Eadie, Chambers, Gardner, & Bell, 1984), and the preferential degradation of more labile plankton derived OM enhanced the apparent importance of terrestrial contributions which contain more resistant constituents (Meyers, 2003). In aquatic environments, suspended particles serve as an important vehicle in the transfer of OM from sources to sediments. Thus through the comparison of OM composition between sediments and suspended particles, early degradation processes of OM can be assessed (Canuel et al., 1995; Jaffé, Cabrera, Hajje, & Carvajal-Chitty, 1996; Saliot, Mejanelle, Scribe, Fillaux, Pepe, & Jabaud et al. , 2001), allowing for a better source strength interpretation of sedimentary OM.

While lipids in sediments and soils from south Florida estuaries have been reported (e.g. Hernandez, Mead, Peralba, & Jaffé, 2001; Jaffé et al., 2001; Mead et al., 2005; Neto, Mead, Louda, & Jaffé, 2006; Xu, Jaffé, Wachnicka, & Gaiser, 2006a; Xu, Mead, & Jaffé, 2006b), little is known about the sources and transport of particulate organic matter (POM). Here, we assess the spatial and temporal variation in the composition and origin of POM in Florida Bay through the measurement of several classes of lipid biomarkers such as aliphatic hydrocarbons, sterols, aliphatic alcohols, fatty acids, highly branched isoprenoids (HBIs) and terrestrially derived triterpenoids from the Florida Bay estuary, South Florida.

2. Sampling and methodology

2.1 Study area

Florida Bay is a large, subtropical body of water with a surface area of ~2200 km² and a mean depth of less than 2m (Corbett, Chanton, Burnett, Dillon, Rutkowski, & Fourqurean, 1999). Florida Bay experiences a warmer wet season from May to October accounting for 75% of the annual precipitation and a cooler dry season from November to April (Fourqurean, & Robbins, 1999). Benthic communities of Florida Bay consist mainly of dense seagrass prairies where *Thalassia testudinum* is the dominant species (Zieman et al., 1989). Mangrove forests dominated by *Rhizophora mangle* (red mangrove) grow along the sea-land interface between the Bay and the Florida Everglades wetlands. In addition, mangrove tree islands are common throughout the northeastern (NE) and central sections of the Bay, but are very sparse in the southwestern (SW) section, and represent an additional source of sedimentary OM to the Bay.

2.2. Sample collection

Water samples were collected during two wet seasons (October of 2003 and 2004) and two dry seasons (March 2004 and April 2005) at five locations corresponding to the Florida Coastal Everglades Long Term Ecological Research project's (FCE-LTER; <http://fcelter.fiu.edu/>) Taylor Slough/Pan Handle sampling transect (TS/Ph 7, 8, 9, 10 and 11; Fig. 1). The samples will subsequently be referred to as W1 and W2 for the wet season (October 2003 and 2004, respectively) or D1 and D2 for the dry season (March 2004 and April 2005, respectively) followed by their corresponding site number (7, 8, 9, 10 and 11). For example, W1-7 indicates a sample from TS/Ph 7 during the wet season of 2003. For

each sample, 50 liters of waters were collected at 20cm depth and kept in clean polycarbonate bottles (Nalgene) until transport to the laboratory (within 8 hours of collection). Samples were kept on ice and immediately vacuum filtered through pre-combusted glass-fiber filters with 0.7 μ m nominal pore size (GF/F). The filters with particulate material were freeze-dried at -40°C under the vacuum condition ($<100 \times 10^{-3}$ mBar). After that, dried samples were kept at -20°C until analysis.

2.3. Chemical analysis

The analytical procedure used in this study has been described by Jaffé et al. (2001). Briefly, dried filters were Soxhlet extracted with pure methylene chloride (Optima, Fisher, USA) for 24 hours. Activated copper was added to remove any elemental sulfur from the organic extracts. Total extractable organics were saponified with 1.0N KOH and separated into neutral and acid fractions. The neutral fraction was further separated by silica gel adsorption chromatography. Known amounts of squalane and per D phenanthrene were added as internal standards for aliphatic hydrocarbons and other fractions, respectively. The % error of the compound quantifications was of the order of 5-8%. Prior to analysis by GC-MS, the acid fraction was derivatized with freshly distilled diazomethane and the aliphatic alcohol and sterol fractions were derivatized with bis(trimethylsilyl)trifluoroacetamide (BSTFA) after addition of 10 μ l pyridine.

A HP 5973 GC-MS system was used for the analysis and operated in electron impact (EI) ionization mode at 70eV. The capillary column was a DB5 MS (30m, 0.25mm i.d., 0.25 μ m). The GC oven was programmed from 60 to 180°C at a rate of 6.0°C/min after 1 min at the initial temperature; then increased to 315°C at a rate of 3°C/min and kept at the

maximum temperature for 15 min. The identification of individual compounds was performed by comparison of chromatographic retention time and mass spectra of authentic standards and those previously reported in the literature.

3. Results and discussion

Several lipid compound classes such as *n*-alkanes, sterols, fatty acids and aliphatic *n*-alcohols as well as highly branched isoprenoids (HBIs) and pentacyclic triterpenoids were examined. All concentrations are expressed in ng per liter (ng/L).

3.1. Aliphatic hydrocarbons

A homologous series of *n*-alkanes ranging from C₁₅ to C₃₃ was observed and displayed a strong odd-over-even carbon number predominance, reflected by their high carbon preference index values (CPI; Fig. 2A and Table 1). Their total concentrations varied from 37 to 349ng/L (Table 1). The C₁₅ and C₁₇ homologues dominated the *n*-alkane distribution accounting for up to 81% of the total *n*-alkanes, while the C₂₉ and C₃₁ *n*-alkanes, indicative of terrestrially derived OM, were only minor components (0~24%). This distribution suggested a primary influence of phytoplankton (Cranwell, Eglinton, & Robinson, 1987). All POM samples except D2-7 featured an average carbon-chain length (ACL) of *n*-alkanes from 16.4 to 21.8, also reflecting strong microbial (planktonic) OM inputs.

Whereas surface sediments have a seagrass-derived lipid signature (Table 2; Xu et al., 2006b), POM was characterized by predominant phytoplankton-derived compounds. This difference may reflect (1) the preferential degradation of short-chain (microbial and

algal) *n*-alkanes relative to mid- (seagrasses) and long-chain (mangrove) counterparts during early diagenesis of POM and surface sediment OM; and (2) an unimportant contribution of seagrass- and mangrove-derived in resuspended OM relative to planktonic POM in the water column.

Along the NE to SW transect, the percentage of ($C_{29}+C_{31}$) *n*-alkanes gradually decreased from 1.5 to 0.3%, 13 to 2.7%, 8.2 to 0% and 24 to 2.7% for the W1, D1, W2 and D2 series, respectively, suggesting a consistent decrease in the source strength of terrestrially-derived OM toward offshore areas. The ratio of long/short ($>C_{20}/\leq C_{20}$) *n*-alkanes, an indicator of terrestrial vs. marine OM, generally decreased from the NE (site 7 and 8) to the central section (site 9) and then increased to the SW bay (site 10, 11), likely as a result of a strong influence of seagrasses in off-shore areas, which are characterized by high amounts of C_{23} and C_{25} *n*-alkanes (Jaffé et al., 2001; Mead et al., 2005). This data is consistent with previous reports that planktonic primary productivity is the highest in the central section of the Bay (Philips, Lynch, & Badylak, 1995).

The *n*-alkane based *Paq* value, defined as $(C_{23}+C_{25})/(C_{23}+C_{25}+C_{29}+C_{31})$ *n*-alkanes, was proposed as a means to distinguish between non-emergent and emergent plant derived OM (Ficken, Li, Swain, & Eglinton, 2000). Mead et al. (2005) reported *Paq* values for seagrasses, *Rhizophora mangle* and *Cladium* sp. (sawgrass) as 0.90-1.00, 0.24 and 0.10, respectively. This distinct difference makes the *Paq* a good proxy to estimate seagrass inputs in Florida Bay. The *Paq* value of POM markedly increased from site 7 (as low as 0.30) to site 11 (up to 1.0) (Fig. 3), reflecting spatial variations in the strength of seagrass vs. terrestrial (e.g. mangrove) OM. It should be noted that phytoplankton contributions are not reflected in the *Paq* proxy.

The POM samples from the dry season (D1 and D2) were enriched in terrestrial components relative to those from the wet season (W1 and W2), as reflected by several *n*-alkane based proxies. The percentage of high molecular weight *n*-alkanes ($C_{29}+C_{31}$) ranged from 3 to 24% during the dry season vs. 0 to 8% during the wet season. The sample D2-7 showed the lowest *Paq* value (0.3), the highest ACL value (25.0) and the most abundant long-chain *n*-alkanes (24%). The terrestrial “background” signal is apparently suppressed by the higher autochthonous primary production in the summer months in Florida Bay (wet season).

Several C_{20} , C_{25} and C_{30} highly branched isoprenoids (HBIs) were detected in the aliphatic hydrocarbon fractions (Fig. 2A). The C_{20} HBI compounds consisted of the saturated alkane namely 2, 6, 10-trimethyl-7-(3-methylbutyl)dodecane ($C_{20:0}$; Retention index; RI_{DB5} 1693) and its corresponding monounsaturated alkene ($C_{20:1_{DB5}}$ 1685) (Jaffé et al., 2001). The C_{20} HBI was present at concentrations as high as 433ng/L while that of the C_{25} HBIs reached 100ng/L. The C_{30} HBIs, however, were only detectable in the D2 series at a concentration below 10ng/L. Cyanobacteria have been suggested as a potential source for the C_{20} HBIs, but direct evidence for this is still lacking (Jaffé et al., 2001; Kenig, Sinninghe Damsté, Kock-van Dalen, Rijpstra, Huc, & de Leeuw, 1995). In contrast, the biosynthesis of C_{25} HBIs by some diatoms has been firmly established (Sinninghe Damsté, Muyzer, Abbas, Rampen, Massé, & Allard et al., 2004; Volkman, Barrett, & Dunstan, 1994). The C_{30} HBIs are also biosynthesized by some marine diatoms (Sinninghe Damsté et al., 2004; Volkman et al., 1994).

Regarding the spatial distribution, C_{20} HBIs were generally more abundant at the NE and central Bay relative to the SW Bay especially at TS/Ph 11. This distribution reflects

the mixed contribution of cyanobacterial mats and periphyton mats which are abundant in the NE and central Bay and in the freshwater Everglades, respectively (Jaffé et al., 2001). The C₂₅ HBIs were most abundant in the central Bay POM, as in the surface sediments (Xu et al., 2006b), reflecting autochthonous marine diatom inputs, while the absence of C₃₀ HBIs in most POM samples may be a reflection of weak source strengths or fast degradation of those polyunsaturated compounds.

3.2 Sterols

The concentrations of C₂₆-C₃₀ sterols ranged from 91 to 3040ng/L (Fig. 2D and Table 1). Among them, the principal compounds were cholest-5-en-3β-ol (C₂₇Δ⁵; 10-48%), 24-methylcholesta-5,22-dien-3β-ol (C₂₈Δ^{5,22}; 2-20%), 24-ethylcholest-5,22-dien-3β-ol (C₂₉Δ^{5,22}; 2-72%) and 24-ethylcholest-5-en-3β-ol (C₂₉Δ⁵; 4-16%). Figure 4 shows the relative abundance of C₂₆-C₃₀ sterols in the POM samples. C₂₇, C₂₈ and C₂₉ sterols were abundant, but the C₂₆ and C₃₀ compounds were minor or undetectable. The only C₂₆ sterol, 24-norcholesta-5,22*E*-dien-3β-ol (C₂₆Δ^{5,22}) accounted for < 3% of total sterols. The C₂₇ sterols were dominated by C₂₇Δ⁵ (10 to 48%) followed by cholesta-5,22-dien-3β-ol (C₂₇Δ^{5,22}), cholestan-3β-ol (C₂₇Δ⁰) and cholest-22-en-3β-ol (C₂₇Δ²²). C₂₈ sterols were composed of C₂₈Δ^{5,22}, 24-methylcholesta-22*E*-en-3β-ol (C₂₈Δ²²), 24-methylcholesta-5,24(28)-dien-3β-ol (C₂₈Δ^{5,24(28)}), 24-methylcholest-5-en-3β-ol (C₂₈Δ⁵) and 24-methylcholestan-3β-ol (C₂₈Δ⁰). Among them, C₂₈Δ⁵ (2-8%) and C₂₈Δ^{5,22} (2-20%) were the most abundant in most POM samples except for W1-11 which had C₂₈Δ^{5,24(28)} at a concentration of 445ng/L. The percentages of C₂₉ sterols were relatively constant and accounted for ~ 25% of the POM samples, except for the W1 series where this value

markedly decreased from site 7 to site 11 (Fig. 4). Several C_{30} sterols such as 4α -methyl-24-ethyl- $5\alpha(H)$ -cholest-22(E)-en- 3β -ol ($C_{30}\Delta^{22}$), $4\alpha,23,24$ -trimethyl- $5\alpha(H)$ -cholest-22(E)-en- 3β -ol ($C_{30}\Delta^{22}$), $4\alpha,23,24$ -trimethyl- $5\alpha(H)$ -cholestan- 3β -ol ($C_{30}\Delta^0$) accounted for 0.2-7% of the total sterols.

While the low abundance of C_{26} sterols may reflect minimal inputs of OM from marine animals (de Leeuw, Rijpstra, Schenck, & Volkman, 1983; Jiang, O'Leary, Volkman, Zhang, Jia, & Yu et al., 1994), the relatively high percentages of $C_{27}\Delta^5$ (10-48%) suggested an important contribution from zooplankton-derived OM, although an origin from other organisms is possible (Volkman, 1986). C_{28} sterols were relatively abundant in most POM samples, indicating a significant input from phytoplankton especially diatoms (Volkman, 1986), although these sterols have also been found to be abundant in seagrasses (Jaffé et al., 2001).

C_{29} sterols have been widely used as biomarkers of terrestrial OM (Huang, & Meinschein, 1976; 1978), but they are also biosynthesized by some microalgae (Volkman, 1986; Volkman, Barrett, Blackburn, Mansour, Sikes, & Gelin, 1998). In Florida Bay, $C_{29}\Delta^5$ in seagrasses accounts for ~ 75% of total sterols (Jaffé et al., 2001). Therefore, the high abundance of C_{29} sterols does not necessarily reflect a large terrestrially-derived OM input in Florida Bay, which is corroborated by the dominance of short-chain n -alkanes and the high Paq values (>0.81). The $C_{28}\Delta^5:C_{29}\Delta^{5,22}:C_{29}\Delta^5$ ratio of POM fell in the range of 1:0.76:1.24 to 1:9.93:0.90, while higher plants have this value in the order of 1:1.4:4.0 to 1:1.6:6.6 (Gogou, & Stephanou, 2004; Volkman, 1986). A significantly enhanced abundance of $C_{28}\Delta^5$ and $C_{29}\Delta^{5,22}$ relative to $C_{29}\Delta^5$ in POM suggests an important OM input from marine algae. The relative abundance of the C_{27} vs. C_{29} sterols did not show

pronounced variations along the NE to SW transect for D1, W2 and D2 series (1 to 3), again suggesting seagrass as an important POM source. As a biomarker for dinoflagellates (de Leeuw et al., 1983), $C_{30}\Delta^{22}$ (dinosterol) remained at low abundance in the POM, suggesting only minor contributions from these organisms. The presence of only minor amounts of stanols in the POM samples suggests that this OM is fairly fresh and has not undergone significant degradation (Jaffé et al., 1996; Wakeham, & Lee, 1989).

3.3 Fatty acids

The concentration of fatty acids was as high as 2580ng/L (Table 1). The *n*-fatty acid homologues displayed a strong even-over-odd carbon predominance (Fig. 2B). Short-chain fatty acids (SCFAs; $\leq C_{20}$) are mainly derived from planktonic and microbial organisms, while long-chain fatty acids (LCFAs; $>C_{20}$) are produced by higher plants including seagrasses (Jaffé et al., 2001). Therefore, the high SCFAs/LCFAs values observed for POM (up to 47) suggest a primary marine planktonic influence, in agreement with the data from *n*-alkanes. However, seagrasses, commonly covered by epiphytic organisms, cannot be excluded as an important source as they have a strong predominance at C_{16} followed by C_{24} in their fatty acids distributions (Jaffé et al., 2001). The long chain fatty acids observed here are most likely derived from mixed inputs of seagrasses and mangroves. Fatty acid homologues maximized at *n*- C_{14} (300ng/l) for W1-11, a distribution typical of phytoplankton.

The relative abundance of monounsaturated fatty acids (MUFAs) varied from 7% to 27%, indicating an important microbial contribution to the POM. Compared to the abundance of polyunsaturated fatty acids (PUFAs) in surface sediments (0.7-13%; Table 2;

Xu et al., 2006b), the POM samples showed markedly elevated concentrations of these constituents (7-57%), which can be attributed to the preferential degradation of MUFAs and PUFAs relative to their saturated counterparts during early diagenesis (Meyers, 2003). The proportion of branched fatty acids in the POM (0.6-4%) was significantly lower than that in surface sediments (5-15%; Table 2; Xu et al., 2006b), reflecting a more active bacterial reworking at the sediment-water interface (Gogou, & Stephanou, 2004) and a reduced contribution of these compounds from sediment resuspension. Branched fatty acids, principally *iso*- and *anteiso*-C₁₅ and C₁₇, are found to be abundant in bacteria (Budge, & Parrish, 1998). Wakeham and Beier (1991) proposed a bacterial indicator value of 6% to be rather low and not indicative of a large contribution of fatty acids from bacteria.

Some fatty acid based proxies can reflect a shift in phytoplankton taxonomic classes such as diatoms and dinoflagellates. The ratios of C_{16:1}/C_{16:0}, $\Sigma C_{16}/\Sigma C_{18}$ and C_{20:5}/C_{22:6} have been proposed as indicators of diatoms (Budge, & Parrish, 1998; Ramos, Parrish, Quibuyen, & Abrajano, 2003). For the Florida Bay POM, the ratio of C_{16:1}/C_{16:0} remained below 1.0 while the $\Sigma C_{16}/\Sigma C_{18}$ was generally higher than 1.0 and had the maximum value of 3.2 (Table 1). This apparent disagreement between the two diatom indicators is most likely attributed to an interference of seagrass derived OM (possibly from epiphytes) in which the C_{16:0} fatty acid is dominant (Jaffé et al., 2001). Therefore, such proxies involve non-source specific biomarkers, and hence their applicability can be limited. The ratio of C_{20:5}/C_{22:6} was as high as 12, indicating an important contribution from diatoms.

Seasonally, the percentages of (MUFAs+PUFAs) in the POM remained at relatively constant, high levels, suggesting consistently important contributions of phytoplankton

derived OM. The fatty acids also showed higher percentages of long-chain homologues during the dry season, as did the *n*-alkanes.

3.4 *n*-Alkanols

The *n*-alkanol distribution exhibited a strong even over odd carbon number predominance with concentrations ranging from 16 to 1220ng/L (Table 1 and Fig. 2C). Terrestrial plants produce long chain *n*-alkanols from C₂₂ to C₃₂ (Sever, & Parker, 1969), while short chain ($\leq$ C₂₀) *n*-alkanols are characteristic of microbial OM inputs (Robinson, Cranwell, Finlay, & Eglinton, 1984). The *n*-alkanol distributions in most POM samples showed maxima at short chain homologues (C₁₂-C₁₈), again consistent with a predominant algal and microbial OM origin. However, a few of the dry season samples (i.e., D1-7, D1-8 and D2-7) had significant levels of C₂₈ or C₃₀ *n*-alkanols (Table 1), suggesting important higher plant inputs (Jaffé et al., 2001). This supports the interpretation of the *n*-alkane and fatty acid distributions as having higher relative contributions from terrestrial-higher plant OM in the dry season. The ratio of long chain/short chain *n*-alkanols markedly decreased (Table 1), further confirming a decreasing trend of terrestrial influence toward offshore. In regards to seasonality, the dry season samples had higher values of long/short chain *n*-alkanols (0.2-10) relative to the wet season samples (0.0-1.4), suggesting more terrestrial OM components for the former, while wet season OM sources seem controlled by increased planktonic primary productivity.

3.5 Higher plant derived pentacyclic triterpenoids

Pentacyclic triterpenoids such as taraxerol, β -amyrin and lupeol are widely found in higher plants, especially in trees, but rarely in other organisms. Hence they are often used as biomarkers of terrestrial inputs (Brassell, Eglinton, & Maxwell, 1983; ten Haven, & Rullkotter, 1988; Versteegh, Schefuß, Dupont, Marret, Sinninghe Damsté, & Jansen, 2004). Previous studies have reported taraxerol concentration of up to 1.4 mg/g in mangrove leaves (Versteegh et al., 2004) but it is absent in the other two main higher plant biomass components of the Everglades ecosystem, namely seagrass and sawgrass (Xu et al., 2006b). Several pentacyclic triterpenoids such as taraxerol, β -amyrin, germanicol and lupeol were detected in the POM. Overall, their concentrations rapidly decreased from site 7 to 11 (Table 1), suggesting reduced transport from the NE mangrove fringe to the Bay and/or the sparse distribution of mangrove tree islands in SW Florida Bay. The abundance of these triterpenoids was found to be higher during the dry season (D1 and D2) relative to the wet season (W1 and W2). For example, taraxerol, a specific biomarker of mangroves (Versteegh et al., 2004), was as high as 1210ng/L for the D1 series and 109ng/L for the D2 series, but only 19ng/L for the W1 series and 102ng/L for the W2 (Table 1). These results are in agreement with data on source variations for dissolved organic matter (DOM) in Florida Bay (Maie, Boyer, Yang, & Jaffé, 2006).

3.6 Principal components analysis (PCA)

PCA is a multivariate statistical technique, useful in extracting information from complex organic geochemical data (e.g. Hernandez et al., 2001; Mead et al., 2005; Mudge, East, Bebianno, & Barreira, 1998). Figure 5 shows the PCA score plot for POM in Florida Bay where the first principal component (PC 1) accounted for 39% and the second principal

component (PC 2) accounted for 27% of the variance. The biomarker concentration data and indices used in this analysis were for the taraxerol, sterols, *Paq*, *n*-alkanes and fatty acid distributions.

The variability among the samples seems to be controlled mainly by OM source and season, and neither the PCs properly define these variables independently. However, PC1 seems to more strongly control the special variables separating the POM samples into two groups, namely near-shore (empty diamonds) and offshore areas (filled diamonds) with the exception of samples W1-8 and D2-8. In contrast, PC2 seems to control the seasonal component and roughly separates the D and W series along its axis (with some exceptions). However, analyzing the sample clusters in more detail, one can observe that the W1 samples are clearly distinct from W2 samples, except for samples from site 11 which both fall in the W1 cluster. Even sample W1-8 falls into the W1 cluster with a high positive correlation with PC1 although station 8 is assumed to contain high relative contributions from terrestrial sources. The main difference between the W1 and W2 sampling is that rainfall for the W1 season was about twice that for the W2 season. The average daily precipitation at the Flamingo Ranger Station close to Florida Bay was 0.81mm during October 2003 (W1) and 0.39mm during October 2004 (W2); (see <http://fcelter.fiu.edu>). Based on this, it seems that higher rainfall resulted in increased planktonic/seagrass primary productivity probably due to atmospheric phosphorus inputs into this highly P-limited environment (Sutula, Perez, Reyes, Childers, Davis, & Day et al., 2003). This result also suggests that potential seasonal variations in POC composition are not primarily driven by hydrology (POM transport) but by primary productivity. Samples W1-11 and W2-11 were

co-located in the PCA plot since this site experiences active P-exchange with waters from the Gulf of Mexico and as such is less sensitive to atmospheric P-inputs.

With regards to the D samples, the drier D1 season compared to D2 (0.04mm during March 2004 (D1) and 0.22mm during April 2005 (D2)) was more controlled by hydrology (POM transport from terrestrial environments) compared to D2. And a consistent shift from more terrestrial (negative correlation with PC1) to more marine (positive correlation with PC1) was observed for the nearshore sites (7, 8 and 9), but as expected not for sites 10 and 11. These data again suggest that POM in Florida Bay is primarily derived from marine (planktonic/seagrass) OM where primary productivity during the wet season is an important ecological driver. In contrast, the dry season exerts a stronger terrestrial OM influence on the nearshore sites, probably as a result of a relative increase in terrestrial (mangrove) OM transport in comparison with marine biomass primary productivity.

4. Summary

Autochthonous/marine sources were the principal contributors to POM in Florida Bay, while allochthonous/terrestrial OM, although minor, played a more important role mostly in the NE, nearshore sites and during the dry season. Multi-biomarker proxies indicated a shift in OM sources along the NE to SW transect from mangrove influenced to plankton/seagrass dominated marine OM inputs. The higher relative abundance of terrestrially derived POM during the dry season relative to the wet season is most likely an indirect consequence of the seasonal primary productivity changes in the Bay.

Higher percentages of mono- and poly-unsaturated fatty acids and lower proportions of branched fatty acids in the POM samples relative to surface sediments reflect a

preferential degradation of unsaturated compounds during sedimentation and active microbial reworking at the water-sediment interface.

PCA analyses classified the POM samples into two groups according to PC1, namely a nearshore and a more marine influenced group, while PC2 separated the POM samples according to seasonality (hydrology/primary productivity). This study is the first to report seasonal variations in suspended POM compositions in Florida Bay.

Acknowledgements

The authors thank Dr. N. Maie for assistance with sample collection. We also thank the B.R.T. Simoneit and two anonymous reviewers for their helpful comments. This research was supported by the National Science Foundation through the FCE-LTER (DEB-9910514) and the Division of Earth Sciences Geology and Paleontology Program (#0345812). Y.X. thanks FIU for a presidential fellowship during his Ph. D. study. This is SERC contribution 342.

References

- Bianchi, T.S., Mitra, S., & McKee, B.A. (2002). Sources of terrestrially-derived organic carbon in lower Mississippi River and Louisiana shelf sediments: implications for differential sedimentation and transport at the coastal margin. *Marine Chemistry*, 77, 211-223.
- Brassell, S.C., Eglinton, G., & Maxwell, J.R. (1983). The geochemistry of terpenoids and steroids. *Biochemical Society Transactions*, 11, 575-586.
- Budge, S.M., & Parrish, C.C. (1998). Lipid biogeochemistry of plankton, settling matter and sediments in Trinity Bay, Newfoundland. II. Fatty acids. *Organic Geochemistry*, 29, 1547-1559.
- Campbell, S.J., & Miller, C.J. (2002). Shoot and abundance characteristics of the seagrass *Heterozostera tasmanica* in Westernport estuary (south-eastern Australia). *Aquatic Botany*, 73, 33-46.
- Canuel, E.A., Cloern, J.E., Ringelberg, D.B., Guckert, J.B., & Rau, G.H. (1995). Molecular and isotopic tracers used to examine sources of organic matter and its incorporation into the food webs of San Francisco Bay. *Limnology and Oceanography*, 40, 67-81.
- Corbett, D.R., Chanton, J., Burnett, W., Dillon, K., Rutkowski, C., & Fourqurean, J.W. (1999). Patterns of groundwater discharge into Florida Bay. *Limnology and Oceanography*, 44, 1045-1055.
- Cranwell, P.A., Eglinton, G., & Robinson, N. (1987). Lipids of aquatic organisms as potential contributors to lacustrine sediment - II. *Organic Geochemistry*, 11, 513-527.
- de Leeuw, J.W., Rijpstra, W.I.C., Schenck, P.A., & Volkman, J.K. (1983). Free, esterified and residual bound sterols in Black Sea Unit I sediments. *Geochimica et Cosmochimica Acta*, 47, 455-465.
- Eadie, B.J., Chambers, R.L., Gardner, W.S., & Bell, G.E. (1984). Sediment trap studies in Lake Michigan: resuspension and chemical fluxes in the southern basin. *Journal of Great Lakes Research*, 10, 307-321.
- Eglinton, G., & Hamilton, R.J. (1967). Leaf epicuticular waxes. *Science*, 156, 1322-1335.
- Ficken, K.J., Li, B., Swain, D.L., & Eglinton, G. (2000). An *n*-alkane proxy for the sedimentary input of submerged/floating freshwater aquatic macrophytes. *Organic Geochemistry*, 31, 745-749.
- Fourqurean, J.W., & Robbins, M.B. (1999). Florida Bay: a history of recent ecological changes. *Estuaries*, 22, 345-357.
- Gogou, A., & Stephanou, E.G. (2004). Marine organic geochemistry of the Eastern Mediterranean: 2. Polar biomarkers in Cretan Sea surficial sediments. *Marine Chemistry*, 85, 1-25.
- Hernandez, M.E., Mead, R., Peralba, M.C., & Jaffé, R. (2001). Origin and transport of *n*-alkan-2-ones in a subtropical estuary: potential biomarkers for seagrass-derived organic matter. *Organic Geochemistry*, 32, 21-32.
- Huang, W., & Meinschein, W.G. (1976). Sterols as source indicators of organic materials in sediments. *Geochimica et Cosmochimica Acta*, 40, 323-330.

- Huang, W.-Y., & Meinschein, W.G. (1978). Sterols in sediments from Baffin Bay, Texas. *Geochimica et Cosmochimica Acta*, 42, 1391-1396.
- Jaffé, R., Boyer, J.N., Lu, X., Maie, N., Yang, C., Scully, N.M., & Mock, S. (2004). Source characterization of dissolved organic matter in a subtropical mangrove-dominated estuary by fluorescence analysis. *Marine Chemistry*, 84, 195-210.
- Jaffé, R., Cabrera, A., Hajje, N., & Carvajal-Chitty, H. (1996). Organic biogeochemistry of a hypereutrophic tropical, freshwater lake - Part 1: particle associated and dissolved lipids. *Organic Geochemistry*, 25, 227-240.
- Jaffé, R., Mead, R., Hernandez, M.E., Peralba, M.C., & DiGuida, O.A. (2001). Origin and transport of sedimentary organic matter in two subtropical estuaries: a comparative, biomarker-based study. *Organic Geochemistry*, 32, 507-526.
- Jiang, S., O'Leary, T., Volkman, J.K., Zhang, H., Jia, R., Yu, S., Wang, Y., Luan, Z., Sun, Z., & Jiang, R. (1994). Origins and simulated thermal alteration of sterols and keto-alcohols in deep-sea marine sediments of the Okinawa Trough. *Organic Geochemistry*, 21, 415-422.
- Kenig, F., Sinninghe Damsté, J.S., Kock-van Dalen, A.C., Rijpstra, W.I.C., Huc, A.Y., & de Leeuw, J.W. (1995). Occurrence and origin of mono-, di-, and trimethylalkanes in modern and Holocene cyanobacterial mats from Abu Dhabi, United Arab Emirates. *Geochimica et Cosmochimica Acta*, 59, 2999-3015.
- Mead, R., Xu, Y., Chong, J., & Jaffé, R. (2005). Sediment and soil organic matter source assessment as revealed by the molecular distribution and carbon isotopic composition of n-alkanes. *Organic Geochemistry*, 36, 363-370.
- McCallister, S.L., Bauer, J.E., Ducklow, H.W., & Canuel, E.A. (2006). Sources of estuarine dissolved and particulate organic matter: A multi-tracer approach. *Organic Geochemistry*, 37, 454-468.
- Maie, N., Boyer, J.N., Yang, C., & Jaffé, R. (2006). Spatial, geomorphologic, and seasonal variability in CDOM in estuaries of Florida Coastal Everglades. *Hydrobiologia*, in press.
- Meyers, P.A. (2003). Applications of organic geochemistry to paleolimnological reconstructions: a summary of examples from the Laurentian Great Lakes. *Organic Geochemistry*, 34, 261-289.
- Mudge, S.M., East, J.A., Bebianno, M.J., & Barreira, L.A. (1998). Fatty acids in the Ria Formosa Lagoon, Portugal. *Organic Geochemistry*, 29, 963-977.
- Neto, R.R., Mead, R.N., Louda, J.W., & Jaffé, R. (2006). Organic biogeochemistry of detrital flocculent material (Floc) in a subtropical, coastal wetland. *Biogeochemistry*, 77, 283-304.
- Paula, J., Fidalgo Ecosta, P., Martins, A., & Gove, D. (2001). Patterns of abundance of seagrasses and associated infaunal communities at Inhaca Island, Mozambique. *Estuarine, Coastal and Shelf Science*, 53, 307-318.
- Philips, E.J., Lynch, T.C., & Badylak, S. (1995). Chlorophyll a, tripton, color, and light availability in a shallow tropical inner-shelf lagoon, Florida Bay, USA. *Marine Ecology Progress Series*, 127, 223-234.
- Prartono, T., & Wolff, G.A. (1998). Organic geochemistry of lacustrine sediments: a record of the changing trophic status of Rostherne Mere, U.K. *Organic Geochemistry*, 28, 729-747.

- 468 Ramos, C.S., Parrish, C.C., Quibuyen, T.A.O., & Abrajano, T.A. (2003). Molecular and
469 carbon isotopic variations in lipids in rapidly settling particles during a spring
470 phytoplankton bloom. *Organic Geochemistry*, 34, 195-207.
- 471 Robinson, N., Cranwell, P.A., Finlay, B.J., & Eglinton, G. (1984). Lipids of aquatic
472 organisms as potential contributors to lacustrine sediments. *Organic Geochemistry*,
473 6, 143-152.
- 474 Saliot, A., Mejanelle, L., Scribe, P., Fillaux, J., Pepe, C., Jabaud, A., & Dagaut, J. (2001).
475 Particulate organic carbon, sterols, fatty acids and pigments in the Amazon River
476 system. *Biogeochemistry*, 53, 79-103.
- 477 Sever, J., & Parker, P.L. (1969). Fatty alcohols (normal and isoprenoid) in sediments.
478 *Science*, 164, 1052-1054.
- 479 Sinninghe Damsté, J.S., Muyzer, G., Abbas, B., Rampen, S.W., Massé, G., Allard, W.G.,
480 Belt, S.T., Robert, J.-M., Rowland, S.J., Moldowan, J.M., Barbanti, S.M., Fago, F.J.,
481 Denisevich, P., Dahl, J., Trinidad, L.A.F., & Schouten, S. (2004). The rise of
482 rhizosolenid diatoms. *Science*, 304, 584-587.
- 483 Sutula, M.A., Perez, B.C., Reyes, E., Childers, D.L., Davis, S., Day, J., John W., Rudnick,
484 D., Sklar, F. (2003). Factors affecting spatial and temporal variability in material
485 exchange between the Southern Everglades wetlands and Florida Bay (USA).
486 *Estuarine, Coastal and Shelf Science*, 57, 757-781.
- 487 ten Haven, H.L., & Rullkötter, J. (1988). The diagenetic fate of taraxer-14-ene and
488 oleanene isomers. *Geochimica et Cosmochimica Acta*, 52, 2543-2548.
- 489 Versteegh, G.J.M., Schefuß, E., Dupont, L., Marret, F., Sinninghe Damsté, J.S., & Jansen,
490 J.H.F. (2004). Taraxerol and *Rhizophora* pollen as proxies for tracking past
491 mangrove ecosystems. *Geochimica et Cosmochimica Acta*, 68, 411-422.
- 492 Volkman, J.K. (1986). A review of sterol markers for marine and terrigenous organic
493 matter. *Organic Geochemistry*, 9, 83-99.
- 494 Volkman, J.K., Barrett, S.M., Blackburn, S.I., Mansour, M.P., Sikes, E.L., & Gelin, F.
495 (1998). Microalgal biomarkers: A review of recent research developments. *Organic*
496 *Geochemistry*, 29, 1163-1179.
- 497 Volkman, J.K., Barrett, S.M., & Dunstan, G.A. (1994). C₂₅ and C₃₀ highly branched
498 isoprenoid alkenes in laboratory cultures of two marine diatoms. *Organic*
499 *Geochemistry*, 21, 407-414.
- 500 Volkman, J.K., Rohjans, D., Rullkötter, J., Scholz-Bottcher, B.M., & Liebezeit, G. (2000).
501 Sources and diagenesis of organic matter in tidal flat sediments from the German
502 Wadden Sea. *Continental Shelf Research*, 20, 1139-1158.
- 503 Wakeham, S.G., & Beier, J.A. (1991). Fatty acid and sterol biomarkers as indicators of
504 particulate matter source and alteration processes in the Black Sea. *Deep-Sea*
505 *Research*, 38, S943-S968.
- 506 Wakeham, S.G., & Lee, C. (1989). Organic geochemistry of particulate matter in the ocean:
507 The role of particles in oceanic sedimentary cycles. *Organic Geochemistry*, 14, 83-
508 96.
- 509 Xu, Y., Jaffé, R., Wachnicka, A., & Gaiser, E.E. (2006a). Occurrence of C₂₅ highly
510 branched isoprenoids (HBIs) in Florida Bay: paleoenvironmental indicators of
511 diatom-derived organic matter inputs. *Organic Geochemistry*, 37, 847-859.

- 512 Xu, Y., Mead, R., & Jaffé, R. (2006b). A molecular marker based assessment of
513 sedimentary organic matter sources and distributions in Florida Bay. *Hydrobiologia*,
514 569, 179-192.
- 515 Zieman, J.C., Fourqurean, J.W., & Iverson, R.L. (1989). Distribution, abundance and
516 productivity of seagrasses and macroalgae in Florida Bay. *Bulletin of Marine*
517 *Science*, 44, 292-311.
518

Figure and table captions:

Fig. 1: Geographic locations of particulate organic matter sampling sites in Florida Bay

Fig. 2: Representative GC/MS chromatograms of (A) *n*-alkanes, (B) fatty acids, (C) *n*-alcohols and (D) sterols in particulate organic matter from Florida Bay

Fig. 3: The *Paq* distributions in particulate organic matter from Florida Bay.

$Paq = (C_{23}+C_{25})/(C_{23}+C_{25}+C_{29}+C_{31})$ *n*-alkanes

Fig. 4: Relative abundance of C₂₆, C₂₇, C₂₈, C₂₉ and C₃₀ sterols in particulate organic matter from Florida Bay

Fig. 5: Biomarker-based score plot of the first two principal components (PC-1 and PC-2) for suspended particles from Florida Bay

Table 1: Concentration of *n*-alkanes, *n*-alkanols, triterpenoids, fatty acids, HBIs and sterols in Florida Bay suspended particles during dry season (D) and wet season (W)

Table 2: Comparison of biomarker proxies between POM and surface sediments in Florida Bay

Figure 1

553

554

555 Fig. 2

Fig. 3

Fig. 4

Fig. 5

580

Table 1 Concentration of *n*-alkanes, *n*-alkanols, triterpenoids, fatty acids, HBIs and sterols in Florida Bay suspended particulates during dry season (D) and wet season (W)

	W1				D1				W2					D2				
	8	9	10	11	7	8	9	10	7	8	9	10	11	7	8	9	10	11
<i>n</i> -Alkanes																		
Total	119	338	349	281	316	266	37	95	43	163	76	75	38	128	95	73.7	214	60
C ₂₉ +C ₃₁	1.8	3	1	1	41	12	1	3	4	10	2	0.2	0	29	8	5	6	4
Paq	0.8	0.8	0.9	1.0	0.4	0.6	0.8	0.8	0.3	0.6	0.7	1.0	1.0	0.3	0.6	0.5	0.6	0.8
Cmax	C ₁₇	C ₁₇	C ₁₅	C ₂₃	C ₁₅	C ₁₅	C ₁₅	C ₁₅	C ₁₅	C ₁₅	C ₁₅	C ₂₁	C ₁₅	C ₁₇	C ₁₇	C ₁₅	C ₁₅	C ₁₅
Short	96	293	326	140	195	212	22	63	33	103	51	50	21	34	45	50	177	23
Long	23	46	23	141	121	55	15	33	10	60	26	54	17	86	50	23	36	35
Long/short	4.1	6.3	14	1.0	1.6	3.8	1.5	1.9	3.2	1.7	2.0	0.9	1.2	0.4	0.9	2.2	5	0.7
ACL	18.5	17.9	16.5	20.6	20.1	17.5	19.7	20.6	18.2	19.6	19.4	20.5	19.4	25	21.8	19.3	17.6	21.5
CPI	7.1	3.4	5.4	1.6	7.8	10.9	2.1	1.9	7.1	3.0	1.6	1.9	2.2	2.4	1.4	5.1	11.3	3.8
%(C ₁₅ +C ₁₇)	69	61	76	11	57	75	32	31	68	41	33	12	31	24	15	63	81	33
%(C ₂₉ +C ₃₁)	1.5	0.9	0.3	0.3	13	4.7	3.2	2.7	8.1	6.0	2.8	0.2	0.0	24	8.8	6.2	2.7	6.8
<i>n</i> -Alkanols																		
Total	16	59	48	29	1220	406	41	63	122	218	80	203	155	143	130	41	177	30
Cmax	C ₁₄	C ₁₂	C ₁₈	C ₁₂	C ₃₀	C ₂₈	C ₁₈	C ₁₈	C ₁₈	C ₁₂	C ₁₂	C ₁₂	C ₁₂	C ₂₈	C ₁₄	C ₁₈	C ₁₈	C ₁₈
Long/short	0.3	0.3	0.0	0.0	10	7.8	1.5	0.2	0.0	1.4	0.1	0.1	0.0	3.9	0.9	1.0	0.5	0.5
Triterpenoids																		
Taraxerol	7	19	0.0	0.0	1210	181	9	8.4	102	30	0.0	0.0	0.0	109	21	2	1	0.0
β-amyrin	0.0	9.9	0.0	0.0	518	97	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	5	0.0	0.0	0.0
Germanicol	0.0	0.0	0.0	0.0	131	29	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	0.0	0.0	0.0
Lupeol	7.0	0.0	0.0	0.0	231	39	0.0	0.0	18	7.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 1 continued

Fatty acids

Total	153	40	165	601	2580	1063	1230	1980	598	214	270	535	454	1370	1824	1800	3411	1130
Short	149	40	162	587	2000	832	1180	1800	543	161	246	495	436	662	752	869	1140	577
Long	3.7	0.0	3.4	14	577	231	44	184	55	53	24	41	18	202	185	174	223	72
Short/long	39.8	nd	47.4	41.1	3.5	3.6	26.7	9.7	10.0	3.0	10.5	12.2	23.9	3.3	4.1	5.0	5.1	8.0
MUFAs (%)	26	21	27	16	20	18	23	23	15	7.0	11	17	20	11	16	18	23	24
PUFAs (%)	10	0.9	7.5	11	6.5	8.6	9.6	9.0	3.3	0.2	0.9	1.5	1.2	22	21	20	35	16
Br-FAs (%)	3.5	2.3	1.7	1.2	2.1	1.2	1.1	1.5	0.8	0.6	0.7	2.0	1.7	3.9	3.1	3.5	2.3	3.0
C16:1/C16:0	0.6	0.3	0.3	0.8	0.3	0.3	0.4	0.5	0.2	0.1	0.1	0.2	0.2	0.5	0.4	0.5	0.7	0.3
C16/C18	2.1	3.2	2.0	2.8	1.6	1.7	1.9	1.7	2.2	2.0	1.7	2.1	2.3	2.0	1.9	1.7	0.9	1.0
A/B	8.7	nd	3.1	0.6	3.9	3.2	5.5	7.0	4.1	nd	nd	nd	nd	4.1	5.4	11.9	2.8	1.3
HBIs																		
C ₂₀ HBIs	154	282	148	7.1	404	191	58	133	40	52	77	17	1.1	410	71	145	433	4.3
C ₂₅ HBIs	13	41	91	0.0	35	22	14	56	3.0	16	24	0.0	0.0	56	33	9.8	100	0.0
C ₃₀ HBIs	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	10.3	nd	nd	nd	nd
Sterols																		
Total	180	91	526	800	3040	2290	1137	1090	1380	1460	1040	1180	161	596	451	430	1500	309

All concentrations are expressed in ng/L. C_{max}: the maximum compounds in homologues.

Short: total concentrations of short-chain homologues ($\leq C_{20}$). Long: total concentrations of long-chain homologues ($>C_{20}$). ACL = average chain length of the homologues. CPI = carbon preference index. MUFAs = mono-unsaturated fatty acids. PUFAs = poly-unsaturated fatty acids. Br-FAs = branched-C₁₅ + C₁₇ fatty acids indicative of bacterial contributions. A/B = the ratio of C_{20:5} over C_{22:6} fatty acids. nd = not detected

Table 2

	TS/Ph 7		TS/Ph 8		TS/Ph 9		TS/Ph 10		TS/Ph 11	
<i>n</i> -alkane	POM	Sed.	POM	Sed.	POM	Sed.	POM	Sed.	POM	Sed.
<i>Paq</i>	0.33	0.44	0.65	na	0.7	0.53	0.83	0.63	0.94	0.83
Cmax	C ₁₅	C ₂₉	C ₁₅	na	C ₁₅	C ₂₅	C ₁₅	C ₂₅	C ₂₃	C ₂₅
ACL	21.1	26.6	19.4	na	19.1	25.7	18.8	24.5	20.5	21.4
%(<i>C</i> ₁₅ + <i>C</i> ₁₇)	50	2.6	67	na	47	7.4	50	9.7	25	13
%(<i>C</i> ₂₉ + <i>C</i> ₃₁)	16	29	5.3	na	3.3	23	1.5	10	2.4	4.6
Fatty acids										
PUFA%	11	0.7	10	na	8	4.4	13	8.2	9.2	13
MUFA%	15	4.9	17	na	18	6.4	23	11	20	37
BrFA%	2.3	7.3	2.1	na	1.9	5.2	1.9	8	2	15
Sterol										
<i>C</i> ₂₇ %	44	19	40	na	36	37	47	38	41	42
<i>C</i> ₂₈ %	23	17	18	na	23	17	22	19	41	22
<i>C</i> ₂₉ %	28	64	38	na	38	46	29	43	15	36
<i>n</i> -alkanol										
Long/short	4.6	10	2.6	na	0.7	50	0.2	6.3	0.1	1.8

na = not analyzed

*The data on sediments are from Xu et al., 2006b while data on POM are the average values from this study.