Differential gene expression in the brain of in response to exposure to polychlorinated biphenyls (PCBs)

Bowen Li, Chonggang Wang, Kai Ye, Ang Yu, Yixin Chen, Zhenghong Zuo

► To cite this version:

Bowen Li, Chonggang Wang, Kai Ye, Ang Yu, Yixin Chen, et al.. Differential gene expression in the brain of in response to exposure to polychlorinated biphenyls (PCBs). *Marine Environmental Research*, 2008, 66 (5), pp.548. 10.1016/j.marenvres.2008.09.001 . hal-00563053

HAL Id: hal-00563053

<https://hal.science/hal-00563053>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Differential gene expression in the brain of *Sebastiscus marmoratus* in response to exposure to polychlorinated biphenyls (PCBs)

Bowen Li, Chonggang Wang, Kai Ye, Ang Yu, Yixin Chen, Zhenghong Zuo

PII: S0141-1136(08)00205-5
DOI: [10.1016/j.marenvres.2008.09.001](https://doi.org/10.1016/j.marenvres.2008.09.001)
Reference: MERE 3287

To appear in: *Marine Environmental Research*

Received Date: 30 April 2008
Revised Date: 30 July 2008
Accepted Date: 5 September 2008

Please cite this article as: Li, B., Wang, C., Ye, K., Yu, A., Chen, Y., Zuo, Z., Differential gene expression in the brain of *Sebastiscus marmoratus* in response to exposure to polychlorinated biphenyls (PCBs), *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.09.001](https://doi.org/10.1016/j.marenvres.2008.09.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Differential gene expression in the brain of *Sebastiscus marmoratus* in
response to exposure to polychlorinated biphenyls (PCBs)**

Bowen Li^a, Chonggang Wang^{a,b}, Kai Ye^a, Ang Yu^b, Yixin Chen^a, Zhenghong Zuo^{a,b*}

^a Key Laboratory of the Ministry of Education for Cell Biology and Tumor Cell Engineering, School of Life Sciences,

Xiamen University, Xiamen City, 361005, PR China

^b State Key Laboratory of Marine Environmental Science (Xiamen University), Xiamen City, 361005, PR China

* Corresponding author telephone: +86-592-2187353; fax: +86-592-2187230;

E-mail: zuozhenghong@xmu.edu.cn

Abstract

Effects of exposure to polychlorinated biphenyls (PCBs) on *Sebastiscus marmoratus* were investigated using a suppression subtractive hybridization method. A total of 108 gene sequences were identified as having the potential for being differentially expressed, and 45 could be identified with homologous database sequences. Functions with which they were associated included long-term potentiation and neurotransmitter release, neuroendocrine, mitosis and cell proliferation, energy-related metabolism, general metabolism, signal protein, hemopoiesis system, immune system, and structure. The expression of 17 of these genes was analyzed in the brain using real time fluorescent quantitative PCR. The present study provided a basis for studying the response of fish to PCB exposure and allowed the characterization of new potential neurotoxicol biomarkers of PCB contamination in seawater.

Keywords: *polychlorinated biphenyls, neurotoxicity, Sebastiscus marmoratus, suppression subtractive hybridization library, real time PCR*

1. Introduction

Polychlorinated biphenyls (PCBs) are a class of industrial compounds consisting of paired phenyl rings with various degrees of chlorination (Tilson and Kodavanti, 1997). Although their production was banned in the 1970s, this was after more than a billion kilograms were produced, and so they remain ubiquitous, persistent environmental contaminants that are routinely found in samples of human and animal tissues (Fisher, 1999). The central nervous system is one of the target organs for PCBs (Altmann et al., 2001), and PCBs are known to be neurotoxic, inducing neurochemical and behavioral disruptions in humans and experimental animals (Brouwer et al., 1995). Cognitive/motor deficits were found among children who had accidental perinatal exposure to PCBs in cooking oil (Rogan and Gladen, 1992).

It has been proved that disruption of the hypothalamus-pituitary-interrenal axis by PCBs involves brain glucocorticoid receptor down-regulation (Aluru et al., 2004). The effects of PCBs on brain development may be attributable, at least in part, to their ability to reduce circulating levels of thyroid hormone. However, the developmental effects of PCB exposure are not simply a function of PCB-induced hypothyroidism (Zoeller et al., 2000). A high dose of PCBs appears to result in neuronal loss, as reflected by a decrease in glucocorticoid receptor (GR), heat shock protein 90 (hsp90), and heat shock protein 70 (hsp70) expression, leading to the disruption of the negative feedback regulation of cortisol during stress (Aluru et al., 2004). In recent years, learning deficits and endocrine disrupting actions have been observed whereby the non-coplanar congener exhibits neurotoxicities via different mechanisms with coplanar PCB (Zhou et al., 2004).

52 However, the neurotoxic mechanism of PCBs is complex and actually still unclear.
53 The aim of this study was to identify the transcripts which were up- or down-regulated by
54 PCBs in the brain of the marine fish *Sebastiscus marmoratus* using suppression PCR
55 subtractive hybridization (SSH). This is a highly effective genome-wide approach that
56 enriches for differentially expressed mRNA transcripts by generating subtracted cDNA
57 libraries (Alonso and Leong, 2002; Diatchenko et al., 1996; Straub et al., 2004).
58 Differential expression of some of the genes identified by SSH was confirmed by real time
59 quantitative PCR. This method is very sensitive for the detection and quantification of gene
60 expression levels, in particular for low abundance mRNA (Feng et al., 2007).

61

62 **2. Materials and methods**

63 **2.1. Chemicals**

64 Aroclor 1254, a commercial PCB mixture was purchased from Sigma-Aldrich
65 (Sigma-Aldrich Co., USA), and a PCR-Select cDNA Subtraction Kit was obtained from BD
66 Biosciences (BD Biosciences, UK). All other chemicals were of analytical grade and were
67 obtained from commercial sources.

68 **2.2. Treatments and sample collection**

69 All animal experiments were conducted according to the research protocols approved
70 by Xiamen University Institutional Animal Care and Use Committee. Female *Sebastiscus*
71 *marmoratus* weighing 23-27 g, were captured from the coastal area of Xiamen city, Fujian
72 Province, China. Before the exposure experiment, the fish were acclimated in tanks (25 fish
73 in each tank) containing 50 L of aerated sand-filtered seawater, with natural photoperiod

for 7 days. Fish were dosed via single intraperitoneal injection with PCBs (3 mg/kg body weight in corn oil) or with only corn oil as a control. After 3 days, fish were randomly sampled from each group. The isolated brains were frozen in liquid N₂ and immediately stored at -80 °C until analysis.

2.3. Suppression subtractive hybridization

For the SSH procedure (PCR-Select cDNA Subtraction Kit, BD Biosciences, UK) mRNA was isolated from fish brains (RNAiso Reagent and *OligotexTM-dT30<Super>* mRNA Purification Kit, TaKaRa Biotechnology (Dalian), PR. China) and two libraries constructed in a pMD18-T vector (TaKaRa Biotechnology (Dalian), PR. China), enriched for genes which had been up-regulated (forward subtracted) or down-regulated (reverse subtracted) in response to PCBs. For each subtracted cDNA library, 150 random white colonies were selected and PCR amplifications were performed with nested PCR (primer sequences: forward 5'-TCG AGC GGC CGC CCG GGC AGG T-3', reverse 5'-AGC GTG GTC GCG GCC GAG GT-3'). Candidate positive clones from the subtracted cDNA library were selected for sequencing. The sequence data were analyzed using the Vec Screen program (NCBI) for vector masking, and all adaptor sequences were removed. Each sequence of the isolated clones was used for homology searches in GenBank by BLASTN (NCBI)

2.4. Real time fluorescent quantitative PCR

The real time fluorescent quantitative PCR analysis used total RNA isolated from single-brain pools (n=6). SYBR green I was used as the fluorescent dye for real time quantitative PCR on a Rotorgene 3000 (ABI, USA). The housekeeping gene of beta-actin

was used as the endogenous assay control (primer sequences: forward 5'-AGG GAA ATC GTG CGT G-3', reverse 5'-ATG ATG CTG TTG TAG GTG GT-3'). PCR reactions were performed in triplicate for each cDNA sample.

The Relative Expression Software Tool - Multiple Condition Solver (REST-MCS ©-version 2) was used to calculate the relative expression of target genes mRNA in real time fluorescent quantitative PCR using Pair Wise Fixed Reallocation Randomization Test© (Pfaffl et al., 2002). The mathematical model used is based on the mean crossing point deviation between sample and control group of target gene, normalized by the mean crossing point deviation of reference genes. Specific amplification efficiencies are included in the correction of the quantification ratio. Significant differences between groups were determined by ANOVA and secondary testing was done by using Tukey's LSD.

3. Results and discussion

PCR screening of colonies for recombinants was done with insert-specific primers in parallel with the generation of archived, 96-well format, bacterial stock cultures. Colonies containing inserts were identified using agarose gel electrophoresis of the PCR products. Eighty clones of forward subtraction and 28 of reverse subtraction were selected and sequenced. A total of 108 expressed gene sequences were identified by BLAST searches. Seventy-nine transcripts were obtained and analyzed by BLASTN (nucleic acid) and TBLASTX (translated protein query search) and a selection are shown in Table 1. The homology search revealed that 45 clones contained substantial sequence homologies ($E\text{-value} < 1e^{-6}$) with known gene fragments or full-length cDNAs (most belong to

118 *Tetraodon nigroviridis*) and 34 possibly novel genes, showing no significant homologies
 119 with any known sequences in the GenBank.

120 The clones could be arranged into a number of distinct functional groups
 121 representative of protective mechanisms associated with responses to PCB exposure,
 122 including genes involved in PCB metabolism, such as *cytochrome b* and *cytochrome c*
 123 (Goksoyr and Forlin, 1992); some correlated with neural functions, such as *snap25a*
 124 (Roberts et al., 1998), *rtn1* (De Craene et al., 2006); and of mechanisms of energy
 125 metabolism and cellular regulation, while a number were cryptic (classifications are given
 126 in Table 1).

127 **Positions of Table 1**

128 The real time quantitative PCR primers for 20 candidate ESTs were designed, and 17
 129 were used for quantitative PCR (Table 2). The differential expression of 17 ESTs included
 130 10 which encoded known proteins likely to be involved in PCB neurotoxicity, 5 which
 131 belong to function-unknown cDNA/mRNA, and 2 were potentially novel gene transcripts.
 132 The results showed the presence of differentially expressed genes in the brains of the
 133 PCB-treated fish compared with the control group (Figure 1).

134 **Position of Table 2**

135 **Position of Figure 1**

136 Long-term potentiation (LTP) is a persistent synaptic enhancement induced by
 137 high-frequency stimulation of afferents and has been suggested to be an important
 138 component of the cellular basis of certain forms of memory and learning (Collingridge,
 139 1987). LTP has been introduced in neurotoxicological studies on PCBs, and PCB-induced

140 LTP deficits reveal that there are multiple factors have to be considered besides the
 141 N-methyl-D-aspartate (NMDA) receptor (Altmann et al., 2001; Gilbert et al., 2000). In the
 142 present study, EST 13 and EST 14 contained substantial sequence homologies with genes
 143 of the α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor β subunit
 144 and calmodulin respectively, which correspond to LTP (Sandkuhler, 2007). It has been
 145 reported that LTP deficits responding to PCBs can only be partly related to the reduction of
 146 binding sites for the NMDA receptor (Altmann et al., 2001). Our result suggested that the
 147 AMPA receptor was also induced by PCB-treatment. The down-regulation of these genes
 148 could induce LTP deficits, which may be one likely mechanism for neurochemical and
 149 behavioral disruptions responding to PCBs. EST 9 is homologous with the gene of 25 kDa
 150 synaptosomal-associated protein (SNAP-25), which is a presynaptic protein involved
 151 in neurotransmitter exocytosis at the nerve terminal (Roberts et al., 1998). At the
 152 neuronal synapse, the central role of SNAP-25 is evident from the impairment of
 153 neuronal secretion by botulinum neurotoxin A/E, a toxin that proteolytically
 154 inactivates SNAP-25 (Hayashi et al., 1994). Moreover, loss of SNAP-25 expression
 155 in mice results in embryonic lethality late in gestation with evoked
 156 neurotransmitter release absent at the neuromuscular junction and central
 157 synapses (Washbourne et al., 2002). This study showed that the expression of
 158 SNAP-25 was decreased after PCB-treatment, which is likely to have induced the
 159 disruption of synaptic plasticity and LTP, resulting in the inordinateness of central
 160 neural system functions. EST 1 is homologous to a gene of Tob (the transducer of
 161 ErbB-2), which is well known as a negative regulator in peripheral cells (Matsuda et al.,

1996). The elevated expression level of Tob responding to PCB-treatment suggested that Tob might be another potential core gene involved in the neurotoxicity of PCBs.

4. Conclusions

In conclusion, our results identified a number of ESTs suitable for further development as environmental biomarkers, most of which had novel sequences and although some remained cryptic, those putatively identified were informative regarding the potential mechanisms of PCB neurotoxicity. However, further investigation is required to understand the effects of PCB-induced neurotoxicity in marine fish.

Acknowledgements

This work was supported by the National Natural Science Foundation (40606027) of China and by Fund (MEL0606) of the State Key Laboratory of Marine Environmental Science, Xiamen University, China. Professor John Hodgkiss is thanked for his assistance with English.

References

- Alonso, M. and Leong, J.A., 2002. Suppressive subtraction libraries to identify interferon-inducible genes in fish. *Mar Biotechnol* (NY) 4, 74-80.
- Altmann, L., Mundy, W.R., Ward, T.R., Fastabend, A. and Lilienthal, H., 2001. Developmental exposure of rats to a reconstituted PCB mixture or aroclor 1254: effects on long-term potentiation and [3H]MK-801 binding in occipital cortex and hippocampus. *Toxicol Sci* 61, 321-330.
- Aluru, N., Jorgensen, E.H., Maule, A.G. and Vijayan, M.M., 2004. PCB disruption of the hypothalamus-pituitary-interrenal axis involves brain glucocorticoid receptor downregulation in anadromous Arctic charr. *Am J Physiol Regul Integr Comp Physiol* 287, R787-793.
- Brouwer, A., Ahlborg, U.G., Van den Berg, M., Birnbaum, L.S., Boersma, E.R., Bosveld, B., Denison, M.S., Gray, L.E., Hagmar, L., Holene, E. and et al., 1995. Functional aspects of developmental toxicity of

- polyhalogenated aromatic hydrocarbons in experimental animals and human infants. *Eur J Pharmacol* 293, 1-40.
- Collingridge, G., 1987. Synaptic plasticity. The role of NMDA receptors in learning and memory. *Nature* 330, 604-605.
- De Craene, J.O., Coleman, J., Estrada de Martin, P., Pypaert, M., Anderson, S., Yates, J.R., 3rd, Ferro-Novick, S. and Novick, P., 2006. Rtn1p is involved in structuring the cortical endoplasmic reticulum. *Mol Biol Cell* 17, 3009-3020.
- Diatchenko, L., Lau, Y.F., Campbell, A.P., Chenchik, A., Moqadam, F., Huang, B., Lukyanov, S., Lukyanov, K., Gurskaya, N., Sverdlov, E.D. and Siebert, P.D., 1996. Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proc Natl Acad Sci U S A* 93, 6025-6030.
- Feng, L., Miao, W. and Wu, Y., 2007. Differentially expressed genes of *Tetrahymena thermophila* in response to tributyltin (TBT) identified by suppression subtractive hybridization and real time quantitative PCR. *Aquat Toxicol* 81, 99-105.
- Fisher, B.E., 1999. Most unwanted. *Environ Health Perspect* 107, A18-23.
- Gilbert, M.E., Mundy, W.R. and Crofton, K.M., 2000. Spatial learning and long-term potentiation in the dentate gyrus of the hippocampus in animals developmentally exposed to Aroclor 1254. *Toxicol Sci* 57, 102-111.
- Goksoyr, A. and Forlin, L., 1992. The Cytochrome P-450 System in Fish, *Aquatic Toxicology and Environmental Monitoring Aquatic Toxicology* 22, 25.
- Hayashi, T., McMahon, H., Yamasaki, S., Binz, T., Hata, Y., Sudhof, T.C. and Niemann, H., 1994. Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on assembly. *EMBO J* 13, 5051-5061.
- Matsuda, S., Kawamura-Tsuzuku, J., Ohsugi, M., Yoshida, M., Emi, M., Nakamura, Y., Onda, M., Yoshida, Y., Nishiyama, A. and Yamamoto, T., 1996. Tob, a novel protein that interacts with p185erbB2, is associated with anti-proliferative activity. *Oncogene* 12, 705-713.
- Pfaffl, M.W., Horgan, G.W., Dempfle, L. Relative Expression Software Tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res.* 30, e36, 2002.
- Roberts, L.A., Morris, B.J. and O'Shaughnessy, C.T., 1998. Involvement of two isoforms of SNAP-25 in the expression of long-term potentiation in the rat hippocampus. *Neuroreport* 9, 33-36.
- Rogan, W.J. and Gladen, B.C., 1992. Neurotoxicology of PCBs and related compounds. *Neurotoxicology* 13, 27-35.
- Sandkuhler, J., 2007. Understanding LTP in pain pathways. *Mol Pain* 3, 9.
- Straub, P.F., Higham, M.L., Tanguy, A., Landau, B.J., Phoel, W.C., Hales, L.S., Jr. and Thwing, T.K., 2004. Suppression subtractive hybridization cDNA libraries to identify differentially expressed genes from contrasting fish habitats. *Mar Biotechnol (NY)* 6, 386-399.
- Tilson, H.A. and Kodavanti, P.R.S., 1997. Neurochemical effects of polychlorinated biphenyls: An overview and identification of research needs. *Neurotoxicology* 18, 727-743.
- Washbourne, P., Thompson, P.M., Carta, M., Costa, E.T., Mathews, J.R., Lopez-Bendito, G., Molnar, Z., Becher, M.W., Valenzuela, C.F., Partridge, L.D. and Wilson, M.C., 2002. Genetic ablation of the t-SNARE SNAP-25 distinguishes mechanisms of neuroexocytosis. *Nat Neurosci* 5, 19-26.
- Zhou, J., Qin, Z., Cong, L. and Xu, X., 2004. Research progress of the endocrine disrupting activities of polychlorinated biphenyls. *Chinese Science Bulletin* 49, 6.
- Zoeller, R.T., Dowling, A.L. and Vas, A.A., 2000. Developmental exposure to polychlorinated biphenyls

233 exerts thyroid hormone-like effects on the expression of RC3/neurogranin and myelin basic protein
234 messenger ribonucleic acids in the developing rat brain. *Endocrinology* 141, 181-189.

ACCEPTED MANUSCRIPT

235 Table 1 Characterisation of ESTs isolated from polychlorinated biphenyls-treated
236 *Sebastiscus marmoratus*.

237

238 Table 2 Primers used for real time quantitative PCR.

239

240 Figure 1. Figure 1. Gene expression profile of genes that were affected by PCBs: The
241 values were gene absolute regulation calculated by REST-MCS ©- version 2 from
242 duplicate PCR determinations of three independent experiments for each set of conditions
243 tested (PCB-treated and the control). * & ** indicate expression levels significantly
244 different from the control ($p < 0.05$ & $p < 0.01$ respectively) by Tukey's LSD.

245

Table 1
Characterisation of ESTs isolated from polychlorinated biphenyls-treated *Sebastiscus marmoratus*

Homologous gene & species	EST #	Insert (bp)	E-value	Change	GenBank Access No.
<u>LTP and neurotransmitter release related</u>					
<i>Danio rerio</i> SNAP 25a	9	271	7e ⁻⁴⁷	↓	<u>FF403915</u>
<i>Carassius auratus</i> kainate receptor (AMPA)beta subunit	13	384	4e ⁻⁶³	↓	<u>FF403841</u>
<i>O. latipes</i> calmodulin	14	541	2e ⁻¹⁶²	↓	<u>FF403842</u>
<i>Scophthalmus maximus</i> clone sba441 ependymin	20	486	2e ⁻³¹	↓	<u>FF403849</u>
<u>Neuroendocrine</u>					
<i>Gasterosteus aculeatus</i> RTN1	8	321	7e ⁻¹¹²	↓	<u>FF403914</u>
<i>Paralichthys olivaceus</i> carboxypeptidase H	22	587	2e ⁻¹³²	↓	<u>FF403851</u>
<u>Mitosis and Cell proliferation related</u>					
<i>Gillichthys mirabilis</i> transducer of ERBB-2(Tob)	1	305	4e ⁻³⁰	↑	<u>FF403837</u>
<i>Danio rerio</i> dynein cytoplasmic 1 heavy chain 1	6	470	5e ⁻²⁵	↓	<u>FF403892</u>
<i>Oreochromis mossambicus</i> stathmin-like mRNA	23	745	0.0	↓	<u>FF403852</u>
<u>Energy-related metabolism</u>					
<i>Scombridae</i> gen. sp. cytochrome c oxidase subunit VI b precursor	4	268	5e ⁻⁶⁰	↑	<u>FF403870</u>
<i>Sebastes maliger</i> cytochrome b	5	326	1e ⁻⁹¹	↑	<u>FF403881</u>
<i>Rhabdosargus sarba</i> sodium potassium ATPase α subunit	15	754	0.0	↓	<u>FF403843</u>
<i>Helicolenus hilgendorfi</i> ATPase 6	26	165	2e ⁻⁵²	↑	<u>FF403855</u>
<i>Kyphosus vaigiensis</i> NADH dehydrogenase subunit 2	31	272	7e ⁻³⁹	↑	<u>FF403861</u>
<u>General metabolism</u>					
<i>Danio rerio</i> proteasome subunit Y	32	246	5e ⁻³²	↓	<u>FF403862</u>
<i>Sphyræna argentea</i> lactate dehydrogenase-A	52	487	3e ⁻⁴⁷	↑	<u>FF403884</u>
<i>Danio rerio</i> glutamic-oxaloacetic transaminase 1, soluble	56	439	8e ⁻⁹²	↓	<u>FF403888</u>
<u>Signal proteins</u>					
<i>Danio rerio</i> zinc finger protein zic1	10	379	4e ⁻¹⁰⁷	↓	<u>FF403838</u>
<i>Sparus aurata</i> V-Fos transformation effector-like protein	11	468	0.0	↓	<u>FF403839</u>
<i>Danio rerio</i> suppression of tumorigenicity 13 (colon carcinoma) (Hsp70 interacting protein)	21	687	5e ⁻¹²²	↓	<u>FF403850</u>
<i>Siniperca chuatsi</i> clone C311 eukaryotic translation elongation factor -1 gamma	34	270	5e ⁻⁶³	↑	<u>FF403864</u>
<i>Danio rerio</i> zinc finger protein zic1	36	197	1e ⁻⁴¹	↓	<u>FF403866</u>
<i>Homo sapiens</i> zinc finger protein 238 (ZNF238), transcript variant 1	74	310	1e ⁻¹⁸	↓	<u>FF403908</u>

(continued on next page)

249 Table 1 (continued)

Homologous gene & species	EST #	Insert (bp)	E-value	Change	GenBank Access No.
<u>Immune system</u>					
<i>Epinephelus akaara</i> clone 004 MHC class I alpha antigen	25	353	5e ⁻¹¹	↑	<u>FF403854</u>
<u>Hemopoiesis system</u>					
<i>Oryzias latipes</i> embryonic alpha-type globin	12	371	3e ⁻⁵⁹	↓	<u>FF403840</u>
<u>Ribosomal proteins</u>					
<i>Platichthys flesus</i> 60S ribosomal protein L28	24	308	1e ⁻¹²⁵	↑	<u>FF403853</u>
<i>Paralichthys olivaceus</i> ribosomal protein L17	28	456	1e ⁻¹⁴⁵	↑	<u>FF403857</u>
<i>Siniperca chuatsi</i> clone C426 60S ribosomal protein L35	29	180	6e ⁻⁷¹	↑	<u>FF403858</u>
<i>Pagrus major</i> 60S ribosomal protein L24	30	346	6e ⁻⁸⁸	↓	<u>FF403860</u>
<i>Solea senegalensis</i> ribosomal protein L35a	51	394	3e ⁻¹²²	↑	<u>FF403883</u>
<i>Siniperca chuatsi</i> clone C123 ribosomal protein L7	57	561	0.0	↓	<u>FF403889</u>
<i>Pagrus major</i> ribosomal protein L37	65	259	2e ⁻⁷⁶	↑	<u>FF403898</u>
<u>Identified genes but function unknown</u>					
<i>Tetraodon nigroviridis</i> full-length cDNA	7	428	1e ⁻⁶¹	↓	<u>FF403903</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	16	342	3e ⁻²⁵	↓	<u>FF403844</u>
<i>Gasterosteus aculeatus</i> clone CNB134-E09 mRNA	17	467	1e ⁻⁵²	↓	<u>FF403845</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	18	262	6e ⁻²¹	↓	<u>FF403846</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	19	300	2e ⁻¹⁶	↓	<u>FF403847</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	27	410	4e ⁻¹²⁷	↑	<u>FF403856</u>
<i>Gasterosteus aculeatus</i> clone CFW195-D 03 mRNA	33	220	1e ⁻⁷	↑	<u>FF403863</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	38	474	2e ⁻¹⁰	↑	<u>FF403868</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	49	111	3e ⁻¹⁰	↑	<u>FF403880</u>
<i>Gasterosteus aculeatus</i> clone CGX01-B09 mRNA	54	514	3e ⁻⁶⁰	↓	<u>FF403886</u>
<i>Gasterosteus aculeatus</i> clone CNB24-E09 mRNA	62	335	7e ⁻⁰⁹	↑	<u>FF403895</u>
<i>Tetraodon nigroviridis</i> full-length cDNA	73	453	3e ⁻¹⁵	↑	<u>FF403907</u>
<i>Gasterosteus aculeatus</i> clone CNB134-E09 mRNA	78	264	5e ⁻²⁹	↓	<u>FF403912</u>
<u>Unidentified genes</u>					
No homologous region	2	423	—	↑	<u>FF403848</u>
No homologous region	3	453	—	↑	<u>FF403859</u>
No homologous region	35	392	—	↑	<u>FF403865</u>
No homologous region	37	191	—	↑	<u>FF403867</u>
No homologous region	39	328	—	↑	<u>FF403869</u>
No homologous region	40	548	—	↑	<u>FF403871</u>
No homologous region	41	424	—	↓	<u>FF403872</u>
No homologous region	42	496	—	↓	<u>FF403873</u>
No homologous region	43	627	—	↓	<u>FF403874</u>

(continued on next page)

250 Table 1 (*continued*)

Homologous gene & species	EST #	Insert (bp)	E- value	Change	GenBank Access No.
No homologous region	44	267	—	↓	<u>FF403875</u>
No homologous region	45	595	—	↓	<u>FF403876</u>
No homologous region	46	756	—	↓	<u>FF403877</u>
No homologous region	47	767	—	↓	<u>FF403878</u>
No homologous region	48	611	—	↓	<u>FF403879</u>
No homologous region	50	179	—	↑	<u>FF403882</u>
No homologous region	53	1143	—	↑	<u>FF403885</u>
No homologous region	55	654	—	↓	<u>FF403887</u>
No homologous region	58	234	—	↓	<u>FF403890</u>
No homologous region	59	139	—	↓	<u>FF403891</u>
No homologous region	60	315	—	↓	<u>FF403893</u>
No homologous region	61	510	—	↓	<u>FF403894</u>
No homologous region	63	409	—	↑	<u>FF403896</u>
No homologous region	64	246	—	↑	<u>FF403897</u>
No homologous region	66	332	—	↑	<u>FF403899</u>
No homologous region	67	931	—	↓	<u>FF403900</u>
No homologous region	68	612	—	↓	<u>FF403901</u>
No homologous region	69	268	—	↓	<u>FF403902</u>
No homologous region	70	240	—	↓	<u>FF403904</u>
No homologous region	71	495	—	↓	<u>FF403905</u>
No homologous region	72	519	—	↓	<u>FF403906</u>
No homologous region	75	402	—	↓	<u>FF403909</u>
No homologous region	76	342	—	↓	<u>FF403910</u>
No homologous region	77	555	—	↓	<u>FF403911</u>
No homologous region	79	338	—	↓	<u>FF403913</u>

251

252 Table 2
 253 Primers used for real time quantitative PCR
 254

EST #	GenBank Access No.	Forward Primer(5'→3')	Reverse Primer(5'→3')	Tm(°C)
1	<u>FF403837</u>	ACTATAATCCTTGAAAGTCCAGT	AACTTCAGTCTGAGCAACATG	53
2	<u>FF403848</u>	ACCTGAGGGACAGAAACG	TCCCAGAGCCACAACAC	53
3	<u>FF403859</u>	GAAATATCACAAGCGTGGC	GCTCATCTGAATCCCTTTTC	53
6	<u>FF403892</u>	ATAATCACCTACATTATTGCT	TACCATCGGTCAACATTCTC	53
7	<u>FF403903</u>	AGAAAGAATACCCGGACAG	GCGATGACGTGGTCACA	53
8	<u>FF403914</u>	ATACAAAAGCAAAAACCCTAC	GCGAAATGCCACAGA	53
9	<u>FF403915</u>	GCAAAATGTCCACCACTTC	GCACACCAAATCGCCA	53
10	<u>FF403838</u>	GTGCTACTCAAACCTGGACCT-	AATATTTGCGATGAATGGAC	53
11	<u>FF403839</u>	GAAACCCACGCAGAACAG	AGCTCATCACCGAGGATG	53
12	<u>FF403840</u>	ATCTTGAAGTTGGCAGGG	ACTGGAAGGACTTGAACCC	53
13	<u>FF403841</u>	ACGGGCTTCTTTGATTTC	GAATCTGTTCTGGCTCACTC	53
14	<u>FF403842</u>	CACAGACAGCGAGGAGGA	CGCCGTCATCATTTGTAC	53
15	<u>FF403843</u>	AAGAAGTCCATCGCCTACA	GGTTCCTGGGCTGTCTC	53
16	<u>FF403844</u>	TTTTGGGGGATTTTATGC	CACAGAAGATACCTTGGACAG	53
17	<u>FF403845</u>	CCTCATCATTCGTAGTCAGC	CACTTGGATAGGGGTTGG	53
18	<u>FF403846</u>	GAGAAGAGGGCTTTTAGAGAG	TTGATAGCAGTTGTAATTGTCTG	53
19	<u>FF403847</u>	TTGAATCACTTCCTCCTCTG	GCCAAGGCTATACCTGCT	53

255

Figure 1. Gene expression profile of genes that were affected by PCBs.

The values were gene absolute regulation calculated by REST-MCS ©- version 2 from duplicate PCR determinations of three independent experiments for each set of conditions tested (PCB-treated and the control). * & ** indicate expression levels significantly different from the control ($p < 0.05$ & $p < 0.01$ respectively) by Tukey's LSD.