Implications of mycosporine-like amino acid and antioxidant defenses in UV-B radiation tolerance for the algae species and

Tse-Min Lee, Chia-Tai Shiu

► To cite this version:

Tse-Min Lee, Chia-Tai Shiu. Implications of mycosporine-like amino acid and antioxidant defenses in UV-B radiation tolerance for the algae species and. *Marine Environmental Research*, 2008, 67 (1), pp.8. 10.1016/j.marenvres.2008.09.006 . hal-00563058

HAL Id: hal-00563058

<https://hal.science/hal-00563058>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Implications of mycosporine-like amino acid and antioxidant defenses in UV-B radiation tolerance for the algae species *Ptercladiella capillacea* and *Gelidium amansii*

Tse-Min Lee, Chia-Tai Shiu

PII: S0141-1136(08)00215-8

DOI: [10.1016/j.marenvres.2008.09.006](https://doi.org/10.1016/j.marenvres.2008.09.006)

Reference: MERE 3291

To appear in: *Marine Environmental Research*

Received Date: 15 June 2008

Revised Date: 1 September 2008

Accepted Date: 27 September 2008

Please cite this article as: Lee, T-M., Shiu, C-T., Implications of mycosporine-like amino acid and antioxidant defenses in UV-B radiation tolerance for the algae species *Ptercladiella capillacea* and *Gelidium amansii*, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.09.006](https://doi.org/10.1016/j.marenvres.2008.09.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Implications of mycosporine-like amino acid and antioxidant defenses in UV-B
radiation tolerance for the algae species *Ptercladiella capillacea* and *Gelidium*
amansii

Tse-Min Lee and Chia-Tai Shiu

Institute of Marine Biology, National Sun Yat-sen University, Kaohsiung, Taiwan 80424, Republic of China

*The Kuroshio Research Group of the Asia-Pacific Ocean Research Center, National Sun Yat-sen University,
Kaohsiung, Taiwan 80424, Republic of China*

Corresponding author: Tse-Min Lee; tel: 886-7-5252000 ext. 5110; fax: 886-7-5251509; e-mail:

tmllee@mail.nsysu.edu.tw

Running title: MAA and UV-B-induced oxidative responses

Abstract

Ultraviolet-B (UV-B) radiation (0.5, 1.0, 1.5, 3.0 W m⁻²) induced higher H₂O₂ production and lipid peroxidation in alga *Gelidium amansii* inhabiting in lower subtidal regions than upper subtidal alga *Ptercladiella capillacea*. Compared to *G. amansii*, mycosporine-like amino acid (MAA) concentration in *P. capillacea* was higher and can be increased by 0.5-1.0 W m⁻² UV-B, while carotenoid concentration was lower but also increased by 1.5-3.0 W m⁻² UV-B. UV-B increased ascorbate concentration, but to a higher degree in *P. capillacea*. UV-B decreased glutathione concentration, but to a higher degree in *G. amansii*. UV-B increased ascorbate peroxidase (APX) and glutathione reductase (GR) activities in *P. capillacea* but decreased them in *G. amansii*. UV-B increased superoxide dismutase and catalase activities, but to a higher degree in *G. amansii*. So, *G. amansii* suffered greater oxidative stress from UV-B radiation. *Ptercladiella capillacea* can effectively reduce UV-B sensitivity by increasing sunscreen ability and antioxidant defense capacity.

Keywords: Antioxidant; carotenoid; *Gelidium amansii*; *Ptercladiella capillacea*; UV-B radiation

1. Introduction

The stratospheric ozone layer is vital to life on Earth because it is the principal agent that absorbs ultraviolet radiation in the Earth's atmosphere. The depletion of the stratospheric ozone layer due to anthropogenic and natural causes since 1979 has led to increasing levels of solar ultraviolet-B (UV-B) radiation (280-320 nm) reaching the Earth's surface, especially in the Arctic, Antarctic and temperate regions (Seckmeyer and McKenzie, 1992; Kerr and McElroy, 1993; Russell et al., 1996), where UV-B and UV-A (320-400 nm) radiation are capable of penetrating the water column to an ecologically-significant depth (Calkins and Thordardottir, 1982; Whitehead et

al., 2000).

Since UV-B radiation damages DNA, algae evolved counteract mechanisms for the repairing of UV-induced damage of DNA by photoreactivation and excision repair (Britt, 1995; Kim and Sancar, 1995). Most algae synthesize mycosporine-like amino acids (MAAs) as an effective UV sunscreen compound that absorbs harmful UV radiation and emits heat or a fluorescence emission to prevent UV-induced photodamage (Riegger and Robinson, 1997; Dunlap and Shick, 1998; Karsten et al., 1998). MAAs are water-soluble substances characterized by a cyclohexenone or cyclohexenimine chromophore conjugated with the nitrogen substituent of an amino acid or its imino alcohol, having absorption maxima ranging from 310 to 360 nm (Nakamura et al., 1982). The MAA content shows a negative correlation with water depth, but the ability for MAA synthesis in response to environmental changes is variable among algae (Hoyer et al., 2001). MAAs also act as antioxidants, for example, mycosporine-glycine shows a moderate antioxidant potential (Dunlap and Yamamoto, 1995).

UV-B radiation has been shown to reduce the photosynthetic electron transport and dark reaction activity (Brandle, 1977; Foyer and Harbinson, 1994), and thus leads to excess ROS formation (Langebartels et al., 2000). As a result, the cellular homeostasis of ROS production is disrupted and causes oxidative stress, that is, oxidative damage to macromolecules (Foyer et al., 1994; Rijstenbil, 2002). Induction of ROS over-production and oxidative stress by UV-B radiation has been observed in microalgae and macroalgae (Tschiersch and Ohmann, 1993; Foyer et al., 1994; Lesser, 1996; Malanga et al., 1997; He et al., 2002; Rijstenbil, 2002; Shiu and Lee, 2005).

The increases in carotenoids, detoxifying enzymes and antioxidants provide UV-B protection by scavenging ROS (Mittler and Tel-Or, 1991; Middleton and Teramura, 1993). Ascorbate (AsA), glutathione (GSH), α -tocopherol, and carotenoids are considered effective antioxidants (Noctor and Foyer, 1998; Smirnoff and Wheeler, 2000). Carotenoids play an essential photoprotective role by quenching triplet state chlorophyll molecules and scavenging singlet oxygen and other toxic oxygen species formed within the chloroplast (Young, 1991). To scavenge ROS enzymatically, superoxide

dismutase (SOD; EC 1.15.1.1) is the first line of detoxifying enzymes that converts $O_2^{\cdot -}$ to a less toxic ROS, H_2O_2 . Then, ascorbate peroxidase (APX; EC 1.11.1.11) degrades H_2O_2 via AsA oxidation and oxidized AsA, such as dehydroascorbate (DHA), is reduced to regenerate AsA by dehydroascorbate reductase (DHAR; EC 1.8.5.1) using GSH as the electron donor (Asada, 1999). Oxidized GSH, GSSG, is reduced to GSH by glutathione reductase (GR; EC 1.6.4.2) utilizing reducing equivalents from NAD(P)H. Catalase (CAT; EC 1.11.1.6) is also responsible for H_2O_2 removal (Willenken et al., 1997).

At low latitudes, ambient UV-B radiation that is high due to the high solar angle and the relatively small amount of stratospheric ozone and can reach more deeply into oligotrophic tropical water column because of the relatively low quantities of dissolved and particulate organic matter (Baker et al., 1980). It is expected that seaweeds from tropical regions encounter higher UV radiation than those occurring in temperate and Polar regions even under the normal ozone layer (Häder, 1993). In subtropical waters, the coastline along northeastern Taiwan is dominated by the rhodophyte *Pterocladia capillacea* (S.G. Gmelin) Santelices & Hommersand in the upper subtidal region (Fig. 1A) while another rhodophyte *Gelidium amansii* (J.V. Lamouroux) J.V. Lamouroux is abundant in the lower subtidal region (Fig. 1C). These two algae belong to the Family Gelidiaceae but different genus (Gelidiales, Rhodophyta). *Pterocladia capillacea* is ascribed to the genus *Pterocladia* and *G. amansii* is ascribed to the genus *Gelidium*. Our surveys during 2004-2005 in four sites along northeastern Taiwan coast (see 2.1. Algal materials and treatments) showed that *P. capillacea* was significantly abundant in the subtidal region of 1-m water depth while *G. amansii* was abundant in the regions of 3-m and 5-m water depth ($p < 0.0001$). Besides, they appeared significant temporal variation ($p < 0.0001$); *P. capillacea* was abundant in the summer (September, 2004) while *G. amansii* was absent in the summer. It is therefore interesting for us to understand the factors influencing the temporal and spatial differences between the two Gelidiaceae algae. We hypothesize that one of factors driving this differential distribution is the sensitivity to high UV/solar radiation, especially in the summer. The MAA contents of *P. capillacea* (Fig. 1B) were

Figure 1

significantly higher than those of *G. amansii* (Fig. 1D), especially in the summer. It is known that physiological responses of macroalgae to UV-B are related to their zonation pattern. A survey of the UV tolerance of marine rhodophyte from tropical habitats showed that UV exposure caused less pronounced effects in intertidally-occurring species than in subtidal species (van de Poll et al., 2003). The Arctic marine macroalgal species from eulittoral and upper sublittoral regions had higher antioxidant enzyme activities upon exposure to UV radiation than those species from lower sublittoral regions (Aguilera et al., 2002). We therefore propose that differences in UV-B tolerance in term of UV screening capacity and antioxidant defense ability are one of the mechanisms that forces different habitation patterns for *P. capillacea* and *G. amansii* in northeastern Taiwan. In the attempts to elucidate the presence of MAA in the regulation of carotenoid synthesis and antioxidant defense responses in algae differing in UV-B tolerance and their roles in mitigating UV-B/solar radiation toxicity, algal thalli were sampled in the field and immediately transferred to the laboratory for UV-B tolerance experiment and the contents of MAA, carotenoid and antioxidant, and the activities of antioxidant enzymes were examined.

2. Materials and methods

2.1. Algal materials and treatments

In the attempt to determine temporal and spatial changes in the abundance and MAA contents of *P. capillacea* and *G. amansii* in northeastern Taiwan, four sampling sites (Lung-tung, Her-mei, Mo-auo, and Shea-cherng) and four sampling times (June 16-22, 2004, September 21-30, 2004, March 17-23, 2005, and June 3-11, 2005) were set. For each sampling site, three 50-m sampling lines parallel to coastal line were set on 1-m, 3-m and 5-m water depth (MHWS), respectively (a sampling line for each water depth) and 4 random sampling stations were set on each line for the determination of algal abundance (biomass). To estimate algal biomass in each sampling station,

algae in a 50×50 cm² quadrat were scraped and *P. capillacea* and *G. amansii* were weighed and calculated as g wet weight (w. wt.) m⁻². After weighting, algae were fixed in liquid nitrogen and stored in a -70°C freezer until analysis of MAAs.

For laboratory experiments, *P. capillacea* and *G. amansii* of 10-cm plant height were collected in September, 2004 and August, 2005 from the intertidal regions at Her-mei and the subtidal regions at Mo-auo in northeastern Taiwan, respectively. Plants were transported to the laboratory, washed with natural seawater to remove attached sands and incubated at 22-26°C for 14 days in 35‰ (parts per thousand) nutrient-enriched artificial seawater (ASW) containing 403.5 mM NaCl, 10 mM KCl, 10 mM CaCl₂, 30 mM MgSO₄, 10 mM Tris-HCl (pH 8.0), N- (NH₄⁺ and NO₃⁻) and P- (PO₄⁻³) free Provasoli nutrient solutions (Provasoli, 1968), 3 mM NaHCO₃, 400 μM NaNO₃, and 20 μM Na₂HPO₄. The light time was 14 h (photosynthetically active radiation (PAR, 400-700 nm) of 100-120 μmol photons m⁻² s⁻¹ in the absence of algae, achieved by 20 W cool-fluorescent lamps (FL20D, China Electric Apparatus Ltd., Taiwan, ROC)). Before UV-B radiation, thallus segments were cultured in a Petri dish (internal diameter: 30 cm) containing 2 L of 35‰ nutrient-enriched ASW for one day (14 h light/10 h dark) incubation (150 μmol photons m⁻² s⁻¹ PAR provided by 40 W cool-fluorescent lamps (FL40D, China Electric Apparatus Ltd., Taiwan)). Thallus segments of 0.15 g wet weight (w. wt.) were transferred to a Petri dish (2.5 cm in diameter) containing 5 mL of 35‰ nutrient-enriched ASW for UV-B radiation (280-320 nm) (0, 0.5, 1.0, 1.5, and 3.0 W m⁻²), achieved by exposure to UV-B lamps (TLD 15W/08, UVP, Japan) covered with cellulose acetate film (to cut off UVC) of 0.75 mm thickness under 150 μmol photons m⁻² s⁻¹ PAR provided by 40 W cool-fluorescent lamps (FL40D, China Electric Apparatus Ltd., Taiwan). After 14 h of ‘UV-B + PAR’ and ‘PAR’ treatments, thalli were sampled for further analysis. The biologically weighted UV-B radiation over 14 h according to generalized plant action spectrum (normalized to 300 nm; Caldwell, 1971) for low-, medium- and high-UV-B radiation were 0.5-1.0 W m⁻² (25.67-50.17 kJ/m²/14 h), 1.5 W m⁻² (75.83 kJ/m²/14 h) and 3.0 W m⁻² (149.33 kJ/m²/14 h), respectively. Thallus segments were fixed in liquid nitrogen and stored in a -70 °C freezer until analysis. The samples for

MAA analysis and antioxidant enzyme activity were subjected to lyophilization and stored in -70 °C freezer. Each Petri dish was a replicate; there were 3 replicates for each treatment.

2.2. Determination of lipid peroxidation and H_2O_2 production

Thalli of 0.05 g w. wt. were ground in liquid nitrogen and 1 mL of 5% (w/v) trichloroacetic acid was added. After centrifugation at $12,000 \times g$ for 10 min at 4°C, the supernatant was collected. The thiobarbituric acid reacting substance (TBARS) resulting from reaction with thiobarbituric acid (TBA) was determined as lipid peroxidation according to Health and Packer (1968). TBARS content was calculated based on the value of $A_{532}-A_{600}$ with an extinction coefficient of $155 \text{ mM}^{-1} \text{ cm}^{-1}$.

H_2O_2 content was determined by peroxidase-mediated decomposition of H_2O_2 (Okuda et al., 1991). The pH of 0.2 mL of supernatant was adjusted to 7.5 by adding 11.51 μL of 4 M KOH and then centrifuged at $12,000 \times g$ for 1 min at 4°C. The supernatant was collected and applied to a 1 mL column of Amberlite IRA-410, and residual H_2O_2 was washed out with 800 μL of distilled water. H_2O_2 content in the eluent was determined within 10 min after elution. Four hundred μL of 12.5 mM 3-dimethylaminobenzoic acid (DMAB), 0.4 mL of 10 mM 3-methyl-2-benzothiazoline hydrazone (MBTH) and finally 0.02 mL of 0.25 U mL^{-1} horseradish peroxidase (Sigma, USA) were added and measured at 590 nm for 3 min. A H_2O_2 standard curve was determined following the same procedure.

2.3. Extraction and determination of MAA

MAA was extracted and analyzed according to Carreto et al. (2005) with one modification. A sample of 0.005 g dry weight (d. wt.) was homogenized in liquid nitrogen and extracted with 1 mL of 25% (v/v) methanol for 2 h in a 45 °C water bath. The extract was centrifuged at $5,000 \times g$ for 5

min and the supernatant was collected. The pellet was re-extracted twice following the same procedure. Then, the pooled extracts were evaporated to dryness under vacuum (Speed Vac Concentrator SVA 100H, Savant Instruments Inc., Holbrook, NY, USA) and re-dissolved in 500 μL of 25% methanol (v/v). After being centrifuged at $15,000 \times g$ for 5 min, the sample was used for HPLC analysis on a Merck RP-8 column (5 μm , 250 mm $\times$ 4 mm I.D.). The mobile phase consisted of 1% methanol (v/v) run at a flow rate of 0.8 mL min⁻¹. MAAs were detected at 330 nm. Identification was confirmed by spectra, retention time (RT) and by co-chromatography with standards, extracted from the algae *Laurencia papillosa* (C. Agardh) Greville and *Porphyra umbilicalis* (L.) Kützinger obtained from Taiwan (supplementary data). Molar extinction coefficients were obtained from Karsten et al. (1998). MAA concentration was expressed as $\mu\text{mol g}^{-1}$ d. wt.

2.4. Determination of ascorbate and glutathione concentrations

Thalli of 0.02 g w. wt. were ground in liquid nitrogen and 1 mL of 5% (w/v) trichloroacetic acid (TCA) was added. After centrifugation at $12,000 \times g$ for 10 min at 4 °C, the supernatant was collected for determination of its antioxidant content. The measurement of reduced (AsA) and oxidized (DHA) ascorbate concentration were performed by a modification of the method of Hodges et al. (1996). Total ascorbate was determined in 1 mL mixture containing 200 μL TCA extract, 50 mM potassium phosphate buffer (pH 7.4), 3 mM EDTA and 1 mM dithiothreitol (DTT). The mixture was incubated at 25 °C for 10 min, 100 μL of N-ethylmaleimide, 400 μL of 0.61 M TCA, 400 μL of 0.8 M orthophosphoric acid and 400 μL of α, α' -bipyridyl were then added. Finally, 200 μL of FeCl_3 was added and the reaction mixture was incubated in a 55 °C water bath for 10 min and the absorbance was detected at A_{525} . AsA concentrations were determined by adding distilled water instead of DTT and N-ethylmaleimide and then following the above method. Total AsA and AsA concentrations were estimated from a standard curve of 0-40 nmole L-AsA. DHA concentration was calculated by subtraction of AsA concentration from total AsA concentration.

Total glutathione was determined by the method of Griffiths (1980). The concentration of glutathione (reduced form) was estimated from a standard curve of 0-20 nmole GSH. After the removal of GSH by 2-vinylpyridine derivatization, GSSG concentration was determined and GSH concentration was calculated by the subtraction of GSSG concentration from total glutathione concentration.

2.5. Determination of enzyme activity

Enzyme activity was determined according to Shiu and Lee (2005). Lyophilized thalli of 0.006 g d. wt. were homogenized in liquid nitrogen and 0.6 mL of 0.1 M sodium phosphate buffer (pH 6.8) containing 1% (w/v) polyvinylpolypyrrolidone (PVPP) and then 0.25% (v/v) Triton X-100 was added. After centrifugation at $12,000 \times g$ for 10 min at 4 °C, the supernatant was used in activity assays for SOD, CAT and GR. For the APX assay, thalli of 0.005 g d. wt. were homogenized in liquid nitrogen and 0.6 mL of extraction buffer (0.1 M sodium phosphate buffer (pH 6.8) containing 1% (w/v) PVPP, 0.25% (v/v) Triton X-100 and 0.5 mM L-ascorbate) was added. The supernatant after centrifugation at $12,000 \times g$ for 10 min at 4 °C was used for the APX activity assay. Protein content was determined according to Bradford (1976) with bovine serum albumin as the standard.

CAT activity was measured at A_{420} for H_2O_2 decomposition rate using $40 \text{ mM}^{-1} \text{ cm}^{-1}$ as the extinction coefficient. SOD activity was determined by the photochemical inhibition of nitro blue tetrazolium. APX activity was determined by the decrease in ascorbate concentration, which was detected at A_{290} (extinction coefficient: $2.8 \text{ mM}^{-1} \text{ cm}^{-1}$). GR activity was monitored by A_{340} for β -NADPH oxidization as a measure of GSSG reduction.

2.6. Chemicals and statistical analysis

Chemicals were purchased from Merck (Germany) or Sigma (USA). Statistics were analyzed

by SAS (SAS v 8.1, NC, USA). The results presented were the mean of 3 duplicates with a Petri dish as a duplicate (mean $\pm$ SD, $n=3$) and the significance of treatments were tested by 1-way analysis of variance (ANOVA). The biomass and total MAA concentrations for the field surveys were analyzed by 3-way ANOVA (time, site, and water depth). The significance of the differences between means was analyzed by Duncan's new multiple range test followed by significance threshold ANOVA value of $p < 0.05$.

3. Results

Figure 2

3.1. TBARS and H_2O_2 concentration

UV-B radiation did not affect the TBARS content of *P. capillacea* ($p = 0.0672$) (Fig. 2A). UV-B radiation $\geq 1.5 \text{ W m}^{-2}$ increased the TBARS content of *G. amansii* ($p = 0.0254$) (Fig. 2A). H_2O_2 concentration in *P. capillacea* and *G. amansii* increased as UV-B radiation increased ($p < 0.05$); the increase was higher in *G. amansii* than *P. capillacea* (Fig. 2B).

3.2. MAA and carotenoid concentration

Figure 3

The amount of MAA is shown in Fig. 3. Shinorine, porphyra-334, and palythanol can be detected in both algae. Total MAA concentration was higher in *P. capillacea* than in *G. amansii*. Porphyra-334 was the dominant MAA in *P. capillacea* and shinorine was the dominant MAA in *G. amansii*. The amounts of shinorine, porphyra-334, and palythanol in *P. capillacea* were increased by UV-B radiation. A linear increase was observed as UV-B radiation increased from $0.5\text{-}1.0 \text{ W m}^{-2}$ followed by a decline as UV-B radiation $\geq 1.5 \text{ W m}^{-2}$. The amount of MAA in *G. amansii* was decreased by UV-B radiation.

The carotenoid concentration for the control thalli was 2-fold higher in *G. amansii* than in *P.*

capillacea (Fig. 4). The carotenoid concentrations of *P. capillacea* did not change in the range of 0.5-1.0 W m⁻² UV-B radiation, but increased as UV-B radiation increased from 1.5-3.0 W m⁻² (Fig. 4). The carotenoid concentration of *G. amansii* increased slightly from 0.5-1.0 W m⁻² UV-B radiation followed by a decrease as UV-B radiation increased from 1.5-3.0 W m⁻² (Fig. 4).

Figure 4

3.3. Ascorbate and glutathione concentration

Figure 5

UV-B radiation increased AsA concentration ($p = 0.0131$) in *P. capillacea* (Fig. 5A) but did not affect its DHA concentration ($p = 0.1141$) (Fig. 5B). UV-B radiation increased AsA ($p = 0.0234$) and DHA ($p = 0.0130$) concentrations of *G. amansii* (Fig. 5). AsA/DHA ratios in *P. capillacea* increased as UV-B radiation increased, while those of *G. amansii* increased linearly as UV-B radiation increased from 0-1.5 W m⁻² but decreased at 3.0 W m⁻² due to an increase in DHA concentration.

GSH concentration in *P. capillacea* was not affected by UV-B radiation ($p = 0.1582$) (Fig. 5C) while GSSG concentration was decreased by UV-B radiation ($p = 0.0249$) (Fig. 5D). The GSH concentration of *G. amansii* was decreased by UV-B radiation ($p = 0.0435$) (Fig. 5C), but GSSG concentration was increased by UV-B radiation in the range of 0-1.5 W m⁻², followed by a drop at 3.0 W m⁻² ($p = 0.0009$) (Fig. 5D). GSH/GSSG ratios in *P. capillacea* and *G. amansii* were decreased by UV-B radiation. However, the extent of the decrease was greater in *G. amansii*.

3.4. Enzyme activities

SOD activities of *P. capillacea* ($p < 0.0001$) and *G. amansii* ($p = 0.0040$) increased at UV-B radiation ≥ 1.0 W m⁻² and 0.5 W m⁻², respectively, and reached a maximum at UV-B radiation ≥ 1.5 -3.0 W m⁻² (Fig. 6A). The increase in SOD activity caused by UV-B radiation was greater in *G. amansii* than in *P. capillacea*.

CAT activity in *P. capillacea* increased as UV-B radiation increased from 0.5-1.0 W m⁻² and then slightly decreased ($p = 0.0025$), while CAT activity in *G. amansii* was increased by UV-B radiation ≥ 1.5 W m⁻² ($p = 0.0051$) (Fig. 6B). The APX activity of *P. capillacea* was increased by UV-B radiation and reached a maximum at 1.0 W m⁻² UV-B radiation, followed by a slight decrease to a level that was still higher than the control ($p = 0.0052$) (Fig. 6C). In contrast, the APX activity in *G. amansii* decreased linearly as UV-B radiation increased ($p = 0.0408$) (Fig. 6C). GR activity in *P. capillacea* increased linearly as UV-B radiation increased ($p = 0.0062$) while that of *G. amansii* increased to a peak at 1.0 W m⁻², followed a sharp decline to a level below the control when exposed to 1.5-3.0 W m⁻² ($p = 0.0408$) (Fig. 6D).

Figure 6

4. Discussion

This study shows that UV-B radiation causes oxidative stress in *P. capillacea* and *G. amansii*. The over-production of ROS and the induction of oxidative stress by UV-B radiation have been found in several algae and cyanobacteria (Tschiersch and Ohmann, 1993; Foyer et al., 1994; Lesser, 1996; Malanga et al., 1997; He et al., 2002; Rijstenbil, 2002; Shiu and Lee, 2005). The higher levels of lipid peroxidation and H₂O₂ production in *G. amansii* in response to UV-B radiation evidence that *G. amansii* which inhabits the lower subtidal regions was relatively more susceptible to UV-B radiation than *P. capillacea* inhabiting the upper subtidal regions. A survey of the UV tolerance of rhodophytes from tropical habitats showed that UV exposure caused less pronounced effects in intertidally occurring species than in subtidal species (van de Poll et al., 2003). It suggests that algae in the upper subtidal regions have developed mechanisms to cope with high levels of UV/solar radiation. Arctic marine macroalgal species from the eulittoral and upper sublittoral regions exhibit higher antioxidant enzyme activity upon exposure to UV radiation than species from lower sublittoral regions (Aguilera et al., 2002). A study of UV radiation effects on photosynthetic performance, growth and the capability to synthesize MAAs of the aeroterrestrial green algae

Stichococcus sp. and *Chlorella luteoviridis* that form biofilms on building facades, and compared with the responses of green algae from soil (*Myrmecia incisa*) and from brackish water (*Desmodesmus subspicatus*) shows that the biosynthesis and accumulation of MAAs under UV radiation resulted in the reduced sensitivity of growth and photosynthesis to UV radiation, which consequently may enhance survival in an environmentally harsh habitat (Karsten et al., 2007). It is believed that *P. capillacea*, from the upper subtidal regions, has higher amounts of UV absorbing/screening compounds as well as other mechanisms related to UV-B radiation resistance than *G. amansii* from the subtidal regions.

Algae are known to synthesize MAAs as an effective UV sunscreen to prevent UV-induced photodamage (Riegger and Robinson, 1997; Dunlap and Shick, 1998; Karsten et al., 1998). But, the ability to synthesize MAAs in response to environmental changes is different for each algae species. In the present study, the data of Fig. 3 have shown that *P. capillacea* contained 2-fold higher MAA amounts than *G. amansii*, and MAA accumulation was induced by UV-B radiation only in *P. capillacea* but not in *G. amansii*. A similar result was reported by Hoyer et al. (2001) in a study of the distribution pattern of MAA and responses to environmental radiations in Antarctic macroalgae. MAA content was found to be negatively correlated with water depth, while MAA content was not changed after transplantation from deep waters to shallow waters. They have identified 3 physiologically different groups of algae in terms of MAA values: 1. species with no capability for MAA biosynthesis; 2. species with a basic MAA content which is adjusted relative to changes in environmental radiation; 3. species with constant high MAA content irrespective of environmental conditions. Evidently, *P. capillacea* belongs to the group 2 and *G. amansii* is a species similar to group 1. Since MAA is a primary sunscreen compound against UV-B radiation, it can be suggested that *G. amansii* with less MAA is more susceptible to UV-B radiation when compared to *P. capillacea*.

Carotenoid is involved in UV-B radiation defense in *P. capillacea*. As MAA concentration decreased when exposed to 1.5-3.0 W m⁻² UV-B radiation, carotenoid concentration in *P.*

1 *capillacea* increased (see Figs. 3 and 4). *Gelidium amansii* did not accumulate carotenoid in
 2 response to UV-B radiation; it linearly decreased when exposed to 0.5-3.0 W m⁻² UV-B radiation.
 3 Carotenoid performs a photoprotective role by quenching triplet state chlorophyll molecules and
 4 scavenging singlet oxygen and other toxic oxygen species formed within chloroplasts (Young,
 5 1991). The induction of carotenoid and detoxifying enzymes or radical quenchers and antioxidants,
 6 as well as the repair of UV-induced DNA damage, are modulated to counteract the toxicity of UV
 7 radiation (Singh et al., 2008). By increasing the capacity for carotenoid synthesis through the
 8 transformation of tobacco plants with a heterologous carotenoid gene encoding β -carotene
 9 hydroxylase from *Erwinia uredovora* under constitutive promoter control for the conversion of
 10 β -carotene into zeaxanthin, Gotz et al. (2002) suggest that the enhancement of zeaxanthin formation
 11 in the presence of functional xanthophyll cycle contributes to UV protection and the prevention of
 12 UV damage. Thus, carotenoids are accumulated in *P. capillacea* to avoid photodamage when it fails
 13 to accumulate MAA in response to 1.5-3.0 W m⁻² UV-B radiation. As a result, UV-B-induced
 14 photodamage in *P. capillacea* is alleviated.

15 Higher antioxidant concentrations and antioxidant enzyme activities are associated with higher
 16 stress tolerance in algae (Butow et al., 1994; Collén and Davison, 1999a, b). However, the present
 17 results based on the data of Fig. 5 do not support the above view. We found that higher antioxidant
 18 regeneration ability is associated with higher UV-B tolerance. As compared to *G. amansii*, *P.*
 19 *capillacea* had higher AsA and GSH regeneration ability in response to UV-B radiation. The
 20 magnitude of the UV-B-induced increase of AsA concentration in *P. capillacea* was also higher
 21 than that of *G. amansii*.

22 The increases of SOD and CAT activities by UV-B radiation were greater in *G. amansii* than in
 23 *P. capillacea*. The response of SOD, the first line of defense against ROS in plants (Alscher et al.,
 24 2002), to UV-B stress depends on the algal species and the exposure time. SOD activity in the green
 25 macroalga *Ulva fasciata* was increased by UV-B radiation (Shiu et al., 2005). UV-B radiation also
 26 increased SOD activity in symbiotic dinoflagellate (Lesser and Shick, 1989) but longterm UV-B

exposure decreased SOD activity in *Chlorella vulgaris* (Malanga and Puntarulo, 1997). Arctic macroalgae (*Monostroma arcticum*, *Acrosiphonia penicilliformis*, *Coccotylus truncates*, *Phycodrys rubens*, *Palmaria palmate*, *Devaleraea ramentacea*) show less SOD induction by UV-A+UV-B, and it is even depressed in some species (Aguilera et al., 2002). The induction of CAT by UV-B radiation was also observed in land plants such as maize (Bold and Scandalios, 1997) and sunflower (Costa et al., 2002). Because of the low affinity of CAT for H₂O₂ but the high affinity of APX for H₂O₂, the higher induction of CAT activity accompanied by depression of APX activity and higher H₂O₂ production in *G. amansii* seems to indicate that *G. amansii* faces greater oxidative stress upon exposure to UV-B radiation than does *P. capillacea*.

APX and GR contribute to H₂O₂ detoxification in *P. capillacea*. GR activity was increased by UV-B radiation to regenerate GSH needed for the reduction of DHA to AsA that is the substrate of APX. In contrast, a decrease in GR activity along with a significant GSSG accumulation in *G. amansii* demonstrates that the regeneration of GSH in *G. amansii* was inhibited by UV-B radiation. Accordingly, the regeneration of AsA in *G. amansii* was relatively low than that of *P. capillacea*, and the action of APX on H₂O₂ decomposition concomitantly decreased. So, the capacity for scavenging ROS became lower in *G. amansii* than in *P. capillacea* on exposure to UV-B radiation.

A comparative study of antioxidant enzyme activity among Arctic macroalgae exposed to UV radiation showed that algal tolerance to UV-induced oxidative stress was correlated with high SOD, APX, and CAT activities (Aguilera et al., 2002). The present study delineated that APX and GR activities are positively correlated with UV-B radiation tolerance in these 2 red algae, but SOD and CAT activities are negatively correlated with UV-B tolerance. It seems that the strategy for UV-B defense is different for subtropical algae compared to Arctic macroalgae.

In conclusion, MAA is involved in the regulation of oxidative responses in red macroalgae in subtropical waters in northeastern Taiwan. As MAA accumulates, UV-B is screened and ROS production is reduced. Higher MAA concentration and the ability to accumulate MAA are primarily used by the upper subtidal alga *P. capillacea* for effective screening of UV-B radiation, and when

MAA accumulation ability decreased, carotenoid levels were increased. The lower subtidal alga *G. amansii*, whose MAA concentration and antioxidant defense capacity were low, suffers severe oxidative stress from UV-B radiation. It seems to suggest that either *G. amansii* inhabits lower subtidal regions because of less MAA concentration or it produces less MAAs because in the lower level it does not need them. It needs more work to prove it.

Acknowledgements

Financial support was provided by the National Science Council (NSC94-2311-B-110-004) and the Council of Agriculture (94AS-9.4.1-FB-e1), Executive Yuan, Taiwan, Republic of China.

References

- Aguilera, J., Dummermuth, A., Karsten, U., Schriek, R., Wiencke, C., 2002. Enzymatic defenses against photooxidative stress induced by ultraviolet radiation in Arctic marine macroalgae. *Polar Biol.* 25, 432-441.
- Alscher, R.G., Erturk, N., Heath, L.S., 2002. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J. Exp. Bot.* 53, 1331-1341.
- Asada, K., 1999. The water-water cycle in chloroplasts: scavenging of active oxygen and dissipation of excess photons. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 50, 601-639.
- Baker, K.S., Smith, R.C., Green, A.E.S., 1980. Middle ultraviolet radiation reaching the ocean surface. *Photochem. Photobiol.* 32, 367-374.
- Boldt, R., Scandalios, J.G., 1997. Influence of UV-light on the expression of the *Cat2* and *Cat3* catalase genes in maize. *Free Rad. Biol. Med.* 23, 505-514.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principal of protein-dye binding. *Analyt. Biochem.* 72, 248-254.

- 1 Brandle, J.R., Campbell, W.F., Sisson, W.B., Martin, M.M., 1977. Net photosynthesis, electron
2 transport capacity, and ultrastructure of *Pisum sativum* L. exposed to ultraviolet-B radiation.
3 Plant Physiol. 60, 165-169.
- 4 Britt, A.B., 1995. Repair of DNA damage induced by ultraviolet radiation, Plant Physiol. 108,
5 891-896.
- 6 Butow, B., Wynne, D., Tel-Or, E., 1994. Response of catalase activity to environmental stress in the
7 freshwater dinoflagellate *Peridinium gatunense*. J. Phycol. 30, 17-22.
- 8 Calkins, J., Thordardottir, T., 1982. Penetration of solar UV-B into waters of Iceland. In: Calkins, J.
9 (ED.), The role of solar ultraviolet radiation in marine ecosystems, Plenum Press, New York, pp.
10 309-321.
- 11 Carreto, J.I., Carignan, M.O., Montoya, N.G., 2005. A high-resolution reverse-phase liquid
12 chromatography method for the analysis of mycosporine-like amino acids (MAAs) in marine
13 organisms. Mar. Biol. 146, 237-252.
- 14 Collén, J., Davison, I.R., 1999a. Reactive oxygen production and damage in intertidal *Fucus* spp.
15 (Phaeophyceae). J. Phycol. 32, 54-61.
- 16 Collén, J., Davison, I.R., 1999b. Reactive oxygen metabolism in intertidal *Fucus* spp.
17 (Phaeophyceae). J. Phycol. 35, 62-69.
- 18 Costa, H., Gallego, S.M., Tomaro, M.L., 2002. Effect of UV-B radiation on antioxidant defense
19 system in sunflower cotyledons. Plant Sci. 162, 939-945.
- 20 Dunlap, W.C., Shick, J.M., 1998 Ultraviolet radiation-absorbing mycosporine-like amino acids in
21 coral reef organism: a biochemical and environmental perspective. J. Phycol. 34, 418-430.
- 22 Dunlap, W.C., Yamamoto, Y., 1995. Small-molecule antioxidants in marine organisms: antioxidant
23 activity of mycosporine-glycine. Comp. Biochem. Physiol. 112B, 105-114.
- 24 Foyer, C.H., Harbinson, J., 1994. Oxygen metabolism and the regulation of photosynthetic electron
25 transport. In: Foyer, C.H., Mullineaux, P.M. (EDs.), Causes of photooxidative stress and
26 amelioration of defense systems in plants, CRC Press, Boca Raton, FL, pp. 1-42.

- 1 Foyer, C.H., Lelandais, M., Kunert, K.J., 1994. Photooxidative stress in plants. *Physiol. Plant.* 92,
2 696-717.
- 3 Gotz, T., Sandmann, G., Romer, S., 2002. Expression of a bacterial carotene hydroxylase gene (crtZ)
4 enhances UV tolerance in tobacco. *Plant Mol. Biol.* 50, 129-142.
- 5 Griffiths, O.W., 1980. Determination of glutathione and glutathione disulphide using glutathione
6 reductase and 2-vinylpyridine. *Analyt. Biochem.* 106, 207-212.
- 7 Häder, D.P., 1993. Risk for enhanced solar ultraviolet radiation for aquatic ecosystems. *Prog.*
8 *Phycol. Res.* 9, 1-45.
- 9 He, Y.Y., Klisch, M., Häder, D.P., 2002. Adaptation of cyanobacteria to UV-B stress correlated
10 with oxidative stress and oxidative damage. *Photochem. Photobiol.* 76, 188-196.
- 11 Health, R.L., Tallman, G., 1968. Photoperoxidation in isolated chloroplasts. I. Kinetics and
12 stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.* 125, 189-198.
- 13 Hodges, D.M., Andrews, C.J., Johnson, D.A., Hamilton, R.I., 1996. Antioxidant compound
14 responses to chilling stress in differentially sensitive inbred maize lines. *Physiol. Plant.* 98,
15 685-692.
- 16 Hoyer, K., Karsten, U., Sawall, T., Wiencke, C., 2001. Photoprotective substances in Antarctic
17 macroalgae and their variation with respect to depth distribution, different tissues and
18 developmental stages. *Mar. Ecol. Prog. Ser.* 211: 117-129
- 19 Karsten, U., Karsten, U., Lembecke, S., Schumann, R., 2007. The effects of ultraviolet radiation on
20 photosynthetic performance, growth and sunscreen compounds in aeroterrestrial biofilm algae
21 isolated from building facades. *Planta* 225, 991-1000.
- 22 Karsten, U., Sawall, T., Wiencke, C., 1998. A survey of the distribution of UV-absorbing substances
23 in tropical macroalgae. *Phycol. Res.* 46, 271-279.
- 24 Kerr, J.B., McElroy, C.T., 1993. Evidence for large upward trends of ultraviolet-B radiation linked
25 to ozone depletion. *Science* 262, 1032-1034.
- 26 Kim, S.-T. Sancar, A., 1995. Photorepair of nonadjacent pyrimidine dimmers by DNA photolyase.

- 1 Photochem. Photobiol. 61, 171-174.
- 2 Langebartels, C., Schraudner, M., Heller, W., Ernst, D., Sandermann, H., 2000. Oxidative stress and
- 3 defense reactions in plants exposed to air pollutants and UV-B radiation. In: Inze, D., Van
- 4 Montagu, M. (EDs.), Oxidative stress in plants, Harwood Academic Publishers, Amsterdam, pp.
- 5 105-135.
- 6 Lesser, M.P., 1996. Elevated temperatures and ultraviolet radiation cause oxidative stress and
- 7 inhibit photosynthesis in symbiotic dinoflagellates. Limnol. Oceanogr. 43, 271-283.
- 8 Lesser, M.P., Shick, J.M., 1989. Effects of irradiance and ultraviolet radiation on photoadaptation in
- 9 the zooxanthellae of *Aiptasia pallida*: primary production, photoinhibition and enzymic
- 10 defenses against oxygen toxicity. Mar. Biol. 102, 243-255.
- 11 Malanga, G., Puntarulo, S., 1997. Oxidative damage to chloroplasts from *Chlorella vulgaris*
- 12 exposed to ultraviolet-B radiation. Physiol. Plant. 101, 455-462.
- 13 Middleton, E.M., Teramura, A.H., 1993. The role of flavonol glycosides and carotenoids in
- 14 protecting soybean from ultraviolet-B damage. Plant Physiol. 103, 741-752.
- 15 Mittler, R., Tel-Or, E., 1991. Oxidative stress responses in the unicellular cyanobacterium
- 16 *Synechococcus* PCC7942. Free Rad. Res. Commun. 12, 845-850.
- 17 Nakamura, H., Kobayashi, J., Hirata, Y., 1982. Separation of mycosporine-like amino acids in
- 18 marine organisms using reverse-phase high performance liquid chromatography, J. Chromatogr.
- 19 250, 113-118.
- 20 Noctor, G., Foyer, C.H., 1998. Ascorbate and glutathione: keeping active oxygen under control. Ann.
- 21 Rev. Plant Physiol. Plant Mol. Biol. 49, 249-279.
- 22 Okuda, T., Matsuda, Y., Yamanaka, A., Sagisaka, S., 1991. Abrupt increase in the level of hydrogen
- 23 peroxide in leaves of winter wheat is caused by cold treatment. Plant Physiol. 97, 1265-1267.
- 24 Provasoli, L., 1968. Media and prospects for the cultivation of marine algae. In: Watanabe, A.,
- 25 Hattori, A. (EDs.) Cultures and collections of algae. Proceeding of the U.S., Japan Conference,
- 26 Hakone, Japanese Society of Plant Physiology 1968, pp. 63-75.

- 1 Riegger, L., Robinson, D., 1997. Photoinduction of UV-absorbing compounds in Antarctic diatoms
2 and *Phaeocystis antarctica*. Mar. Ecol. Prog. Ser. 160, 13-25.
- 3 Rijstenbil, J.W., 2002, Assessment of oxidative stress in the planktonic diatom *Thalassiosira*
4 *pseudonana* in response to UVA and UVB radiation. J. Plankton Res. 24, 1277-1288.
- 5 Russell, J.M., Luo, M.Z., Cicerone, R.J., Deaver, L.E., 1996. Satellite confirmation of the
6 dominance of chlorofluorocarbons in the global stratospheric chlorine budget. Nature 379,
7 526-529.
- 8 Seckmeyer, G., McKenzie, R.L., 1992. Elevated ultraviolet radiation in New Zealand (45°S)
9 contrasted with Germany (45°N). Nature 359, 135-137.
- 10 Shiu, C.T., Lee, T.M., 2005. Ultraviolet-B-induced oxidative stress and responses of the
11 ascorbate-glutathione cycle in a marine macroalga *Ulva fasciata*. J. Exp. Bot. 56, 2851-2865.
- 12 Singh, S.P., Kumari, S., Rastogi, R.P., Singh, K.L., Sinha, R.P., 2008. Mycosporine-like amino acids
13 (MAAS): Chemical structure, biosynthesis and significance as UV-absorbing/screening
14 compounds. Indian J. Exp. Biol. 46, 7-17.
- 15 Smirnoff, N., Wheeler, G.L., 2000. Ascorbic acid in plants: biosynthesis and function. Crit. Rev.
16 Plant Sci. 19, 267-290.
- 17 van de Poll, W.H., Bischof, K., Buma, A.G.J., 2003. Habitat related variation in UV tolerance of
18 tropical marine red macrophytes is not temperature dependent. Physiol. Plant. 118, 74-83.
- 19 Whitehead, R.F., De Mora, S.J., Demers, S., 2000. Enhanced UV radiation-a new problem for the
20 marine environment. In: De Mora, S.J., Demers, S., Vernet, M. (Eds.) The effects of UV
21 radiation in the marine environment, Cambridge University Press, Cambridge, pp. 1-34.
- 22 Willekens, H., Chamnongpol, S., Davey, M., Schraudner, M., Langebartels, C., 1997. Catalase is a
23 sink for H₂O₂ and is indispensable for stress defense in C3 plants. EMBO J. 16, 4806-4816.
- 24 Young, A.J., 1991. The photoprotective role of carotenoids in higher plants. Physiol. Plant. 83,
25 702-708.
- 26

Figure Legends

Fig. 1. Temporal (sampling times were 2004-J: June 16-22, 2004, 2004-Sept: September 21-30, 2004, 2005-Mar: March 17-23, 2005, and 2005-Jun: June 3-11, 2005) and spatial (sampling sites were L: Lung-tung, H: Her-mei, M: Mo-auo, and S: Shea-cherng) changes in biomass (A, C) and total MAA contents (B, D) (means, n=4) in *Ptercladiella capillacea* and *Gelidium amansii*.

Fig. 2. Changes in TBARS (A) and H₂O₂ (B) contents (means±SD, n=3) in *Ptercladiella capillacea* (●) and *Gelidium amansii* (▽) in response to UV-B radiation. Different letters indicate significant difference among treatments (large font for *Ptercladiella capillacea* and small font for *Gelidium amansii*).

Fig. 3. The total MAA concentrations of (A, E), shinorine (B, F), porphyra 334 (C, G), and palythanol (D, H) in *Ptercladiella capillacea* (●) and *Gelidium amansii* (▽) in response to UV-B radiation. Data are presented as mean±SD (n=3) and different letters indicate significant difference among treatments (large font for *Ptercladiella capillacea* and small font for *Gelidium amansii*).

Fig. 4. The carotenoid concentration in *Ptercladiella capillacea* (●) and *Gelidium amansii* (▽) in response to UV-B radiation. Data are presented as mean±SD (n=3) and different letters indicate significant difference among treatments.

Fig. 5. The AsA (A), DHA (B), GSH (C), and GSSG (D) concentrations in *Ptercladiella capillacea* (●) and *Gelidium amansii* (▽) in response to UV-B radiation. Data are presented as mean±SD (n=3)

and different letters indicate significant difference among treatments (large font for *Ptercladiella capillacea* and small font for *Gelidium amansii*).

Fig. 6. Effects of UV-B radiation on SOD (A), CAT (B), APX (C), and GR (D) activities (mean $\pm$ SD, n=3) in *Ptercladiella capillacea* (●) and *Gelidium amansii* (▽). Different letters indicate difference among treatments (large font for *Ptercladiella capillacea* and small font for *Gelidium amansii*).

Supplementary data:

Fig. 1. HPLC profile of MAAs of *Ptercladiella capillacea* and *Gelidium amansii* in response to UV-B radiation.