Accepted Manuscript

Effects of Benzo[*a*]pyrene on DNA Damage and Histological alterations in Gonad of Scallop *Chlamys farreri*

Miao Jing-jing, Pan Lu-qing, Liu Jing, Zhang Lin

PII: S0141-1136(08)00234-1

DOI: [10.1016/j.marenvres.2008.10.006](https://doi.org/10.1016/j.marenvres.2008.10.006)

Reference: MERE 3298

To appear in: *Marine Environmental Research*

Received Date: 29 April 2008

Revised Date: 20 October 2008

Accepted Date: 23 October 2008

Please cite this article as: Jing-jing, M., Lu-qing, P., Jing, L., Lin, Z., Effects of Benzo[*a*]pyrene on DNA Damage and Histological alterations in Gonad of Scallop *Chlamys farreri*, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.10.006](https://doi.org/10.1016/j.marenvres.2008.10.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Effects of Benzo[a]pyrene on DNA Damage and Histological alterations in Gonad of Scallop *Chlamys farreri*

Miao Jing-jing, Pan Lu-qing*, Liu Jing, Zhang Lin

The Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, No. 5, Yushan Road, Qingdao, China

Abstract

To investigate the biological impact of polycyclic aromatic hydrocarbons (PAHs) on reproductive system, scallops *Chlamys farreri* were continuously exposed to benzo[a]pyrene (BaP) (0.5, 3, 10 $\mu\text{g L}^{-1}$) during 15 days. DNA damage and histological alterations were examined in female gonad. DNA strand break levels significantly increased after 12 h exposure, and remained high and significantly different from control values until the end of the exposure. In the ovaries of the scallops exposed to 10 $\mu\text{g L}^{-1}$ BaP for 10 days, histological analysis showed that the cytoplasm of the oocytes in the outer layer of the ovaries became sparse, and the nuclear membranes were obscure. Damaging effects on ovaries and oocytes were more severe after 15 days exposure. Degenerating oocytes increased and the connective tissue of the ovary envelopes became loose. Electron microscopic examination revealed ultrastructural aspects of degenerating oocyte and degenerated oocyte after 15 days exposure.

Keywords: Benzo[a]pyrene; Scallop; DNA strand break; Genotoxicity; Endocrine;

Histopathology; Ultrastructure; Gonad; Oocyte

*Corresponding author. Tel.: +86 532 8203 2963; fax: +86 532 8289 4024.

E-mail address: panlq@ouc.edu.cn (L. Pan).

1. Introduction

In recent years, special attention has been given to the disruption of reproduction by organic compounds and the possible consequences on wildlife populations, because reproductive systems appear to be particularly vulnerable to environmental contaminants (Harries et al., 1997; Kovacs et al., 1997). Several studies have demonstrated that endocrine disturbances due to xenobiotics in wildlife include sex changes in marine gastropods (Horiguchi et al., 2002), delay in oocyte maturation and sperm release and perturbation of the neuroendocrine control of spawning in mussels (Binelli et al., 2004; Gagné et al., 2004). Field observations of reproductively abnormal organisms and populations decline in polluted sites stimulated major research efforts to understand stressor-induced changes of some of the reproductive parameters within an organism that could be valuable as early warning indicators of the potential long-term hazard of environmental deterioration at the population level (Thomas, 1990).

Bivalves are particularly at risk to aquatic xenobiotic chemicals because they are stationary and filter relatively high volume of water, including suspended particles. Many different effects have been recorded in bivalves exposed to chemical contaminants. Among them, histological alterations have been widely employed as a method of assessing the disruption of reproduction systems (Ketata et al., 2008). Binelli et al. (2004) reported degenerating oocytes in relation to spawning delay in the zebra mussel (*Dreissena polymorpha*) exposed to DDT. Numerous degenerating ovarian follicles and atretic oocytes have also been observed in female mussels (*Mytilus edulis*) after exposure to bisphenol A (50 µg L⁻¹), BDE-47 (1 µg L⁻¹), North Sea oil (0.5 mg L⁻¹) and a mixture of North Sea oil (0.5

43 mg L⁻¹), alkyphenols (0.1 mg L⁻¹) and polycyclic aromatic hydrocarbons (PAHs) (0.1 mg L⁻¹)
 44 for three weeks (Aarab et al., 2004; Aarab et al., 2006). These results indicated dramatic toxic
 45 effects on reproductive tissues in relation to environmental toxicants. PAHs, owing to their
 46 highly mutagenic and carcinogenic features and widespread distributions in the global
 47 environment, are considered as one of the major pollutants affecting marine organisms
 48 (Menzie et al., 1992). In addition to genetic toxicity, PAHs are also listed by the US
 49 Environmental Protection Agency (EPA) as compounds capable of disrupting the endocrine
 50 system of animals. However, few laboratory studies of PAHs toxicity on the reproductive
 51 system of bivalves have been carried out.

52 In the present study we used benzo[*a*]pyrene (BaP), a powerful carcinogen that is
 53 included in the 16 PAHs designated as primary contaminants, as a representative of PAHs. We
 54 investigated the female gonad damage induced by BaP exposure, including DNA damage and
 55 histological perturbations in Zhikong scallop *Chlamys farreri*. The objective of our work was
 56 to determine the basic characteristics of the endpoints of genotoxic effects and endocrine
 57 disruption of BaP, and to compare the temporal and dose response relationship between BaP
 58 exposure and the toxic effects. The study may provide preliminary information on evaluation
 59 of gonadal status under the stress of PAHs.

60

61 **2. Materials and methods**

62 *2.1 Scallops*

63 Scallops (*C. farreri*) used for this experiment were rope-cultivated scallops, aged 2 years
 64 and with shell length of 5.8±0.8 cm, collected from Taiping Bay (Huanghai Sea, Qingdao,

China) on April 10th, 2006. The sunken ventral portion of the anterior ear of the right shell of the scallop *C. farreri* provides an aperture when the two shells are closed. Female scallops were identified after gently stimulating the aperture of the scallop to make it gape widely using forceps, and observing the colour of the gonad. Mature female scallops have a red gonad, whereas a male gonad is white. Female scallops held in aquaria (1 L water per scallop) were acclimated to laboratory conditions for 7 days in filtered seawater pumped from Taiping Bay. The sea water was continuously aerated, and salinity, temperature and pH were respectively maintained at 31‰, 17(±1)°C and 8.1. The water was renewed every 24 hours. The scallops were fed daily with dried powder of *Spirulina platensis* (30 mg for each individual).

2.2 Experimental design

The background concentration of BaP in seawater from Taiping Bay was analysed by HPLC before the experiment.. The effects of three waterborne BaP concentrations were studied (0.5, 3 and 10 µg L⁻¹). BaP was first dissolved in acetone, and then added to seawater to achieve final acetone concentration of 0.0025%. Control groups included a seawater control and an acetone control. There were three aquaria used as replicates per group. Experimental conditions (salinity, pH, temperature, scallop density and feeding) were the same as those used for acclimation, and the exposure media were renewed daily. The exposure experiment lasted for 15 days. During this period, we determined the BaP concentration of the exposed groups every day before renewing the water during the experiments. The analyzed BaP concentrations of each group were 0.52±0.07 µg L⁻¹, 2.93±0.13 µg L⁻¹, 9.15±0.11 µg L⁻¹. Female gonads of scallops were excised and frozen immediately at -80°C at days 0, 0.5, 1, 3, 6, 10 and 15 for the subsequent examination of DNA

87 integrity, two to three scallops were sampled per replicate. Female gonads from sea water control,
 88 acetone control and $10 \mu\text{g L}^{-1}$ BaP treatment groups were sampled at day 10 and day 15 for
 89 histological evaluation, and sampled for ultrastructural evaluation at day 15. Gonad tissues from
 90 six scallops were sampled as 6 replicates. No mortalities were observed during the exposure
 91 experiment.

92 *2.3 Alkaline unwinding assay*

93 DNA was extracted from female gonads by mashing the tissues from two to three scallops
 94 with plastic sticks and homogenizing the tissue in 3 mL TE buffer (50 mM Tris, 100 mM
 95 EDTA, pH 8.0). 0.45 mL lysate was transferred into 1.5 mL Eppendorf tubes. Then 0.05 mL
 96 10% SDS (Sodium dodecyl sulfate) and $3 \mu\text{L } 20 \text{ g L}^{-1}$ proteinase K were added to it and the
 97 mixture was incubated for 5 h at 55°C . An equal volume of buffered
 98 phenol/chloroform/isoamyl alcohol (PCI) (25:24:1, v/v/v, pH 8.0) was then added to the
 99 sample. The sample was gently mixed for 15 min and then centrifuged for 10 min at 13 000
 100 rpm at 4°C . The aqueous layer was transferred to a new tube and digested with $0.8 \mu\text{L}$ Rnase
 101 (10 mg mL^{-1}) for 30 min at 37°C and the digesta was extracted twice with PCI. DNA was
 102 precipitated from the resulting aqueous layer by adding 2 volumes of absolute ethanol and
 103 1/10 volume of 3 M sodium acetate, pH 5.2. The samples were allowed to settle for 2 h at
 104 -20°C , and then centrifuged for 15 min at 13 000 rpm. The resulting pellet was rinsed with 1
 105 mL 70% ethanol and air-dried, then dissolved in 1 mL of TE buffer (10 mM Tris, 1 mM
 106 EDTA, pH 8.0). The amount of DNA was quantified using a UV/Visible spectrophotometer
 107 (Ultro spec 2100 pro, Amersham Biosciences, Sweden).

108 The alkaline unwinding assay used in the study was adapted from Ching et al (2001). In

the assay, the rate of transition of double-stranded DNA (dsDNA) to single-stranded DNA (ssDNA) under pre-defined alkaline denaturing conditions was proportional to the number of breaks in the phosphodiester backbone, and thus was used as a measure of DNA integrity (Daniel et al., 1985). The amounts of various types of DNA were quantified by measuring the varying degrees of fluorescence resulting from a DNA-binding dye, bisbenzamide. After reaction with the dye, the fluorescence of dsDNA is double that of the ssDNA (Cesarone et al., 1979).

The DNA sample was diluted and divided into three equal portions for fluorescence determination of dsDNA, ssDNA and alkaline unwound DNA (auDNA). The fluorescence of the initial DNA or dsDNA was determined by placing 100 μL of the DNA sample, 100 μL of 25 mM NaCl and 2 μL of 0.5% SDS in a pre-chilled test tube, followed by the addition of 3 mL 0.2 M potassium phosphate (pH 6.9), and 3 μL bisbenzamide (1 mg mL⁻¹). The contents were mixed and allowed to react in darkness for 15 min. The fluorescence of the sample was measured using a spectrofluorimeter (Molecular Spectroscopy LS 55 from Instruments, P.E., MA, USA) with an excitation wavelength of 360 nm and an emission wavelength of 450 nm. The fluorescence of ssDNA was determined as above but using a DNA sample that had already been boiled at 80°C for 30 min to completely unwind the DNA. auDNA samples were made by subjecting initial DNA samples to alkaline treatment, i.e. 50 μL of 50 mM NaOH was rapidly mixed with 100 μL of DNA sample in a pre-chilled test tube and incubated on ice in darkness for 30 min, followed by a rapid addition and mixing of 50 μL of 50 mM HCl and 2 μL of 0.5% SDS then the mixture was forcefully passed through a 21G needle several times. The fluorescence of the alkaline unwound DNA sample was measured as described above.

131 The ratio between double-stranded DNA to total DNA (F value) was determined as
 132 follows: $F \text{ value} = (\text{auDNA} - \text{ssDNA})/(\text{dsDNA} - \text{ssDNA})$

133 *2.4 Histological evaluation*

134 The female gonads were excised and fixed with Bouin's fixative for 18 hours. Following
 135 dehydration through an ascending ethanol series, the tissues were embedded in paraffin,
 136 sectioned at 5 μm , stained with Ehrlich's haematoxylin and eosin (H&E), and observed with
 137 an Olympus BX-41 microscope (Olympus, Japan).

138 For transmission electron microscopy (TEM), tissues were prepared as follows: selected
 139 female gonads of scallops were cut at 1 mm^3 , fixed in 2.5% glutaraldehyde in phosphate
 140 buffer (pH 7.2) at 4°C, postfixed in 1% osmium tetroxide, dehydrated in an ethanol series,
 141 and embedded in Epon 812. Ultrathin sections were cut on an LKB ultramicrotome, stained
 142 with uranyl acetate and lead citrate. Sections were examined in a Hitachi H-7000 transmission
 143 electron microscope (Hitachi, Japan).

144 For quantifying the oocyte degeneration, the degenerating oocytes were counted in the
 145 histological sections of each ovarian piece. The number of the degenerating oocytes was
 146 determined in a 90,000 μm^2 field at a magnification of x400.

147 *2.5 Statistical analysis*

148 All values were expressed as mean $\pm$ SD of three replicates in one representative
 149 experiment. Experiments were performed three times to confirm the results. The analyses
 150 were carried out using SPSS software (Version 11.5). Data were compared with one-way
 151 ANOVA, and Newman-Keuls posthoc test when appropriate. The α level used was 0.05. All
 152 data were checked for normality and homogeneity of the variances, and transformed when

153 necessary.

154 **3. Results**

155 *3.1 DNA strand breaks*

156 The results of mean F values of each individual treatment group are presented in Figure
 157 1. No significant difference in F values was observed over the 15-day exposure period for the
 158 control and acetone control groups ($P > 0.05$). F values of control and acetone control groups,
 159 respectively, were 0.640 ± 0.084 and 0.633 ± 0.050 . F values of samples from 3 and $10 \mu\text{g L}^{-1}$
 160 treatment groups showed marked decreases (0.516 ± 0.014 and 0.462 ± 0.063 , respectively) (P
 161 < 0.05) after 12 h of exposure. While the $0.5 \mu\text{g L}^{-1}$ BaP treatment groups showed marked
 162 decreases in mean F values (0.495 ± 0.074 , $P < 0.05$) after 1 day exposure. The F values
 163 remained low and significantly different from the control values ($P < 0.05$) until day 15 for the
 164 $0.5 \mu\text{g L}^{-1}$ (0.369 ± 0.024), $3 \mu\text{g L}^{-1}$ (0.259 ± 0.035) and $10 \mu\text{g L}^{-1}$ (0.270 ± 0.020) BaP treatment
 165 groups. There were no significant differences between the F values of $3 \mu\text{g L}^{-1}$ and $10 \mu\text{g L}^{-1}$
 166 BaP exposed groups from day 6 to day 15 ($P > 0.05$).

167 *3.2 Histological evaluation*

168 Histological and ultrastructural studies of female gonads from both seawater control and
 169 acetone control scallops showed the same histological features. The female gonad
 170 development of *C. farreri* can be divided into five stages: proliferation, growth, maturation,
 171 spawning and recovery. Histological sections of female gonad displayed numerous ovarian
 172 follicles filled with mature oocytes, the interspace among follicles disappeared, indicating that
 173 the female gonads of scallops were at the late pre-spawning stage (Figure 2a, 2b). Electron
 174 microscopic examination showed that most of the oocytes were at the vitellogenic stage.

175 Oolemma of oocytes formed microvilli, and the oocytes' ooplasm was rich in mitochondria,
 176 cisternae of rough surfaced endoplasmic reticulum and ribosomes, vast amounts of yolk
 177 bodies and cortical granules. Nucleoli were visible in nucleus, nuclear membrane
 178 protuberated at some site, and the euchromatins uniformly distributed in nucleus (Figure 3a,
 179 3b).

180 Histological analysis of the ovaries indicated that after 10 and 15 days exposure of $10\ \mu\text{g}$
 181 L^{-1} BaP degenerating oocytes increased especially in the outer layer of the ovaries. The
 182 degenerating oocytes appeared irregular or polyhedric, and were deformed by compression of
 183 the oocyte (Figure 2c, 2d, 3c, 3f). Cytoplasm of the degenerating oocytes were sparse, and the
 184 nuclear membranes were obscure compared with the control. More severe damaging effects
 185 on ovaries and oocytes were observed after 15 days exposure. Degenerating oocytes increased,
 186 meanwhile the connective tissue of the ovary envelopes became loose in most of the ovaries
 187 observed (Figure 2c, 2d). Oocyte count indicated that degenerating oocytes increased
 188 significantly ($P < 0.05$) after 10 days and 15 days exposure compared with control (Table 1).
 189 Ultrastructural sections of female gonad from $10\ \mu\text{g L}^{-1}$ BaP treatment group after 15 days
 190 exposure displayed degenerating oocyte and degenerated oocyte. The degenerating and
 191 degenerated oocytes appeared irregular or polyhedric, and were deformed by compression of
 192 the oocyte (Figure 2c, 2d, 3c, 3f). During the gradual disintegration of the oocytes, numerous
 193 vacuoles appeared in the ooplasm of oocytes. A few lysosomes (phagosomes) and vacuoles
 194 also appeared in the cytoplasm of the follicle cells which were attached to the degenerating
 195 oocyte. (Figure 3c, 3d, 3e). In degenerated oocytes, endoplasmic reticulum, mitochondria, and
 196 the quantity of yolk bodies decreased, resulting in the electron density reduction of the

197 ooplasm. The microvilli were shed and numerous glycogen particles occurred in cytoplasm.
 198 Nuclear membranes were anomalous, and the distribution of euchromatins in nucleus was not
 199 uniform (Figure 3f).

200 **4. Discussion**

201 BaP is a genotoxic agent capable of binding to DNA via CYP transformation to arene
 202 oxides, forming DNA adducts that are more effective sites for the production of strand breaks.
 203 It may also increase the presence of reactive oxygen species (ROS) via redox cycling, which
 204 can be responsible for lipid peroxidation processes and result in DNA strand break eventually
 205 (Martinez and Livingstone, 1995; Canova et al., 1998). The micronucleus frequency observed
 206 for adult oysters *Crassostrea gigas* contaminated for 48 h with BaP (0.05, 0.5, 1 and 500 μg
 207 L^{-1}) showed the clastogenic action of BaP and a dose dependent effect. Siu et al. (2003, 2004)
 208 and Ching et al. (2001) studied BaP (0.3, 3, 30 $\mu\text{g L}^{-1}$) effects on DNA strand breaks of
 209 haemocytes and hepatopancreas in mussels *Perna viridis*, and the results suggested that the
 210 response of DNA alteration upon BaP exposure may be tissue-specific and threshold
 211 dependent in the species. In a previous study, Pan et al. (2007) reported that the *F* values in
 212 gill and digestive gland of 0.5 and 3 $\mu\text{g L}^{-1}$ BaP treatment groups decreased until day 6 and
 213 then showed a recovery. This was considered to be due to an early elicitation of the DNA
 214 repair system. While, in this study, DNA strand breaks of gonads from 0.5, 3 and 10 $\mu\text{g L}^{-1}$
 215 treatment groups showed marked decrease after 12 h or 1 day exposure, and no recoveries
 216 appeared until the end, indicating a decreased capacity for DNA repair activity. The result
 217 corresponds with the findings of Gagné et al. (2004) which suggests that ripe mussels whose
 218 oocytes were meiose-arrested might have reduced DNA synthesis/repair activity during the

219 mature stage. However, further studies about the relationship between DNA repair activity of
220 oocytes and the gonad mature status are needed. In addition, a dose-dependent relationship
221 was observed between the DNA strand break levels and the BaP exposure dose (from 0.5 to 3
222 $\mu\text{g L}^{-1}$), while the DNA strand break levels had no marked difference between 3 and 10 $\mu\text{g L}^{-1}$
223 BaP from day 6 to day 15, indicating that the toxic effects on DNA in gonads may be
224 threshold dependent.

225 Microscopic observations in the present work showed time course of the BaP toxic
226 effects towards ovaries of scallops. During the 15 days exposure, degenerating oocytes
227 significantly increased in ovaries (Table 1). Ultrastructure of the ovaries was studied to find
228 out more detailed effects. Results indicated that oocytes were at the vitellogenesis stage and
229 yolk granules were accumulating. After 15 days exposure, degenerating oocytes increased in
230 the treatment group and ultrastructural sections displayed degenerated oocytes with reduced
231 density of the ooplasm and yolk granules, which could be the reason for the obscure nuclear
232 membrane observed under the microscopic. As a result of reductions in yolk bodies and cell
233 organelles in degenerated oocytes, and numerous glycogen particles appeared in cytoplasm,
234 there was a reduction in the electron density of the ooplasm, and the chromatin distributed
235 around the nuclear membrane. Consequently, the electron density of cytoplasm and nucleus
236 became similar and made the nuclear membrane appear obscure (Figure 2c, 2d, 3f). Oocyte
237 degeneration is a continuous process throughout the development of oocyte, and it seems to
238 be most important for vitellogenic oocytes (Gaulejac et al., 1995; Chung et al., 2007).
239 Degenerated oocytes may be observed in the ovaries amongst the normal oocytes at any stage
240 of development. It may be caused by overproduction of oocytes in spawners as a result of

241 environmental stress (Leino and Maccormik, 1997). Oocyte degeneration due to xenobiotic
242 and heavy metal contamination was reported in several bivalve species (Maung and Tyler,
243 1982; Rasmussen et al., 1983). Lowe and Pipe reported a high level of gamete atresia in *M.*
244 *edulis* treated with hydrocarbons, and presumed that the degeneration was due to the direct
245 action of these chemicals on gametes by the destabilization of lysosomes and yolk granules
246 which contain lysosomal enzymes (1986). The pathological effects in gill and digestive gland
247 of *C. farreri* exposed to 10 µg L⁻¹ BaP were also reported (Miao et al., 2007). Considering the
248 results of DNA strand break in this work, it was suggested that the oocyte degeneration can be
249 considered to be subsequences of the effect of BaP reactive intermediates on the
250 biomacromolecules, which caused DNA, protein and the cell membrane system damages.

251 On the other hand, BaP is also a compound capable of disrupting the endocrine system of
252 animals. Fong et al. (1994) observed that methiothepin, a serotonin (5-hydroxytryptamine,
253 5HT) receptor antagonist, caused oocyte degeneration, with morphological aspects similar to
254 those observed in this work. Degenerating oocytes in relation to spawning delay have also
255 been reported in association with endocrine disruption induced by DDT in zebra mussel
256 *Dreissena polymorpha* (Binelli et al., 2004). In our results, ultrastructure of the ovaries was
257 studied to find out more detailed effects. The ultrastructure characteristics of degenerating and
258 degenerated oocytes displayed the same aspects with the natural interdigitation phenomena
259 reported in the marine bivalve *Pina nobilis* (Gaulejac et al., 1995) and *Cyclina sinensis*
260 (Chung et al., 2007). Meanwhile, it was reported in *C. sinensis* that follicle cells appear to
261 play an integral role in oocyte degeneration (Chung et al., 2007), which were also observed as
262 a common phenomenon in our results (Figure 3d). Furthermore the ultrastructure observation

indicated that quantity of yolk granules in exposed samples decreased significantly during the oocyte degenerating which suggested that oocyte degeneration caused by PAHs exposure could be a reason for vitellogenin-like protein levels variation reported in mussels *M. edulis* and *D. polymorpha* (Aarab et al., 2004; Lafontaine et al., 2000). A long-term and low-dose effect of PAHs needs to be further studied to examine the endocrine effects and the mechanisms of PAH impact on bivalve reproductive systems.

5. Conclusion

The results of the present study confirmed that BaP exposure resulted in DNA damage and perturbations in histological structure of female gonad in scallops. According to the results, the effect of BaP reactive intermediates on biomacromolecules, which caused DNA and cell membrane system damage in scallop female gonad, were corroborated. The experiment indicated that DNA damage in gonad was more sensitive than other tissues, and gonad DNA strand break could be a potentially suitable biomarker as early warning indicator for toxicity of PAHs. However, the mechanism of BaP effects needs to be studied further to understand the phenomenon of increased oocyte degeneration.

Acknowledgement

We wish to express our sincere gratitude to all the staff of the Laboratory of Aquatic Environmental Physiology for the organization of the exposure experiment. This work was supported by Science and Technology Development Project of Qingdao City of China (06-2-2-21-jch) and Science Foundation of The Key Laboratory of Mariculture, Ministry of

285 Education, Ocean University of China.

286

287 **References**

288 Aarab, N., Lemaire, S., Unruh, E., Hansen, P.D., Larsen, B.K. & Andersen, O.K., et al., 2006.

289 Preliminary study of responses in mussel *Mytilus edulis* exposed to bisphenol A, diallyl
290 phthalate and tetrabromodiphenyl ether. *Aquatic Toxicology* 78, 86-92.

291 Aarab, N., Minier, C., Lemaire, S., Unruh, E., Hansen, P.D. & Larsen, B.K. et al., 2004.

292 Biochemical and histological responses in mussel (*Mytilus edulis*) exposed to North Sea
293 oil and to a mixture of North Sea oil and alkylphenols. *Marine Environmental Research*
294 58, 437-441.

295 Binelli, A., Bachetta, R., Mantecca, P., Ricciardi, F., Provini, A., Vailati, G., 2004. DDT in
296 zebra mussels from Lake Maggiore (N. Italy): level of contamination and endocrine
297 disruptions. *Aquatic Toxicology* 69, 175-188.

298 Canova, S., Degan, P., Peters, L. D., Livingstone, D. R., Voltan, R., Venier, P., 1998. Tissue
299 dose, DNA adducts, oxidative DNA damage and CYP1A-immunopositive proteins in
300 mussels exposed to waterborne benzo[a]pyrene. *Mutation Research* 399, 17-30.

301 Cesarone, D. F., Bolognesi, C., Santi, L., 1979. Improved microfluorometric DNA
302 determination in biological material using 33258 Hoechst. *Analytical Biochemistry* 100,
303 188-197.

304 Ching, E. W. K., Siu, W. H. L., Lam, P. K. S., Xu, L., Zhang, Y., Richardson, B., Wu, R. S. S.,
305 2001. DNA Adduct formation and DNA strand breaks in green-lipped mussels (*Perna*
306 *viridis*) exposed to benzo[a]pyrene: dose- and time- dependent relationships. *Marine*

- 307 Pollution Bulletin 42, 603- 610.
- 308 Chung, E., Koh, C. H., Park, G. M., 2007. Oogenesis, oocyte degeneration and sexual
 309 maturation in female *Cyclina sinensis* (Gmelin, 1971) (Bivalvia: Veneridae) in Korea.
 310 Integrative Biosciences 11, 191-198.
- 311 Daniel, F. B., Haas, D. L., Santi, L., 1985. Quantitation of chemically induced DNA strand
 312 breaks in human cells via an alkaline unwinding assay. Analytical Biochemistry 144,
 313 390- 402.
- 314 Fong, P.P., Hardege, J.D., Ram, J.L., 1994. Long-lasting, sex-specific inhibition of
 315 serotonin-induced spawning by methiothepin in the zebra mussel, *Dreissena polymorpha*
 316 (Pallas). Journal of Experimental Zoology 270, 314–320.
- 317 Gagné, F., Blaise, C., Hellou, J., 2004. Endocrine disruption and health effects of caged
 318 mussels, *Elliptio complanata*, placed downstream from a primary-treatedmunicipal
 319 effluent plume for one year. Comparative Biochemistry and Physiology Part C 138,
 320 33-44.
- 321 Gaulejac, B. D., Henry, M., Vicente, N., 1995. An ultrastructural study of gametogenesis of
 322 the marine bivalve *Pinna nobilis* (Linnaeus 1758) I. Oogenesis. Journal of Mollusc Study
 323 61, 375-392.
- 324 Harries, J.E., Sheahan, D.A., Jobling, S., Matthiessen, P., Neall, P., Sumpter, J.P., et al., 1997.
 325 Estrogenic activity in five United Kingdom rivers detected by measurement of
 326 vitellogenesis in caged male trout. Environmental Toxicological Chemistry 16, 534-542.
- 327 Horiguchi, T., Kojima, M., Kaya, M., Matsuo, T., Shiraishi, H. & Morita, M. et al., 2002.
 328 Tributyltin and triphenyltin induce spermatogenesis in ovary of female abalone, *Haliotis*

- 329 *gigantea*. Marine Environmental Research 54, 679-684.
- 330 Ketata, I., Denier, X., Hamza-Chaffai, A., Minier, C., 2008. Endocrine-related reproductive
- 331 effects in mollusks. Comparative Biochemistry and Physiology Part C 147, 261-270.
- 332 Kovacs, T.G., Voss, R.H., Megraw, S.R., Martel, P.H., 1997. Perspectives on Canadian field
- 333 studies examining the potential of pulp and paper mill effluent to affect fish reproduction.
- 334 Journal of Toxicology and Environmental Health 51, 305-352.
- 335 Lafontaine, Y., Gagné, F., Blaise, C., 2000. Biomarkers in zebra mussels (*Dreissena*
- 336 *polymorpha*) for the assessment and monitoring of water quality of the St Lawrence
- 337 River (Canada). Aquatic Toxicology 50, 51-71.
- 338 Leino, R. L. and McCormik, J. H., 1997. Reproductive characteristics of ruffe,
- 339 *Gymnocephalus cernuus*, in the St. Louis River estuary on western Lake Superior: A
- 340 histological examination of the ovaries over one annual cycle. Canadian Journal of
- 341 Fisheries and Aquatic Sciences 54, 256-263.
- 342 Lowe, V.L., Pipe, R.K., 1986. Hydrocarbon exposure in mussels: a quantitative study of the
- 343 responses in the reproductive and nutrient storage cell systems. Aquatic Toxicology 8,
- 344 265-272.
- 345 Martinez, P. G. and Livingstone, D. R., 1995. Benzo[a]pyrene –dione-stimulated oxyradical
- 346 production by microsomes of digestive gland of the common mussel, *Mytilus edulis* L..
- 347 Marine Environmental Research 39, 185-189.
- 348 Maung Myint, U., Tyler, P.A., 1982. Effects of temperature, nutritive and metal stressors on
- 349 the reproductive biology of *Mytilus edulis*. Marine Biology 67, 209–223.
- 350 Menzie, C. A., Potocki, B. B., Santodonato, J., 1992. Exposure to carcinogenic polycyclic

- 351 aromatic hydrocarbons in the environment. Environmental Science and Technology 26,
352 1278-1284.
- 353 Miao, J-J, Pan, L-Q., Wang, J., 2007. Effects of B[a]P on micro and ultrastructure of gill and
354 digestive gland of *Chlamys farreri*. Acta Scientiae Circumstantiae 27(9), 1497-1503.
- 355 Pan, L-Q., Miao, J-J., Wang, J., Liu, J., 2007. AHH activity, tissue dose and DNA damage in
356 different tissues of the scallop *Chlamys farreri* exposed to benzo[a]pyrene,
357 Environmental Pollution 153, 192-198.
- 358 Rasmussen, L.P.D., Hage, E., Karlog, O., 1983. Light and electron microscopic studies of the
359 acute chronic toxic effects of N-nitroso compounds on the marine mussel, *Mytilus*
360 *edulis* (L.). 1. N-nitrosodimethylamine. Aquatic Toxicology 3, 285-299.
- 361 Siu, W.H.L., Cao, J., Jack, R. W., Wu., R. S. S., Richardson, B. J., Xu, L., Lam., R. K. S.,
362 2004. Application of the comet and micronucleus assays to the detection of B[a]P
363 genotoxicity in haemocytes of the green-lipped mussels (*Perna viridis*). Aquatic
364 Toxicology 66, 381-392.
- 365 Siu, W.H.L., Hung, C.L.H., Wong, H.L., et al., 2003. Exposure and time dependent DNA
366 strand breakage in hepatopancreas of green-lipped mussels (*Perna viridis*) exposed to
367 Aroclor 1254, and mixtures of B[a]P and Aroclor 1254. Marine Pollution Bulletin 46,
368 1285e1293.
- 369 Thomas, P., 1990. Molecular and biochemical responses of fish to stressors and their potential
370 use in environmental monitoring. American Fisheries Society Symposium 8, 9-28.

371 Table 1

372 Effect of Benzo[a]pyrene on oocyte degeneration in mature ovarian tissues prepared from scallop
 373 (*Chlamys farreri*)

374

Treatment	Rate of degenerating oocyte (%)		
	0 day	10 days	15 days
Seawater control	2.33 ± 1.55	2 ± 0.57	2.33 ± 1.52
Acetone control		1.67 ± 0.33	2 ± 1.15
$10 \mu\text{g L}^{-1}$ BaP		9.67 ± 2.08	25.67 ± 4.04

375 Values are means $\pm$ S.E. (n=6)

376 **Figure Captions**

377 Fig.1. Variations in the ratio between double-stranded DNA to total DNA (F value) (mean $\pm$
378 SD) in mature female gonads of scallop (*Chlamys farreri*) groups exposed to various
379 concentrations of BaP over a 15-day period (n=3).

380

381 Fig.2. Effects of BaP in female scallop *Chlamys farreri* gonad section. 2a and 2b - control; 2c
382 - exposed to 10 $\mu\text{g L}^{-1}$ BaP for 10 days. Arrows indicate the sparse cytoplasts of oocytes; 2d -
383 exposed to 10 $\mu\text{g L}^{-1}$ BaP for 15 days. Ovarian follicles filled with atretic oocytes. Arrows
384 indicate the sparse cytoplasts of oocytes. (CT- connective tissue; DO-
385 degenerating/degenerated oocyte; FC- follicle cell; Fw- follicular wall; MO- mature oocyte;
386 N- nucleus; Nu-nucleolus) (Paraffin sections, 5 μm -thick, hematoxylin-eosin-saffron.)

387

388 Fig.3. Effects of BaP on ultrastructure of scallop *Chlamys farreri* oocytes. 3a and 3b -control;
389 3c, 3d, 3e and 3f - exposed to 10 $\mu\text{g L}^{-1}$ BaP for 15 days. 3c showed perivitelline space of
390 several degenerating oocyte. 3d showed part of a degenerating oocyte and a follicle cell. 3e
391 showed a degenerating oocyte with numerous vacuoles and modified mitochondrion. 3f
392 showed a degenerated oocyte with large quantity of glycogen in the cytoplasm. (CG- cortical
393 granules; DO- degenerating oocyte; FC- follicle cell; Fw- follicular wall; GP- glycogen
394 particle; L- lysosome; Mi-mitochondrion; Mv-microvillus; N- nucleus; Nu-nucleolus;
395 RER-rough endoplasmic reticulum; Va-vacuoles; Yg-yolk granules) (Epon 812 embedded,
396 stained with uranyl acetate and lead citrate.)

397

398

Figures

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

501
502
503
504
505
506
507
508
509
510
511
512
513
514