Accepted Manuscript

Title: Antibiotic resistance of bacterial biofilms

Authors: Niels Høiby, Thomas Bjarnsholt, Michael Givskov,
Søren Molin, Oana Ciofu

PII: S0924-8579(10)00009-9
DOI: doi:10.1016/j.ijantimicag.2009.12.011
Reference: ANTAGE 3212

To appear in: *International Journal of Antimicrobial Agents*

Received date: 13-12-2009

Accepted date: 15-12-2009

Please cite this article as: Høiby N, Bjarnsholt T, Givskov M, Molin S, Ciofu O, Antibiotic resistance of bacterial biofilms, *International Journal of Antimicrobial Agents* (2008), doi:10.1016/j.ijantimicag.2009.12.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Antibiotic resistance of bacterial biofilms

Niels Høiby^{a, b*}, Thomas Bjarnsholt^{a, b}, Michael Givskov^b, Søren Molin^c and Oana Ciofu^b

^aDepartment of Clinical Microbiology 9301, Juliane Maries Vej 22, Rigshospitalet, 2100 Copenhagen, Denmark

^bDepartment of Bacteriology, Institute of International Health, Immunology and Microbiology, University of Copenhagen, Denmark.

^cBiocentrum, Danish Technical University, Lyngby, Denmark

*Corresponding author. Tel.: +4535457788; fax: +4535456412.

E-mail address: hoiby@hoibyniels.dk (N. Høiby)

Keywords:

- Biofilm
- Antibiotic resistance
- Antibiotic tolerance
- Beta-lactamase
- Efflux pumps
- Mutators
- Pseudomonas aeruginosa
- Cystic fibrosis
- Foreign body infections

ABSTRACT

A biofilm is a structured consortium of bacteria embedded in a self-produced polymer matrix consisting of polysaccharide, protein and DNA. Bacterial biofilms cause chronic infections because they show increased tolerance to antibiotics, disinfective chemicals and resist phagocytosis and other components of the defense system of the body. The persistence of e.g. staphylococcal infections related to foreign bodies is due to biofilm formation. Likewise, chronic *Pseudomonas aeruginosa* lung infections in cystic fibrosis patients are caused by biofilm growing mucoid strains. Characteristically, gradients of nutrients and oxygen exist from the top to the bottom of biofilms and these gradients are associated to decrease of bacterial metabolic activity and increase of doubling times of the bacterial cells and these more or less dormant cells are therefore responsible for some of the tolerance to antibiotics. Biofilm growth is associated with increased level of mutations and with quorum sensing regulated mechanisms. Conventional resistance mechanisms such as chromosomal beta-lactamase, up-regulated efflux pumps and mutations of antibiotic target molecules in the bacteria also contribute to the survival of biofilms. Biofilm can be prevented by early aggressive antibiotic prophylaxis or therapy and they can be treated by chronic suppressive therapy. A promising strategy may be the use of enzymes which can dissolve the biofilm matrix (e.g. DNase, alginate lyase) and quorum sensing inhibitors, which increases biofilm susceptibility to antibiotics.

1. Introduction

Biofilm growing bacteria cause chronic infections (1) which are characterized by persisting inflammation and tissue damage (2). Chronic infections, including foreign

body infections, are infections which 1) persist in spite of antibiotic therapy and the innate and adaptive immune- and inflammatory response of the host, and 2) which, in contrast to colonization, are characterized by immune response and persisting pathology (Table 1).

2. The occurrence and architecture of bacterial biofilms

Foreign body infections are characterized by biofilm growth of the bacteria on the outer and/or inner surface of the foreign body (Table 2). Biofilm growth also occurs on natural surfaces such as teeth (3), heart valves (endocarditis), (4) in the lungs of cystic fibrosis (CF) patients causing chronic bronchopneumonia (2), in the middle ear in patients with persistent otitis media (5), in chronic rhinosinusitis (6), in chronic osteomyelitis and prosthetic joint infections (7) (8) (9), in intravenous catheters and stents (10) and in chronic wounds (11) (12)(Fig. 1 A & B). The microbes in biofilms are kept together by a self-produced biopolymeric matrix. The matrix contains polysaccharides, proteins and DNA originating from the microbes and the bacterial consortium can consist of one or more species living in a sociomicrobiological way (1, 13) (2, 14). The matrix is important since it provides structural stability and protection to the biofilm. The development of bacterial biofilms over time has been intensively studied *in vitro* by confocal scanning laser microscopy employing *green fluorescent protein (gfp)*-tagged bacteria. This technique has been combined with advanced *in silico* image-analysing to produce 3-D images of biofilm (15, 16) (17). As an example, *P. aeruginosa* produce a mature *in vitro* biofilm in 5-7 days (Fig. 2).

The development of an *in vitro* biofilm is initiated by planktonic (freely moving) bacteria which are attached reversible to a surface, which may be conditioned of a layer of e.g. proteins (a pellicle) (3) (18). At this stage, the bacteria

are still susceptible to antibiotics and this is in accordance with the success of the peroperative antibiotic prophylaxis for e.g. alloplastic surgery. The next step is irreversible binding to the surface within the next few hours and multiplication of the bacteria which forms microcolonies on the surface and begin to produce a polymer matrix around the microcolonies (18). The biofilm grows thickness (up to 50 μm) and under *in vitro* conditions, mushroom-like or tower-like structures are often observed in the mature biofilm. At that stage, the biofilm show maximum tolerance (= resistance) to antibiotics. Subsequently follows a stage where focal areas of the biofilm dissolves and the liberated bacterial cells can then spread to other location where new biofilms can be formed. This liberation process may be caused by bacteriophage activity within the biofilm (19). The mature biofilm may contain water-filled channels and thereby resemble primitive, multicellular organisms. Motile bacteria can use type-IV pili to mount or climb a biofilm formed by other bacteria and colonize the top of the biofilm resembling a hat (16). Important properties of biofilm growing bacteria are different from those of planktonic growing bacteria and this has significant diagnostic and therapeutic consequences. The bacteria appear different in biofilm infections since they are located close to each other in aggregates surrounded by the self-produced matrix. In clinical specimens (biopsies, pus, sputum) biofilm can often be recognized by light microscopy although precise identification of all the bacteria within a biofilm can only be done by DNA hybridization techniques and identification of the components of the biofilm matrix require specialized staining techniques (2). Traditional sampling techniques may not be sufficient to culture biofilm growing bacteria sticking to a surface unless the bacteria are released by ultrasonic pre-treatment (20). The ordinary culture techniques, however, reveals only the properties of planktonically growing bacteria, and e.g. antibiotic susceptibility

testing therefore gives misleading results which do not reflect the increased resistance of the bacteria when living in biofilms. The minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) of antibiotics to biofilm growing bacteria may be up to 100 – 1000 fold higher compared to planktonic bacteria (21) (22, 23). Methods to test biofilm growing bacteria have therefore been developed but their clinical relevance as regards prediction of clinical successful therapy awaits confirmation (22, 24, 25).

3. Stationary phase physiology, low oxygen concentration and slow growth.

Inspection of environmental as well as in vitro biofilms has revealed that oxygen concentration may be high at the surface but low in the center of biofilms where anaerobic conditions may be present (26). Likewise, growth, protein synthesis and metabolic activity is stratified in biofilms i.e. high level of activity at the surface and low level and slow growth or no growth in the center and this is one of the explanations for the reduced susceptibility of biofilms to antibiotics (27) (28). Very slow *in situ* growth rates of biofilms of *P. aeruginosa* have been measured in sputum of CF patients (average doubling time 2-3 hours and presence of a significant number of cells in stationary growth phase) (29). Monotherapy with antibiotics like beta-lactams, which are only active against dividing *P. aeruginosa* cells, are therefore not very efficient to eradicate biofilm infections (21).

4. Mutators

The mutation frequency of biofilm-growing bacteria is significantly increased compared to planktonically growing isogenic bacteria (30) and there is an increased horizontal gene transmission in biofilms (31). These physiological conditions may explain why biofilm growing bacteria easily become multi-drug resistant by means

of traditional resistance mechanisms against beta-lactam antibiotics, aminoglycosides and fluoroquinolones which are detected by routine susceptibility testing in the clinical microbiology laboratory where planktonic bacterial growth is investigated. Thus, bacterial cells in the biofilms may simultaneously produce enzymes that degrade antibiotics, have antibiotic targets of low-affinity and overexpresses efflux pumps which have a broad spectrum of substrates. Achievement of multiple mutations in a bacterial population size of 10^8 - 10^{10} / ml sputum as is attained under infection of the CF lung (32) imply the presence of a hypermutable bacterial subpopulation and the presence of high percentages of hypermutable *P. aeruginosa* isolates associated with antibiotic resistance have actually been found in CF patients (33) (34).

The hypermutable phenotype of CF *P. aeruginosa* isolates is due to alterations in the genes of the DNA repair systems of either the mismatch repair system (MMR) which involves *mutS*, *mutL* and *uvrD* or of the DNA oxidative lesions repair system (GO) which involves *mutT*, *mutY* and *mutM* (35, 36). It has been shown that mutations in either of the two systems determine emergence of antibiotic resistant isolates, especially due to selection of isolates expressing multidrug efflux-pumps (36, 37). An increased production of endogenous reactive oxygen species and a deficient anti-oxidant system (38) (39) determine an imbalance between oxidative burden and anti-oxidant defenses leading to oxidative stress in biofilms. The oxidative stress is considered to cause enhanced mutability in biofilms (30, 40) . Recent data suggest that microcolony structures, due to endogenous oxidative stress, are specific sites within the biofilms for enhanced genetic adaptation and evolutionary change (40). In addition, Boles and Singh (41) showed that the endogenous oxidative stress in biofilms promote antibiotic resistance and that addition of anti-oxidants reduced the

occurrence of diversity in biofilms. We have previously shown that oxidative stress is linked to the occurrence of hypermutable *P. aeruginosa* strains in CF patients (34).

In addition to their endogenous oxidative stress, the biofilm-growing bacteria in the CF airways are exposed to reactive oxygen species (ROS) from the activated polymorphonuclear leukocytes (PMN)(34). We have recently shown that the hypoxic environment in the CF sputum is due to the consumption of oxygen by PMNs which liberate reactive oxygen species which can react with the biofilm-embedded bacterial cells (42), thus creating an unique environment with low-oxygen tension filled with reactive oxygen species.

The hypermutability of bacteria in biofilms promotes the emergence of mutations conferring antibiotic resistance which will be selected for by the repeated antibiotic courses administered in order to maintain the lung function of CF patients.

Development of resistance to all classes of antibiotics during the chronic lung infection in CF has been documented (43). Resistance to beta-lactam antibiotics occur due to mutations in regulatory genes of the beta-lactamase production leading to the occurrence of isolates with stable or partially-stable derepressed production of AmpC beta-lactamase(44). Resistance to ciprofloxacin of CF *P. aeruginosa* isolates was shown to be mediated by mutations in *gyrA* and alterations in the two efflux systems MexCD-OprJ and MexEF-OprN and resistance to tobramycin was overexpression of the MexXY-OprM multi-drug efflux pump (45) (46). Resistance to colistin was shown to occur due to mutations in the *pmr* system involved in the LPS structure (47).

5. Chromosomal beta-lactamase and biofilm matrix components

The overproduction of chromosomally encoded AmpC cephalosporinase is considered the main mechanism of resistance of CF isolates of *P. aeruginosa* to β -lactam antibiotics (48). The most common phenotype of beta-lactamase production in CF isolates is the partially derepressed phenotype with high basal levels of beta-lactamase that can be induced further to higher levels in the presence of beta-lactam antibiotics (44). The role of this beta-lactamase phenotype is important especially for the resistance to beta-lactam antibiotics acting as strong inducers (carbapenems like imipenem). However, not all beta-lactams are strong inducers and the overexpression of the MexAB-OprM efflux pumps may play, together with beta-lactamases, an important role in the resistance to poor inducers (e.g. piperacillin). Totally derepressed beta-lactamase production is encountered in 2.5% of clinical CF isolates (44) and is responsible for the resistance to both poor and strong inducer beta-lactam antibiotics, independent of the presence of efflux pumps overexpression (49) (50)(51). We have found an insertion sequence (IS 1669) inactivating the *ampD* gene in several resistant clinical *P. aeruginosa* isolates with constitutive high expression of chromosomal beta-lactamase (51).

The diffusion barrier in biofilms (52) plays a role for biofilm resistance of *P. aeruginosa* that overproduce beta-lactamase due to the presence in the biofilm matrix of beta-lactamases which will hydrolyze the beta-lactam antibiotics before reaching the bacterial cells (53) (54). Giwercman (55) showed that imipenem and piperacillin were able to induce β -lactamase production in *P. aeruginosa* biofilms. Nichols (56) predicted from mathematical models that the biofilm would not afford protection

208 against diffusion of beta-lactam antibiotics into the bacteria embedded in the biofilm
209 as long as the level of chromosomal beta-lactamase is low. However bacteria
210 expressing high level of chromosomal beta-lactamase growing in biofilms would be
211 exposed to reduced concentration of beta-lactam antibiotics due to accumulation of
212 the enzyme in the polysaccharide matrix. The extracellular β -lactamase would
213 inactivate the antibiotic as it penetrates, thereby protecting the deeper-lying cells.

214 The source of β -lactamase in biofilms has been considered to be from a layer of lysed
215 bacteria due to exposure to an antibiotic, with release of defensive enzymes into the
216 extracellular space. We have shown that the source of β -lactamase in biofilm may
217 also be the membrane vesicles (MVs) containing beta-lactamase liberated by resistant
218 *P. aeruginosa* bacteria (57)(58)(54) and we have shown, that high level of free
219 chromosomal beta-lactamase is present in CF sputum (58). We have also shown that
220 strong inducers like imipenem will induce the beta-lactamase through all the bacterial
221 layers while poorer inducers like ceftazidime will influence just the superficial layers
222 of the biofilm, probably due to the inactivation of the antibiotic by beta-lactamase (59,
223 60)(61)(58) (Figure 3).

224 The protective role played by beta-lactamase in impairing the penetration of beta-
225 lactams in the biofilm can be seen in Fig. 3A. Treatment with ceftazidime of a
226 biofilm formed by a *P. aeruginosa* CF strain with stable derepressed levels of beta-
227 lactamase due to an insertion sequence in *ampD* (*P. aeruginosa ampD*⁻) killed very
228 few bacterial cells (dead bacteria in red) (Figure 3B) in contrast to the complemented
229 strain with low level of beta-lactamase (Fig. 3 C). However, addition of aztreonam
230 improved the efficacy of ceftazidime treatment of the biofilm (Figure 3D, probably
231 because aztreonam acts as a beta-lactamase inhibitor (58). In addition, meropenem, a

beta-lactamase stable beta-lactam showed good *in vitro* efficacy in the treatment of *P. aeruginosa* biofilms (22) (61). Treatment with ceftazidime of a biofilm formed by the same strain expressing basal levels of beta-lactamase due to complementation with the wild-type *ampD* (*P. aeruginosa ampD*+) led to eradication of the biofilm (Figure 3C).

The matrix of the biofilm may also be part of the resistance mechanisms to antibiotics since e.g. sub-MIC concentrations of beta-lactam antibiotics induce increased alginate synthesis in *P. aeruginosa* biofilms (Fig. 4) (62) (63) and also enhance the biofilm matrix of some slime-producing coagulase-negative staphylococci (64)(65). Originally, it has been thought that tolerance of the biofilms to aminoglycosides was due to a transport limitation due to the binding of these positively charged antibiotics to the negatively charged exopolysaccharide matrix but the repeated dosing of antibiotics during therapy probably leads to saturation of the binding sites (66) (67) (68). As previously discussed, the oxygen limitation and the metabolic rates are probably more important factors which contribute to the tolerance of biofilms to aminoglycosides and ciprofloxacin (69). However, in the respiratory zone of the CF lung with poor access to aminoglycoside aerosols, where the antibiotic concentration is low delayed penetration of the aminoglycosides through thick biofilms may play a role in the tolerance of biofilms to aminoglycosides (2). Recently, it has been shown that administration of DNase and alginate lyase enhanced the activity of tobramycin in biofilms by dissolving the biofilm matrix (70).

6. Tolerance, adaptive resistance and efflux pumps

Colistin is only active against the non-dividing central part of *P. aeruginosa* biofilms *in vitro* (Fig. 5A), whereas the superficial, metabolic active part of the biofilm

become tolerant due to upregulation of PmrA-PmrB two-component regulatory systems involved in the adaptive resistance to cationic peptides leading to addition of aminoarabinose to lipid A of LPS (22, 71, 72). Since the metabolic active surface layer of the biofilm is susceptible to ciprofloxacin (Fig. 5B) in contrast to the dormant central part of the biofilm, combination therapy with this drug and colistin was able to kill all cells in the biofilm *in vitro* (Fig. 5C)(71). The clinical efficacy has been demonstrated of this combination therapy for the early eradication treatment of *P. aeruginosa* in CF patients (73).

Tolerance of biofilms to tobramycin is also mediated by low metabolic activity but the high cell density that results in accumulation of extracellular signalling molecules is probably important, as it has been shown that tolerance to tobramycin of *P. aeruginosa* strain PAO1 biofilm is quorum sensing mediated (Fig. 6)(74) (see below). In addition, a non-specific mechanism for the tolerance of the metabolic active cells to colistin was shown to be up-regulation of the efflux pump MexAB-OprM (71). Furthermore, increased efflux pump activity due to mutations has been shown to be a major resistance mechanism against aminoglycoside antibiotics and fluorquinolones in *P. aeruginosa* from CF patients (46) (45).

7. High cell density and quorum sensing

Bacteria communicate by means of synthesising and reacting on signal molecules (75-78). The term QS indicates that this system permits bacteria to sense when a critical number (concentration) of bacteria are present in a limited space in the environment and respond by activating certain genes which then produce e.g. virulence factors such as enzymes or toxins. The QS molecules are small peptides in many Gram-

positive bacteria whereas the most well-described QS molecules in Gram-negative bacteria are N-acyl-L-homoserinelactones (AHL) (78). For *P. aeruginosa* QS regulates the production of virulence factors such as extracellular enzymes and cellular lysins (e.g. rhamnolipid), which are important for the pathogenesis of infections where it functions as a protective shield against phagocytes (79, 80) (81). QS may also have influence on the development of the biofilm (82) and QS have been shown to determine the tolerance of *P. aeruginosa* biofilms against antibiotic therapy and against the innate inflammatory response dominated by polymorphonuclear leucocytes (PMNs) (83). The connections between QS and biofilms has been named sociomicrobiology (84).

8. Quorum sensing inhibitors (QSI)

Much of our knowledge about QS originate from experiments with QS knock-out mutants and from use of naturally occurring and artificially syntetized QSI compounds (85)(86). Screening for QSI in nature has identified many QSI compounds (87). These naturally occurring QSI compounds can be synthesized and their structure modified and used to inhibit QS *in vivo* in experimental animal infections (85). Since it has been shown, that bacteria used for experimental animal biofilm infections actually communicate *in vivo* (88) and also in e.g. CF patients with chronic *P. aeruginosa* lung infection QSI may be used to treat the infection (89). Interestingly, some macrolide antibiotics like azithromycin (90) but also other antibiotics like ceftazidime and ciprofloxacin (91) inhibit QS in *P. aeruginosa* at sub-MIC concentrations leading to inhibition of the virulence of these bacteria although they cannot inhibit their growth at obtainable concentrations *in vivo*. Controlled

clinical trials using azithromycin to treat the chronic *P. aeruginosa* lung infection in both CF children and adults have shown significant improvement of their lung function (92) (93) (94). Most CF patients with chronic *P. aeruginosa* lung infections are therefore now treated continuously with azithromycin (95). An expected side-effect has, however, been development of resistance to macrolides in other pathogenic bacteria like *S. aureus* in CF patients (96). It would therefore be desirable to develop QSI without conventional growth-inhibiting or bacteriocidal activity (78). One example of such naturally occurring QSI is found in garlic extract, which *in vitro* and *in vivo* have been able to render otherwise resistant *P. aeruginosa* biofilms susceptible to antibiotic therapy and to PMN activity and the consequence is eradication of the biofilm both as regards antibiotic therapy and PMN activity which dominates the inflammatory response in CF patients (Fig. 7) (97). According to current knowledge, QSI resistance can only occur due to mutations, which renders the QS deficient bacteria unable to produce virulence factors (98) i.e. the bacteria become non-virulent similar to the result of QSI therapy. If this holds true, then resistance problems against conventional antibiotics as we face today will not be a clinical problem.

Foreign body infections constitute an steadily increasing medical problem and comprise e.g. intravenous catheters, intrauterine catheters, naso-laryngeal tubes, stents (Table 2, Fig. 1A), alloplastic materials, hydrocephalus shunts and artificial hearts (99). If foreign bodies become colonized with biofilm-forming bacteria the result is most often chronic inflammation around the foreign body which either has to be replaced or treated with sometimes life-long antibiotic suppressive therapy, although early therapy may sometimes lead to eradication of the condition. Antibiotic coated foreign bodies like e.g. catheters, vascular prosthesis have been introduced to prevent biofilm formation and they are quite efficient (100) (101). It is, however, desirable to

develop other compounds due to the risk of development of bacterial resistance and QSI are strong candidates. QSI have been shown synergistically to improve the weak effects of antibiotics and PMNs on biofilm growing bacteria *in vitro* and *in vivo* in animal experiments leading to elimination of biofilms (Fig. 7) (102). These results have lead to further development of QSI as pharmaceutical compounds for patients who are subject to implantation of foreign bodies.

9. Prophylaxis and treatment of *P. aeruginosa* biofilms in CF lungs – perspectives for other biofilm infections?

The currently used methods for preventing chronic *P. aeruginosa* biofilms in CF lungs are 1) prevention of cross-infection from other already chronically infected CF patients by isolation techniques and hygienic measures (103), 2) early aggressive eradication therapy of intermittent colonization by means of oral ciprofloxacin and nebulized colistin for 3 weeks or even better for 3 months or by using nebulized tobramycin as monotherapy (104), 3) daily nebulized DNase (Pulmozyme) (105). These 3 methods, which are combined in most CF centers, are successful and cost-efficient and has completely changed the epidemiology of chronic *P. aeruginosa* lung infection in CF patients from being very common in CF children to being predominantly a problem for adult patients and no problems of resistance to the antibiotics have been recorded (73, 106, 107). Early aggressive eradication therapy has also been shown to superior in an animal model of *P. aeruginosa* infection in CF (108). Furthermore, although vaccines against *P. aeruginosa* have been developed and undergone clinical trials, they have not been further developed due to the success of the early, aggressive eradication therapy (109).

The recommended method for treatment of chronic *P. aeruginosa* biofilm

infection is chronic suppressive antibiotic therapy (2)(110) (Fig. 8) which is started when the chronic infection is diagnosed (continuous colonization at the monthly bacteriological examination for 6 months and/or increased level of antibodies against *P. aeruginosa* (111)(112) . The chronic suppressive therapy consists of daily nebulized colistin or tobramycin for the rest of the patient's life combined with either regular 2-week courses every 3 months of intravenous anti-pseudomonas antibiotics (combination therapy of 2 antibiotics: tobramycin or colistin + ceftazidime, or piperacillin/tazobactam, or carbapenem, or aztreonam, or ciprofloxacin) or ad hoc intravenous therapy when clinical deteriorations occur (110). Additionally, DNase is inhaled every day to reduce the viscosity of the DNA-containing sputum (105), and daily oral azithromycin is given to the patients as mentioned above (95). The chronic suppressive therapy (maintenance therapy) has successfully been able to maintain the pulmonary function or slow the decline of the pulmonary function and prolong the life of the patients for many years (113). The side-effects of the maintenance therapy is high level of conventional resistance mechanisms in the persisting strains and high level of allergy to the beta-lactam antibiotics (114) (43). Similar principles, a) systemic or local prophylactic use of antibiotics to prevent biofilm formation, b) early aggressive eradication therapy to eradicate planktonic growth or early biofilm formation and c) chronic suppressive antibiotic therapy to maintain the function of a inserted foreign body are gradually being introduced in other area of biofilm infection (100) (8) (9). Possibly other approaches taken advantage of the use of QSI may further improve the management of biofilm infections.

Funding

None.

Conflict of interest statement

None to declare.

References

1. Costerton W, Veeh R, Shirtliff M, Pasmore M, Post C, Ehrlich G. The application of biofilm science to the study and control of chronic bacterial infections. *J Clin Invest.* 2003;112(10):1466-77.
2. Bjarnsholt T, Jensen PØ, Fiandaca MJ, Pedersen J, Hansen CR, Andersen CB, et al. *Pseudomonas aeruginosa* biofilms in the respiratory tract of cystic fibrosis patients. *Pediatric Pulmonology.* 2009;44:547-58.
3. Kolenbrander P. E., Palmer Jr. R. J. In: (ed) eAP, Washington, D.C. Human oral bacterial biofilms. In: Ghannoum M., A. OTG, editors. *Microbial biofilms.* Washington D.C.: ASM Press; 2004.
4. Høiby N, Döring G, Schiøtz PO. The role of immune complexes in the pathogenesis of bacterial infections. *Ann Rev Microbiol.* 1986;40:29-53.
5. Hall-Stoodley L, Hu FZ, Gieseke A, Nistico A, Nguyen L, Hayes J, et al. Direct detection of bacterial biofilms on the middle-ear mucosa of children with chronic otitis media. *Journal of the American Medical Association.* 2006;296:202-11.
6. Sanderson A.R., Leid J.G., D. H. Bacterial biofilms on the sinus mucosa of human subjects with chronic rhinosinusitis. *Laryngoscope.* 2006;116:1121-6.
7. Gristina A.G., Oga M., Webb L.X., C.D H. Adherent bacterial colonization in the pathogenesis of osteomyelitis. *Science.* 1985;228:990-3.
8. Trampuz A., W. Z. Prosthetic joint infections: update in diagnosis and treatment. *Swiss Medical Weekly.* 2005;135:243-51.
9. Del Pozo J.L., R. P. Infection associated with prosthetic joints. *new England Journal of Medicine.* 2009;361:787-94.
10. Taconelli E., Smith G., Hieke K., Lafuma A., P. B. Epidemiology, medical outcomes and costs of catheter-related bloodstream infection in intensive care units of four European countries: literature- and registry-based estimates. *Journal of Hospital Infection.* 2009;72:97-103.
11. Kirketerp-Møller K., Jensen P.Ø., Fazli M., Madsen K.G., Pedersen J., Moser C., et al. Distribution, and ecology of bacteria in chronic wounds. *Journal of Clinical Microbiology.* 2008;46:2717-22.

12. Bjarnsholt T., Kirketerp-Møller K., Jensen P.Ø., Madsen K.G., Phipps R., Krogfelt K., et al. Why chronic wounds won't heal: a novel hypothesis. *Wound Repair and Regeneration*. 2008;16:2-10.
13. Wingender J, Strathmann M, Rode A, Leis A, Flemming HC. Isolation and biochemical characterization of extracellular polymeric substances from *Pseudomonas aeruginosa*. *Microbial Growth In Biofilms*, 2001. p. 302-14.
14. Whitchurch CB, TolkerNielsen T, Ragas PC, Mattick JS. Extracellular DNA required for bacterial biofilm formation. *Science*. 2002;295(5559):1487.
15. Sternberg C, Christensen BB, Johansen T, Nielsen AT, Andersen JB, Givskov M, et al. Distribution of bacterial growth activity in flow-chamber biofilms. *Appl Environ Microbiol*. 1999;65(9):4108-17.
16. Klausen M, Heydorn A, Ragas P, Lambertsen L, AaesJorgensen A, Molin S, et al. Biofilm formation by *Pseudomonas aeruginosa* wild type, flagella and type IV pili mutants. *Mol Microbiol*. 2003;48(6):1511-24.
17. Heydorn A, Ersboll B, Kato J, Hentzer M, Parsek MR, TolkerNielsen T, et al. Statistical analysis of *Pseudomonas aeruginosa* biofilm development: Impact of mutations in genes involved in twitching motility, cell-to-cell signaling, and stationary-phase sigma factor expression. *Appl Environ Microbiol*. 2002;68(4):2008-17.
18. Marshall KC. Biofilms: An overview of bacterial adhesion, activity, and control at surfaces. *ASM News*. 1992;58(4):202-7.
19. Webb JS, Thompson LS, James S, Charlton T, TolkerNielsen T, Koch B, et al. Cell death in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol*. 2003;185(15):4585-92.
20. Kobayashi N., Bauer T.W., Tuohy M.J., Fujishiro T., G.W. P. Brief ultrasonication improves detection of biofilm-formative bacteria around a metal implant. *Clinical Orthopaedic Related Research*. 2007;475:210-3.
21. Anwar H, Costerton JW. Enhanced Activity of Combination of Tobramycin and Piperacillin for Eradication of Sessile Biofilm Cells of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 1990;34(9):1666-71.
22. Moskowitz SM, Foster JM, Emerson J, Burns JL. Clinically feasible biofilm susceptibility assay for isolates of *Pseudomonas aeruginosa* from patients with cystic fibrosis. *J Clin Microbiol*. 2004;42(5):1915-22.
23. Bjarnsholt T., Kirketerp-Møller K., Kristiansen S., Phipps R., Nielsen A.K., Jensen P.Ø., et al. Silver against *Pseudomonas aeruginosa* biofilms. *APMIS*. 2007;115:921-8.
24. Anwar H, Dasgupta MK, Costerton JW. Testing the Susceptibility of Bacteria in Biofilms to Antibacterial Agents. *Antimicrob Agents Chemother*. 1990;34(11):2043-6.
25. Domingue G, Ellis B, Dasgupta M, Costerton JW. Testing antimicrobial susceptibilities of adherent bacteria by a method that incorporates guidelines of the National Committee for Clinical Laboratory Standards. *J Clin Microbiol*. 1994;32(10):2564-8.
26. Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappinscott HM. Microbial biofilms. *Annu Rev Microbiol*. 1995;49:711-45.
27. Werner E, Roe F, Bugnicourt A, Franklin MJ, Heydorn A, Molin S, et al. Stratified growth in *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol*. 2004;70(10):6188-96.

28. Keren I, Kaldalu N, Spoering A, Wang YP, Lewis K. Persister cells and tolerance to antimicrobials. *Fems Microbiol Lett.* 2004;230(1):13-8.
29. Yang L, Haagenzen JAJ, Jelsbak L, Johansen HK, Sternberg C, Høiby N, et al. In situ growth rates and biofilm development of *Pseudomonas aeruginosa* populations in chronic lung infection. *Journal of Bacteriology.* 2008;190:2767-76.
30. Driffield K, Miller K, Bostock M, O'Neill AJ, Chopra I. Increased mutability of *Pseudomonas aeruginosa* in biofilms. *Journal of Antimicrobial Chemotherapy.* 2008;61:1053-6.
31. Molin S, Tolker-Nielsen T. Gene transfer occurs with enhanced efficiency in biofilms and induces enhanced stabilisation of the biofilm structure. *Curr Opin Biotechnol.* 2003;14(3):255-61.
32. Høiby N, Johansen HK, Moser C, Song ZJ, Ciofu O, Kharazmi A. *Pseudomonas aeruginosa* and the biofilm mode of growth. *Microbes and Infection.* 2001;3:1-13.
33. Oliver A, Canton R, Campo P, Baquero F, Blazquez J. High frequency of hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science.* 2000;288(5469):1251-3.
34. Ciofu O, Riis B, Pressler T, Poulsen HE, Høiby N. Occurrence of hypermutable *P. aeruginosa* in cystic fibrosis patients is associated with the oxidative stress caused by chronic lung inflammation. *Antimicrobial Agents and Chemotherapy.* 2005;49:2276-82.
35. Oliver A, Sanchez JM, Blazquez J. Characterization of the GO system of *Pseudomonas aeruginosa*. *Fems Microbiol Lett.* 2002;217(1):31-5.
36. Mandsberg LF, Ciofu O, Kirkby N, Christiansen LE, Poulsen HE, Høiby N. Antibiotic resistance in *P. aeruginosa* strains with increased mutation frequency due to inactivation of the DNA oxidative repair system. *Antimicrobial Agents and Chemotherapy.* 2009;53:2483-91.
37. Macia MD, Blanquer D, Togores B, Saulea J, Perez JL, Oliver A. Hypermutation is a key factor in development of multiple-antimicrobial resistance in *Pseudomonas aeruginosa* strains causing chronic lung infections. *Antimicrob Agents Chemother.* 2005;49(8):3382-6.
38. Hassett D.J., Ma J.F., Elkins J.G., McDermott T.R., Ochsner U.A., West S.E.H., et al. Quorum sensing in *Pseudomonas aeruginosa* controls expression of catalase and superoxide dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. *Mol Microbiol.* 1999;34(5):1082-93.
39. Mai-Prochnow A., Lucas-Elio P., Egan S, Thomas T, Webb J, Sanchez-Amat, A., Kjellerberg S. Hydrogen peroxide linked to lysine oxidase activity facilitates biofilm differentiation and dispersal in several gram-negative bacteria. *Journal of Bacteriology.* 2008;190:5493-501.
40. Conibear T.C., Collins S.L., J.S. W. Role of mutation in *Pseudomonas aeruginosa* biofilm development. *PLoS.* 2009;14(7):6289.
41. Boles B.R., P.K. S. Endogenous oxidative stress produces diversity and adaptability in biofilm communities. *Proceedings of the National Academy of Science USA.* 2008;105(34):12503-8.
42. Kolpen M, Hansen CR, Bjarnsholt T, Moser C, Christensen LD, van Gennip M, et al. Ongoing respiratory burst of polymorphonuclear leukocytes accelerates oxygen depletion in sputum from patients with chronic *Pseudomonas aeruginosa* lung infection. *Thorax.* 2009, in press.

43. Ciofu O, Giwercman B, Pedersen SS, Hoiby N. Development of antibiotic resistance in *Pseudomonas aeruginosa* during two decades of antipseudomonal treatment at the Danish CF center. *APMIS*. 1994;102(9):674-80.
44. Ciofu O. *Pseudomonas aeruginosa* chromosomal beta-lactamase in patients with cystic fibrosis and chronic lung infection - Mechanism of antibiotic resistance and target of the humoral immune response. *Apmis*. 2003;111:4-47.
45. Jalal S, Ciofu O, Høiby N, Gotoh N, Wretling B. Molecular mechanisms of fluoroquinolone resistance in *Pseudomonas aeruginosa* isolates from cystic fibrosis patients (Vol 44, pg 710, 2000). *Antimicrob Agents Chemother*. 2000;44(5):1410.
46. Islam S, Oh H, Jalal S, Ciofu O, Høiby N, Wretling B. Chromosomal resistance mechanisms for aminoglycosides in *Pseudomonas aeruginosa* cystic fibrosis isolates. *Clinical Microbiology and Infection*. 2009;15:60-6.
47. Johansen HK, Moskowitz SM, Ciofu O, Pressler T, Høiby N. Spread of colistin-resistant non-mucoid *Pseudomonas aeruginosa* among chronically infected Danish cystic fibrosis patients. *Journal of Cystic Fibrosis*. 2008;7:391-7.
48. Giwercman B, Lambert PA, Rosdahl VT, Shand GH, Hoiby N. Rapid Emergence of Resistance in *Pseudomonas-Aeruginosa* in Cystic Fibrosis Patients Due to In Vivo Selection of Stable Partially Derepressed Beta-Lactamase Producing Strains. *J Antimicrob Chemother*. 1990;26(2):247-59.
49. Nakae T, Nakajima A, Ono T, Saito K, Yoneyama H. Resistance to beta-lactam antibiotics in *Pseudomonas aeruginosa* due to interplay between the MexAB-OprM efflux pump and beta-lactamase. *Antimicrob Agents Chemother*. 1999;43(5):1301-3.
50. Nakae T, Nakajima A, Ono T, Saito K, Yoneyama H. Resistance to beta-lactam antibiotics in *Pseudomonas aeruginosa* due to interplay between the MexAB-OprM efflux pump and beta-lactamase. *Antimicrob Agents Chemother*. 1999 May;43(5):1301-3.
51. Bagge N, Ciofu O, Hentzer M, Campbell JIA, Givskov M, Hoiby N. Constitutive high expression of chromosomal beta-lactamase in *Pseudomonas aeruginosa* caused by a new insertion sequence (IS1669) located in *ampD*. *Antimicrob Agents Chemother*. 2002;46(11):3406-11.
52. deBeer D, Stoodley P, Lewandowski Z. Measurement of local diffusion coefficients in biofilms by microinjection and confocal microscopy. *Biotechnol Bioeng*. 1997;53(2):151-8.
53. Dibdin GH, Assinder SJ, Nichols WW, Lambert PA. Mathematical model of beta-lactam penetration into a biofilm of *Pseudomonas aeruginosa* while undergoing simultaneous inactivation by released beta-lactamases. *J Antimicrob Chemother*. 1996;38(5):757-69.
54. Ciofu O, Beveridge TJ, Kadurugamuwa J, Walther Rasmussen J, Hoiby N. Chromosomal beta-lactamase is packaged into membrane vesicles and secreted from *Pseudomonas aeruginosa*. *J Antimicrob Chemother*. 2000;45(1):9-13.
55. Giwercman B, Jensen ET, Hoiby N, Kharazmi A, Costerton JW. Induction of beta-Lactamase Production in *Pseudomonas-Aeruginosa* Biofilm. *Antimicrob Agents Chemother*. 1991;35(5):1008-10.
56. Nichols W.W., Evans M.J., Slack M.P., H.L. W. The penetration of antibiotics into aggregates of mucoid and non-mucoid *Pseudomonas aeruginosa*. *Journal of General Microbiology* 1989;135:1291-303.

- 559 57. Ciofu O, Beveridge TJ, Kadurugamuwa J, Walther-Rasmussen J, Hoiby N.
560 Chromosomal beta-lactamase is packaged into membrane vesicles and secreted
561 from *Pseudomonas aeruginosa*. *J Antimicrob Chemother*. 2000;45(1):9-13.
- 562 58. Giwercman B, Meyer C, Lambert PA, Reinert C, Hoiby N. High-Level beta-
563 Lactamase Activity in Sputum Samples from Cystic Fibrosis Patients During
564 Antipseudomonal Treatment. *Antimicrob Agents Chemother*. 1992;36(1):71-6.
- 565 59. Bagge N, Hentzer M, Andersen JB, Ciofu O, Givskov M, Hoiby N. Dynamics
566 and spatial distribution of beta-lactamase expression in *Pseudomonas*
567 *aeruginosa* biofilms. *Antimicrob Agents Chemother*. 2004 Apr;48(4):1168-74.
- 568 60. Bagge N, Hentzer M, Andersen JB, Ciofu O, Givskov M, Hoiby N. Dynamics
569 and spatial distribution of beta-lactamase expression in *Pseudomonas*
570 *aeruginosa* biofilms. *Antimicrob Agents Chemother*. 2004;48(4):1168-74.
- 571 61. Hill D., Rose B., Pajkos A., Robinson M., Bye P., Bell S., et al. Antibiotic
572 susceptibilities of *Pseudomonas aeruginosa* isolates derived from patients with
573 cystic fibrosis under aerobic, anaerobic, and biofilm conditions. *Journal of*
574 *Clinical Microbiology*. 2005;43:5085-90.
- 575 62. Bagge N, Schuster M, Hentzer M, Ciofu O, Givskov M, Greenberg EP, et al.
576 *Pseudomonas aeruginosa* biofilms exposed to imipenem exhibit changes in
577 global gene expression and beta-lactamase and alginate production. *Antimicrob*
578 *Agents Chemother*. 2004;48(4):1175-87.
- 579 63. Wood L.F., Leech A.J., D.E. O. Cell wall-inhibitory antibiotics activate the
580 alginate biosynthesis operon in *Pseudomonas aeruginosa*: roles of sigma (AlgT)
581 and the AlgW and Prc proteases. *Molecular Microbiology*. 2006;62:412-26.
- 582 64. Dunne WM. Effects of Subinhibitory Concentrations of Vancomycin or
583 Cefamandole on Biofilm Production by Coagulase-Negative Staphylococci.
584 *Antimicrobial Agents and Chemotherapy*. 1990 34(3):390-3.
- 585 65. Permin H, Koch C, Høiby N, Christensen HO, Møller AF, Møller S.
586 Ceftazidime treatment of chronic *Pseudomonas aeruginosa* respiratory tract
587 infection in cystic fibrosis. *J Antimicrob Chemother*. 1983;12, supplementum
588 A:313-23.
- 589 66. Anwar H, Strap JL, Costerton JW. Establishment of aging biofilms: possible
590 mechanism of bacterial resistance to antimicrobial therapy. *Antimicrobial Agents*
591 *and Chemotherapy*. 1992;36:1347-51.
- 592 67. Levy J, Smith, A.L., Kenny, M.A., Ramsey, B., & Schoenknecht, F.D.
593 Bioactivity of Gentamicin in purulent sputum from patients with cystic fibrosis
594 or bronchiectasis: Comparison with activity in serum. *J Infect Dis*.
595 1983;148:1069-76.
- 596 68. Hunt BE, Weber A, Berger A, Ramsey B, Smith AL. Macromolecular
597 mechanisms of sputum inhibition of tobramycin activity. *Antimicrob Agents*
598 *Chemother*. 1995;39(1):34-9.
- 599 69. Walters MC, Roe F, Bugnicourt A, Franklin MJ, Stewart PS. Contributions
600 of antibiotic penetration, oxygen limitation, and low metabolic activity to
601 tolerance of *Pseudomonas aeruginosa* biofilms to ciprofloxacin and tobramycin.
602 *Antimicrob Agents Chemother*. 2003;47(1):317-23.
- 603 70. Alipour M., Suntres Z. E., A. O. Importance of DNase and alginate lyase for
604 enhancing free and liposome encapsulated aminoglycoside activity against
605 *Pseudomonas aeruginosa*. *Journal of Antimicrobial Chemother* apy.
606 2009;64:317-25.

- 607 71. Pamp SJ, Gjermansen M, Johansen HK, Tolker-Nielsen T. Tolerance to the
608 antimicrobial peptide colistin in *Pseudomonas aeruginosa* biofilms is linked to
609 metabolically active cells, and depends on the *prm* and *mexAB-oprM* genes.
610 *Molecular Microbiology*. 2008;68:223-40.
- 611 72. Haagensen J, Klausen M, Ernst RK, Miller SI, Folkesson A, Tolker-Nielsen
612 T, et al. Differentiation and distribution of colistin- and sodium dodecyl sulfate-
613 tolerant cells in *Pseudomonas aeruginosa* biofilms. *Journal of Bacteriology*.
614 2007;189:28-37.
- 615 73. Hansen CR, Pressler T, Høiby N, . Early aggressive eradication therapy for
616 intermittent *Pseudomonas aeruginosa* airway colonization in cystic fibrosis
617 patients: 15 years experience. *Journal of Cystic Fibrosis*. 2008;7:523-30.
- 618 74. Bjarnsholt T, Jensen P-Ø, Burmølle M, Hentzer M, Haagensen JAJ, Hougen
619 HP, et al. *Pseudomonas aeruginosa* tolerance to tobramycin, hydrogen peroxide
620 and polymorphonuclear leukocytes is quorum-sensing dependent *Microbiology*.
621 2005;151:373-83.
- 622 75. Greenberg EP. Quorum sensing in Gram-negative bacteria. *ASMNews*.
623 1997;63(7):371-7.
- 624 76. Lindum PW, Anthoni U, Christophersen C, Eberl L, Molin S, M G. N-acyl-L-
625 homoserine lactone autoinducers contro production of an extracellular
626 lipopeptide biosurfactant required for swarming motility of *Serratia liquefaciens*
627 MG1. *J Bact*. 1998;180(23):6384-8.
- 628 77. Williams P, Camara M, Hardman A, Swift S, Milton D, Hope VJ, et al.
629 Quorum sensing and the population-dependent control of virulence. *Phil Trans*
630 *Roy Soc London B*. 2000;355(1397):667-80.
- 631 78. Bjarnsholt T, Givskov M. The role of quorum sensing in the pathogenicity
632 of the cunning aggressor *Pseudomonas aeruginosa* *Analytical and Bioanalytical*
633 *Chemistry*. 2007;387:409-14.
- 634 79. Jensen PØ, Bjarnsholt T, Phipps R, Rasmussen TB, Calum H, Christoffersen
635 L, et al. Rapid necrotic killing of polymorphonuclear leukocytes is caused by
636 quorum-sensing-controlled production of rhamnolipid by *Pseudomonas*
637 *aeruginosa*. *Microbiology*. 2007;153:1329-38.
- 638 80. Gennip Mv, Christensen LD, Alhede M, Phipps R, Jensen PØ,
639 Christophersen L, et al. Inactivation of the *rhlA* genes in *Pseudomonas*
640 *aeruginosa* prevents rhamnolipid production, disabling the protection against
641 polymorphonuclear leukocytes. *APMIS*. 2009;117:537-46.
- 642 81. Alhede M., Bjarnsholt T., Jensen P.Ø., Phipps R.K., Moser C., Christophersen
643 L., et al. *Pseudomonas aeruginosa* recognizes and responds aggressively to the
644 presence of polymorphonuclear leukocytes. *Microbiology*. 2009.
- 645 82. Davies DG, Parsek MR, Pearson JP, Iglewski BH, Costerton JW, Greenberg
646 EP. The involvement of cell-to-cell signals in the development of a bacterial
647 biofilm. *Science*. 1998;280(5361):295-8.
- 648 83. Bjarnsholt T., Jensen P.Ø., Burmølle M., Hentzer M., Haagensen J.A.,
649 Hougen H.P., et al. *Pseudomonas aeruginosa* tolerance to tobramycin, hydrogen
650 peroxide and polymorphonuclear leukocytes is quorum-sensing dependent.
651 *Microbiology*. 2005
652 ;151:373-83.
- 653 84. Parsek M.R., E.P. G. Sociomicrobiology: the connections between quorum
654 sensing and biofilms. . *Trends in Microbiology*. 2005;13:27-33.

85. Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, et al. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *Embo J*. 2003;22(15):3803-15.
86. Givskov M, de Nys R, Manefield M, Gram L, Maximilien R, Eberl L, et al. Eukaryotic interference with homoserine lactone mediated procaryotic signalling. *Journal of Bacteriology*. 1996;178:6618-22.
87. Rasmussen TB, Bjarnsholt T., Hentzer M., Kristoffersen P., Köte M, Eberl L, et al. Screening for Quorum Sensing Inhibitors Using a Novel Tool – the QSI Selector. *Journal of Bacteriology*. 2005;151:373-83.
88. Wu H, Song ZJ, Hentzer M, Andersen JB, Heydorn A, Mathee K, et al. Detection of N-acylhomoserine lactones in lung tissues of mice infected with *Pseudomonas aeruginosa*. *Microbiology Uk*. 2000;146:2481-93.
89. Hentzer M, Givskov M. Pharmacological Inhibition of Quorum Sensing for Treatment of Chronic Bacterial Infections. *Journal of Clinical Investigation*. 2003;112:1300-7.
90. Hoffmann N, Lee B, Hentzer M, Rasmussen TB, Song Z, Johansen HK, et al. Azithromycin blocks quorum sensing and alginate polymer formation and increases the sensitivity to serum and stationary growth phase killing of *P. aeruginosa* and attenuates chronic *P. aeruginosa* lung infection in *Cftr*^{-/-} mice. *Antimicrobial Agents and Chemotherapy*. 2007(in press).
91. Skindersø M.E., Alhede M., Phipps R., Yang L., Jensen P.O., Rasmussen T.B., et al. Effects of antibiotics on quorum sensing in *Pseudomonas aeruginosa*. *Antimicrobial Agents Chemother*. 2008;52:3548-663.
92. Wolter J, Seeney S, Bell S, Bowler S, Masel P, McCormack J. Effect of long term treatment with azithromycin on disease parameters in cystic fibrosis: a randomized trial. *Thorax*. 2002;57:212-6.
93. Equi A, BalfourLynn IM, Bush A, Rosenthal M. Long term azithromycin in children with cystic fibrosis: a randomised, placebo-controlled crossover trial. *Lancet*. 2002;360(9338):978-84.
94. Saiman L, Marshall BC, MayerHamblett N, Burns JL, Quittner AL, Cibene DA, et al. Azithromycin in patients with cystic fibrosis chronically infected with *Pseudomonas aeruginosa* - A randomized controlled trial. *Jama J Am Med Assn*. 2003;290(13):1749-56.
95. Hansen CR, Pressler T, Koch C, Høiby N. Long-term azithromycin treatment of cystic fibrosis patients with chronic *P. aeruginosa* infection; an observational cohort study. *Journal of Cystic Fibrosis* 2005;4:35-40.
96. Tramper-Stranders GA, Wolfs TFW, Fleer A, Kimpen JLL, Ent CKvd. Maintenance Azithromycin treatment in pediatric patients with cystic fibrosis. Long-term outcomes related to macrolide resistance and pulmonary function. *The Pediatric Infectious Disease Journal*. 2007;26:8-12.
97. Bjarnsholt T, Jensen PØ, Rasmussen TB, Christophersen L, Calun H, Hentzer M, et al. Garlic blocks quorum sensing and promotes rapid clearing of pulmonary *Pseudomonas aeruginosa* infections. *Microbiology*. 2005;151:3873-80.
98. Lee B, Haagensen JAJ, Ciofu O, Andersen JB, Høiby N, Molin S. Heterogeneity of biofilms formed by non-mucoid *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Journal of Clinical Microbiology*. 2005;43:5247-55.

99. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: A common cause of persistent infections. *Science*. 1999 284(5418):1318-22.
100. Stensballe J, Tvede M, Loorns D, Lippert FK, Dahl B, Tønnesen E, et al. Infection risk with nitrofurazone-impregnated urinary catheters in trauma patients. *Ann Intern Med*. 2007;147:285-93.
101. Reid G. Biofilms in infectious disease and on medical devices. *Int J Antimicrobial Agents*. 1999;11(3-4):223-6.
102. Jensen PØ, Bjarnsholt T, Phipps R, Rasmussen TB, Calum H, Christoffersen L, et al. Rapid necrotic killing of polymorphonuclear leukocytes is caused by quorum-sensing-controlled production of rhamnolipid by *Pseudomonas aeruginosa* *Microbiology*. 2007;153:1329-38.
103. Høiby N, Pedersen SS. Estimated risk of cross-infection with *Pseudomonas aeruginosa* in Danish Cystic Fibrosis patients. *Acta Paediatr Scand*. 1989;78:395-404.
104. Döring G, Høiby N. Early intervention and prevention of lung disease in cystic fibrosis: a European consensus. *Journal of Cystic Fibrosis*. 2004;3:67-91.
105. Frederiksen B, Pressler T, Hansen A, Koch C, 2006. HNAIP-. Effect of aerosolised rhDNase (Pulmozyme®) on pulmonary colonization in patients with cystic fibrosis. *Acta Paediatrica*. 2006;95:1070-4.
106. Baumann U, Stocklossa C, Greiner W, Graf von der Schulenburg J-M, von der Hardt H. Cost of care and clinical condition in paediatric cystic fibrosis patients. *Journal of Cystic Fibrosis*. 2003;2:84-90.
107. Frederiksen B, Koch C, Høiby N. The changing epidemiology of *Pseudomonas aeruginosa* infection in Danish cystic fibrosis patients, 1974-1995. *Pediatric Pulmonology*. 1999;28:159-66.
108. van Gennip M, Moser C, Christensen L.D., Bjarnsholt T, Calum H, Jensen P.Ø., et al. Augmented effect of early antibiotic therapy in mice with experimental lung infections due to sequentially adapted mucoid strains of *Pseudomonas aeruginosa*. *Journal of Antimicrobial Chemotherapy*. 2009;64:1241-50.
109. Döring G, Pier GB. Vaccines and immunotherapy against *Pseudomonas aeruginosa*. *Vaccine*. 2008;26:1011-24.
110. Döring G, Conway SP, Heijerman HGM, Hodson ME, Høiby N, Smyth A, et al. Antibiotic therapy against *Pseudomonas aeruginosa* in cystic fibrosis: a European consensus. *Eur Resp J*. 2000;16(4):749-67.
111. Høiby N, Flensburg EW, Beck B, Friis B, Jacobsen L, Jacobsen SV. *Pseudomonas aeruginosa* infection in cystic fibrosis. Diagnostic and prognostic significance of *Pseudomonas aeruginosa* precipitins determined by means of crossed immunoelectrophoresis. *Scand J Resp Dis*. 1977;58:65-79.
112. Proesmans M, Balinska-Miskiewicz W, Dupont L, Bossuyt X, Verhaegen J, Høiby N, et al. Evaluating the "Leeds criteria" for *Pseudomonas aeruginosa* infection in a cystic fibrosis centre. *European Respiratory Journal*. 2006;27:937-43.
113. Frederiksen B, Lanng S, Koch C, Høiby N. Improved survival in the Danish cystic fibrosis centre : results of aggressive treatment. *Pediatr Pulmonol*. 1996;21:153-8.
114. Koch C, Hjelt K, Pedersen SS, Jensen ET, Jensen T, Lanng S, et al. Retrospective Clinical Study of Hypersensitivity Reactions to Aztreonam and Six Other beta-Lactam Antibiotics in Cystic Fibrosis Patients Receiving Multiple Treatment Courses. *Rev Infect Dis*. 1991;13(S7):S608-S11.

115. Givskov M. Jamming the command language of bacteria: a new approach
to the control of bacterial infections [Doctoral Thesis]: Danish Technical
University; 2005.
116. Ciofu O, Høiby N. Cystic fibrosis - coping with resistance. . In: Gould I, van
der Meer J, editors. Antibiotic Policies – fighting resistance. New York: Springer
Verlag; 2007. p. 149-74.
117. Bagge N, Schuster M, Hentzer M, Ciofu O, Givskov M, Greenberg EP, et al.
Pseudomonas aeruginosa biofilms exposed to imipenem exhibit changes in
global gene expression and beta-lactamase and alginate production. Antimicrob
Agents Chemother. 2004 Apr;48(4):1175-87.

Table 1. Some general features of biofilm infections in humans compared to acute planktonic infections and superficial colonization/normal flora on skin and mucosal membranes. The bold fonts indicate biofilm specific features.

Features of biofilm infections	Necessary condition for biofilm infections	Sufficient condition for biofilm infections	Also found in acute planktonic infections	Also found in colonization/ normal flora on skin and mucosal membranes
Aggregates of bacteria embedded in a self-produced polymer matrix	Yes	Yes	No	No/Yes
Tolerant to clinical relevant PK/PD dosing of antibiotics in spite of susceptibility of planktonic cells	Yes	Yes	No	No/Yes
Tolerant to both innate and adaptive immune response	Yes	Yes	No	No/Yes - unknown (s-IgA)
Inflammation	Yes	No	Yes	No
Biofilm-specific antigens	No and Yes - seldom – e.g. <i>Pseudomonas aeruginosa</i> alginate	No and Yes - seldom – e.g. <i>Pseudomonas aeruginosa</i> alginate	No	No
Antibody response	Yes - after some weeks	No	Yes - after some weeks	No
Chronic infections	Yes	Yes	No	No
Foreign body associated infections	No	Yes	No but yes the first day of infection	No
Located on surfaces	No	No	Yes	Yes
Localized infection	Yes	No	Yes	Yes
Focus for spreading or local exacerbation	Yes	No	Yes	Yes

Table 2. Natural and pathogenic biofilms on human tissue and foreign bodies.

'Organ A' with normal flora	Connection via foreign bodies	'Organ B' without normal flora
Skin	→	blood, peritoneum, middle ear
Mouth	→	teeth
Pharynx	→	bronchi, lungs,
Duodenum	→	bile tract, pancreas
Urethra	→	urine bladder
Vagina	→	uterus
'Air in operation room'*	→	alloplastic, cerebrospinal shunt
No symptoms	→	pathologi

*most frequently coagulase negative staphylococci, which occur as biofilms on detached epidermal cells.

Antibiotic resistance of bacterial biofilms

Niels Høiby ^{a,b,*}, Thomas Bjarnsholt ^{a,b}, Michael Givskov ^b, Søren Molin ^c,
Oana Ciofu ^b

^a *Department of Clinical Microbiology 9301, Juliane Mariesvej 22,
Rigshospitalet, 2100 Copenhagen, Denmark*

^b *Department of Bacteriology, Institute of International Health, Immunology
and Microbiology, University of Copenhagen, Denmark*

^c *BioCentrum, Danish Technical University, Lyngby, Denmark*

ARTICLE INFO

Article history:

Received 13 December 2009

Accepted 15 December 2009

Keywords:

Biofilm

Antibiotic resistance

Antibiotic tolerance

β-Lactamase

Efflux pumps

Mutators

Pseudomonas aeruginosa

Cystic fibrosis

Foreign-body infections

* Corresponding author. Tel.: +45 3545 7788; fax: +45 3545 6412.

E-mail address: hoiby@hoibyniels.dk (N. Høiby).

Accepted Manuscript

ABSTRACT

A biofilm is a structured consortium of bacteria embedded in a self-produced polymer matrix consisting of polysaccharide, protein and DNA. Bacterial biofilms cause chronic infections because they show increased tolerance to antibiotics and disinfectant chemicals as well as resisting phagocytosis and other components of the body's defence system. The persistence of, for example, staphylococcal infections related to foreign bodies is due to biofilm formation. Likewise, chronic *Pseudomonas aeruginosa* lung infection in cystic fibrosis patients is caused by biofilm-growing mucoid strains. Characteristically, gradients of nutrients and oxygen exist from the top to the bottom of biofilms and these gradients are associated with decreased bacterial metabolic activity and increased doubling times of the bacterial cells; it is these more or less dormant cells that are responsible for some of the tolerance to antibiotics. Biofilm growth is associated with an increased level of mutations as well as with quorum-sensing-regulated mechanisms. Conventional resistance mechanisms such as chromosomal β -lactamase, upregulated efflux pumps and mutations in antibiotic target molecules in bacteria also contribute to the survival of biofilms. Biofilms can be prevented by early aggressive antibiotic prophylaxis or therapy and they can be treated by chronic suppressive therapy. A promising strategy may be the use of enzymes that can dissolve the biofilm matrix (e.g. DNase and alginate lyase) as well as quorum-sensing inhibitors that increase biofilm susceptibility to antibiotics.

1. Introduction

Biofilm-growing bacteria cause chronic infections [1] characterised by persistent inflammation and tissue damage [2]. Chronic infections, including foreign-body infections, are infections that (i) persist despite antibiotic therapy and the innate and adaptive immune and inflammatory responses of the host and (ii) in contrast to colonisation, are characterised by an immune response and persisting pathology (Table 1).

2. Occurrence and architecture of bacterial biofilms

Foreign-body infections are characterised by biofilm growth of bacteria on the outer and/or inner surface of the foreign body (Table 2). Biofilm growth also occurs on natural surfaces such as teeth [3], heart valves (endocarditis) [4], in the lungs of cystic fibrosis (CF) patients causing chronic bronchopneumonia [2], in the middle ear in patients with persistent otitis media [5], in chronic rhinosinusitis [6], in chronic osteomyelitis and prosthetic joint infections [7–9], in intravenous (i.v.) catheters and stents [10] and in chronic wounds [11,12] (Fig. 1). The microbes in biofilms are kept together by a self-produced biopolymer matrix. The matrix contains polysaccharides, proteins and DNA originating from the microbes, and the bacterial consortium can consist of one or more species living in sociomicrobiological way [1,2,14,15]. The matrix is important since it provides structural stability and protection to the biofilm. Development of bacterial biofilms over time has been intensively studied in vitro by confocal scanning laser microscopy employing green fluorescent protein (GFP)-tagged bacteria. This technique has been combined with

advanced in silico image analysis to produce three-dimensional images of the biofilm [16–18]. As an example, *Pseudomonas aeruginosa* produces a mature in vitro biofilm in 5–7 days (Fig. 2).

Development of an in vitro biofilm is initiated by planktonic (freely moving) bacteria that reversibly attach to a surface, which may be covered by a layer of, for example, proteins (a pellicle) [3,20]. At this stage, the bacteria are still susceptible to antibiotics and this is in accordance with the success of perioperative antibiotic prophylaxis, e.g. for alloplastic surgery. The next step is irreversible binding to the surface within the next few hours and multiplication of the bacteria, which form microcolonies on the surface and begin to produce a polymer matrix around the microcolonies [20]. The biofilm grows in thickness (up to 50 μm) and under in vitro conditions mushroom-like or tower-like structures are often observed in the mature biofilm. At that stage, the biofilm shows maximum tolerance (= resistance) to antibiotics.

Subsequently, a stage follows where focal areas of the biofilm dissolve and the liberated bacterial cells can then spread to another location where new biofilms can be formed. This liberation process may be caused by bacteriophage activity within the biofilm [21]. The mature biofilm may contain water-filled channels and thereby resemble primitive, multicellular organisms. Motile bacteria can use type 1V pili to mount or climb a biofilm formed by other bacteria and colonise the top of the biofilm, resembling a hat [17]. Important properties of biofilm-growing bacteria are different from those of planktonic bacteria and this has significant diagnostic and therapeutic consequences. The bacteria appear different in biofilm infections since they

are located close to each other in aggregates surrounded by the self-produced matrix. In clinical specimens (biopsies, pus, sputum), biofilm can often be recognised by light microscopy, although precise identification of all the bacteria within a biofilm can only be done by DNA hybridisation techniques, and identification of the components of the biofilm matrix require specialised staining techniques [2]. Traditional sampling techniques may not be sufficient to culture biofilm-growing bacteria sticking to a surface unless the bacteria are released by ultrasonic pre-treatment [22]. Ordinary culture techniques, however, reveal only the properties of planktonically growing bacteria and, for example, antibiotic susceptibility testing therefore gives misleading results that do not reflect the increased resistance of the bacteria living in biofilms. The minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) of antibiotics to biofilm-growing bacteria may be up to 100–1000-fold higher compared with planktonic bacteria [23–25]. Methods to test biofilm-growing bacteria have therefore been developed, but their clinical relevance with regard to prediction of clinically successful therapy awaits confirmation [24,26,27].

3. Stationary-phase physiology, low oxygen concentration and slow growth

Inspection of environmental as well as in vitro biofilms has revealed that the oxygen concentration may be high at the surface but low in the centre of the biofilm where anaerobic conditions may be present [28]. Likewise, growth, protein synthesis and metabolic activity is stratified in biofilms, i.e. a high level

of activity at the surface and a low level and slow or no growth in the centre, and this is one of the explanations for the reduced susceptibility of biofilms to antibiotics [29,30]. Very slow in situ growth rates of *P. aeruginosa* biofilms have been measured in the sputum of CF patients (average doubling time 2–3 h and the presence of a significant number of cells in stationary growth phase) [31]. Monotherapy with antibiotics such as β -lactams, which are only active against dividing *P. aeruginosa* cells, are therefore not very efficient at eradicating biofilm infections [23].

4. Mutators

The mutation frequency of biofilm-growing bacteria is significantly increased compared with planktonically growing isogenic bacteria [32] and there is increased horizontal gene transmission in biofilms [33]. These physiological conditions may explain why biofilm-growing bacteria easily become multidrug resistant by means of traditional resistance mechanisms against β -lactam antibiotics, aminoglycosides and fluoroquinolones, which are detected by routine susceptibility testing in the clinical microbiology laboratory where planktonic bacterial growth is investigated. Thus, bacterial cells in biofilms may simultaneously produce enzymes that degrade antibiotics, have antibiotic targets of low affinity and overexpress efflux pumps that have a broad range of substrates. Achievement of multiple mutations in a bacterial population size of 10^8 – 10^{10} /mL of sputum, as attained in infection of the CF lung [34], implies the presence of a hypermutable bacterial subpopulation, and the presence of

high percentages of hypermutable *P. aeruginosa* isolates associated with antibiotic resistance has actually been found in CF patients [35,36].

The hypermutable phenotype of CF *P. aeruginosa* isolates is due to alterations in genes of the DNA repair systems of either the mismatch repair system (MMR), which involves *mutS*, *mutL* and *uvrD*, or the DNA oxidative lesions repair system (GO), which involves *mutT*, *mutY* and *mutM* [37,38]. It has been shown that mutations in either of the two systems determine the emergence of antibiotic-resistant isolates, especially due to selection of isolates expressing multidrug efflux pumps [38,39]. Increased production of endogenous reactive oxygen species (ROS) and a deficient antioxidant system [40,41] determine an imbalance between oxidative burden and antioxidant defences leading to oxidative stress in biofilms. This oxidative stress is considered to cause enhanced mutability in biofilms [32,42]. Recent data suggest that microcolony structures, due to endogenous oxidative stress, are specific sites within biofilms for enhanced genetic adaptation and evolutionary change [42]. In addition, Boles and Singh [43] showed that the endogenous oxidative stress in biofilms promotes antibiotic resistance and that addition of antioxidants reduced the occurrence of diversity in biofilms. We have previously shown that oxidative stress is linked to the occurrence of hypermutable *P. aeruginosa* strains in CF patients [36].

In addition to their endogenous oxidative stress, biofilm-growing bacteria in the CF airways are exposed to ROS from activated polymorphonuclear leukocytes (PMNs) [36]. We have recently shown that the hypoxic

environment in CF sputum is due to the consumption of oxygen by PMNs which liberate ROS that can react with the biofilm-embedded bacterial cells [44], thus creating an unique environment with low oxygen tension filled with ROS.

The hypermutability of bacteria in biofilms promotes the emergence of mutations conferring antibiotic resistance, which will be selected for by the repeated antibiotic courses administered in order to maintain the lung function of CF patients. Development of resistance to all classes of antibiotics during chronic lung infection in CF has been documented [45]. Resistance to β -lactam antibiotics occurs due to mutations in the regulatory genes of β -lactamase production leading to the occurrence of isolates with stable or partially stable derepressed production of AmpC β -lactamase [46]. Resistance to ciprofloxacin of CF *P. aeruginosa* isolates was shown to be mediated by mutations in *gyrA* and alterations in two efflux systems (MexCD-OprJ and MexEF-OprN), and resistance to tobramycin was due to overexpression of the MexXY-OprM multidrug efflux pump [47,48]. Resistance to colistin was shown to occur due to mutations in the *pmr* system involved in the lipopolysaccharide (LPS) structure [49].

5. Chromosomal β -lactamase and biofilm matrix components

Overproduction of chromosomally encoded AmpC cephalosporinase is considered the main mechanism of resistance of CF *P. aeruginosa* isolates to β -lactam antibiotics [50]. The most common β -lactamase production

phenotype in CF isolates is the partially derepressed phenotype with high basal levels of β -lactamase that can be further induced to higher levels in the presence of β -lactam antibiotics [46]. The role of this β -lactamase phenotype is especially important for resistance to β -lactam antibiotics acting as strong inducers (carbapenems such as imipenem). However, not all β -lactams are strong inducers and overexpression of the MexAB-OprM efflux pump may, together with β -lactamases, play an important role in resistance to poor inducers (e.g. piperacillin). Totally derepressed β -lactamase production is encountered in 2.5% of clinical CF isolates [46] and is responsible for resistance both to poor- and strong-inducer β -lactam antibiotics, independent of the presence of efflux pump overexpression [51,52]. We have found an insertion sequence (IS 1669) inactivating the *ampD* gene in several resistant clinical *P. aeruginosa* isolates with constitutive high expression of chromosomal β -lactamase [52].

The diffusion barrier in biofilms [53] plays a role in biofilm resistance of *P. aeruginosa* that overproduce β -lactamase owing to the presence in the biofilm matrix of β -lactamases that will hydrolyse the β -lactam antibiotics before reaching the bacterial cells [54,55]. Giwercman et al. [56] showed that imipenem and piperacillin were able to induce β -lactamase production in *P. aeruginosa* biofilms. Nichols et al. [57] predicted from mathematical models that the biofilm would not afford protection against diffusion of β -lactam antibiotics into the bacteria embedded in the biofilm as long as the level of chromosomal β -lactamase was low. However, bacteria expressing a high

level of chromosomal β -lactamase growing in biofilms would be exposed to a reduced concentration of β -lactam antibiotics owing to accumulation of the enzyme in the polysaccharide matrix. The extracellular β -lactamase would inactivate the antibiotic as it penetrates, thereby protecting the deeper-lying cells.

The source of β -lactamase in biofilms has been considered to be from a layer of lysed bacteria owing to exposure to an antibiotic, with release of defensive enzymes into the extracellular space. We have shown that the source of β -lactamase in biofilm may also be the membrane vesicles containing β -lactamase liberated by resistant *P. aeruginosa* bacteria [55,58] and we have shown that a high level of free chromosomal β -lactamase is present in CF sputum [58]. We have also shown that strong inducers such as imipenem will induce the β -lactamase through all the bacterial layers, whilst poorer inducers such as ceftazidime will influence just the superficial layers of the biofilm, probably due to inactivation of the antibiotic by β -lactamase [58–60] (Fig. 3).

The protective role played by β -lactamases in impairing the penetration of β -lactams in the biofilm can be seen in Fig. 3A. Treatment with ceftazidime of a biofilm formed by a *P. aeruginosa* CF strain with stable derepressed levels of β -lactamase due to an insertion sequence in *ampD* (*P. aeruginosa ampD⁻*) killed very few bacterial cells (dead bacteria in red) (Fig. 3B) in contrast to the complemented strain with a low level of β -lactamase (*P. aeruginosa ampD⁺*) (Fig. 3C). However, addition of aztreonam improved the efficacy of ceftazidime treatment of the biofilm (Fig. 3D), probably because aztreonam

acts as a β -lactamase inhibitor [58]. In addition, meropenem, a β -lactamase-stable β -lactam, showed good in vitro efficacy in the treatment of *P. aeruginosa* biofilms [24,60]. Treatment with ceftazidime of a biofilm formed by the same strain expressing basal levels of β -lactamase due to complementation with the wild-type *ampD* (*P. aeruginosa ampD*⁺) led to eradication of the biofilm (Fig. 3C).

The matrix of the biofilm may also be part of the resistance mechanism to antibiotics since, for example, sub-MIC concentrations of β -lactam antibiotics induce increased alginate synthesis in *P. aeruginosa* biofilms (Fig. 4) [62,63] and also enhance the biofilm matrix of some slime-producing coagulase-negative staphylococci [64,65]. Originally, it was thought that tolerance of biofilms to aminoglycosides was the result of transport limitation due to the binding of these positively charged antibiotics to the negatively charged exopolysaccharide matrix, but the repeated dosing of antibiotics during therapy probably leads to saturation of the binding sites [66–68]. As discussed previously, oxygen limitation and the metabolic rate are probably more important factors contributing to the tolerance of biofilms to aminoglycosides and ciprofloxacin [69]. However, in the respiratory zone of the CF lung with poor access to aminoglycoside aerosols, where the antibiotic concentration is low, delayed penetration of aminoglycosides through thick biofilms may play a role in the tolerance of biofilms to these aminoglycosides [2]. Recently, it has been shown that administration of DNase and alginate lyase enhanced the activity of tobramycin in biofilms by dissolving the biofilm matrix [70].

6. Tolerance, adaptive resistance and efflux pumps

Colistin is only antimicrobial active against the non-dividing central part of *P. aeruginosa* biofilms in vitro (Fig. 5A), whereas the superficial, metabolically active part of the biofilm becomes tolerant due to upregulation of the PmrA-PmrB two-component regulatory system involved in adaptive resistance to cationic peptides leading to addition of aminoarabinose to lipid A of LPS [24,71,72]. Since the metabolically active surface layer of the biofilm is susceptible to ciprofloxacin (Fig. 5B), in contrast to the dormant central part of the biofilm, combination therapy with ciprofloxacin and colistin was able to kill all cells in the biofilm in vitro (Fig. 5C) [71]. Clinical efficacy has been demonstrated with this combination therapy for the early eradication treatment of *P. aeruginosa* in CF patients [73].

Tolerance of biofilms to tobramycin is also mediated by low metabolic activity, but the high cell density that results in accumulation of extracellular signalling molecules is probably important, as it has been shown that tolerance to tobramycin of *P. aeruginosa* strain PAO1 biofilm is quorum-sensing-mediated (Fig. 6) [74] (see below). In addition, a non-specific mechanism for the tolerance of the metabolically active cells to colistin was shown to be upregulation of the MexAB-OprM efflux pump [71]. Furthermore, increased efflux pump activity due to mutations has been shown to be a major resistance mechanism against aminoglycoside antibiotics and fluoroquinolones in *P. aeruginosa* from CF patients [47,48].

7. High cell density and quorum sensing (QS)

Bacteria communicate by means of synthesising and reacting on signal molecules [75–78]. The term QS indicates that this system allows bacteria to sense when a critical number (concentration) of bacteria are present in a limited space in the environment and respond by activating certain genes that then produce, for example, virulence factors such as enzymes or toxins. The QS molecules are small peptides in many Gram-positive bacteria, whereas the most well described QS molecules in Gram-negative bacteria are *N*-acyl-L-homoserine lactones [78]. For *P. aeruginosa*, QS regulates the production of virulence factors such as extracellular enzymes and cellular lysins (e.g. rhamnolipid), which are important for the pathogenesis of infections where it functions as a protective shield against phagocytes [79–81]. QS may also influence the development of the biofilm [82] and QS has been shown to determine the tolerance of *P. aeruginosa* biofilms to antibiotic therapy and to the innate inflammatory response dominated by PMNs [74]. The connection between QS and biofilms has been named sociomicrobiology [83].

8. Quorum-sensing inhibitors (QSIs)

Much of our knowledge about QS originates from experiments with QS knock-out mutants and from the use of naturally occurring and artificially synthesised QSI compounds [84,85]. Screening for QSIs in nature has identified many QSI compounds [86]. These naturally occurring QSI compounds can be synthesised and their structure modified and used to inhibit QS in vivo in experimental animal infections [84]. Since it has been shown that bacteria

used for experimental animal biofilm infections actually communicate in vivo [87] and also in, for example, CF patients with chronic *P. aeruginosa* lung infection, QSIs may be used to treat these infections [88]. Interestingly, some macrolide antibiotics such as azithromycin [89], but also other antibiotics such as ceftazidime and ciprofloxacin [90], inhibit QS in *P. aeruginosa* at sub-MIC concentrations, leading to inhibition of the virulence of these bacteria even though they cannot inhibit their growth at concentrations obtainable in vivo. Controlled clinical trials using azithromycin to treat chronic *P. aeruginosa* lung infection both in CF children and adults have shown significant improvement of their lung function [91–93]. Most CF patients with chronic *P. aeruginosa* lung infections are therefore now treated continuously with azithromycin [94]. An expected side effect, however, has been the development of resistance to macrolides in other pathogenic bacteria such as *Staphylococcus aureus* in CF patients [95]. It would therefore be desirable to develop QSIs without conventional growth-inhibiting or bactericidal activity [78]. One example of such a naturally occurring QSI is found in garlic extract, which in vitro and in vivo has been able to render otherwise resistant *P. aeruginosa* biofilms susceptible to antibiotic therapy and to PMN activity and the consequence is eradication of the biofilm both with regard to antibiotic therapy and PMN activity which dominates the inflammatory response in CF patients (Fig. 7) [96]. According to current knowledge, QSI resistance can only occur due to mutations, which render the QS-deficient bacteria unable to produce virulence factors [97], i.e. the bacteria become non-virulent, similar to the result of QSI therapy. If this holds true, then resistance problems against conventional antibiotics as we face today will not be a clinical problem.

Foreign-body infections constitute a steadily increasing medical problem and comprise, for example, infections of i.v. catheters, intrauterine catheters, nasolaryngeal tubes, stents (Table 2; Fig. 1A), alloplastic materials, hydrocephalus shunts and artificial hearts [98]. If foreign bodies become colonised with biofilm-forming bacteria the result is most often chronic inflammation around the foreign body that either has to be replaced or treated with sometimes life-long antibiotic suppressive therapy, although early therapy may sometimes lead to eradication of the condition. Antibiotic-coated foreign bodies, e.g. catheters and vascular prostheses, have been introduced to prevent biofilm formation and they are quite efficient [99,100]. However, it is desirable to develop other compounds owing to the risk of development of bacterial resistance, and QSIs are strong candidates. QSIs have been shown to improve synergistically the weak effects of antibiotics and PMNs on biofilm-growing bacteria in vitro and in vivo in animal experiments, leading to elimination of biofilms (Fig. 7) [79]. These results have led to further development of QSIs as pharmaceutical compounds for patients who are subject to implantation of foreign bodies.

9. Prophylaxis and treatment of *Pseudomonas aeruginosa* biofilms in cystic fibrosis lungs: perspectives for other biofilm infections?

The currently used methods for preventing chronic *P. aeruginosa* biofilms in CF lungs are (i) prevention of cross-infection from other already chronically

infected CF patients by isolation techniques and hygienic measures [101], (ii) early aggressive eradication therapy of intermittent colonisation by means of oral ciprofloxacin and nebulised colistin for 3 weeks or, even better, for 3 months or by using nebulised tobramycin as monotherapy [102] and (iii) daily nebulised DNase (Pulmozyme[®]) [103]. These three methods, which are combined in most CF centres, are successful and cost efficient and have completely changed the epidemiology of chronic *P. aeruginosa* lung infection in CF patients from being very common in CF children to being predominantly a problem for adult patients, and no problems of resistance to the antibiotics have been recorded [73,104,105]. Early aggressive eradication therapy has also been shown to be superior in an animal model of *P. aeruginosa* infection in CF [106]. Furthermore, although vaccines against *P. aeruginosa* have been developed and have undergone clinical trials, they have not been further developed owing to the success of early aggressive eradication therapy [107].

The recommended method for treatment of chronic *P. aeruginosa* biofilm infection is chronic suppressive antibiotic therapy [2,108] (Fig. 8), which is started when the chronic infection is diagnosed (continuous colonisation at the monthly bacteriological examination for 6 months and/or increased level of antibodies against *P. aeruginosa*) [109,110]. Chronic suppressive therapy consists of daily nebulised colistin or tobramycin for the rest of the patient's life combined with either regular 2-week courses every 3 months of i.v. antipseudomonal antibiotics (combination therapy of two antibiotics: tobramycin or colistin + ceftazidime, or piperacillin/tazobactam, or carbapenem, or aztreonam, or ciprofloxacin) or ad hoc i.v. therapy when

clinical deterioration occurs [108]. Additionally, DNase is inhaled every day to reduce the viscosity of the DNA-containing sputum [103], and daily oral azithromycin is given to patients as mentioned above [94]. Chronic suppressive therapy (maintenance therapy) has successfully been able to maintain pulmonary function or slow the decline in pulmonary function and prolong the life of patients for many years [111]. The side effects of maintenance therapy is a high level of conventional resistance mechanisms in the persisting strains and a high level of allergy to β -lactam antibiotics [45,112]. Similar principles, e.g. (a) systemic or local prophylactic use of antibiotics to prevent biofilm formation, (b) early aggressive eradication therapy to eradicate planktonic growth or early biofilm formation and (c) chronic suppressive antibiotic therapy to maintain the function of an inserted foreign body, are gradually being introduced in other areas of biofilm infection [8,9,99]. Possibly other approaches taking advantage of the use of QSIs may further improve the management of biofilm infections.

Funding

None.

Conflicts of interest

None declared.

Ethical approval

Not required.

References

- [1] Costerton W, Veeh R, Shirtliff M, Pasmore M, Post C, Ehrlich G. The application of biofilm science to the study and control of chronic bacterial infections. *J Clin Invest* 2003;112:1466–77.
- [2] Bjarnsholt T, Jensen PØ, Fiandaca MJ, Pedersen J, Hansen CR, Andersen CB, et al. *Pseudomonas aeruginosa* biofilms in the respiratory tract of cystic fibrosis patients. *Pediatr Pulmonol* 2009;44:547–58.
- [3] Kolenbrander PE, Palmer RJ Jr. Human oral bacterial biofilms. In: Ghannoum MA, O'Toole GA, editors. *Microbial biofilms*. Washington, DC: ASM Press; 2004.
- [4] Høiby N, Döring G, Schiøtz PO. The role of immune complexes in the pathogenesis of bacterial infections. *Annu Rev Microbiol* 1986;40:29–53.
- [5] Hall-Stoodley L, Hu FZ, Gieseke A, Nistico A, Nguyen L, Hayes J, et al. Direct detection of bacterial biofilms on the middle-ear mucosa of children with chronic otitis media. *JAMA* 2006;296:202–11.
- [6] Sanderson AR, Leid JG, Hunsaker D. Bacterial biofilms on the sinus mucosa of human subjects with chronic rhinosinusitis. *Laryngoscope* 2006;116:1121–6.
- [7] Gristina AG, Oga M, Webb LX, Hobgood CD. Adherent bacterial colonization in the pathogenesis of osteomyelitis. *Science* 1985;228:990–3.
- [8] Trampuz A, Zimmerli W. Prosthetic joint infections: update in diagnosis and treatment. *Swiss Med Wkly* 2005;135:243–51.
- [9] Del Pozo JL, Patel R. Infection associated with prosthetic joints. *N Engl J Med* 2009;361:787–94.

- [10] Taconelli E, Smith G, Hieke K, Lafuma A, Bastide P. Epidemiology, medical outcomes and costs of catheter-related bloodstream infection in intensive care units of four European countries: literature- and registry-based estimates. *J Hosp Infect* 2009;72:97–103.
- [11] Kirketerp-Møller K, Jensen PØ, Fazli M, Madsen KG, Pedersen J, Moser C, et al. Distribution, and ecology of bacteria in chronic wounds. *J Clin Microbiol* 2008;46:2717–22.
- [12] Bjarnsholt T, Kirketerp-Møller K, Jensen PØ, Madsen KG, Phipps R, Krogfelt K, et al. Why chronic wounds won't heal: a novel hypothesis. *Wound Repair Regen* 2008;16:2–10.
- [13] Høiby N, Johansen HK, Ciofu O, Jensen PØ, Bjarnsholt T, Givskov M. Foreign body infections—biofilms and quorum sensing [in Danish]. *Ugeskr Laeger* 2007;169:4163–6.
- [14] Wingender J, Strathmann M, Rode A, Leis A, Flemming HC. Isolation and biochemical characterization of extracellular polymeric substances from *Pseudomonas aeruginosa*. *Methods Enzymol* 2001;336:302–14.
- [15] Whitchurch CB, Tolker-Nielsen T, Ragas PC, Mattick JS. Extracellular DNA required for bacterial biofilm formation. *Science* 2002;295:1487.
- [16] Sternberg C, Christensen BB, Johansen T, Nielsen AT, Andersen JB, Givskov M, et al. Distribution of bacterial growth activity in flow-chamber biofilms. *Appl Environ Microbiol* 1999;65:4108–17.
- [17] Klausen M, Heydorn A, Ragas P, Lambertsen L, Aaes-Jorgensen A, Molin S, et al. Biofilm formation by *Pseudomonas aeruginosa* wild type, flagella and type IV pili mutants. *Mol Microbiol* 2003;48:1511–24.

- [18] Heydorn A, Ersboll B, Kato J, Hentzer M, Parsek MR, Tolker-Nielsen T, et al. Statistical analysis of *Pseudomonas aeruginosa* biofilm development: impact of mutations in genes involved in twitching motility, cell-to-cell signaling, and stationary-phase sigma factor expression. *Appl Environ Microbiol* 2002;68:2008–17.
- [19] Givskov M. *Jamming the command language of bacteria: a new approach to the control of bacterial infections* [Doctoral thesis]. Danish Technical University; 2005.
- [20] Marshall KC. Biofilms: an overview of bacterial adhesion, activity, and control at surfaces. *ASM News* 1992;58:202–7.
- [21] Webb JS, Thompson LS, James S, Charlton T, Tolker-Nielsen T, Koch B, et al. Cell death in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* 2003;185:4585–92.
- [22] Kobayashi N, Bauer TW, Tuohy MJ, Fujishiro T, Procop GW. Brief ultrasonication improves detection of biofilm-formative bacteria around a metal implant. *Clin Orthop Relat Res* 2007;475:210–3.
- [23] Anwar H, Costerton JW. Enhanced activity of combination of tobramycin and piperacillin for eradication of sessile biofilm cells of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1990;34:1666–71.
- [24] Moskowitz SM, Foster JM, Emerson J, Burns JL. Clinically feasible biofilm susceptibility assay for isolates of *Pseudomonas aeruginosa* from patients with cystic fibrosis. *J Clin Microbiol* 2004;42:1915–22.

- [25] Bjarnsholt T, Kirketerp-Møller K, Kristiansen S, Phipps R, Nielsen AK, Jensen PØ, et al. Silver against *Pseudomonas aeruginosa* biofilms. *APMIS* 2007;115:921–8.
- [26] Anwar H, Dasgupta MK, Costerton JW. Testing the susceptibility of bacteria in biofilms to antibacterial agents. *Antimicrob Agents Chemother* 1990;34:2043–6.
- [27] Domingue G, Ellis B, Dasgupta M, Costerton JW. Testing antimicrobial susceptibilities of adherent bacteria by a method that incorporates guidelines of the National Committee for Clinical Laboratory Standards. *J Clin Microbiol* 1994;32:2564–8.
- [28] Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappinscott HM. Microbial biofilms. *Annu Rev Microbiol* 1995;49:711–45.
- [29] Werner E, Roe F, Bugnicourt A, Franklin MJ, Heydorn A, Molin S, et al. Stratified growth in *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol* 2004;70:6188–96.
- [30] Keren I, Kaldalu N, Spoering A, Wang YP, Lewis K. Persister cells and tolerance to antimicrobials. *FEMS Microbiol Lett* 2004;230:13–8.
- [31] Yang L, Haagensen JAJ, Jelsbak L, Johansen HK, Sternberg C, Høiby N, et al. In situ growth rates and biofilm development of *Pseudomonas aeruginosa* populations in chronic lung infection. *J Bacteriol* 2008;190:2767–76.
- [32] Driffield K, Miller K, Bostock M, O'Neill AJ, Chopra I. Increased mutability of *Pseudomonas aeruginosa* in biofilms. *J Antimicrob Chemother* 2008;61:1053–6.

- [33] Molin S, Tolker-Nielsen T. Gene transfer occurs with enhanced efficiency in biofilms and induces enhanced stabilisation of the biofilm structure. *Curr Opin Biotechnol* 2003;14:255–61.
- [34] Høiby N, Johansen HK, Moser C, Song ZJ, Ciofu O, Kharazmi A. *Pseudomonas aeruginosa* and the biofilm mode of growth. *Microbes Infect* 2001;3:1–13.
- [35] Oliver A, Canton R, Campo P, Baquero F, Blazquez J. High frequency of hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science* 2000;288:1251–3.
- [36] Ciofu O, Riis B, Pressler T, Poulsen HE, Høiby N. Occurrence of hypermutable *Pseudomonas aeruginosa* in cystic fibrosis patients is associated with the oxidative stress caused by chronic lung inflammation. *Antimicrob Agents Chemother* 2005;49:2276–82.
- [37] Oliver A, Sanchez JM, Blazquez J. Characterization of the GO system of *Pseudomonas aeruginosa*. *FEMS Microbiol Lett* 2002;217:31–5.
- [38] Mandsberg LF, Ciofu O, Kirkby N, Christiansen LE, Poulsen HE, Høiby N. Antibiotic resistance in *Pseudomonas aeruginosa* strains with increased mutation frequency due to inactivation of the DNA oxidative repair system. *Antimicrob Agents Chemother* 2009;53:2483–91.
- [39] Macia MD, Blanquer D, Togores B, Sauleda J, Perez JL, Oliver A. Hypermutation is a key factor in development of multiple-antimicrobial resistance in *Pseudomonas aeruginosa* strains causing chronic lung infections. *Antimicrob Agents Chemother* 2005;49:3382–6.
- [40] Hassett DJ, Ma JF, Elkins JG, McDermott TR, Ochsner UA, West SEH, et al. Quorum sensing in *Pseudomonas aeruginosa* controls expression of

- catalase and superoxide dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. *Mol Microbiol* 1999;34:1082–93.
- [41] Mai-Prochnow A, Lucas-Elio P, Egan S, Thomas T, Webb J, Sanchez-Amat A, et al. Hydrogen peroxide linked to lysine oxidase activity facilitates biofilm differentiation and dispersal in several Gram-negative bacteria. *J Bacteriol* 2008;190:5493–501.
- [42] Conibear TC, Collins SL, Webb JS. Role of mutation in *Pseudomonas aeruginosa* biofilm development. *PLoS One* 2009;14:e6289.
- [43] Boles BR, Singh PK. Endogenous oxidative stress produces diversity and adaptability in biofilm communities. *Proc Natl Acad Sci USA* 2008;105:12503–8.
- [44] Kolpen M, Hansen CR, Bjarnsholt T, Moser C, Christensen LD, van Gennip M, et al. Polymorphonuclear leucocytes consume oxygen in sputum from chronic *Pseudomonas aeruginosa* pneumonia in cystic fibrosis. *Thorax* 2010;65:57–62.
- [45] Ciofu O, Giwercman B, Pedersen SS, Hoiby N. Development of antibiotic resistance in *Pseudomonas aeruginosa* during two decades of antipseudomonal treatment at the Danish CF Center. *APMIS* 1994;102:674–80.
- [46] Ciofu O. *Pseudomonas aeruginosa* chromosomal β -lactamase in patients with cystic fibrosis and chronic lung infection—mechanism of antibiotic resistance and target of the humoral immune response. *APMIS* 2003;111:4–47.
- [47] Jalal S, Ciofu O, Høiby N, Gotoh N, Wretling B. Molecular mechanisms of fluoroquinolone resistance in *Pseudomonas aeruginosa* isolates from

- cystic fibrosis patients. *Antimicrob Agents Chemother* 2000;44:710–2.
- Erratum in: *Antimicrob Agents Chemother* 2000;44:1410.
- [48] Islam S, Oh H, Jalal S, Ciofu O, Høiby N, Wretling B. Chromosomal mechanisms of aminoglycoside resistance in *Pseudomonas aeruginosa* isolates from cystic fibrosis patients. *Clin Microbiol Infect* 2009;15:60–6.
- [49] Johansen HK, Moskowitz SM, Ciofu O, Pressler T, Høiby N. Spread of colistin-resistant non-mucoid *Pseudomonas aeruginosa* among chronically infected Danish cystic fibrosis patients. *J Cyst Fibros* 2008;7:391–7.
- [50] Giwercman B, Lambert PA, Rosdahl VT, Shand GH, Hoiby N. Rapid emergence of resistance in *Pseudomonas aeruginosa* in cystic fibrosis patients due to in-vivo selection of stable partially derepressed β -lactamase producing strains. *J Antimicrob Chemother* 1990;26:247–59.
- [51] Nakae T, Nakajima A, Ono T, Saito K, Yoneyama H. Resistance to β -lactam antibiotics in *Pseudomonas aeruginosa* due to interplay between the MexAB-OprM efflux pump and β -lactamase. *Antimicrob Agents Chemother* 1999;43:1301–3.
- [52] Bagge N, Ciofu O, Hentzer M, Campbell JIA, Givskov M, Hoiby N. Constitutive high expression of chromosomal β -lactamase in *Pseudomonas aeruginosa* caused by a new insertion sequence (IS 1669) located in *ampD*. *Antimicrob Agents Chemother* 2002;46:3406–11.
- [53] de Beer D, Stoodley P, Lewandowski Z. Measurement of local diffusion coefficients in biofilms by microinjection and confocal microscopy. *Biotechnol Bioeng* 1997;53:151–8.
- [54] Dibdin GH, Assinder SJ, Nichols WW, Lambert PA. Mathematical model of β -lactam penetration into a biofilm of *Pseudomonas aeruginosa*

while undergoing simultaneous inactivation by released β -lactamases. J Antimicrob Chemother 1996;38:757–69.

- [55] Ciofu O, Beveridge TJ, Kadurugamuwa J, Walther-Rasmussen J, Hoiby N. Chromosomal β -lactamase is packaged into membrane vesicles and secreted from *Pseudomonas aeruginosa*. J Antimicrob Chemother 2000;45:9–13.
- [56] Giwercman B, Jensen ET, Hoiby N, Kharazmi A, Costerton JW. Induction of β -lactamase production in *Pseudomonas aeruginosa* biofilm. Antimicrob Agents Chemother 1991;35:1008–10.
- [57] Nichols WW, Evans MJ, Slack MP, Walmsley HL. The penetration of antibiotics into aggregates of mucoid and non-mucoid *Pseudomonas aeruginosa*. J Gen Microbiol 1989;135:1291–303.
- [58] Giwercman B, Meyer C, Lambert PA, Reinert C, Hoiby N. High-level β -lactamase activity in sputum samples from cystic fibrosis patients during antipseudomonal treatment. Antimicrob Agents Chemother 1992;36:71–6.
- [59] Bagge N, Hentzer M, Andersen JB, Ciofu O, Givskov M, Hoiby N. Dynamics and spatial distribution of β -lactamase expression in *Pseudomonas aeruginosa* biofilms. Antimicrob Agents Chemother 2004;48:1168–74.
- [60] Hill D, Rose B, Pajkos A, Robinson M, Bye P, Bell S, et al. Antibiotic susceptibilities of *Pseudomonas aeruginosa* isolates derived from patients with cystic fibrosis under aerobic, anaerobic, and biofilm conditions. J Clin Microbiol 2005;43:5085–90.

- [61] Ciofu O, Høiby N. Cystic fibrosis—coping with resistance. In: Gould I, van der Meer J, editors. *Antibiotic policies—fighting resistance*. New York, NY: Springer Verlag; 2007. p. 149–74.
- [62] Bagge N, Schuster M, Hentzer M, Ciofu O, Givskov M, Greenberg EP, et al. *Pseudomonas aeruginosa* biofilms exposed to imipenem exhibit changes in global gene expression and β -lactamase and alginate production. *Antimicrob Agents Chemother* 2004;48:1175–87.
- [63] Wood LF, Leech AJ, Ohman DE. Cell wall-inhibitory antibiotics activate the alginate biosynthesis operon in *Pseudomonas aeruginosa*: roles of sigma (AlgT) and the AlgW and Prc proteases. *Mol Microbiol* 2006;62:412–26.
- [64] Dunne WM. Effects of subinhibitory concentrations of vancomycin or cefamandole on biofilm production by coagulase-negative staphylococci. *Antimicrob Agents Chemother* 1990;34:390–3.
- [65] Permin H, Koch C, Høiby N, Christensen HO, Møller AF, Møller S. Ceftazidime treatment of chronic *Pseudomonas aeruginosa* respiratory tract infection in cystic fibrosis. *J Antimicrob Chemother* 1983;12(Suppl A):313–23.
- [66] Anwar H, Strap JL, Costerton JW. Establishment of aging biofilms: possible mechanism of bacterial resistance to antimicrobial therapy. *Antimicrob Agents Chemother* 1992;36:1347–51.
- [67] Levy J, Smith AL, Kenny MA, Ramsey B, Schoenknecht FD. Bioactivity of gentamicin in purulent sputum from patients with cystic fibrosis or bronchiectasis: comparison with activity in serum. *J Infect Dis* 1983;148:1069–76.

- [68] Hunt BE, Weber A, Berger A, Ramsey B, Smith AL. Macromolecular mechanisms of sputum inhibition of tobramycin activity. *Antimicrob Agents Chemother* 1995;39:34–9.
- [69] Walters MC, Roe F, Bugnicourt A, Franklin MJ, Stewart PS. Contributions of antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of *Pseudomonas aeruginosa* biofilms to ciprofloxacin and tobramycin. *Antimicrob Agents Chemother* 2003;47:317–23.
- [70] Alipour M, Suntres ZE, Omri A. Importance of DNase and alginate lyase for enhancing free and liposome encapsulated aminoglycoside activity against *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2009;64:317–25.
- [71] Pamp SJ, Gjermansen M, Johansen HK, Tolker-Nielsen T. Tolerance to the antimicrobial peptide colistin in *Pseudomonas aeruginosa* biofilms is linked to metabolically active cells, and depends on the *prm* and *mexAB-oprM* genes. *Mol Microbiol* 2008;68:223–40.
- [72] Haagensen J, Klausen M, Ernst RK, Miller SI, Folkesson A, Tolker-Nielsen T, et al. Differentiation and distribution of colistin- and sodium dodecyl sulfate-tolerant cells in *Pseudomonas aeruginosa* biofilms. *J Bacteriol* 2007;189:28–37.
- [73] Hansen CR, Pressler T, Høiby N. Early aggressive eradication therapy for intermittent *Pseudomonas aeruginosa* airway colonization in cystic fibrosis patients: 15 years experience. *J Cyst Fibros* 2008;7:523–30.
- [74] Bjarnsholt T, Jensen P-Ø, Burmølle M, Hentzer M, Haagensen JAJ, Hougen HP, et al. *Pseudomonas aeruginosa* tolerance to tobramycin,

hydrogen peroxide and polymorphonuclear leukocytes is quorum-sensing dependent. *Microbiology* 2005;151:373–83.

- [75] Greenberg EP. Quorum sensing in Gram-negative bacteria. *ASM News* 1997;63:371–7.
- [76] Lindum PW, Anthoni U, Christophersen C, Eberl L, Molin S, Givskov M. *N*-acyl-L-homoserine lactone autoinducers control production of an extracellular lipopeptide biosurfactant required for swarming motility of *Serratia liquefaciens* MG1. *J Bacteriol* 1998;180:6384–8.
- [77] Williams P, Camara M, Hardman A, Swift S, Milton D, Hope VJ, et al. Quorum sensing and the population-dependent control of virulence. *Philos Trans R Soc Lond B Biol Sci* 2000;355:667–80.
- [78] Bjarnsholt T, Givskov M. The role of quorum sensing in the pathogenicity of the cunning aggressor *Pseudomonas aeruginosa*. *Anal Bioanal Chem* 2007;387:409–14.
- [79] Jensen PØ, Bjarnsholt T, Phipps R, Rasmussen TB, Calum H, Christoffersen L, et al. Rapid necrotic killing of polymorphonuclear leukocytes is caused by quorum-sensing-controlled production of rhamnolipid by *Pseudomonas aeruginosa*. *Microbiology* 2007;153:1329–38.
- [80] Van Gennip M, Christensen LD, Alhede M, Phipps R, Jensen PØ, Christophersen L, et al. Inactivation of the *rhIA* gene in *Pseudomonas aeruginosa* prevents rhamnolipid production, disabling the protection against polymorphonuclear leukocytes. *APMIS* 2009;117:537–46.
- [81] Alhede M, Bjarnsholt T, Jensen PØ, Phipps RK, Moser C, Christophersen L, et al. *Pseudomonas aeruginosa* recognizes and

responds aggressively to the presence of polymorphonuclear leukocytes.

Microbiology 2009;155:3500–8.

- [82] Davies DG, Parsek MR, Pearson JP, Iglewski BH, Costerton JW, Greenberg EP. The involvement of cell-to-cell signals in the development of a bacterial biofilm. Science 1998;280:295–8.
- [83] Parsek MR, Greenberg EP. Sociomicrobiology: the connections between quorum sensing and biofilms. Trends Microbiol 2005;13:27–33.
- [84] Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, et al. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. EMBO J 2003;22:3803–15.
- [85] Givskov M, de Nys R, Manefield M, Gram L, Maximilien R, Eberl L, et al. Eukaryotic interference with homoserine lactone-mediated prokaryotic signalling. J Bacteriol 1996;178:6618–22.
- [86] Rasmussen TB, Bjarnsholt T, Skindersoe ME, Hentzer M, Kristoffersen P, Kte M, et al. Screening for quorum-sensing inhibitors (QSI) by use of a novel genetic system, the QSI selector. J Bacteriol 2005;187:1799–814.
- [87] Wu H, Song ZJ, Hentzer M, Andersen JB, Heydorn A, Mathee K, et al. Detection of *N*-acylhomoserine lactones in lung tissues of mice infected with *Pseudomonas aeruginosa*. Microbiology 2000;146:2481–93.
- [88] Hentzer M, Givskov M. Pharmacological inhibition of quorum sensing for treatment of chronic bacterial infections. J Clin Invest 2003;112:1300–7.
- [89] Hoffmann N, Lee B, Hentzer M, Rasmussen TB, Song Z, Johansen HK, et al. Azithromycin blocks quorum sensing and alginate polymer formation and increases the sensitivity to serum and stationary-growth-

phase killing of *Pseudomonas aeruginosa* and attenuates chronic *P. aeruginosa* lung infection in *Cftr*^{-/-} mice. *Antimicrob Agents Chemother* 2007;51:3677–87.

- [90] Skindersø ME, Alhede M, Phipps R, Yang L, Jensen PO, Rasmussen TB, et al. Effects of antibiotics on quorum sensing in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2008;52:3548–663.
- [91] Wolter J, Seeney S, Bell S, Bowler S, Masel P, McCormack J. Effect of long-term treatment with azithromycin on disease parameters in cystic fibrosis: a randomized trial. *Thorax* 2002;57:212–6.
- [92] Equi A, Balfour-Lynn IM, Bush A, Rosenthal M. Long term azithromycin in children with cystic fibrosis: a randomised, placebo-controlled crossover trial. *Lancet* 2002;360:978–84.
- [93] Saiman L, Marshall BC, Mayer-Hamblett N, Burns JL, Quittner AL, Cibene DA, et al. Azithromycin in patients with cystic fibrosis chronically infected with *Pseudomonas aeruginosa*: a randomized controlled trial. *JAMA* 2003;290:1749–56.
- [94] Hansen CR, Pressler T, Koch C, Høiby N. Long-term azithromycin treatment of cystic fibrosis patients with chronic *Pseudomonas aeruginosa* infection; an observational cohort study. *J Cyst Fibros* 2005;4:35–40.
- [95] Tramper-Stranders GA, Wolfs TFW, Fleer A, Kimpen JLL, van der Ent CK. Maintenance azithromycin treatment in pediatric patients with cystic fibrosis: long-term outcomes related to macrolide resistance and pulmonary function. *Pediatr Infect Dis J* 2007;26:8–12.
- [96] Bjarnsholt T, Jensen PØ, Rasmussen TB, Christophersen L, Calun H, Hentzer M, et al. Garlic blocks quorum sensing and promotes rapid

clearing of pulmonary *Pseudomonas aeruginosa* infections. Microbiology 2005;151:3873–80.

- [97] Lee B, Haagensen JAJ, Ciofu O, Andersen JB, Høiby N, Molin S. Heterogeneity of biofilms formed by non-mucoid *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. J Clin Microbiol 2005;43:5247–55.
- [98] Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. Science 1999;284:1318–22.
- [99] Stensballe J, Tvede M, Loorns D, Lippert FK, Dahl B, Tønnesen E, et al. Infection risk with nitrofurazone-impregnated urinary catheters in trauma patients. Ann Intern Med 2007;147:285–93.
- [100] Reid G. Biofilms in infectious disease and on medical devices. Int J Antimicrob Agents 1999;11:223–6.
- [101] Høiby N, Pedersen SS. Estimated risk of cross-infection with *Pseudomonas aeruginosa* in Danish cystic fibrosis patients. Acta Paediatr Scand 1989;78:395–404.
- [102] Döring G, Høiby N. Early intervention and prevention of lung disease in cystic fibrosis: a European consensus. J Cyst Fibros 2004;3:67–91.
- [103] Frederiksen B, Pressler T, Hansen A, Koch C, Høiby N. Effect of aerosolised rhDNase (Pulmozyme®) on pulmonary colonization in patients with cystic fibrosis. Acta Paediatr 2006;95:1070–4.
- [104] Baumann U, Stocklossa C, Greiner W, von der Schulenburg JM, von der Hardt H. Cost of care and clinical condition in paediatric cystic fibrosis patients. J Cyst Fibros 2003;2:84–90.

- [105] Frederiksen B, Koch C, Høiby N. The changing epidemiology of *Pseudomonas aeruginosa* infection in Danish cystic fibrosis patients, 1974–1995. *Pediatr Pulmonol* 1999;28:159–66.
- [106] van Gennip M, Moser C, Christensen LD, Bjarnsholt T, Calum H, Jensen PØ, et al. Augmented effect of early antibiotic therapy in mice with experimental lung infections due to sequentially adapted mucoid strains of *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2009;64:1241–50.
- [107] Döring G, Pier GB. Vaccines and immunotherapy against *Pseudomonas aeruginosa*. *Vaccine* 2008;26:1011–24.
- [108] Döring G, Conway SP, Heijerman HGM, Hodson ME, Høiby N, Smyth A, et al. Antibiotic therapy against *Pseudomonas aeruginosa* in cystic fibrosis: a European consensus. *Eur Respir J* 2000;16:749–67.
- [109] Høiby N, Flensburg EW, Beck B, Friis B, Jacobsen L, Jacobsen SV. *Pseudomonas aeruginosa* infection in cystic fibrosis. Diagnostic and prognostic significance of *Pseudomonas aeruginosa* precipitins determined by means of crossed immunoelectrophoresis. *Scand J Respir Dis* 1977;58:65–79.
- [110] Proesmans M, Balinska-Miskiewicz W, Dupont L, Bossuyt X, Verhaegen J, Høiby N, et al. Evaluating the 'Leeds criteria' for *Pseudomonas aeruginosa* infection in a cystic fibrosis centre. *Eur Respir J* 2006;27:937–43.
- [111] Frederiksen B, Lanng S, Koch C, Høiby N. Improved survival in the Danish cystic fibrosis centre: results of aggressive treatment. *Pediatr Pulmonol* 1996;21:153–8.

[112] Koch C, Hjelt K, Pedersen SS, Jensen ET, Jensen T, Lanng S, et al.

Retrospective clinical study of hypersensitivity reactions to aztreonam and six other β -lactam antibiotics in cystic fibrosis patients receiving multiple treatment courses. Rev Infect Dis 1991;13(Suppl 7):S608–11.

Accepted Manuscript

Fig. 1. (A) Example of a biofilm on a biliary stent, which became the focus of repeated incidents of sepsis that were ultimately lethal. DNA typing (pulsed-field gel electrophoresis) showed that it was the same clonal type of *Escherichia coli* in the biofilm and in the blood. Despite relevant antibiotic therapy, it was not possible to eradicate the biliary focus on the stent and the second incident of sepsis was lethal. Microphotos: Gram and methylene blue staining, magnification $\times 100$ and $\times 1000$. Reproduced with permission from [13]. (B) Chronically infected wound with microcolonies of *Pseudomonas aeruginosa* surrounded but not penetrated by polymorphonuclear leukocytes (PMNs). Bacteria are identified using fluorescence in situ hybridisation (FISH) using a fluorescein-labelled peptide nucleic acid specific for *P. aeruginosa*. PMNs were stained by 4',6-diamidino-2-phenylindole (DAPI). Reproduced from [11] with permission.

Fig. 2. Time course of formation of a *Pseudomonas aeruginosa* biofilm based on in vitro experiments with green fluorescent protein-tagged *P. aeruginosa*, grown in a flow cell and examined using confocal scanning laser microscopy. QS (quorum sensing) indicates cell-to-cell communication, and EPS is the hydrated extracellular biofilm matrix. The thickness of the biofilm varies between 25 μm and 100 μm and is determined by the balance between growth and liberation of bacteria. Reproduced from [19] with permission.

Fig. 3. (A) Induction of β -lactamase in *Pseudomonas aeruginosa* biofilm. *Pseudomonas aeruginosa* PAO1 expressing green fluorescent protein (GFP) when the promoter of the AmpC β -lactamase is induced (*PampC-gfp*): 6-day-

old biofilm exposed to 100 µg/mL ceftazidime for 4 h. Detection level of the monitor, 10 µg/mL ceftazidime. Reproduced from [61] with permission. (B) Treatment of *P. aeruginosa* biofilm with a β-lactam. *Pseudomonas aeruginosa ampD⁻* (levels of AmpC β-lactamase: basal 1050 mU; induced 4255 mU) expressing GFP as a tag. Seven-day-old biofilm treated with 10× minimal inhibitory concentration (MIC) of ceftazidime. Propidium iodide (PI) was added after Day 6 to monitor continuously the killing of the biofilm by ceftazidime. Reproduced from [61] with permission. (C) Treatment of *P. aeruginosa* biofilm with a β-lactam. *Pseudomonas aeruginosa ampD⁺* (levels of AmpC β-lactamase: basal 3 mU; induced 175 mU) expressing GFP as a tag. Seven-day-old biofilm treated with 10× MIC of ceftazidime. PI was added after Day 6 to monitor continuously the killing of the biofilm by ceftazidime. Reproduced from [61] with permission. (D) Treatment of *P. aeruginosa* biofilm with a combination of ceftazidime and aztreonam. *Pseudomonas aeruginosa ampD⁻* (levels of AmpC β-lactamase: basal 1050 mU; induced 4255 mU) expressing GFP as a tag. Seven-day-old biofilm treated with a combination of ceftazidime and aztreonam (10× MICs). PI was added after Day 6 to monitor continuously the killing of the biofilm by ceftazidime. Reproduced from [61] with permission.

Fig. 4. Induction of alginate in *Pseudomonas aeruginosa* biofilms treated with sub-minimal inhibitory concentrations (MICs) of imipenem [62]. (A) *Pseudomonas aeruginosa* PAO1 not exposed to antibiotics; (B) PDO300 (a PAO1 derivative constitutively expressing alginate) not exposed to antibiotics; (C) PAO1 exposed to imipenem for 18 h; and (D) PAO1 biofilm exposed to imipenem for 37 h. Alginate is stained green by concanavalin A-conjugated

fluorescein isothiocyanate (conA-FITC). Reproduced from [62] with permission.

Fig. 5. (A) Treatment of *Pseudomonas aeruginosa* biofilm with colistin.

Pseudomonas aeruginosa PAO1 expressing green fluorescent protein (GFP) as a tag was grown as a biofilm in a flow chamber for 4 days. Propidium iodide (PI) was added after Day 4 to monitor continuously the killing of the biofilm by colistin. Image shows the biofilm after 2 days of treatment with 25 $\mu\text{g/mL}$ colistin. Reproduced from [71] with permission. (B) Treatment of *P.*

aeruginosa biofilm with ciprofloxacin. *Pseudomonas aeruginosa* PAO1 expressing GFP as a tag was grown as a biofilm in a flow chamber for 4 days and was treated for 2 days with 10 $\mu\text{g/mL}$ ciprofloxacin. PI was added after Day 4 to monitor continuously the killing of the biofilm by ciprofloxacin. Red staining shows that ciprofloxacin kills the bacteria located at the surface of the biofilm. Reproduced from [71] with permission. (C) Treatment of *P.*

aeruginosa biofilm with a combination of ciprofloxacin and colistin.

Pseudomonas aeruginosa PAO1 expressing GFP as a tag was grown as a biofilm in a flow chamber for 4 days. PI was added after Day 4 to monitor continuously the killing of the biofilm by ciprofloxacin and colistin. The image shows the biofilm after 2 days of treatment with 10 $\mu\text{g/mL}$ ciprofloxacin and 25 $\mu\text{g/mL}$ colistin. Reproduced from [71] with permission.

Fig. 6. Treatment of *Pseudomonas aeruginosa* biofilm with tobramycin. Wild-type PAO1 and $\Delta\text{lasRrhIR}$ mutant, both expressing green fluorescent protein as a tag, were grown as biofilms in flow chambers for 3 days. On Day 3,

tobramycin 10 $\mu\text{g/mL}$ and 20 $\mu\text{g/mL}$ was added. Propidium iodide (PI) was added after Day 3 to monitor continuously the killing of the biofilm by tobramycin. Image shows the biofilm after 48 h of treatment: (a) untreated wild-type; (b) 10 $\mu\text{g/mL}$ tobramycin-treated wild-type; (c) 20 $\mu\text{g/mL}$ tobramycin-treated wild-type; (d) untreated $\Delta\text{lasRrhIR}$ mutant; (e) 10 $\mu\text{g/mL}$ tobramycin-treated $\Delta\text{lasRrhIR}$ mutant; and (f) 20 $\mu\text{g/mL}$ tobramycin-treated $\Delta\text{lasRrhIR}$ mutant. Reproduced with permission from [74].

Fig. 7. Four-day-old biofilm formed by green fluorescent protein-tagged *Pseudomonas aeruginosa* cultured in (A,B) the absence and (C,D) the presence of 2% garlic extract. Biofilms in B and D were treated on Day 3 with 340 $\mu\text{g/mL}$ tobramycin for 24 h. Biofilms were then stained with LIVE/DEAD BacLight™ Bacterial Viability Kit, where dead bacteria are red and living bacteria are green. It is seen that in the presence of garlic extract + tobramycin can kill the bacteria in the biofilm whereas they survive if tobramycin or garlic is used alone. Reproduced from [74] with permission.

Fig. 8. Gram stain ($\times 1000$) of an explanted lung with a biofilm of *Pseudomonas aeruginosa* surrounded by numerous polymorphonuclear leukocytes. The patient is a 41-year-old cystic fibrosis male who has suffered from chronic mucoid *P. aeruginosa* lung infection for 28 years and has been treated with 114 courses of 2-week antipseudomonal antibiotic treatment (chronic suppressive maintenance therapy, total: 1 kg tobramycin, 10 kg β -lactam antibiotics and 1 kg colistin). He had developed 46 precipitating antibodies against *P. aeruginosa* (normal, 0–1).

Table 1

Some general features of biofilm infections in humans compared with acute planktonic infections and superficial colonisation/normal flora on skin and mucosal membranes. Bold indicates biofilm-specific features

Features of biofilm infections	Necessary condition for biofilm infection	Sufficient condition for biofilm infection	Also found in acute planktonic infection	Also found in colonisation/normal flora on skin and mucosal membranes
Aggregates of bacteria embedded in a self-produced polymer matrix	Yes	Yes	No	No/Yes
Tolerant to clinically relevant PK/PD dosing of antibiotics despite susceptibility of planktonic cells	Yes	Yes	No	No/Yes

Tolerant both to innate and adaptive immune responses	Yes	Yes	No	No/Yes—unknown (sIgA)
Inflammation	Yes	No	Yes	No
Biofilm-specific antigens	No and yes—seldom, e.g. <i>Pseudomonas aeruginosa</i> alginate	No and yes—seldom, e.g. <i>P. aeruginosa</i> alginate	No	No
Antibody response	Yes—after some weeks	No	Yes—after some weeks	No
Chronic infections	Yes	Yes	No	No
Foreign-body-associated infections	No	Yes	No, but yes the first day of infection	No
Located on surfaces	No	No	Yes	Yes
Localised infection	Yes	No	Yes	Yes
Focus for spreading or local exacerbation	Yes	No	Yes	Yes

PK/PD, pharmacokinetic/pharmacodynamic; sIgA, serum immunoglobulin A.

Table 2

Natural and pathogenic biofilms on human tissue and foreign bodies

'Organ A' with normal flora	Connection via foreign bodies	'Organ B' without normal flora
Skin	→	blood, peritoneum, middle ear
Mouth	→	teeth
Pharynx	→	bronchi, lungs
Duodenum	→	bile tract, pancreas
Urethra	→	urine, bladder
Vagina	→	uterus
'Air in operation room' ^a	→	alloplastic, cerebrospinal shunt
No symptoms	→	pathology

^a Most frequently, coagulase-negative staphylococci, which occur as biofilms on detached epidermal cells.

Fig. 1A.

Fig. 1B.

Figure 1. *A.* Example of a biofilm on a biliary stent which became focus of repeated incidents of sepsis which ultimately was lethal. DNA typing (pulsed field gel electrophoresis) showed that it was the same clonal type of *E. coli* in the biofilm and in the blood. In spite of relevant antibiotic therapy it was not possible to eradicate the biliary focus on the stent and the second incident of sepsis was lethal. Microphotos: Gram- and methylene blue staining, magnification x 100 and x 1000. Reproduced with permission from Høiby et al. *Ugeskrift for Læger* 169:4163-4166, 2007. *B* Chronic infected wound with microcolonies of *P. aeruginosa* surrounded but not penetrated by polymorphonuclear leukocytes. The bacteria are identified by use of fluorescence in situ hybridization (FISH) using a fluorescein-labelled (FITC) peptide nucleic acid (PNA) specific for *P. aeruginosa*. The polymorphonuclear leukocytes were stained by DAPI. Reproduced from (11) with permission.

Figure 2. Time-course of formation of a *P. aeruginosa* biofilm based on *in vitro* experiments with *green fluorescent protein*-tagged *P. aeruginosa*, which grow in a *flow-cell* and is examined with a confocal scanning laser microscope. QS indicates cell-to-cell communication and EPS is hydrated extracellular biofilm matrix. The thickness of the biofilm varies between 25 – 100 μm and is determined by the balance between growth and lysis of the bacteria. Reproduced from (115) with permission.

Figure 3A. Induction of beta-lactamase in *P. aeruginosa* biofilm. *P. aeruginosa* PAO1 expressing green fluorescent protein (gfp) when the promoter of the AmpC beta-lactamase is induced (*PampC-gfp*): 6 days old biofilm exposed to 100 $\mu\text{g/ml}$ ceftazidime for 4 h. Detection level of the monitor: 10 $\mu\text{g/ml}$ ceftazidime.

Reproduced from (116) with permission.

Figure 3B. Treatment of *P. aeruginosa* biofilm with beta-lactams. *P. aeruginosa* *ampD*⁻ (levels AmpC beta-lactamase (mU) :1050 basal, 4255 induced) expressing green fluorescent protein (gfp) as a tag. 7 days old biofilm treated with 10 times MIC of ceftazidime. Propidium iodide was added after day 6 to continuously monitor the killing of the biofilm by ceftazidime. Reproduced from (116) with permission.

Figure 3C. Treatment of *P. aeruginosa* biofilm with beta-lactams. *P. aeruginosa* *ampD*⁺ (levels AmpC beta-lactamase (mU): basal 3, induced 175) expressing green fluorescent protein (gfp) as a tag. 7 days old biofilm treated with 10 times MIC of ceftazidime. Propidium iodide was added after day 6 to continuously monitor the killing of the biofilm by ceftazidime. Reroduced from (116) with permission.

Figure 3D. Treatment of *P. aeruginosa* biofilms with a combination of ceftazidime and aztreonam. *P. aeruginosa ampD⁻* (levels AmpC beta-lactamase (mU) :1050 basal, 4255 induced) expressing green fluorescent protein (gfp) as a tag. 7 days old biofilm treated with a combination of ceftazidime and aztreonam (10 times MICs). Propidium iodide was added after day 6 to continuously monitor the killing of the biofilm by ceftazidime. Reproduced from (116) with permission.

Figure 4 Induction of alginate in *P. aeruginosa* biofilms treated with sub-MIC concentrations of imipenem (62) A. *P. aeruginosa* PAO1 not exposed to antibiotics; B. PDO300 (a PAO1 derivative constitutively expressing alginate) not exposed to antibiotics; C. PAO1 exposed to imipenem for 18 hours; D. PAO1 biofilm exposed to imipenem for 37 hours. Alginate is stained green by conA-FITC. Reproduced from (117) with permission.

Figure 5A. Treatment of *P. aeruginosa* biofilms with colistin. *P. aeruginosa* PAO1 expressing green fluorescent protein (gfp) as a tag were grown as biofilms in flow-chambers for 4 days. Propidium iodide was added after day 4 to continuously monitor the killing of the biofilm by colistin. The picture shows the biofilm after 2 days of treatment with colistin 25 $\mu\text{g/ml}$. Reproduced from(71) with permission.

Figure 5 B. Treatment of *P. aeruginosa* biofilms with ciprofloxacin. *P. aeruginosa* PAO1 expressing green fluorescent protein (gfp) as a tag was grown as biofilm in a flow-chamber for 4 days and was treated for 2 days with ciprofloxacin 10 $\mu\text{g/ml}$ (A). Propidium iodide was added after day 4 to continuously monitor the killing of the biofilm by ciprofloxacin. Red staining shows that ciprofloxacin kills the bacteria located at the surface of the biofilm. Reproduced from (71)with permission.

Figure 5C Treatment of *P. aeruginosa* biofilm with a combination of ciprofloxacin and colistin. *P. aeruginosa* PAO1 expressing green fluorescent protein (gfp) as a tag was grown as biofilm in flow-chambers for 4 days. Propidium iodide was added after day 4 to continuously monitor the killing of the biofilm by ciprofloxacin and colistin. The present picture shows the biofilm after 2 days of treatment with 10 $\mu\text{g/ml}$ ciprofloxacin and 25 $\mu\text{g/ml}$ colistin. Reproduced from (71) with permission.

Figure 6. Treatment of *P. aeruginosa* biofilm with tobramycin. Wild-type PAO1 and Δ lasRrhIR mutant, both expressing green-fluorescent protein (gfp) as a tag were grown as biofilms in flow-chambers for 3 days. On day 3 tobramycin 10 μ g/ml and 20 μ g/ml were added. Propidium iodide was added after day 3 to continuously monitor the killing of the biofilm by tobramycin. The present pictures show the biofilm after 48 hours of treatment. (a) untreated wild-type, (b) 10 μ g/ml wild-type, (c) 20 μ g/ml wild-type, (d) untreated Δ lasRrhIR mutant, (e) 10 μ g/ml Δ lasRrhIR mutant, (f) 20 μ g/ml Δ lasRrhIR mutant. Reproduced with permission from (74) .

Fig. 7. A 4 day old biofilm formed by *green fluorescent protein*-tagged *P. aeruginosa* cultured in absense (A and B) or presence (C and D) of 2% garlic extract. The biofilms B and D were treated on day 3 with 340 µg/ml tobramycin for 24 h. The biofilms were then stained with LIVE/DEAD *BacLight* bacterial Viability Kit, where dead bacteria are red and living bacteria are green. It is seen that in the presence of garlic extract + tobramycin can kill the bacteria in the biofilm whereas they survive if tobramycin or garlic is used alone. Reproduced from (74) with persission.

Fig. 8. Gram stain (x 1000) of an explanted lung with a biofilm of *P. aeruginosa* surrounded by numerous polymorphonuclear leukocytes. The patient is a 41 year old cystic fibrosis male who has suffered from chronic mucoid *P. aeruginosa* lung infection for 28 years and has been treated with 114 2-weeks anti-pseudomonas antibiotic treatment courses (chronic suppressive maintenance therapy, total: one kg tobramycin, 10 kg betalactam antibiotics and 1 kg colistin). He had developed 46 precipitating antibodies against *P. aeruginosa* (normal: 0-1).