



Seasonal differences in at-sea activity of seabirds underline high energetic demands during the breeding period

Nele Markones, Volker Dierschke, Stefan Garthe

► To cite this version:

Nele Markones, Volker Dierschke, Stefan Garthe. Seasonal differences in at-sea activity of seabirds underline high energetic demands during the breeding period. *Journal für Ornithologie* = *Journal of Ornithology*, 2009, 151 (2), pp.329-336. 10.1007/s10336-009-0459-2 . hal-00568354

HAL Id: hal-00568354

<https://hal.science/hal-00568354>

Submitted on 23 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Nele Markones, Volker Dierschke, and Stefan Garthe

2 **Seasonal differences in at-sea activity of seabirds underline high energetic**
3 **demands during the breeding period**

4
5
6 N. Markones, V. Dierschke and S. Garthe, Research and Technology Centre (FTZ),
7 University of Kiel, Hafentörn 1, 25761 Büsum, Germany

8
9 Corresponding author:

10 N. Markones

11 Research and Technology Centre (FTZ)

12 University of Kiel

13 Hafentörn 1

14 25761 Büsum

15 Germany

16 markones@ftz-west.uni-kiel.de

17 Tel: +49 4834 604 122

18 Fax: +49 4834 604 199

Abstract

We assessed seasonal differences in at-sea activity of Lesser Black-backed Gulls *Larus fuscus*, Black-legged Kittiwakes *Rissa tridactyla* and Common Guillemots *Uria aalge* in the south-eastern North Sea. The three species correspond to different ecological groups with Lesser Black-backed Gulls representing omnivorous generalists, Kittiwakes representing surface-feeding pelagic seabirds and Guillemots representing pursuit-diving pelagic seabirds. Using data from aerial surveys, we differentiated between active (flying or scavenging at fishing vessels) and inactive behaviour (swimming). We estimated activity budgets of all three species for the different seasons and tested for differences in activity between different seasons. All species exhibited significant seasonal differences in activity with highest levels of activity during the breeding season. Numbers of flying auks however were exceptionally low in autumn due to moult and guarding of not-yet fledged chicks at sea. Our results underline the high energetic demands of the breeding season that lead to increased foraging and travelling activity.

Key words

Black-legged Kittiwake, Lesser Black-backed Gull, Common Guillemot, at-sea activity, time-activity budget, seasonality

Introduction

Activity budgets, in combination with knowledge of energetic costs, provide information on resource allocation of seabirds (Goldstein 1990). They may thus be used as indicators of food availability (Cairns 1987) and provide a basis for ecological energetics models (Tasker and Furness 1996). However, information on activity of seabirds at sea is still scarce although the

sea represents the major feeding habitat for seabirds and their exclusive habitat outside the breeding season. Data logger studies have started to fill this gap in our knowledge (e.g. Benvenuti et al. 2001; Garthe et al. 2003; Ropert-Coudert et al. 2004) but usually sample sizes are low and studies have so far mostly covered temporally restricted periods at the order of hours or days. In addition, both interannual variability of extrinsic parameters and individual differences may significantly influence the level of reproductive costs in the breeding season and thus necessitate multi-year comparisons (Golet and Irons 1999). Moreover, logger studies mostly deal with breeding birds due to the need to recapture birds for the detachment of loggers. Due to technological constraints, few studies focusing on activity of birds have so far taken place during the non-breeding season (e.g. Daunt et al. 2006 and see "Outlook"). Bioenergetics models covering the whole year generally extrapolate data of breeding birds and are thus likely to overestimate energy requirements of seabirds (Ellis and Gabrielsen 2002). Hence, data on activity both during and outside the breeding season are needed to minimise errors in bioenergetics models.

So far, data on seabird occurrence and behaviour collected in the course of surveys at sea have seldom been used to gain information on activity of seabirds (but see Camphuysen 1998) although at-sea surveys of seabird abundance have been carried out in many sea areas over the world and studies often cover several decades. In the south-eastern North Sea, the German Bight, Seabirds at Sea surveys have been carried out by ship since 1990 and by aircraft since 2002 (Garthe et al. 2007). Data from aerial surveys are more homogeneous with respect to spatial and temporal effort than ship-based surveys and were thus chosen for the following analyses. Surveys took place throughout the year and activity of seabirds was recorded at least by differentiating between swimming and flying behaviour.

Our study aims to assess reasonable indications of time-activity budgets of selected species for each season in the study area. The null-hypothesis implies no differences in activity levels between different seasons. However, we expect elevated activity levels of breeding birds in the breeding season as parent birds need to maintain self-provisioning and additionally have to raise their young by commuting between the colony and often remote foraging areas at sea (Ricklefs 1984).

We formulate the following specific hypotheses:

(1) Due to the high energetic demands of individuals during the breeding season, we expect a higher amount of actively foraging or travelling birds during the breeding season compared to the other seasons.

(2) Differences in flight activity are expected to be less pronounced in diving species like alcids as flight activity observed may less well represent activity related to foraging. In addition, swimming behaviour includes low proportions of resting (Tremblay et al. 2003). Thus, foraging activity can not be expressed by considering the behaviour “flying”. Nevertheless, the proportion of flying behaviour should be higher due to breeding birds commuting to and from the colony.

Materials and methods

Studies were carried out in the German Bight which is defined here as the area between 53° 21' and 55° 01' N and 05° to 09° E in the south-eastern North Sea. Only breeding species of the study area which forage virtually exclusively at sea were considered appropriate for the analyses. Northern Gannet *Sula bassana*, Northern Fulmar *Fulmarus glacialis* and Razorbill *Alca torda* would meet this criterion, but they are either very scarce or their breeding populations comprise an insignificant share of the total numbers in summer in the German Bight due to large numbers of non-breeding individuals (following Garthe et al. 2007 and

breeding numbers according to Dierschke et al. 2007). Thus, only Lesser Black-backed Gull *Larus fuscus*, Common Guillemot *Uria aalge* (hereafter ‘Guillemot’) and Black-legged Kittiwake *Rissa tridactyla* (hereafter ‘Kittiwake’), whose breeding populations comprised two thirds or more of the total summer populations respectively (following Garthe et al. 2007 and breeding numbers according to Koffijberg et al. 2006 and Dierschke et al. 2007), were considered appropriate for the study. While the latter two occupy a single breeding colony on the small offshore island Helgoland in the study area, Lesser Black-backed Gulls breed in several large colonies along the coast line. During winter, spring and autumn, the proportion of German breeding birds of Kittiwakes and Lesser Black-backed Gulls corresponds to 90% or more of the total population of the two species in the German EEZ of the North Sea (following Garthe et al. 2007 and breeding numbers according to Koffijberg et al. 2006 and Dierschke et al. 2007). On the contrary, the majority of Guillemots observed outside the breeding season belong to populations other than the German breeding population, most probably to the Scottish breeding population. The latter however do not exhibit different phenologies (compare Grunsky-Schöneberg 1998 and Forrester et al. 2007) and thus can be expected not to bias our results on activity patterns. The three species correspond to different ecological groups with Lesser Black-backed Gulls representing omnivorous generalists, Kittiwakes representing surface-feeding pelagic seabirds and Guillemots representing pursuit-diving pelagic seabirds.

Since 2002, data on seabird occurrence in the German Bight have been recorded by aerial surveys following standardised methods described by Kahlert et al. (2000) and Diederichs et al. (2002). Surveys were performed from a high winged twin-engine Partenavia P-68 provided with bubble windows at a flight altitude of 78 m (250 feet) and a cruising speed of circa 185 km/h (100 knots). The occurrence of birds was recorded within 397 m wide transects running in parallel to the flight route of the observation platform. Under good observation conditions,

both sides of the flight route could be covered by two trained observers, resulting in a survey transect of 794 m. All birds were recorded to the second giving details on species, number, and activity. Survey methods included the recording of information on vessel association of species feeding on fishery discards and offal. Surveys sampled a total area of circa 24,600 km², covering large sampling areas during single surveys. Surveys were mostly restricted to the late morning and noon (Table 1).

Due to inferior observation and identification conditions during aerial surveys, it is not possible to identify age classes on the basis of the respective plumages. By applying the ratio of individuals in adult plumage to individuals in immature plumage recorded in ship-based surveys (85:15% for Lesser Black-backed Gulls and 92:8% for kittiwakes during the respective breeding seasons, German Seabirds at Sea database ship v5.07) to aerial surveys, results of all birds from aerial surveys can be assumed to mirror the situation of adult birds and thus mostly breeding individuals of both species. However note that birds in immature plumage comprise first-year and second-year individuals of Kittiwakes and first-year to fourth-year individuals of Lesser Black-backed Gulls respectively. Mean age of first breeding is 4-5 years in Black-legged Kittiwakes (Glutz von Blotzheim and Bauer 1999; Rothery et al. 2002). Thus, birds identified as in adult plumage possibly comprise a substantial proportion of not yet breeding immatures. Immature Guillemots can be identified at sea due to their smaller size only in their first weeks at sea after leaving the colony but can not be identified properly once they are fully grown (proportion of Guillemots identified as immature in summer: 3%, autumn: 16%, winter and spring: <1%, German Seabirds at Sea database ship v5.07). Due to identification difficulties during aerial surveys, we combined data of Guillemot, Razorbill *Alca torda* and “razormot” (Common Guillemot/Razorbill) to obtain an indication of activity budgets of Guillemots. This group is termed as razormots in the following text. We considered this method appropriate as breeding numbers of Razorbills at Heligoland are very

low compared to numbers of Guillemots (18 compared to 2655 apparently occupied nests in 2006; Dierschke et al. 2007). Razorbill numbers comprise less than 5% of Guillemot numbers in the German North Sea during spring, summer and autumn. During winter they make up a share of about 23% compared to numbers of Guillemots (Garthe et al. 2007). However, we assumed that activity budgets of wintering Razorbills and Guillemots do not differ significantly, and are thus confident that the combined data set produces representative results for the Guillemot population of the study area. During rough sea conditions, a flying bird is detected more easily than a swimming bird. This fact influences the results in estimating time budgets. To compensate for the inconspicuousness of swimming auks at sea especially during rough conditions, we thus only included data of relatively calm conditions (sea state <4).

We assessed activity of selected species in all four seasons by distinguishing between flying and swimming individuals and between individuals associated with vessels and those that were not. For the interpretation of our results, we took the proportion of individuals exhibiting a specific behaviour as an indicator for the proportion of time being spent with this behaviour of the respective species in the studied season to get an indication of time-activity budgets (instantaneous sampling; Altmann 1974). We considered flying behaviour and vessel-association to be active (foraging) behaviour. This classification is appropriate in particular for seabirds that mainly forage at the sea surface like gulls and terns and mostly sit on the water surface during resting. Seasons were classified for each species according to timing of breeding, moulting and migration (Table 2). We assessed the ratio of flying and swimming individuals for each season, year and species. Lesser Black-backed Gulls are frequently associated with fishing vessels, feeding on discards and offal. Scavengers at the stern are mostly recorded swimming (pers. observation). We consequently incorporated information on vessel association of single individuals in our analyses of Lesser Black-backed Gulls to test an influence of vessel association on activity as well. Moreover, this method allowed us to

distinguish between swimming birds that were associated with vessels and thus active foragers and those swimmers that were not associated with vessels and thus presumably resting. Very few Kittiwakes (less than 5%, German Seabirds at Sea ship database, version 5.07) and only one Guillemot were recorded in association with vessels. Thus, we ignored this parameter for the latter two species.

It has to be noted that length of potential foraging time per day for individual birds undergoes strong seasonal variation due to changing day length and varying colony attendance of breeding birds. However, we assumed individual length of time available for foraging at sea to be relatively constant between seasons when regarding breeding birds as day length and proportion of time of day spent at the colony are positively correlated such that both effects are offset. For example, day length in summer roughly doubles day length in winter but on the other hand breeders spend approximately 50% of the day during summer at the colony (Tasker and Furness 1996), thus spending approximately the same number of hours at sea under daylight during summer and winter.

Seasonal differences in the ratio of flying and swimming birds and in the proportion of vessel association were tested independent of interannual variability applying a Generalized Linear Mixed Model (GLMM; Faraway 2006) in R Version 2.8.1 (<http://www.r-project.org/>) using the library lme4 (Bates and Sarkar 2007). The model was set as follows: response variable = activity[/vessel association], predictor = breeding/non-breeding period (& vessel association & interaction between both variables), random effect = year, family = binomial. To obtain an estimation of model accuracy we calculated the root mean square error (RMSE). For all species studied we tested whether higher proportions of flying individuals were recorded during breeding season compared to the rest of the year. Furthermore, we tested the influence of breeding season/non-breeding season on vessel association of Lesser Black-backed Gulls.

For the latter species we always omitted the winter data from the analysis because the sum of individuals observed during this season was negligible but could have possibly biased the results. For auks we additionally tested differences in flight activity between autumn and the rest of the year to detect the effect of the moulting and chick-guarding period. Bonferroni correction was applied to account for multiple testing.

Results

Seasonal differences in at-sea activity were evident for all studied species. Analyses revealed significantly higher proportions of flying individuals in summer compared to the rest of the year for all three species.

According to the GLMM, razormots showed significant differences in activity between seasons (Fig. 1, Table 3). The proportion of flying birds was significantly higher during the breeding season and was significantly lower during autumn with only 2 flying individuals of a total of 397 (0.5%). Activity of Kittiwakes differed significantly with season and was highest during summer (Fig. 1, Table 3). Activity of Lesser Black-backed Gulls differed significantly with season and vessel attendance. The proportion of flying individuals was highest during summer and decreased with increasing vessel association (Fig. 1, Table 3). Significant seasonal differences were also recorded for the proportion of vessel association of Lesser Black-backed Gulls with highest values recorded in summer.

Discussion

Methodological aspects

It has to be noted that results on proportions of flying and swimming behaviour can not fully be equated to absolute activity budgets. Considering flying and swimming behaviour only, it

is not possible to differentiate between specific foraging behaviours like plunge diving, surface dipping, etc. over the entire data set. Furthermore, no information can be collected on frequency and length of dive bouts of razormots from a moving observation platform. We analysed at-sea surveys only and thus are not able to incorporate information on length of time spent on land/in the colony and activity at this site. Guillemots and Kittiwakes are truly pelagic seabirds as they do not return to land at night outside the breeding season (Furness and Monaghan 1987). Lesser Black-backed Gulls are not exclusively pelagic concerning their foraging as well as resting behaviour. Individuals of this species usually rest on land throughout the whole year and we assumed that the time spent on land does not differ between breeding season and the rest of the year.

Lesser Black backed Gulls often follow vessels in high numbers. This flocking behaviour complicates quantitative analyses as behaviour of birds in aggregations is not independent. However, as a thorough identification of distinct aggregations at sea is not feasible, this aspect was not included in the analyses.

Timing of surveys with respect to time of day influences results on distribution (Markones et al. 2008) and activity due to the fact that most seabirds exhibit diurnal patterns in activity (Shealer 2002). This problem will be highlighted in the following for the example of the Kittiwake but is assumed to be valid for other species in a similar manner. Breeding Kittiwakes in Scotland showed distinct diurnal differences in activity according to logger data with highest flight activity in the morning between 8 and 13 UTC and during late evening (Daunt et al. 2002). Thus, aerial surveys of the present study covered exclusively periods of highest activity in summer (see Table 1). Consequently, estimated activity budgets are likely to overestimate actual values of the whole day period. General patterns and differences between seasons however should not be influenced by this aspect as the main time of day sampled was very similar between seasons (Tab. 1).

As surveys are carried out visually and thus are restricted to daylight conditions, no information could be incorporated on activity during the night time. Guillemots are thought to be active only during the daytime (Glutz von Blotzheim and Bauer 1999). Studies in the North Sea revealed that Kittiwakes are less active at night both during and outside the breeding season (Garthe and Hüppop 1993, 1996; Daunt et al. 2002) while on the other hand Lesser Black-backed Gulls were frequently foraging at fishing vessels during the night both during and outside the breeding season (Garthe and Hüppop 1993, 1996; Mendel et al., in prep). Studies revealed no information on seasonal differences in diurnal activity. Relatively high RMSE-values of models for Lesser Black-backed Gulls and Kittiwakes (Tab. 3) indicate that factors other than season and vessel association have a strong influence on activity in seabirds. Low RMSE-values of models for Razormots however can be explained by their rather uniform activity patterns comprising high values of swimming behaviour during all seasons.

Seasonal differences in activity

Our results confirmed hypothesis (1) by assessing higher proportions of flying or actively foraging birds in summer compared to the rest of the year for all three species of the different ecological groups. Kittiwakes showed second-highest activity during winter following values of the breeding season (Fig. 1) probably caused by higher thermoregulatory costs and a presumably lower food availability in winter.

Our results support the assumption that breeding birds presumably spend more energy by being engaged in high-costly behaviour like flying and foraging at vessels. It is known that chick-rearing birds increase their working level by 33-50% compared to the non-reproductive period (Drent and Daan 1980). Correspondingly, the summerly proportion of flying Kittiwakes recorded in aerial surveys of our study was nearly 20% higher compared to spring and more than 25% higher compared to autumn. A quarter of all Kittiwakes observed in

summer during aerial surveys in our study was swimming. This value is well in accordance with the proportion of the foraging trip spent on the sea surface recorded by activity loggers for breeding Scottish Kittiwakes (25.0% during daytime, Daunt et al. 2002) and Kittiwakes in Alaska ($21.4\% \pm 15.8\%$, Jodice et al. 2003).

In correspondence to hypothesis (2), activity of razormots was only slightly (but still significantly) elevated in summer. This may be explained by the fact that alcids mainly forage while swimming and the proportion of flying is generally low in this species. Tremblay et al. (2003) found that breeding Guillemots were resting in only 17% of the time they spent at the surface and thus were active in the majority of the time (i.e. preening, swimming actively and interactions with congeners). During the breeding season, flight activity was slightly elevated probably due to movements between foraging area and colony but the difference was very low as can be expected from a mean foraging radius of only 5-10 km only around the single breeding colony in the German Bight on Helgoland (Dierschke et al. 2004). Overall, differences between seasons were very low for this species and thus probably not of real biological relevance.

The recorded low values of flying razormots in autumn correspond to the moulting and chick-guarding period of Guillemots. In correspondence, only 13 individuals out of 1929 (0.7%) were recorded flying during ship-based surveys in autumn (1990-2006, German Seabirds at Sea database ship v5.07). It is striking that virtually no flying individuals were recorded within a period of 76 days (16 July – 30 September) although adults are flightless for a period of only 45-50 days (Glutz von Blotzheim and Bauer 1999). However, Guillemot chicks, exhibiting an intermediate post-hatching development, still need approximately 70 days after leaving the colony to fledge completely (Glutz von Blotzheim and Bauer 1999).

304 To the best of our knowledge, our study is the first to provide information on activity of
305 seabird species in a given study area throughout the year using data of a substantial subset of
306 the whole regional population. The applied method qualified best in assessing activity budgets
307 of surface-feeding species, but was also suited to detect significant differences in activity
308 between seasons in a diving species. We recorded high foraging activity in the breeding
309 season due to high demands, i.e. reproductive costs (self-provisioning plus chick-rearing), and
310 elevated foraging activity in winter presumably due to reduced food supply and high
311 thermoregulatory costs.

312 Results on activity of Kittiwakes correspond well to activity budgets reported in data logger
313 studies. Thus, our analyses of at-sea activity apparently give reasonable estimates of time-
314 activity budgets.

315 The extensive dataset used thus allows the testing of basic theories and contributes to
316 estimates of energy expenditure at the level of seasons. It fills a gap in existing studies of
317 seabird biology as information on activity and energy expenditure is mostly collected in the
318 colony and thus only covers the time period of birds being present in the colony. Data logger
319 studies can give information on both periods, time at the colony and time at sea,
320 simultaneously. However, numbers of samples are mostly small in these studies and generally
321 cover only a short period of a specific breeding season while our study gives an overview
322 over the at-sea activity of an entire region during the whole breeding season and even more on
323 every other season of the year. However, further progress in microtechnology has recently
324 enabled attachments of devices to seabirds the whole-year round producing tracking data over
325 vast ocean areas (e.g. Croxall et al. 2005), and detailed activity data (Catry et al. 2004; Shaffer
326 et al. 2006). Nevertheless, it has to be taken into account that such studies deal with
327 handled birds carrying extra weight and thus may not always produce unbiased results (e.g.
328 Ropert-Coudert and Wilson 2004).

While results on activity in summer lack information on time spent at the colony and thus can not be equated to actual activity budgets, results on activity budgets outside the breeding season can be directly incorporated in models of energy expenditure and food consumption. However, it is recommended to complement results of Lesser Black-backed Gulls by logger studies to obtain data on time spent on land. The data obtained by this methodology may prove important to better understand the energetic demands of birds better understand the energetic demands of birds at year-round, and may further indicate which periods of the year may act as bottlenecks. Also comparisons of activity patterns between different areas may enable us to better identify the importance of certain sea areas.

Zusammenfassung

Saisonale Unterschiede in der Aktivität auf See von Seevögeln

unterstreichen hohe energetische Kosten zur Brutzeit

Wir untersuchten die saisonalen Unterschiede in der Aktivität auf See beobachteter Heringsmöwen *Larus fuscus*, Dreizehenmöwen *Rissa tridactyla* und Trottellummen *Uria aalge* in der südöstlichen Nordsee (Deutsche Bucht). Die drei Arten repräsentieren dabei verschiedene ökologische Gruppen: die Heringsmöwe die der omnivoren Generalisten, die Dreizehenmöwe die der meeresoberflächennah fressenden Pelagen und die Trottellumme die der pelagischen Seevögel, die ihre Nahrung durch Verfolgungstauchen erbeuten.

Dazu analysierten wir Daten, die bei Seevogelerfassungen vom Flugzeug aus erhoben wurden. Wir unterschieden aktives (fliegend oder nahrungssuchend an Fischkuttern) und inaktives Verhalten (schwimmend). Wir berechneten Aktivitätsbudgets aller drei Arten für jede Jahreszeit und testeten Unterschiede in der Aktivität zwischen den verschiedenen Jahreszeiten. Bei allen Arten wurden signifikante Unterschiede in der Aktivität zwischen den

verschiedenen Jahreszeiten festgestellt. Die höchsten Aktivitätsraten traten während der
Brutsaison auf. Außergewöhnlich niedrige Zahlen fliegender Alken wurden im Herbst
festgestellt, die auf die Mauser und das Führen noch flugunfähiger Küken zurückzuführen
sind. Unsere Ergebnisse unterstreichen die hohen Energieanforderungen der Brutsaison, die in
einer erhöhten Nahrungssuch- und Flugaktivität resultieren.

Acknowledgments

Aerial surveys comprise data collected under financial support of the Federal Environmental
Ministry (BMU), the Federal Agency for Nature Conservation (BfN), the Landesbetrieb für
Küstenschutz, Nationalpark und Meeresschutz Schleswig-Holstein (LKN, formerly NPA SH)
and the Niedersächsischer Landesbetrieb für Wasserwirtschaft, Küsten- und Naturschutz
(NLWKN, formerly NLOE). Various observers contributed to data collection, which was
made comfortable by the staff of Sylt Air (Westerland) and FLM Aviation (Kiel). Sven Adler
gave valuable statistical support. John C. Coulson, Bob Furness, Nils Guse and two
anonymous reviewers gave valuable comments on earlier drafts. The analysis was partly
funded by the project ‘Zeitlich-räumliche Variabilität der Seevogel-Vorkommen in der
deutschen Nord- und Ostsee und ihre Bewertung hinsichtlich der Offshore-
Windenergienutzung’ (project 5 of MINOS plus, financed by the Federal Environmental
Ministry).

References

- Altmann J (1974) Observational study of behaviour: sampling methods. Behaviour 49:227-
267
- Ashmole NP (1963) The regulation of numbers of tropical oceanic birds. Ibis 103b:458-473

379 Bates D, Sarkar D (2007) Reference manual lme4: Linear mixed-effects models using S4
 380 classes. <http://mirrors.dotsrc.org/cran/>
 381 Benvenuti S, Dall'Antonia L, Lyngs P (2001) Foraging behaviour and time allocation of
 382 chick-rearing Razorbills *Alca torda* at Græsholmen, central Baltic Sea. *Ibis* 143:402-
 383 412
 384 Cairns DK (1987) Seabirds as indicators of marine food supplies. *Biol Oceanogr* 5:261-271
 385 Camphuysen CJ (1998) Diurnal activity patterns and nocturnal group formation of wintering
 386 Common Murres in the central North Sea. *Colon Waterbirds* 21:406-413
 387 Catry P, Phillips RA, Phalan B, Silk JRD, Croxall JP (2004) Foraging strategies of Grey-
 388 headed Albatrosses *Thalasse chrysostoma*: integration of movements, activity and
 389 feeding events. *Mar Ecol Prog Ser* 280:261-273
 390 Croxall JP, Silk JRD, Phillips RA, Afanasyev V, Briggs DR (2005) Global
 391 circumnavigations: tracking year-round ranges of non-breeding albatrosses. *Science*
 392 307:249-250
 393 Daunt F, Benvenuti S, Harris MP, Dall'Antonia L, Elston DA, Wanless S (2002) Foraging
 394 strategies of the Black-legged Kittiwake *Rissa tridactyla* at a North Sea colony:
 395 evidence for a maximum foraging range. *Mar Ecol Prog Ser* 245:239-247
 396 Daunt F, Afanasyev V, Silk JRD, Wanless S (2006) Extrinsic and intrinsic determinants of
 397 winter foraging and breeding phenology in a temperate seabird. *Behav Ecol Sociobiol*
 398 59:381-388
 399 Diederichs A, Nehls G, Pedersen IK (2002) Flugzeugzählungen zur großflächigen Erfassung
 400 von Seevögeln und marinen Säugern als Grundlage für Umweltverträglichkeitsstudien
 401 im Offshorebereich. *Seevögel* 23:38-46
 402 Dierschke V, Garthe S, Markones N (2004) Aktionsradien Helgoländer Dreizehenmöwen
 403 *Rissa tridactyla* und Trottellummen *Uria aalge* während der Aufzuchtphase.
 404 *Vogelwelt* 125:11-19

405 Dierschke J, Dierschke V, Jachmann F, Stühmer F (2007) Ornithologischer Jahresbericht
 406 2006 für Helgoland. Ornithol Jber 17:1-89
 407 Drent RH, Daan S (1980) The prudent parent: energetic adjustments in avian breeding. Ardea
 408 68:225-252
 409 Ellis HI, Gabrielsen GW (2002) Energetics of free-ranging seabirds. In: Schreiber EA, Burger
 410 J (eds) Biology of marine birds. CRC Press, Boca Raton, pp 359-407
 411 Faraway JJ (2006) Extending the linear model with R: Generalized linear, mixed-effects and
 412 non-parametric models. Chapman and Hall, London
 413 Forrester R, Andrews I, McInerny C, Murray R, McGowan B, Zonfrillo B, Betts M, Jardine
 414 D, Grundy D (2007) The Birds of Scotland. Scottish Ornithologists' Club
 415 Furness RW, Monaghan P (1987) Seabird ecology. Blackie, Glasgow and London
 416 Garthe S, Hüppop O (1993) Gulls and Fulmars following ships and feeding on discards at
 417 night. Ornis Svecica 3:159-161
 418 Garthe S, Hüppop O (1996) Nocturnal scavenging by gulls in the southern North Sea. Colon.
 419 Waterbirds 19:232-241
 420 Garthe S, Benvenuti S, Montevecchi WA (2003) Temporal patterns of foraging activities of
 421 Northern Gannets *Morus bassanus* in the north-west Atlantic. Can J Zool 81:453-461
 422 Garthe S, Sonntag N, Schwemmer P, Dierschke V (2007) Estimation of seabird numbers in
 423 the German North Sea throughout the annual cycle and their biogeographic
 424 importance. Vogelwelt 128:163-178
 425 Glutz von Blotzheim UN, Bauer KM (1999) Handbuch der Vögel Mitteleuropas, Band 8
 426 Charadriiformes (3. Teil). AULA-Verlag, Wiesbaden
 427 Goldstein DL (1990) Energetics of activity and free living in birds. Stud Avian Biol 13:423-
 428 426
 429 Golet GH, Irons DB (1999) Raising young reduces body condition and fat stores in Black-
 430 legged Kittiwakes. Oecologia 120:530-538

431 Grunsky-Schöneberg B (1998) Brutbiologie und Nahrungsökologie der Trottellumme (*Uria*
 432 *aalge* Pont.) auf Helgoland. Ökol Vögel 20:217-274
 433 Jodice PGR, Roby DD, Suryan RM, Irons DB, Kaufman AM, Turco KR, Visser GH (2003)
 434 Variation in energy expenditure among Black-legged Kittiwakes: effects of activity-
 435 specific metabolic rates and activity budgets. Physiol Biochem Zool 76:375–388
 436 Kahlert J, Desholm M, Clausager I, Petersen IK (2000) Environmental impact assessment of
 437 an offshore wind farm at Rødsand: Technical report on birds. Report from NERI
 438 Koffijberg K, Dijkse L, Hälterlein B, Laursen K, Potel P, Südbeck P (2006) Breeding birds
 439 in the Wadden Sea in 2001 - Results of the total survey in 2001 and trends in numbers
 440 between 1991-2001. Common Wadden Sea Secretariat, Trilateral Monitoring and
 441 Assessment Group, Joint Monitoring Group of Breeding Birds in the Wadden Sea,
 442 Wilhelmshaven, Germany
 443 Lack D (1966) Population studies of birds. Oxford University Press, Oxford
 444 Markones N, Garthe S, Dierschke V, Adler S (2008) Small-scale temporal variability of
 445 seabird distribution patterns in the south-eastern North Sea. In: Wollny-Goerke K,
 446 ESKILDSEN K (eds) Marine mammals and seabirds in front of offshore wind energy.
 447 Teubner, Wiesbaden, pp 115-140
 448 Maul AM (1994) Ernährungsweisen und Brutbiologie der Dreizehenmöwe *Rissa tridactyla*
 449 (Linnaeus, 1758) auf Helgoland. Diploma thesis, University of Graz
 450 Prüter J (1989) Phänologie und Nahrungsökologie der Dreizehenmöwen (*Rissa tridactyla*)-
 451 Brutpopulation auf Helgoland. Ökol Vögel 11:189-200
 452 Ricklefs RE (1984) Some considerations on the reproductive energetics of pelagic seabirds.
 453 Stud Avian Biol 8:84-94
 454 Ropert-Coudert Y, Grémillet D, Kato A, Ryan PG, Naito Y, Le Maho Y (2004) A fine-scale
 455 time budget of Cape Gannets provides insights into the foraging strategies of coastal
 456 seabirds. Anim Behav 67:985-992

457 Ropert-Coudert Y, Wilson R (2004) Subjectivity in bio-logging science: do logged data
 458 mislead? Mem Natl Inst Polar Res, Spec. Issue 58:23-33
 459 Rothery P, Harris MP, Wanless S, Shaw DN (2002) Colony size, adult survival rates,
 460 productivity and population projections of Black-legged Kittiwakes *Rissa tridactyla*
 461 on Fair Isle. Atlantic Seabirds 4:17-28
 462 Shaffer SA, Tremblay Y, Weimerskirch H, Scott D, Thompson DR, Sagar PM, Moller H,
 463 Taylor GA, Foley DG, Block BA, Costa DP (2006) Migratory shearwaters integrate
 464 oceanic resources across the Pacific Ocean in an endless summer. Proc Natl Acad Sci
 465 US 103:12799-12802
 466 Shealer DA (2002) Foraging behavior and food of seabirds. In: Schreiber EA, Burger J (eds)
 467 Biology of marine birds. CRC Press, Boca Raton, pp 137-177
 468 Tasker ML, Furness RW (1996) Estimation of food consumption by seabirds in the North
 469 Sea. In: Hunt GLJ, Furness RW (eds) Seabird/fish interactions, with particular
 470 reference to seabirds in the North Sea. ICES Cooperative Research Report No. 216
 471 Tremblay Y, Cherel Y, Oremus M, Tveraa T, Chastel O (2003) Unconventional ventral
 472 attachment of time-depth recorders as a new method for investigating time budget and
 473 diving behaviour of seabirds. J Exp Biol 206:1929-1940
 474

Tables

Table 1 Seasonal survey effort and main day periods sampled by aerial surveys (2002-2006, German Seabirds at Sea database plane v5.06) in the German Bight. Classification of seasons as defined for Black-legged Kittiwakes (see Table 2). Total area surveyed=24,595 km². SD=standard deviation

Season	Mean area surveyed / year [km ²] ± SD	Earliest start time [UTC]	Latest stop time [UTC]	Main time of day sampled [UTC] (>80% of area surveyed)
Winter	1,265 ± 868	08:46	16:13	10-13
Spring	2,071 ± 1,126	06:39	17:54	8-14
Summer	893 ± 1,718	07:04	13:25	8-11
Autumn	690 ± 552	06:15	14:59	7-11

480 **Table 2** Classification of species-specific seasons for the analysis of activity. Spring covers the return to the
 481 breeding site and egg laying while summer comprises the incubation and chick-rearing period in all three species
 482 (Prüter 1989; Maul 1994; Grunsky-Schöneberg 1998; Glutz von Blotzheim and Bauer 1999). Lesser Black-b.
 483 Gull = Lesser Black-backed Gull

	Winter	Spring	Summer	Autumn
Lesser Black-b. Gull	01/11/ - 15/03/	16/03/ - 15/05/	16/05/ - 15/07/	16/07/ - 31/10/
Black-legged Kittiwake	01/11/ - 29/02/	01/03/ - 15/05/	16/05/ - 31/07/	01/08/ - 31/10/
Common Guillemot	01/10/ - 29/02/	01/03/ - 15/04/	16/04/ - 15/07/	16/07/ - 30/09/

484

Table 3 Seasonal variation of seabird activity. Results of GLMMs (for details see Methods). The variable “year” was included as random effect (n = 5 years). *** = significant at 0.001 level, ** = significant at 0.01 level, * = significant at 0.05 level, n.s. = not significant (values Bonferroni-adjusted); Razormot = Common Guillemot/Razorbill

Species	response variable	predictor variable(s)	p	Effect size	z value	RMSE	n birds
Lesser Black-backed Gull	activity	summer/rest of year +	***	0.538	10.765	0.41	15111
		vessel association +	***	-4.249	-11.135		
		interaction (summer/rest of year & vessel association)	***	1.692	4.345		
Lesser Black-backed Gull	vessel association	summer/rest of year	***	1.326	25.979	0.47	15111
Kittiwake	activity	summer/rest of year	***	0.837	11.306	0.48	6061
Razormot	activity	summer/rest of year	**	0.494	3.017	0.15	8176
Razormot	activity	autumn/rest of year	*	-1.827	-2.554	0.15	8176

Figure legends

Fig. 1 Season-specific activity of Lesser Black-backed Gull, Black-legged Kittiwake and “Razormot” (Common Guillemot/Razorbill) in the German Bight as recorded during aerial surveys from 2002-2006. sw=swimming, sw_ves=swimming and associated with fishing vessel, fl=flying, fl_ves=flying and associated with fishing vessel. Activity of Lesser Black-backed Gull in winter is not given due to low sample size

Figures




Fig. 1.