Enzymatic production of β -D-glucose-1-phosphate from trehalose

Journal:	<i>Biotechnology Journal</i>
Manuscript ID:	biot.201000203.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	29-Jul-2010
Complete List of Authors:	Van der Borght, Jef; Ghent University, Centre of Expertise for Industrial Biotechnology and Biocatalysis Desmet, Tom; Ghent University, Centre of Expertise for Industrial Biotechnology and Biocatalysis Soetaert, Wim; Ghent University, Centre of Expertise for Industrial Biotechnology and Biocatalysis
Primary Keywords:	Biocatalysis
Secondary Keywords:	Downstream Processing
Keywords:	Glycosylphosphates

SCHOLARONE™
Manuscripts

Enzymatic production of β -D-glucose-1-phosphate from trehalose

Jef Van der Borght*, Tom Desmet, Wim Soetaert

Center of Expertise for Industrial Biotechnology and Biocatalysis, Department of Biochemical and Microbial Technology, Ghent University, Coupure Links 653, 9000 Ghent, Belgium

* Corresponding author

Key words

Beta-D-glucose-1-phosphate, bioconversion, trehalose phosphorylase, phosphorolysis, downstream processing

Correspondence

Jef Van der Borght

Center of Expertise for Industrial Biotechnology and Biocatalysis

Coupure Links 653

B-9000 Ghent, Belgium

Telephone: +32-9-2649921

Fax: +32-9-2646231

Jef.VanderBorght@UGent.be

List of abbreviations

β Glc1P (β -D-glucose-1-phosphate), Tre (trehalose), Glc (glucose), α Glc1P (α -D-glucose-1-phosphate), α Gal1P (α -D-galactose-1-phosphate), P_i (inorganic phosphate)

Abstract

β -D-glucose-1-phosphate (β Glc1P) is an efficient glucosyl donor for both enzymatic and chemical glycosylation reactions but is currently very costly and not available in large amounts. This paper provides an efficient production method of β Glc1P from trehalose and phosphate using the thermostable trehalose phosphorylase from *Thermoanaerobacter brockii*. At the process temperature of 60 °C, *Escherichia coli* expression host cells are lysed and cell treatment prior to the reaction is therefore not required. In that way, the theoretical maximum yield of 26% could be easily achieved. Two different purification strategies have been compared, *i.e.* either anion exchange chromatography or carbohydrate removal by treatment with trehalase and yeast followed by chemical phosphate precipitation. In a next step, β Glc1P was precipitated with ethanol but this did not induce crystallization, in contrast to what is observed with other glycosylphosphates. After conversion of the product to its cyclohexylammonium salt, however, crystals could be readily obtained. Although both purification methods were quantitative (>99% recovery), a large amount of product (50%) was lost during crystallization. Nevertheless, a production process for crystalline β Glc1P is now available from the cheap substrates trehalose and inorganic phosphate.

1. Introduction

Trehalose phosphorylase (EC 2.4.1.64) catalyzes the reversible phosphorolysis of trehalose (Tre) to β -D-glucose-1-phosphate (β Glc1P) and glucose (Glc) (Fig. 1). It belongs to glycoside hydrolase family 65 together with maltose phosphorylase (EC 2.4.1.8) and kojibiose phosphorylase (EC 2.4.1.230) [1]. These enzymes can be used for the production of β Glc1P from cheap substrates such as maltose or trehalose. Trehalose phosphorylase can be applied for the synthesis of novel disaccharides thanks to its broad acceptor specificity [2, 3]. In these reactions, β Glc1P functions as an activated donor for the α -glucosylation of different monosaccharides. Like other glycosylphosphates, it can also be used as donor for chemical glycosylation reactions [4]. Furthermore, these compounds can have direct applications in anti-inflammatory or immunosuppressive drugs [5] or can be used to manufacture agents with vitamin D-like activity for preventing or curing skin diseases, parathyroid diseases, malignant tumors and bone diseases [6].

These potential applications have drawn considerable interest in the synthesis of glycosylphosphates. Unfortunately, their chemical synthesis suffers from a low yield and lack of anomeric selectivity [7, 8]. In that respect, biocatalytic production processes would be more advantageous. The development of galactokinase variants with modified specificity has been reported but these enzymes require ATP as phosphate donor, which hampers their large-scale application [9]. Glycoside phosphorylases, in contrast, only need inorganic phosphate to produce a glycosylphosphate. In literature, the synthesis of α -glucose-1-phosphate (α Glc1P) from starch [10, 11] or sucrose [12] and that of α -galactose-1-phosphate (α Gal1P) from lactose [13, 14] have previously been reported. A production process for β Glc1P has not yet been described in detail, but is similarly feasible using GH65 disaccharide phosphorylases from cheap substrates

1
2
3 such as trehalose and maltose. Watanabe *et al.* [15] briefly mention the production of β Glc1P
4
5 from trehalose, using typical reaction conditions employed for α Glc1P [16]. In our hands,
6
7 however, a crystalline product could not be obtained using these procedures. Therefore, we
8
9 decided to optimize the production, purification and crystallization of this compound.
10
11

12
13 The use of thermostable enzymes is desired for industrial carbohydrate conversions, as high
14
15 temperatures diminish the odds of contamination. The trehalose phosphorylase of
16
17 *Thermoanaerobacter brockii* is the most stable disaccharide phosphorylase from family GH65
18
19 that is currently available. No maltose phosphorylases have been described that can withstand
20
21 typical process conditions of 60°C for multiple hours. Also, trehalose has become a bulk product
22
23 since the development of an efficient production process from starch by Hayashibara [17],
24
25 making substrate price no longer an issue in the choice of the biocatalyst for β Glc1P production.
26
27
28
29

30
31 The use of a thermostable enzyme also has its benefits for the recovery of a recombinantly
32
33 expressed protein prior to the bioconversion. Indeed, permeabilization and lysis of *E. coli* cells
34
35 can be achieved upon exposure to increased temperatures, which enables the use of a whole
36
37 cell bioconversion [18]. Furthermore, endogenous *E. coli* enzymes are typically inactivated under
38
39 these conditions, which considerably reduces contaminating activities. The production of β Glc1P
40
41 by phosphorolysis of trehalose using both whole *E. coli* cells and cell extracts are compared in
42
43 this paper, as well as various purification methods.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2. Materials and methods

Bacterial strains, plasmids and growth conditions

Escherichia coli BL21(DE3) and XL10 Gold were obtained from Stratagene. *E. coli* CGSC8974 was obtained from the Coli Genetic Stock Center (New Haven, USA). This strain is acid glucose-1-phosphatase (*agp*) negative.

The trehalose phosphorylase (*treP*) gene from *Thermoanaerobacter Brockii* (accession number AB073930) was cloned into the pET15b vector (Novagen) and recombinantly expressed in *E. coli* BL21(DE3). The periplasmic trehalase gene (*treA*) from *E. coli* MG1655 (accession number ECK1185) was cloned into the pTrc99a vector (Amersham Pharmacia Biotech) and recombinantly expressed in *E. coli* XL10 Gold and *E. coli* CGSC8974.

E. coli was routinely grown at 37 °C on LB medium (10 g/L tryptone, 5 g/L yeast extract, 10 g/L NaCl, pH 7.0) supplemented with 0.1 g/L ampicillin.

Trehalose was kindly provided by Cerestar (The Netherlands). All other chemical and medium components were obtained from Sigma or BD Biosciences unless stated otherwise.

Recombinant enzyme production

A glycerol stock of *E. coli* BL21(DE3), transformed with the pET15b-TP vector, was used to inoculate a 5 mL preculture of LB medium supplemented with ampicillin. After overnight growth, 4 mL of culture was used to inoculate 200 mL LB medium supplemented with ampicillin in a 1 L Erlenmeyer. The culture was grown to OD600 0.6 and protein expression was induced by adding IPTG to a final concentration of 0.1 mM. After 6h of protein expression, the culture

(OD₆₀₀ ~ 4.2) was centrifuged and the cell pellet was used either fresh for the bioconversion reaction or frozen at -20°C.

E. coli trehalase was produced in *E. coli* CGSC8974 and XL10 Gold in a similar manner from a glycerol stock. Cell pellets were always frozen at -20°C and lysed before use.

Production of β Glc1P

All bioconversion reactions were performed at 60°C, with a substrate solution containing 200 mM trehalose in 200 mM phosphate buffer pH 7. The effect of a pretreatment of the cells prior to the bioconversion was tested by performing the reaction in parallel with fresh cells, frozen cells and cell extract. These three cell pellets were obtained by centrifuging 25 mL of culture. One of these pellets was lysed by freezing it overnight and subsequently dissolving it in an enzymatic lysis solution (50 mM Tris-HCl, 1 mM EDTA, 4 mM MgCl₂, 50 mM NaCl, 1 mg/mL lysozyme). After 30 min incubation at room temperature, cell debris was removed by centrifugation. To the supernatant, one frozen pellet and one fresh pellet, 50 mL of substrate was added, which was preheated at 60°C. Bioconversion reactions were performed in 50 mL Falcon tubes using an Eppendorf Thermomixer Comfort (Eppendorf AG, Germany) with 750 rpm shaking.

To determine the K_{eq} of the phosphorolysis reaction of trehalose, a frozen cell pellet was lysed with the enzymatic lysis solution. The recombinantly produced trehalose phosphorylase possessed an N-terminal His-tag, due to the cloning in the pET15b plasmid, and was purified using Ni-NTA Spin Columns (Qiagen) with the standard native purification protocol. The purified enzyme was washed, and its buffer was changed to 30 mM MOPS pH 7 using a Microcon YM-30 Centrifugal Filter (Millipore). Bioconversion reactions were performed using substrate solutions containing 5 mM and 10 mM of β Glc1P and glucose in 30 mM MOPS buffer pH 7, in 1.7 mL Eppendorf tubes in a heating block set at 60°C. Both reactions were performed in

triplicate. At regular time-intervals, samples were inactivated for 10 min at 95°C in a heating block and subsequently analyzed by colorimetrically measuring the phosphate concentration with the phosphomolybdate method [19].

For a batch production of β Glc1P, 20 mL of an overnight culture of *E. coli* BL21(DE3) pET15b-TP was used to inoculate 1 L of LB medium with ampicillin in a 2 L Biostat B reactor (B. Braun Biotech Inc., Pennsylvania, USA) and fermentation was performed at 37°C, 400 rpm mixing and 0.8 vvm aeration. When OD600 reached 0.6, IPTG was added to a final concentration of 0.1 mM. After 16 h of protein expression, the culture (OD600 ~ 4.5) was centrifuged and the cell pellets were added to a 5 L Biostat B reactor, containing 2 L of preheated bioconversion substrate. After 2 h, the bioconversion reaction reached its equilibrium concentrations and biomass was removed by filtration through a Seitz EKS filter. The resulting bioconversion product was used to compare different purification methods for β Glc1P.

Purification and crystallization of β Glc1P

Two strategies for purification are compared in this paper. In the first strategy, separation of β Glc1P from phosphate, glucose and trehalose was achieved by anion-exchange chromatography as described by Hokse [20]. Dowex 1-X8 (Alfa Aesar, Germany) in the acetate form in a 77 x 3 cm column and a bed volume of 250 mL was used at room temperature. Elution was done with 250 mM potassium acetate buffer pH 4.5 and fractions corresponding to 1 column volume (250 mL) were collected. After each cycle, the resin was regenerated with 1 M potassium acetate. Elution fractions containing β Glc1P were pooled and concentrated in a rotary evaporator at 60°C. After adjustment of the pH to > 8 with KOH, β Glc1P was precipitated by the addition of ethanol to form a transparent gel after overnight incubation at 4°C. This gel was

resuspended in distilled water and concentrated in a rotary evaporator to remove traces of ethanol. The resulting solution contained the pure potassium salt of β Glc1P.

In the second strategy, β Glc1P was purified by simultaneous removal of trehalose by *E. coli* trehalase (200 mL cell extract / L substrate) and glucose by baker's yeast (20 g/L) at 37°C during 24 h. Phosphate was precipitated by the addition of magnesium acetate and ammonia [16]. Biomass and precipitated phosphate were removed by filtration through a Seitz EKS filter, and β Glc1P was further purified by ethanol precipitation as described earlier.

To obtain a crystalline product, the cyclohexylammonium salt was prepared, as it is known that sugar-phosphates readily crystallize in this form [21-23]. The potassium cations were removed by adding Amberlite IR120H resin (Aldrich) to pH 2.5. The resin was recovered by filtration and the flow-through was titrated to pH 8 with cyclohexylamine. The cyclohexylammonium salt was crystallized by the method of Friedkin [23]. β Glc1P crystals were filtered and dried overnight at 70°C.

The hydrolysis of Glc1P by *E. coli* phosphatases present in the cell extract during carbohydrate removal by trehalase was quantified. A phosphatase positive (XL10 Gold) and a phosphatase negative *E. coli* strain (CGSC8974) were grown in both normal LB medium and LB medium supplemented with 30 mM KH_2PO_4 and 2 g/L glucose. In analogy to the production of recombinant trehalase, an overnight culture was used to inoculate fresh medium. The fresh culture was grown for 6 h after an OD600 of 0.6 was reached and centrifuged. The four cell pellets thus obtained were frozen and lysed by the enzymatic lysis solution. Their phosphatase activity on both 30 mM α Glc1P and β Glc1P in 30 mM MOPS buffer pH 7 was tested at 37°C in 1.7 mL Eppendorf tubes in a heating block. Samples were inactivated for 10 min at 95°C and subsequently analyzed for their phosphate concentration using the phosphomolybdate method [19]. All reactions were performed in triplicate.

Analysis of β Glc1P

All bioconversion reactions were monitored by HPLC analysis. Samples were centrifuged to remove biomass, diluted if necessary in MilliQ water and subsequently filtrated through a Minisart RC-15 filter (pore size 0.45 μ m, Sartorius). The analysis was performed on a Prostar system (Varian) composed of a Prostar 230 Solvent Delivery Module, a cooled (4 °C) Prostar 410 Autosampler, an LaChrom RI detector L-7490 (Merck) and Varian Star 6 Chromatography Workstation software. Chromatographic separation was performed with an Aminex HPx-87H Ion Exchange Column (300 x 7.8 mm, Bio-Rad) at 30 °C. The eluent was 5 mM H₂SO₄ and the flow-rate 0.6 mL/min. The expected retention times were 6.3 min (β Glc1P), 7.3 min (trehalose), 8.1 min (phosphate) and 8.7 min (glucose).

3. Results and discussion

Production of the biocatalysts

The trehalose phosphorylase from *Thermoanaerobacter Brockii* was recombinantly expressed in *E. coli*. Expression host cells can be disrupted completely to release recombinant enzymes and perform reactions of interest with cell-free extracts. These methods range from mechanical such as ultrasonication [24] to enzymatic through the use of lysozyme [25] to chemical methods by using detergents or EDTA and urea [26]. A significant advantage of the use of thermostable enzymes is the possibility to use the exposure to higher temperatures to achieve a similar effect. Thermolysis of *E. coli* to release a recombinant thermostable esterase was previously described by Ren *et al.* [18]. The host cells are lysed, and *E. coli* proteins are denatured while the thermostable enzyme remains active. This makes it possible to perform bioconversion reactions without tedious pretreatment of the cells and without significant background activities from

expression host enzymes. For the biocatalytic production of β Glc1P specifically, this means that since the reaction is performed at 60°C, there is no product degradation by *E. coli* phosphatases. Experiments confirmed that the maximal yield of β Glc1P was not influenced by phosphatase activity during a bioconversion process at these temperatures.

In a second purification strategy, however, a cell extract containing recombinant trehalase is used to hydrolyse unreacted trehalose. Since the optimum temperature of this enzyme is only 37°C, phosphatases remain active which could result in significant product loss. In this case, *E. coli* expression host phosphatase activity had to be avoided in another way. It has been described in literature that *E. coli* phosphatase activity could be lowered significantly by using an *agp*-negative strain like CGSC8974 and growing it in LB medium, supplemented with glucose and phosphate [13, 27, 28]. This effect was quantified and the specificity of *E. coli* phosphatases for α Glc1P and β Glc1P was evaluated (Fig. 2). A large difference is visible between the hydrolysis of α Glc1P and β Glc1P by the phosphatase positive strain. This is most significant when grown on standard LB medium, where the phosphatase activity on β Glc1P is 24 times lower. Both the use of a phosphatase negative strain and the addition of glucose and phosphate to the growth medium are successful in lowering Glc1P breakdown. In total, phosphatase activity on β Glc1P can be lowered 7 times to 0.29 μ M/min. In a 24 h purification, this will lead to hydrolysis of 418 μ M β Glc1P. In a mixture containing 52 mM starting concentration, this corresponds to a loss of less than 1%.

Production of β Glc1P

To determine the maximum yield of conversion, the K_{eq} of the reaction was determined experimentally by performing a bioconversion at 60°C with low substrate concentrations (5 – 10 mM β Glc1P and glucose) in triplicate and monitoring the phosphate concentration until the

reaction stopped. The concentrations of trehalose and phosphate produced were presumed to be equimolar. Indeed, inverting phosphorylases such as TP do not display detectable hydrolytic activity, in contrast to retaining enzymes [29]. The equilibrium constant was determined by calculating $[Glc].[βGlc1P]/[Tre].[P_i]$ and was equal to 0.122. This reflects the high stability of trehalose. The equilibrium constant (K_{eq}) of the phosphorolysis reaction of glucosides, including di-, oligo- and polysaccharides is usually in the range of 0.2 or greater [12]. This implies that the reaction equilibrium of trehalose phosphorolysis lies more towards disaccharide synthesis than for other phosphorolytic reactions. Thus, using maltose could result in a higher yield of $βGlc1P$, but then production could not be performed at the desired reaction temperature of 60°C since no stable maltose phosphorylase is available. Overall, 26% conversion of trehalose and phosphate to $βGlc1P$ and glucose can be achieved. The inefficient use of trehalose is not a problem, since it has recently become a bulk product. The relative yield of $βGlc1P$ compared to trehalose could be increased by using higher phosphate concentrations but that would require the removal of a much higher amount of phosphate in the downstream processing.

Given that heat treatment is sufficient to permeabilize and/or lyse *E. coli* cells, it should be possible to perform the bioconversion reaction at 60°C without preliminary cell lysis. This was tested by comparing the efficiency of three bioconversion conditions using respectively fresh cells, frozen cells and cell extract (Fig. 3). From these results, it is clear that each condition possessed a different lag-phase at the start of the reaction. The reaction with cell extract started immediately and already contained 7.9 mM $βGlc1P$ by the time the first sample was taken (1 min), while it took the frozen cells approximately 8 min and the fresh cells 16 min to reach the same concentration. This shows that freezing the cells made them more sensitive to thermolysis and fresh cells were lysed more gradually. Although the initial velocities were different, all three reactions took equally long to reach the equilibrium concentration of 52 mM $βGlc1P$ from 200

mM trehalose and phosphate, so cell lysis or freezing seems unnecessary prior to the bioconversion.

A batch bioconversion was then performed on a 2 L scale to optimize the purification of β Glc1P. The reaction was performed at 60°C with 200 mM trehalose in 200 mM phosphate buffer pH 7. As biocatalyst, cells from a 1 L fermentation were used, which had an activity of approximately 1 U/mL of culture. After 2 h, the conversion reached its equilibrium and the cells were removed by filtration.

Purification of β Glc1P

Two different purification methods were compared in terms of ease of use and recovery of product. The first method was based on anion exchange chromatography as described by Hokse [20] for the purification of α Glc1P. Negatively charged phosphate and sugar-phosphates interact with the strongly basic groups on the polystyrene-divinylbenzene matrix, while uncharged mono- and disaccharides are not retained on the column. A typical elution profile is presented in Fig. 4. One column volume (250 mL) of bioconversion product, containing 52 mM of β Glc1P and glucose and 148 mM of trehalose and phosphate, was loaded on the resin and eluted with 250 mM potassium acetate pH 4.5. The fractions that contained β Glc1P were pooled and analyzed by HPLC. This showed that all β Glc1P was recovered and no phosphate, glucose or trehalose was present. After each cycle, the resin was regenerated with 1 M potassium acetate. In total 1 L of bioconversion product was purified. The β Glc1P was precipitated by adding ethanol to a final concentration of 80% to remove the acetate. Prior to this precipitation, the solution had been concentrated in a rotary evaporator to minimize the loss of β Glc1P and limit the amount of ethanol needed. It was found that the potassium salt of β Glc1P did not crystallize, but formed a gel instead as will be discussed later. After this gel was dissolved in

1
2
3 distilled water and traces of ethanol were removed in a rotary evaporator, all β Glc1P was
4
5 recovered and highly pure.
6

7
8 An alternative purification method was also investigated that might be more suitable for large-
9
10 scale operations. In literature, baker's yeast *Saccharomyces cerevisiae* is often used to
11
12 metabolize sugar by-products in bioconversion reactions [30-32] while phosphate can be
13
14 removed by chemical precipitation [16]. However, Yoon *et al.* [32] have noted that trehalose is
15
16 only partially hydrolyzed by baker's yeast. This was confirmed in our experiments, which implied
17
18 the need for the addition of a trehalase to assist in carbohydrate removal. The trehalase from *E.*
19
20 *coli* was therefore overexpressed in the phosphatase negative *E. coli* strain CGSC8974 as
21
22 described earlier.
23
24

25
26 Similar to the anion exchange purification method, a 1 L batch of bioconversion product was
27
28 purified using baker's yeast and *E. coli* trehalase at 37°C. No hydrolysis of β Glc1P was observed
29
30 during this process, which means that contaminating phosphatase activity is negligible in both
31
32 enzyme preparations. After 24 h all trehalose and glucose were metabolized with the formation
33
34 of ethanol and some organic acids. Yeast biomass was removed by filtration. Phosphate was
35
36 precipitated from the solution in the form of struvite (magnesium ammonium phosphate) as
37
38 described by Asbhy *et al.* [16]. Struvite can be recovered from the process and has potential
39
40 applications as a slow-release fertilizer [33]. All phosphate precipitated immediately at room
41
42 temperature and was removed by filtration through a Seitz EKS filter. As previously described for
43
44 the other purification method, the β Glc1P solution was precipitated by adding ethanol to a final
45
46 concentration of 80% after concentrating in a rotary evaporator. HPLC analysis of the solution
47
48 showed that acetate was completely removed, but traces (< 1%) of other organic acids
49
50 remained.
51
52
53
54
55
56
57
58
59
60

Overall, both methods were successful in recovering β Glc1P with a high yield (> 99%). Through anion exchange, completely pure product could be obtained, while some traces of organic acids (< 1%) were still present with the alternative strategy. For scale-up of the process, the price and complexity of the procedures also needs to be considered. Although chromatographic separation is very efficient, it is a relatively sophisticated method that requires specialized equipment and careful attention. The high price of the resin is, perhaps, not problematic as it can be regenerated and re-used many times. However, a large amount of waste is generated during the elution and regeneration steps. Furthermore, the product becomes strongly diluted and the remaining phosphate and carbohydrates can not be recovered due to contamination with acetate. The main drawback of the alternative strategy is the need for trehalase as additional enzyme. Although biomass production is generally regarded as cheap, preparation of the enzyme is more labour-intensive. However, no investment in equipment is required in this case, and recovered phosphate can be valorized as highly pure struvite.

For long term storage or to remove the final impurities, crystallization of the product is desirable. Surprisingly, the potassium salt did not crystallize after addition of ethanol as mentioned earlier. This was the case for the β Glc1P purified by both methods and thus was not influenced by product purity. The same was true for the sodium salt, which was obtained by the addition of cation exchange resin to pH 2.5, followed by titration to pH > 8 with sodium hydroxide. Therefore, we decided to convert β Glc1P to its cyclohexylammonium salt by means of cation exchange, as it is known that sugar-phosphates readily crystallize in this form [21-23]. After addition of butanol and diethylether as described by Friedkin [23], small needles indeed formed in the solution. All purified product was pooled and crystallized. The formed β Glc1P crystals were filtered and dried overnight at 70°C, after which 22.9 g were obtained, which corresponds to a yield of approximately 50 %. In contrast to the efficient production and purification methods,

crystallization of the product results in a significant product loss. However, HPLC analysis showed that the remaining β Glc1P was still present in the solution, and could be recovered by performing a second or third harvest.

4. Concluding remarks

Here, an efficient biocatalytic process for the production of β Glc1P from trehalose is described. At the process temperature of 60°C, *E. coli* cells expressing a thermostable trehalose phosphorylase become permeabilized for substrate and product, allowing a whole-cell biotransformation to be performed. In that way, a maximal conversion of 26% can be readily achieved, which is limited only by the reversible nature of the reaction. Next, two purification methods have been optimised that are highly efficient in product recovery (> 99%). On the one hand, anion exchange chromatography results in a very pure product but generates a lot of waste. On the other hand, fermentation of the remaining carbohydrates followed by chemical precipitation of phosphate is a simpler process but results in product that is slightly less pure. Finally, crystallization of β Glc1P was found to be possible only as cyclohexylammonium salt, in contrast to what is observed for other glycosyl phosphates.

Acknowledgements

Research funded by a Ph.D grant (SB71279, SB73279) of the Institute for the Promotion of Innovation through Science and Technology in Flanders (IWT-Vlaanderen).

The authors have declared no conflict of interest.

5. References

- [1] Cantarel, B. L., Coutinho, P. M., Rancurel, C., Bernard, T., *et al.*, The Carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res.* 2009, *37*, D233-D238.
- [2] Aisaka, K., Masuda-Kato, T., Chikamune, T., Kamitori, K., *et al.*, Enzymatic synthesis of novel disaccharides using disaccharide phosphorylases. *J. Biosci. Bioeng.* 2000, *90*, 208-213.
- [3] Chaen, H., Nakada, T., Nishimoto, T., Kuroda, N., *et al.*, Purification and characterization of thermostable trehalose phosphorylase from *Thermoanaerobium Brockii*. *J. Appl. Glycosci.* 1999, *46*, 399-405.
- [4] Plante, O. J., Andrade, R. B., Seeberger, P. H., Synthesis and use of glycosyl phosphates as glycosyl donors. *Org. Lett.* 1999, *1*, 211-214.
- [5] Parish, C. R., Cowden, W. B., Willenborg, D. O., Phosphosugar-based anti-inflammatory and/or immunosuppressive drugs. *US Patent 5506210* 1996.
- [6] Fujinaka, H., Nakamura, J., Murase, D., Souno, H., Kobayashi, H., Use of monosaccharide phosphates for improving the intestinal function. *EU Patent 1477173* 2007.
- [7] Reithel, F. J., β -D-Galactose-1-(barium phosphate). *J. Am. Chem. Soc.* 1945, *67*, 1056-1057.
- [8] Posternak, T., Synthesis of α -D-glucose-1-phosphate and α -D-galactose-1-phosphate. *J. Am. Chem. Soc.* 1950, *72*, 4824-4825.
- [9] Yang, J., Fu, X., Liao, J., Liu, L., Thorson, J. S., Structure-based engineering of *E. coli* galactokinase as a first step toward in vivo glycorandomization. *Chem. Biol.* 2005, *12*, 657-664.
- [10] Bae, J., Lee, D., Kim, D., Cho, S. J., *et al.*, Facile synthesis of glucose-1-phosphate from starch by *Thermus caldophilus* GK24 alpha-glucan phosphorylase. *Process Biochem.* 2005, *40*, 3707-3713.

- [11] Weinhausel, A., Nidetzky, B., Kysela, C., Kulbe, K. D., Application of *Escherichia coli* maltodextrin-phosphorylase for the continuous production of glucose-1-phosphate. *Enzyme Microb. Technol.* 1995, 17, 140-146.
- [12] Goedel, C., Schwarz, A., Minani, A., Nidetzky, B., Recombinant sucrose phosphorylase from *Leuconostoc mesenteroides*: Characterization, kinetic studies of transglucosylation, and application of immobilised enzyme for production of alpha-D-glucose 1-phosphate. *J. Biotechnol.* 2007, 129, 77-86.
- [13] De Groeve, M. R. M., Depreitere, V., Desmet, T., Soetaert, W., Enzymatic production of alpha-d-galactose 1-phosphate by lactose phosphorolysis. *Biotechnol. Lett.* 2009, 31, 1873-1877.
- [14] De Groeve, M. R. M., De Baere, M., Hoflack, L., Desmet, T., *et al.*, Creating lactose phosphorylase enzymes by directed evolution of cellobiose phosphorylase. *Protein Eng. Des. Sel.* 2009, 22, 393-399.
- [15] Watanabe, H., Higashiyama, T., Aga, H., Nishimoto, T., *et al.*, Enzymatic synthesis of a 2-O-[alpha]-d-glucopyranosyl cyclic tetrasaccharide by kojibiose phosphorylase. *Carbohydr. Res.* 2005, 340, 449-454.
- [16] Ashby, J. H., Clarke, H. B., Crook, E. M., Datta, S. P., Thermodynamic quantities for the dissociation equilibria of biologically important compounds. 4. The second acid dissociation of glucose 1-phosphoric acid. *Biochem. J.* 1955, 59, 203-208.
- [17] Maruta, K., Nakada, T., Kubota, M., Chaen, H., *et al.*, Formation of trehalose from maltooligosaccharides by a novel enzymatic system. *Biosci. Biotechnol. Biochem.* 1995, 59, 1829 - 1834.
- [18] Ren, X. D., Yu, D. W., Yu, L., Gao, G., *et al.*, A new study of cell disruption to release recombinant thermostable enzyme from *Escherichia coli* by thermolysis. *J. Biotechnol.* 2007, 129, 668-673.
- [19] Gawronski, J. D., Benson, D. R., Microtiter assay for glutamine synthetase biosynthetic activity using inorganic phosphate detection. *Anal. Biochem.* 2004, 327, 114-118.
- [20] Hokse, H., Purification of alpha-D-glucose-1-phosphate. *Starke* 1983, 35, 101-102.
- [21] MacDonald, D. L., Phosphates and other inorganic esters, in: Pigman, W., Horton, D. (Eds.), *The Carbohydrates. Chemistry and biochemistry.*, Academic Press 1972, pp. 253-277.

- [22] McGilvery, R. W., Cyclohexylammonium fructose-1, 6-diphosphates. *J. Biol. Chem.* 1953, *200*, 835-838.
- [23] Friedkin, M., Desoxyribose-1-phosphate. II. The isolation of crystalline desoxyribose-1-phosphate. *J. Biol. Chem.* 1950, *184*, 449-460.
- [24] Feliu, J. X., Cubarsi, R., Villaverde, A., Optimized release of recombinant proteins by ultrasonication of *E-coli* cells. *Biotechnol. Bioeng.* 1998, *58*, 536-540.
- [25] Salazar, O., Asenjo, J. A., Enzymatic lysis of microbial cells. *Biotechnol. Lett.* 2007, *29*, 985-994.
- [26] Falconer, R. J., Oneill, B. K., Middelberg, A. P. J., Chemical treatment of *Escherichia coli* .1. Extraction of intracellular protein from uninduced cells. *Biotechnol. Bioeng.* 1997, *53*, 453-458.
- [27] Bolton, P. G., Dean, A. C. R., Phosphatase synthesis in *Klebsiella-(Aerobacter) aerogenes* growing in continuous culture. *Biochem. J.* 1972, *127*, 87-96.
- [28] Rossolini, G. M., Thaller, M. C., Pezzi, R., Satta, G., Identification of an *Escherichia-coli* periplasmic acid-phosphatase containing of a 27 KDa-polypeptide component. *FEMS Microbiol. Lett.* 1994, *118*, 167-173.
- [29] Wiesbauer, J., Goedel, C., Schwarz, A., Brecker, L., Nidetzky, B., Substitution of the catalytic acid-base Glu237 by Gln suppresses hydrolysis during glucosylation of phenolic acceptors catalyzed by *Leuconostoc mesenteroides* sucrose phosphorylase. *J. Mol. Catal. B: Enzym.*, *65*, 24-29.
- [30] Nishimoto, M., Kitaoka, M., Practical preparation of lacto-N-biose I, a candidate for the bifidus factor in human milk. *Biosci. Biotechnol. Biochem.* 2007, *71*, 2101-2104.
- [31] Chaen, H., Nakada, T., Mukai, N., Nishimoto, T., *et al.*, Efficient enzymatic synthesis of disaccharide, alpha-D-galactosyl alpha-D-glucoside, by trehalose phosphorylase from *Thermoanaerobacter brockii*. *J. Appl. Glycosci.* 2001, *48*, 135-137.
- [32] Yoon, S. H., Mukerjea, R., Robyt, J. F., Specificity of yeast (*Saccharomyces cerevisiae*) in removing carbohydrates by fermentation. *Carbohydr. Res.* 2003, *338*, 1127-1132.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[33] Shu, L., Schneider, P., Jegatheesan, V., Johnson, J., An economic evaluation of phosphorus recovery as struvite from digester supernatant. *Bioresour. Technol.* 2006, 97, 2211-2216.

For Peer Review

Figure Legends

Figure 1. Reaction scheme of trehalose phosphorylase.

Figure 2. Comparison of phosphatase activity on α Glc1P and β Glc1P in an *agp*-positive (XL10 Gold) and negative (CGSC8974) *E. coli* strain, grown on standard LB medium and LB medium supplemented with phosphate (30 mM KH_2PO_4) and glucose (2 g/L).

Figure 3. Effect of cell pretreatment on the phosphorolysis of trehalose. Cells were centrifuged and either immediately resuspended in preheated substrate, frozen or frozen and lysed with an enzymatic lysis solution before adding preheated substrate (200 mM trehalose in 200 mM phosphate buffer pH 7).

Figure 4. Anion exchange elution profile for the separation of β Glc1P from trehalose, glucose and phosphate. Separation was performed on Dowex 1-X8 resin with 250 mM potassium acetate pH 4.5 as eluent. Each fraction corresponds to one column volume (250 mL).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Reaction scheme of trehalose phosphorylase.
36x8mm (600 x 600 DPI)

E. coli strain and growth medium

Comparison of phosphatase activity on α Glc1P and β Glc1P in an agp-positive (XL10 Gold) and negative (CGSC8974) *E. coli* strain, grown on standard LB medium and LB medium supplemented with phosphate (30 mM KH_2PO_4) and glucose (2 g/L).
70x63mm (600 x 600 DPI)

Effect of cell pretreatment on the phosphorolysis of trehalose. Cells were centrifuged and either immediately resuspended in preheated substrate, frozen or frozen and lysed with an enzymatic lysis solution before adding preheated substrate (200 mM trehalose in 200 mM phosphate buffer pH 7). 60x46mm (600 x 600 DPI)

Anion exchange elution profile for the separation of β Glc1P from trehalose, glucose and phosphate. Separation was performed on Dowex 1-X8 resin with 250 mM potassium acetate pH 4.5 as eluent. Each fraction corresponds to one column volume (250 mL). 60x46mm (600 x 600 DPI)