Additional Findings at Preoperative MRI. A Simple Golden Rule for a Complex Problem?

Francesco Sardanelli

*Università degli Studi di Milano, Dipartimento di Scienze Medico-Chirurgiche, Unità di Radiologia, IRCCS
Policlinico San Donato, San Donato Milanese, Milan, Italy*

Commentary

Short running title:

A Golden Rule for MRI Additional Findings?

Correspondence to:

Prof. Francesco Sardanelli
Università degli Studi di Milano
Dipartimento di Scienze Medico-Chirurgiche
Servizio di Radiologia, IRCCS Policlinico San Donato
Via Morandi 30
20097 San Donato Milanese (MI)
Tel +390252774468
Fax +39025279695
E-mail: f.sardanelli@grupposandonato.it
francesco.sardanelli@unimi.it

Additional Findings at Preoperative MRI. A Simple Golden Rule for a Complex Problem?

Preoperative magnetic resonance imaging (MRI) is a hot topic, a complex problem which probably will remain unresolved for several years.

The starting point is the great body of evidence for the higher sensitivity of MRI compared with mammography and ultrasonography (US) in detecting malignant lesions in the breast harboring the index cancer [1] and in the contralateral breast [2]. However, the discussion about advantages [3] and disadvantages [4] is challenging [5].

Potential advantages of preoperative MRI can be summarized as follows:

1. better short-term patient outcome (reduced re-excision rate; reduced amount of removed tissue due to a tailored surgical strategy);
2. better mid-term patient outcome (reduced rate of ipsilateral local recurrence and contralateral cancer);
3. better long term patient outcome (longer disease free and overall survival);
4. better psychological patient status due to the use of the most sensitive tool for local staging.

However, none of these potential advantages have been clearly demonstrated by high-level studies, i.e. by a large multi-institutional randomized controlled trial (RCT). On the other side, drawbacks of preoperative MRI can be outlined as follows:

1. overtreatment due to false positive MRI findings;
2. overdiagnosis and overtreatment of malignant additional lesions which could have been cured by radiotherapy and/or systemic adjuvant chemotherapy or hormone therapy;
3. delay in definition of surgical strategy due to difficult managing of MRI additional findings, especially those visible only at MRI, hence requiring an MR-guided needle biopsy and/or localization;
4. increase in patient anxiety due to treatment delay and/or uncertainties related to the interpretation and management of additional findings.

Obviously, advantages and disadvantages can be mixed in variable combinations. The problem is hard because we have in our hands a technique surely being the best option for evaluating ipsilateral disease extent and possible contralateral cancers but we are not sure that, using this technique, we have a better treatment for our patients. We could have a worse (i.e., an avoidable more aggressive) treatment. Moreover, patients know that MRI is highly sensitive and self-referred presentation is possible.

Looking at the secondary evidence, meta-analyses found a rate of more aggressive surgery for true positive MRI findings of 11.1% for the ipsilateral breast [1] and 4.1% for the contralateral breast [2]. The rate of MRI-induced ipsilateral wider surgical treatment should be compared with the rate of positive margins after breast conserving treatment (BCT), reported to be from 20% to 40% or more, and that of local recurrences after BCT, usually considered from 5% to 10% at ten years and reported about 9% at 20 years. The rate of MRI-induced contralateral surgery for synchronous cancer should be compared with 0.5-1% annual risk of contralateral breast cancer in women with a previous history of breast cancer [3]. Notably, only ipsilateral recurrences or contralateral cancers which would have appeared in the first years after a conservative treatment might be avoided by pre-operative MRI. Thus, we obtain a relatively balanced result for the contralateral breast: 3-4% of MRI-detected contralateral cancers versus 2-3% cumulative rate of expected contralateral cancers. Conversely, we have an 11% of MRI-induced wider ipsilateral surgery versus and only 2-3% of cumulative rate of expected ipsilateral recurrences [3].

As a consequence, the risk of overdiagnosis and overtreatment by preoperative MRI is probable at least for the ipsilateral breast. However, we must consider that: a) the rate of MRI-detected ipsilateral and contralateral cancers is probably overestimated due to the fact that preoperative MRI has been performed in non-consecutive (selected) patients with a probable higher likelihood of these lesions; b) a publication bias can be hypothesized; c) ipsilateral and/or contralateral overtreatment could be compensated by the reduction of additional surgical interventions needed to achieve free margins; d) considering the role of radiation and systemic therapy, a patient-based perspective should evaluate the combined effect on both breasts. This last point means that MRI could determine an unnecessary ipsilateral excision but also anticipate the diagnosis of a contralateral cancer – or vice versa – avoiding the future second cancer event. For an individual patient, preoperative MRI may determine a spectrum of possibilities from a bilateral advantage to no change of treatment planning to a bilateral overtreatment [3]. Finally, if preoperative MRI would be demonstrated to increase free disease or overall survival, overdiagnosis and overtreatment would change in simple diagnosis and treatment. But this is beyond the current strategic horizon.

A shared consensus was recently reached by an interdisciplinary working group promoted by the European Society of Breast Cancer Specialists (EUSOMA) in considering the following four indications for preoperative MRI [5]:

1. patients newly diagnosed with an invasive lobular cancer;
2. patients at high-risk for breast cancer;
3. patients under 60 years of age with discrepancy in size >1 cm between mammography and US with expected impact on treatment decision;
4. patients eligible for partial breast irradiation on the basis of clinical breast examination and conventional imaging.

My personal view is that other two groups of women could be considered for preoperative MRI [3]:

5. women with mammographically heterogeneously or extremely dense breast; and

6. women with a multifocal or multicentric or bilateral cancer already diagnosed at conventional imaging.

The indication concerning invasive lobular cancer is in agreement with the study recently published by Mann et al in this journal [6]. With the limitation of a retrospective design, they demonstrated a significant reduction in the re-excision rate in the women who did undergo preoperative MRI (9%) versus those who did not (27%) without any increase either of mastectomy rate (48% versus 59%) or mean time to final pathology (40 versus 44 days, respectively) [6]. Thus, the paradigm is inverted showing the possibility to use MRI preoperatively with the aim to reduce the aggressiveness of surgical treatment.

However, all nonrandomized studies on this matter have intrinsic bias and limitations. This is true also for other retrospective studies which reported results in favor [7] or partially in favor [8] of preoperative MRI as well as for retrospective studies which reported results against [9-11] preoperative MRI. Two interesting small studies from a Japanese group showed important advantages of a surgical approach precisely tailored on the basis of preoperative MRI performed in supine surgical position. In one non randomized study [12], they demonstrated significantly reduced excision area and additional excision rate compared to US-guided dye application in the surgical treatment of invasive ductal carcinomas after neoadjuvant chemotherapy. In a second study [13], they randomized 52 patients with localized ductal carcinoma in situ: 24 of them received breast conserving surgery (BCS) using projection and reproduction techniques of surgical-position breast MRI; 28 of them underwent conventional prone-position breast MRI and BCS using mammography-guided hookwire. Average volume of the pathologic specimens in the new technique group was significantly smaller than that in the conventional BCS group (27.5 cm³ versus 57.6 cm³) while the positive margin rate was significantly lower (12.5% versus 39.3%, respectively). These experiences indicate that a precise MRI-based tailored BCS is the way for the future, as is shown also by a recent research on US navigation of breast MRI volumes acquired in supine position [14]. The key point is the translation of a three-dimensional (3D) information from the MR room to the operating theatre.

Preoperative MRI is a typical matter which should be solved by means of large RCTs. This is due to the fact that this use of breast MRI is not for a simple diagnosis (we are already aware of the index cancer). It may seem a paradox, but preoperative MRI is a kind of screening looking for other cancers than the index one. If the results of preoperative MRI impact on treatment, this diagnostic tool should be regarded as a therapy.

Few months ago the results of the COMICE trial came to publication [15]. The investigators enrolled more than 1,800 women with a newly diagnosed breast cancer randomized into two groups, with (n=816) and without (n=807) preoperative MRI. The reoperation rate (primary end-point) was 19% in both groups; the total mastectomy rate was 13% versus 9%, respectively. These results are important. They show the possible consequences of a suboptimal use of preoperative MRI. Several limitations of the study have been already discussed [16]. Importantly, the investigators faced a very slow enrollment by a large number of centers, with many of them enrolling less than 4-5 patients/year, only 2-3 randomized to MRI. The results show no

systematic use of needle biopsy and localization (including MR-guided procedures), with many cases of changed treatment due to unverified false positive findings. Regarding the initial surgical treatment in the MRI arm, the investigators say: “Of the 58 patients who underwent a mastectomy, 32 had an additional biopsy, 11 did not have a biopsy, and data were unavailable for the remaining 15 patients. Of the 16 patients who underwent an avoidable mastectomy, three did and six did not have a biopsy, and data are missing for the remaining seven patients” [15].

In this large RCT, the management of MRI additional findings found has been a key point. In this perspective, an expert use of second look targeted US is essential. This requires not only to be familiar with (supine) breast US but also with (prone) breast MRI as well as with the different lesion location at the two imaging techniques. Six recent studies reported percentages of US correlate for initially MR-detected findings variable from 46% to 82% [17-22]. Pooling their results, we have a total of 1,208 findings, 759 of them with a US correlate (63%). While the mean malignancy rate is 27% (range, 22%-29%), that of lesions with a US-correlate is 32%, (range, 27%-36%), and that of lesion without a US correlate is 18% (range, 10%-28%), showing a variability of results probably depending also on the specific experience and expertise for targeted US examination. Thus, MR-guided procedures are necessary only in a minority of cases, those where targeted US fails to find the MRI finding. However, the rate of malignancy of MRI findings without US correlate does not go below 10% also when the rate of US detection reaches 82% [17]

In this context, in the current issue of this journal, Lotte E. Elshof et al [23] report their experience on 690 consecutive patients with 698 pathology-proven index cancers planned for BCT based on clinical examination and conventional imaging who underwent preoperative breast MRI. MRI additional findings were prospectively managed without using MR-guided procedures. They defined additional findings located within a 3-cm 3D space including the index tumor as “multifocal”, those located outside that space as “multicentric”, those in the other breast as contralateral. Multifocal findings were not sent to targeted US and these findings typically led to BCS with larger excision to include the additional findings. Multicentric and contralateral findings were sent to targeted US. If they were found, needle biopsy was performed. If malignant disease was confirmed over a region too large to allow cosmetically acceptable conserving treatment, mastectomy was planned. The key rule of this approach is as follows: *If pathology proof could not be obtained (i.e., if targeted US failed to find the lesion), the therapy plan was not changed and follow-up was advised.*

At preoperative MRI, 141 additional findings were detected in 121/690 patients (18%). MRI additional findings without pathology proof – named unidentified bright objects (UBOs) by the authors – were found in 81/690 patients (12%). Importantly, of 141 additional findings, 44 multicentric and contralateral findings (31%) were followed up at least with conventional imaging (40 of them being MRI BI-RADS 3). The median follow up was 55 months (range, 22-103 months). In none of the patients these findings turned out to be detected as suspicious and confirmed to be malignant (only four additional findings were lost to

follow-up). Multifocal lesions were significantly more often malignant than multicentric and contralateral lesions.

This approach resulted in a rate of change of surgical planning inferior to 10%, composed of mastectomy for 24 breasts in 23/690 patients (3.3%), wider ipsilateral excisions in 40/690 patients (5.8%), and contralateral excision in 3/690 (0.4%). Of 40 wider ipsilateral excisions, 32 (80%) were pathologically confirmed after surgery.

This experience contains a golden rule and a possible misleading message, the latter already acknowledged by the authors. The *golden rule* is: Do not convert a BCS to mastectomy on the basis of MRI additional finding(s) not pathologically verified to be malignant. The possible *misleading message* is: The use of MR-guidance is not necessary in the preoperative setting.

To perform mastectomy or contralateral surgery for MRI additional findings without needle biopsy verification of malignancy should be regarded as malpractice. Moreover, MR-guided biopsy should be integrated in clinical practice as it was for needle biopsy under stereotactic guidance. A large experience in MR-guided biopsy is reported in the literature, including a large prospective multi-institutional study of 538 lesions [24], many recent single-center experiences [25-31], also evaluating problems and limitations [32-35], as well as the results of a multidisciplinary consensus meeting on the use of vacuum-assisted MR-guided biopsy [36]. The mean time needed for an MR-guided procedure is not different than that of a stereotactic biopsy in the film-screen era: about 45 minutes [37]. Using MR-guidance, the rate of unnecessary ipsilateral wider excisions reported by Elshof et al [23] – 8/40 (20%) – could have been drastically reduced. Also MR-guidance has limitations and some findings can be difficult or impossible to be reached. In these cases, the *golden rule* by Elshof et al should be applied. Notably, the application of this rule demonstrated that multicentric and contralateral findings without US correlate were not detected as malignant at follow up. However, we do not know how many of them were benign and how many of them were malignant but cured by radiation or systemic therapy.

The general recommendations of the EUSOMA interdisciplinary working group still appear to be valid:

1. irrespective of whether the clinical team routinely uses preoperative MRI or not, women newly diagnosed with breast cancer should always be informed of the potential risks and benefits of preoperative MRI if this is under consideration prior to therapy;
2. results of preoperative MRI should be interpreted taking into account clinical breast examination as well as mammography and US (whenever mammography and US are indicated);
3. MRI findings with impact on patient treatment should be verified by percutaneous biopsy whenever possible;

4. lesions visible on MRI alone require MR-guidance for needle biopsy with pathological assessment and, if needed, presurgical localization, implying the availability of specialized equipment and personnel;
5. the total treatment delay due to preoperative MRI and possible workup should be no longer than 1 month;
6. possible changes in therapeutic planning resulting from the findings of preoperative MRI should be decided by a multidisciplinary team [5].

At any rate, the use of preoperative MRI remains an open issue and high-quality research using the patient's outcome as primary end-point is still warranted.

References

1. Houssami N, Ciatto S, Macaskill P, et al (2008) Accuracy and surgical impact of magnetic resonance imaging in breast cancer staging: systematic review and meta-analysis in detection of multifocal and multicentric cancer. *J Clin Oncol* 26:3248–3258
2. Brennan ME, Houssami N, Lord S, et al (2009) Magnetic resonance imaging screening of the contralateral breast in women with newly diagnosed breast cancer: systematic review and meta-analysis of incremental cancer detection and impact on surgical management. *J Clin Oncol* 27(33):5640–5649
3. Sardanelli F (2010) Overview of the role of pre-operative breast MRI in the absence of evidence on patient outcomes. *Breast* 19:3–6
4. Solin LJ (2010) Counterview: Pre-operative breast MRI (magnetic resonance imaging) is not recommended for all patients with newly diagnosed breast cancer. *Breast* 19:7–9
5. Sardanelli F, Boetes C, Borisch B, et al (2010) Magnetic resonance imaging of the breast: recommendations from the EUSOMA working group. *Eur J Cancer* 46:1296–1316
6. Mann RM, Loo CE, Wobbes T, et al (2010) The impact of preoperative breast MRI on the re-excision rate in invasive lobular carcinoma of the breast. *Breast Cancer Res Treat* 119:415–422
7. Fischer U, Zachariae O, Baum F, von Heyden D, Funke M, Liersch T (2004) The influence of preoperative MRI of the breasts on recurrence rate in patients with breast cancer. *Eur Radiol* 14:1725–1731.
8. Pengel KE, Loo CE, Teertstra HJ, et al (2009) The impact of preoperative MRI on breast-conserving surgery of invasive cancer: a comparative cohort study. *Breast Cancer Res Treat* 116:161–169
9. Solin LJ, Orel SG, Hwang WT, Harris EE, Schnall MD (2008) Relationship of breast magnetic resonance imaging to outcome after breast-conservation treatment with radiation for women with early-stage invasive breast carcinoma or ductal carcinoma in situ. *J Clin Oncol* 26:386–391
10. Hwang N, Schiller DE, Crystal P, Maki E, McCready DR (2009) Magnetic resonance imaging in the planning of initial lumpectomy for invasive breast carcinoma: its effect on ipsilateral breast tumor recurrence after breast-conservation therapy. *Ann Surg Oncol* 16:3000–3009
11. Bleicher RJ, Ciocca RM, Egleston BL, et al (2009) Association of routine pretreatment magnetic resonance imaging with time to surgery, mastectomy rate, and margin status. *J Am Coll Surg* 209:180–187
12. Nakamura R, Nagashima T, Sakakibara M, et al (2008) Breast-conserving surgery using supine magnetic resonance imaging in breast cancer patients receiving neoadjuvant chemotherapy. *Breast* 17:245–251
13. Sakakibara M, Nagashima T, Sangai T, et al (2008) Breast-conserving surgery using projection and reproduction techniques of surgical-position breast MRI in patients with ductal carcinoma in situ of the breast. *J Am Coll Surg* 207:62–68
14. Alderliesten T, Loo C, Paape A, et al (2010) On the feasibility of MRI-guided navigation to demarcate breast cancer for breast-conserving surgery. *Med Phys* 37:2617–2626
15. Turnbull L, Brown S, Harvey I, et al (2010) Comparative effectiveness of MRI in breast cancer (COMICE) trial: a randomised controlled trial. *Lancet* 375:563–571
16. Morris EA (2010) Should we dispense with preoperative breast MRI? *Lancet* 375:528–530
17. Linda A, Zuiani C, Londero V, Bazzocchi M (2008) Outcome of initially only magnetic resonance mammography-detected findings with and without correlate at second-look sonography: distribution according to patient history of breast cancer and lesion size. *Breast* 17:5–57
18. Demartini WB, Eby PR, Peacock S, Lehman CD (2009) Utility of targeted sonography for breast lesions that were suspicious on MRI. *AJR Am J Roentgenol* 192:1128–1134

19. Destounis S, Arieno A, Somerville PA, et al (2009) Community-based practice experience of unsuspected breast magnetic resonance imaging abnormalities evaluated with second-look sonography. *J Ultrasound Med* 28:1337–1346
20. Meissnitzer M, Dershaw DD, Lee CH, Morris EA (2009) Targeted ultrasound of the breast in women with abnormal MRI findings for whom biopsy has been recommended. *AJR Am J Roentgenol* 193:1025–1029
21. Abe H, Schmidt RA, Shah RN, et al (2010) MR-directed (“Second-Look”) ultrasound examination for breast lesions detected initially on MRI: MR and sonographic findings. *AJR Am J Roentgenol* 194:370–377
22. Carbognin G, Girardi V, Calciolari C, et al (2010) Utility of second-look ultrasound in the management of incidental enhancing lesions detected by breast MR imaging. *Radiol Med* Jun 23 [Epub ahead of print]
23. Elshof LE, Rutgers EJ, Deurloo EE, et al (2010) A practical approach to manage additional lesions at preoperative breast MRI in patients eligible for breast conserving therapy: results. *Breast Cancer Res Treat* xx:xx-xx
24. Perlet C, Heywang-Kobrunner SH, Heinig A, et al (2006) Magnetic resonance-guided, vacuum-assisted breast biopsy: results from a European multicenter study of 538 lesions. *Cancer* 106:982–990
25. Tozaki M, Yamashiro N, Suzuki T, et al (2009) MR-guided vacuum-assisted breast biopsy: is it an essential technique? *Breast Cancer* 16:121–125
26. Han BK, Schnall MD, Orel SG, Rosen M (2008) Outcome of MRI-guided breast biopsy. *AJR Am J Roentgenol* 191:1798–1804
27. El Khouli RH, Macura KJ, Barker PB, et al (2009) MRI-guided vacuum-assisted breast biopsy: a phantom and patient evaluation of targeting accuracy. *J Magn Reson Imaging* 30:424–429
28. Li J, Dershaw DD, Lee CH, Kaplan J, Morris EA (2009) MRI follow-up after concordant, histologically benign diagnosis of breast lesions sampled by MRI-guided biopsy. *AJR Am J Roentgenol* 193:850–855
29. Friedman P, Enis S, Pinyard J (2009) Magnetic resonance imaging-guided vacuum-assisted breast biopsy: an initial experience in a community hospital. *Can Assoc Radiol J* 60:196–200
30. Malhaire C, El Khoury C, Thibault F, et al (2010) Vacuum-assisted biopsies under MR guidance: results of 72 procedures. *Eur Radiol* 20:1554–1562
31. Schrading S, Simon B, Braun M, Wardelmann E, Schild HH, Kuhl CK (2010) MRI-guided breast biopsy: influence of choice of vacuum biopsy system on the mode of biopsy of MRI-only suspicious breast lesions. *AJR Am J Roentgenol* 194:1650–1657
32. Meeuwis C, Peters NH, Mali WP, et al (2007) Targeting difficult accessible breast lesions: MRI-guided needle localization using a freehand technique in a 3.0 T closed bore magnet. *Eur J Radiol* 62: 283–288
33. Liberman L, Holland AE, Marjan D, et al (2007) Underestimation of atypical ductal hyperplasia at MRI-guided 9-gauge vacuum-assisted breast biopsy. *AJR Am J Roentgenol* 188:684–690
34. Lee JM, Kaplan JB, Murray MP, et al (2007) Underestimation of DCIS at MRI-guided vacuum-assisted breast biopsy. *AJR Am J Roentgenol* 189:468–474
35. Lee JM, Kaplan JB, Murray MP, et al (2007) Imaging histologic discordance at MRI-guided 9-gauge vacuum-assisted breast biopsy. *AJR Am J Roentgenol* 189:852–859
36. Heywang-Köbrunner SH, Sinnatamby R, et al; Consensus Group (2009) Interdisciplinary consensus on the uses and technique of MR-guided vacuum-assisted breast biopsy (VAB): results of a European consensus meeting. *Eur J Radiol* 72:289–294
37. Fischer U, Schwethelm L, Baum FT, Luftner-Nagel S, Teubner J (2009) Effort, accuracy and histology of MR-guided vacuum biopsy of suspicious breast lesions--retrospective evaluation after 389 interventions. *Rofo* 181:774–781