Estimate of intake of benzoic acid in the Belgian adult population

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2008-373.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	24-Feb-2009
Complete List of Authors:	Vandevijvere, Stefanie; Scientific Institute of Public Health, Epidemiology Andjelkovic, Mirjana; Scientific Institute of Public Health, Food De Wil, Marc; Scientific Institute of Public Health, Food Vinkx, Christine; Federal Service of Public Health, Food Chain and the Environment Huybrechts, Inge; Ghent University, Public health Van Loco, Joris; Scientific Institute of Public Health, Food van oyen, herman; Scientific Institute of Public Health, Epidemiology Goeyens, Leo; Scientific Institute of Public Health, Food
Methods/Techniques:	Chromatographic analysis, Chromatography - HPLC, Exposure assessment
Additives/Contaminants:	Additives general
Food Types:	Beverages, Confectionary, Fish and fish products, Snack products

SCHOLARONE™
Manuscripts

Estimate of intake of benzoic acid for the Belgian adult population

Vandevijvere S.¹, Andjelkovic M.², De Wil M.², Vinkx C.³, Huybrechts I.⁴, Van Loco J.², , Van Oyen H.¹, Goeyens, L.²

¹Scientific Institute of Public Health, Unit of Epidemiology, Brussels, Belgium

²Scientific Institute of Public Health, Unit of Food, Brussels, Belgium

³Federal Public Service of Health, Food Chain Safety and Environment, Brussels, Belgium

⁴Department of Public Health, Ghent University, Ghent, Belgium

Abstract

An exposure assessment was performed to estimate average daily benzoic acid intake for Belgian adults. Food consumption data were retrieved from the national food consumption survey. In a first step, individual food consumption data were multiplied with the maximum permitted use levels for benzoic acid per food group (Tier 2). In a second step, a label survey to identify the foods where benzoic acid is effectively used as an additive and a literature review of the possible occurrence of benzoic acid as a natural substance, were performed. With this information a refined list of foods was drafted for the quantification of benzoic acid, which was performed by an HPLC method, optimized and validated for this purpose. Individual food consumption data were then multiplied with the actual average concentrations of benzoic acid per food group (Tier 3). Usual intakes were calculated using the Nusser method. The mean benzoic acid intake was 1.58 mg/kg bw/day (Tier 2) and 1.25 mg/kg bw/day (Tier 3). In Tier 2, men exceeded the acceptable daily intake (ADI) of 5 mg/kg bw/day at the 99th percentile. The greatest contributors to the benzoic acid intake were soft drinks. Benzoic acid as a natural substance represents only a small percentage of the total intake. The results show that actual benzoic acid intake is very likely to be below the ADI. However, there is a need to collect national food consumption data for children as they might be more vulnerable for an excessive intake.

Introduction

Benzoic acid (E210) or its sodium (E211), potassium (E212) or calcium (E213) salt, is intentionally added to certain foods as a food preservative or may naturally be present in specific foods. Sodium benzoate is about 200 times more soluble than benzoic acid and is used preferably as one of the main antimicrobial preservatives in foods and beverages. Benzoic acid is used as an antiseptic, antimycotic and antipyretic agent (Suhr and Nielsen 2004).

The temporary acceptable daily intake (ADI) was set at 5 mg/kg body weight by the European Scientific Committee on Food in 1994. The Joint WHO/FAO Expert Committee on Food Additives (JECFA) set an ADI of 5 mg/kg body weight in 1996, which was a confirmation of the evaluation in 1994. Use of benzoic acid is authorised in the European Union in several food groups on the condition of respecting the maximum use levels (Directive 95/2/EC), which are all expressed as the free acid. The European Commission (Directive 95/2/EC) asks the Member States to monitor the consumption and use of food additives, in order to allow evaluation of the consumption trends as well as differences in consumption patterns among various Member States.

The Report from the European Commission on Dietary Food Additive Intake in the European Union (European Commission 2001) describes a harmonized, tiered method to estimate additive intakes. In Tier 1, the theoretical food consumption data are combined with the maximum permitted usage levels for the additive. This conservative estimate was done at the EU level and is not a task for the Member States, as no national data are used. Only for a limited list of additives, national estimates are necessary to refine the estimate for those additives for which in Tier 1 an exceeding of the ADI was observed. In the next step (Tier 2), actual national food consumption data are used and combined with maximum permitted usage levels. Such intake estimate will still overestimate the real intake, but can exclude additives which do not pose a major risk from the most difficult estimate in Tier 3: a combination of actual national food consumption data with actual usage levels of the additive.

A study on the intake of benzoic acid was chosen because of the following elements: (1) high intakes were estimated in other countries; (2) benzoic acid is much less reactive than sulphites or nitrites resulting in residual levels more close to usage levels; (3) feasibility to analyse; (4) benzoic acid is important in soft drinks for which consumption data are high and increasing; (5) there is a long list of food groups in which the additive is authorised and extensions of authorisations are from time to time requested.

In a former study in Belgium, a Tier 2 exposure assessment of benzoic acid was carried out in three different population groups: Flemish preschool children (n=697; age 2-6; 3 days estimated dietary

record), adolescents from a major city in Flanders, Ghent (n=341; age 14-18; 7-day estimated dietary record) and women of reproductive age from Flanders (n=641; age 18-40; 2-day estimated dietary record) (Bilau et al. 2008). The median estimated daily intake was respectively 2.0 mg/kg bw, 1.7 mg/kg bw and 1.92 mg/kg bw. In all age groups the greatest contributor to total benzoic acid intake was the group of non-alcoholic flavoured drinks. At respectively the 92nd percentile (children), the 97th percentile (adolescents) and the 94th percentile (women), the ADI was slightly exceeded. This study showed the need to perform a more refined estimate, using actual levels.

Besides being widely used as a food additive, benzoic acid occurs naturally in different fruits (Hegnaur 1992). Appreciable amounts have been detected in most berries (around 0.05%) (Budavari et al. 1996), in particular in cranberries where they are present in a range from 300 to 1300 mg/kg (Hegnaur 1966). Benzoic acid has also been detected in many animal products, including dairy products, as the result of the lactic acid fermentation (Sauh et al. 1990). It is also produced during the ripening of cheese by the propionic acid fermentation following the lactic acid fermentation (Thierry and Maillard 2002). Naturally occurring benzoic acid has also been found in honey (Steeg and Montag 1987), mushrooms (Beltran-Garcia 1997) and wine (Matejcek et al. 2003). Further, benzoic acid can be used as a plasticizer in the manufacturing of plastic packing materials for yoghurt, butter or margarine (Vermeiren et al. 2002). The latter source of contamination with benzoic acid will not be further discussed in this paper.

The objective of this study was to perform a refined estimate of the benzoic acid intake in the Belgian adult population, using the data from the national food consumption survey of 2004. Another objective was to identify the food groups contributing most to the total benzoic acid intake in Belgium. The national food consumption data and maximum usage levels of benzoic acid were first combined to perform a Tier 2 exposure assessment. Analytical results were obtained for chosen food groups, and then, as far as possible, actual average levels of benzoic acid were combined with consumption data to perform a Tier 3 exposure assessment. Besides taking into account additive use, the refined intake estimate also took into account the natural presence of benzoic acid in certain foods to obtain the best estimate of real intake. In Tier2, as well as in Tier 3, intake of benzoic acid was compared with the ADI,

Materials and methods

Study design

For the Tier 2 exposure assessment, maximum usage levels of benzoic acid in different foods were as far as possible linked to the food consumption data of these specific foods, derived from the Belgian food consumption survey 2004). For some foods in the EU legislation, it is unlikely that they exist on

the Belgian market (for example grape juice for sacramental use or liquid tea concentrates). Other products were not found in the food consumption survey (e.g. food supplements). For the authorised *quantum satis* use as surface treatment of dried meat products, a Tier 2 approach was not possible. The exclusion of some food groups could lead to an underestimation of the intake, but this was supposed to be very limited.

For the Tier 3 exposure assessment, first a food label survey was performed and then samples of different foods, in which benzoic acid is actually added or present as a natural substance, were taken from different supermarkets. In these samples concentrations of benzoic acid were determined with an optimized and validated analytical method. Benzoic acid concentrations in different foods were then linked to the consumption quantity of these foods, calculated from the data of the food consumption survey. For foods for which no analyses were performed (e.g. shrimps, crustaceans and molluscs, which were designated as a less important source of benzoic acid due to its less frequent intake), the maximum permitted level of benzoic acid was used.

Food consumption data

Consumption data from the national food consumption survey 2004 were used to perform the exposure assessments. Aims, design and methods of this survey are described elsewhere (De Vriese et al. 2005). The target population comprised all Belgian inhabitants of 15 years or older. The sample included 3245 participants randomly selected from the National Register, using a multi-stage stratified procedure.

Information on dietary intake was collected by two non-consecutive 24-hour recalls in combination with a food frequency questionnaire. During the 24-hour recall interviews the respondent reported the quantity of all foods and beverages consumed during the preceding day. The 24-hour recall was carried out using EPIC-SOFT software (Slimani and Valsta 2002). This program allows obtaining very detailed information about the foods consumed and the recipes used in a standardized way.

3083 participants completed two 24-hour recalls of which 1537 women and 1546 men. Participants were categorized into four age groups: 15-18 years (n=760), 19-59 years (n=830), 60-74 years (n=789) and 75 years or older (n=704).

Benzoic acid concentration of selected foods

Before the development of the sampling plan for the collection of the samples to be analysed, a food label survey was performed to identify the food products where benzoic acid is actually used as an additive. Five supermarket chains with good market share in Belgium were visited. Based on the regulation (Directive 95/2/EC), several food groups have been surveyed, 1245 items in total. Products

were selected according to the indications on the labels concerning addition of benzoic acid. Although this was not an absolutely exhaustive survey of all existing categories of food, it was attempted to cover the most important national and marketplace's brands. The final goal of the survey was to get an overview of the general manufacturing practices regarding benzoic acid application as an additive in Belgium.

Further, a literature review of the possible occurrence of benzoic acid as a natural substance was performed. After this first selection of foods, a refined list of foods was drafted for the quantification of benzoic acid. The food consumption survey 2004 was used to evaluate the various channels used for food provisions (supermarkets), the frequencies of consumption of different foods and the preparations of foods. The Tier 2 exposure assessment provided an indicative list of the most important foods contributing to benzoic acid intake in the Belgian population. The comparison of the Tier 2 acquired list with the refined list from the label survey and knowledge of foods in which benzoic acid may be present as a natural substance, permitted the development of the sampling plan for the collection of the food samples to be analysed.

A great effort was put into ensuring representative concentration data of benzoic acid in foods on the Belgian market. The sampling plan included two categories: food items with benzoic acid used as an additive and those where benzoic acid is likely to be present as natural substance. In total for both categories 58 samples (28 and 30, respectively to the category) were collected and analysed. The results of the analyses were grouped to ensure a clear data presentation. The grouping was performed according to the EU directive 95/2/EC as follows: desserts, confectionary, prepared salads, non emulsified sauces, and non-alcoholic drinks. Additionally, two food groups were included, namely dairy products (cheese, yoghurt, and dairy flavoured with fruits) and products based on fruits (berries, grapes, apples, pears and tropical fruits). Collection of samples was performed from August to September 2007 in 5 major supermarket chains with a good market share. These 5 food distributors are visited by approximately 75% of the Belgian population in search for their food provisioning (CRIOC - OIVO 2007). Prepared salads were additionally taken from sandwich delivery snack bars. If the analyses were not conducted on the same day, the sample was stored deep frozen for a maximum time of one month.

Extraction from and determination of benzoic acid in different matrices was done by a modified method described in ISO/FDIS 22855:2007(E) (International Organisation for Standardisation 2007) where benzoic acid was quantified by a high performance liquid chromatography method optimized and validated for this purpose. This proposed method is based on a formerly used method (Küppers and Jans 1988).

Sample preparation and extraction of benzoic acid was adjusted to the state (liquid or solid) of the sample. In this manner liquid samples were homogenised prior to opening of the package while carbonated drinks were degassed in an ultrasonic bath for 3-5 min. A sample (10ml) was transferred into a 50ml conical tube where 10ml of methanol-water (80:20, v:v) solution of 3-chlorobenzoic acid (75mg/l) was added. The solution pH was adjusted with a droplet of 36.5% HCl. In the following step the solution was vortex mixed and centrifuged for 5 minutes at 7000rpm. From this 1ml was used for further purification.

Solid samples (200g) were homogenised with an Ultra-Turrax mixer. Out of that 10g±0.1g was weighed and transferred into a 50ml conical tube. 10 ml of methanol-water (80:20, v:v) solution of 3-chlorobenzoic acid (75mg/l) was added. After adding a droplet of 36.5% HCl, the tube was closed and vortex mixed (non fatty solid partially hydrophilic samples, e.g. jams and gums) or mixed with the Ultra-Turrax (fatty solid hydrophobic samples, e.g. cheese, fish, and crustaceans) until dissolution into the aqueous phase was obtained. If necessary, the sample was placed in a warm (45-60°C) water bath for 30 min with frequent agitation. In the following step the solution was vortex mixed, centrifuged for 5 min at 7000 rpm and cooled to room temperature. From this 1ml was used for further purification.

All prepared samples were subjected to purification through SPE C18ec (Chromabond, 500mg, 3ml) which was conditioned by 1 cartridge volume methanol followed by 1 cartridge volume HCl 0.1M. Next, 1ml of the sample was applied (at the flow rate 0.4ml/min) and the cartridge was successively washed twice with 2ml of 0.1M HCl. and additionally washed with 1ml hexane where before and after this step the cartridge was left to dry for 3 or 1 min (respectively). The extract was eluted with 3ml methanol and concentrated under nitrogen at 40°C to approximately 1ml.

Finally the extracts were filtered through PTFE 0.45µm and injected into HPLC. HPLC analysis was conducted using an Agilent 1100 Series (Waldbronn, Germany) equipped with a quaternary pump, an auto sampler and DAD detector operated by ChemStation software (version B.01.03) A LiChrospher RP Select B (5µm; 250x4.6mm) HPLC column with Lichrospher RP Select B pre-column was used throughout the study.

The separation was obtained by gradient elution with three mobile phases. Mobile phase A (ammonium acetate aqueous solution, pH 3.9, 0.05M) was kept constant at 10% while mobile phase B (acetonitrile) and C (MiliQ water) were kept constant first 10 min: 25% B and 65% C; then the gradient changed 10-13min 60%B and 30%C and kept constant for 7 min; 20-21min: 25% B and

65% C and the system was equilibrated for 4 min. The flow rate was 1ml/min at 35°C. The detection was carried out by DAD detector at 235 and 255nm.

Identification of benzoic acid was done by comparing the retention time and peak shape to that of reference standard whereas quantification was performed according to the calibration curve ($y=1.48x+0.03$; $R^2=0.99$) plotted as the respond ratio of benzoic acid and 3-chlorobenzoic acid versus concentration of benzoic acid in the range 0-250mg/l for nine calibration points (0; 0.2; 0.5; 1; 5; 10; 25; 50; 100; 250mg/l).

The method was validated on 3 different matrices: sugared liquids, emulsified solids and dairy products (mayonnaise and plain yoghurt respectively). The limit of detection (LOD) and limit of quantification (LOQ) for benzoic acid were 2.5 and 5 mg/l respectively. The intra-day and inter-day precision of the method showed a coefficient of variation (%CV) between 0.25 and 16% for spiked samples (taking $\leq 10\%$ as acceptable limit) and the recovery of the method was found to be between 87 to 113%.. The precision of the method was dependent on the matrix and the concentration. For both non-alcoholic flavoured drinks (soft drinks) and dairy products (yoghurt) the lowest level (10 mg/l and 5 mg/kg respectively) resulted in higher %CV's (12-16 %) whereas CV's for determination of benzoic acid in fatty samples (mayonnaise) were low at all three levels.

Statistical analysis

Only respondents with two completed 24-hour recall interviews were included in the analyses (n= 3083; 1546 men and 1537 women).

The individual intake of benzoic acid from a certain food product was estimated by the following equation:

$$y_i (mg / kgbw / day) = \frac{c_i \times x_i}{bw_i} ,$$

where y_i is the intake of benzoic acid by individual i from a particular food (in mg per kg bodyweight per interview day), c_i is the concentration of benzoic acid in that food (mg per kg), x_i is the consumption of a certain food by individual i (kg) and bw_i is the self-reported body weight of individual i (kg). To estimate the total intake of benzoic acid per food group or per day, individual daily intakes of benzoic acid from different foods were added up.

As mentioned before, two approaches were used for the calculation of benzoic acid intake. In the Tier 2 approach, c_i represents the maximum allowed concentration in each food. The maximum permitted levels of benzoic acid used in different foods are listed in the European Directive 95/2/EC (Table 1) (

Directive 95/2/EC). In the Tier 3 approach, c_i represents the actual average concentration of benzoic acid that is observed in a particular food.

The usual intake distribution for benzoic acid was estimated with the Nusser method (Nusser et al. 1996) using the C-side software (Iowa State University 1996). Several statistical methods are available to estimate usual intake distributions with the correct mean, variance and skewness. These statistical procedures adjust for within-person or day-to-day variability. Of all different statistical procedures, the Nusser method (Nusser et al., 1996) is highly recommended because it eliminates the intra-individual variance and additionally transforms the data to obtain approximately normally distributed data. The method is suitable to estimate usual intake distributions in a population both for normally and non-normally distributed foods and nutrients. The usual intake distribution was weighted and adjusted for the age and sex distribution of the Belgian population and adjusted for day of the week and season.

Results

The label survey revealed that in biscuits, cakes and confectionary no benzoic acid is used on the Belgian market, although it is allowed. In the category of emulsified sauces the difference between regular and light sauces was remarkable. Of 55 light emulsified sauces, 19 contained benzoic acid (35%) whereas only 10% of the regular products had the additive indicated on the label. From the results it appeared that benzoic acid may more often be used in light sauces. Similarly, in relation to the fat percentage of the product, it was observed that benzoic acid was more often used in products containing less than 60% fat.

The concentration of benzoic acid in samples suspected to contain benzoic acid as a natural substance, was under official limits. Red and white wine, fruit juices and syrups, in particular those from berries, did not contain benzoic acid above the LOQ. Among dairy products, the highest amount of benzoic acid was detected in cottage cheese. The average amounts in yoghurts were around 25 mg/kg which was comparable to the amounts detected in different yoghurts collected from stores in Belgium in 1982 (Guyot et al. 1998; Sieber et al. 1995; Van Renterghem 1982). On the other hand, the amount of the additive in all analysed soft drinks (coke and lemonade) was between 100 mg/l and the maximum permitted level of 150 mg/l. As a result, the use of actual levels instead of maximum levels did not reduce the intake estimate dramatically (Table 1). Intake estimates of consumption of food groups can be found in Table 2.

The Tier 2 approach, which is the most conservative approach, revealed a mean usual daily benzoic acid intake of 1.58 mg/kg bw/day (Table 3; Figure 1). At the 97.5th percentile, the usual intake of benzoic acid was 3.97 mg/kg bw/day which was still below the ADI. The most important contributor

to the intake of benzoic acid was the group of non-alcoholic flavoured beverages (31.07%). Other important contributors at the upper end of the intake distribution were non-emulsified sauces (16.16%), vegetables in vinegar, brine or oil (13.24%), prepared salads (8.08%), semi-preserved fish products (6.90%) and shrimps, molluscs and crustaceans (6.84%).

The Tier 3 approach is more accurate than the Tier 2 since it takes into account the actual average concentrations of benzoic acid in different foods and it also considers the presence of benzoic acid as a natural substance. Nevertheless the results of Tier 3 were comparable to those of Tier 2 (Table 4, Figure 2), where no exceeding of the ADI was observed. Non-alcoholic flavoured drinks (34.28%) like coke and lemonade were again the most important contributors to the intake of benzoic acid. Benzoic acid present as a natural substance in food only contributed to a very small percentage (2.7%) to the total benzoic acid intake (Table 4).

It was observed that the benzoic acid intake in men was higher than in women (Figure 1; Figure 2). Mean usual benzoic acid intake in men (Tier 3) was 1.41 mg/kg bw/day. For women mean usual benzoic acid intake (Tier 3) was 1.22 mg/kg bw/day. Men consumed more non-alcoholic flavoured beverages than women (288 ± 307 g/day versus 177 ± 198 g/day for women), which were the main contributors to the benzoic acid intake. Men only exceeded the acceptable daily intake at the 99th percentile for the Tier 2 estimation at an intake of 5.18 mg/kg bw/day (while women only reached 4.22 mg/kg bw/day at the 99th percentile). However, men did not exceed the ADI at the 99th percentile for the Tier 3 estimation where the intake was 4.64 mg/kg bw/day (while 4.10 mg/kg bw/day for women).

Discussion

The food label survey showed that in some products, like confectionary, although allowed, no benzoic acid is used in Belgium. Remarkable was that benzoic acid is used more frequently in light sauces than in regular sauces. A possible explanation could be the higher water content of these products which in turn require higher levels of benzoic acid to exert a sufficient preservative effect. The analyses of the food samples for quantification of benzoic acid showed that no products with higher levels of benzoic acid than allowed were present in selected food stores in Belgium.

In the most conservative method (Tier 2), both mean usual benzoic acid intake and the usual benzoic acid intake at the higher percentiles were found to be lower than the ADI. When a Tier 3 exposure assessment was performed, the mean usual intake of benzoic acid and the usual intake of benzoic acid at the higher percentiles were also lower than the ADI, which means there is no direct concern of an excessive benzoic acid intake in the Belgian adult population. Benzoic acid intake in this population

may be regarded as safe and does not need further consideration if dietary patterns do not change tremendously. However, there are some minor concerns which may be raised.

The most important contributor to the benzoic acid intake in the Belgian adult population is the food group of non-alcoholic flavoured drinks. People might be loyal to one brand and hence not consume an average concentration. Using maximum levels in the calculation might better protect consumers. As explained above, some food groups in which use of benzoic acid is authorised were not taken into account. Drugs and cosmetics were not taken into account either.

In the Belgian population 15 years and older, there is no direct threat to consumers regarding benzoic acid intake. However, there may be special groups which are at risk for exceeding the ADI. First, men seem to have a higher intake of benzoic acid than women, due to a higher intake of soft drinks. Men exceed the ADI at the 99th percentile in a Tier 2 exposure assessment. Second, soft drinks are usually related to children's preferences and children are particularly vulnerable to food safety issues. On the other hand sodium benzoate will be phased out as additive in soft drinks, resulting in lower intake of benzoic acid in the near future (Nyman et al. 2008). Moreover it is remarkable that no benzoic acid is used in Belgium in cakes, biscuits and confectionary, food groups which are frequently consumed by children.

Unfortunately, the lack of national data about children's consumption habits posed a limitation to the study. The EU Scientific committee on Food has recommended that special attention should be given to additive intake in children, since the intake is expressed on a body weight basis and may be markedly higher than that of adults.

Benzoic acid has also been linked to hyperactivity in children when mixed with other artificial food colours (McCann et al., 2007). In Flemish preschool children, the calculated (Tier 2) intake estimate was only slightly higher than in adolescents or adult women (Bilau et al., 2008).

The intake as well as the pattern of benzoic acid in the Belgian population was comparable with that in some other countries but somewhat higher. The average intake of benzoic acid in Australia and New Zealand was between 10 and 15% of the ADI for the consumers older than 2 years whereas for the 95th percentile the intake was between 50% and 65% of the ADI. The most exposed population group were children between 2 and 12 years. In this age group, benzoic acid exposure was estimated to be about 1.2-1.4 times the ADI (with higher intake for boys than for girls). The major intake for the average population came from soft drinks (55%), regular cordial (21%) and orange juice (18%) (Food Standards Australia New Zealand 2005).

An intake survey of different food additives in Finland (Pentilla et al. 1988) identified fruit and berry juices, soft drinks and semi-preserved fish as the major sources of intake of benzoic acid which was estimated to be 40 mg/day representing 13% of the acceptable daily intake. Those results were obtained from the food balance sheets of the Government statistical office and actual concentration in 1400 foods products was used.

At last, there are some important methodological limitations that should be taken into account when interpreting the findings of this study. First, it is very difficult to exactly match the food groups or items defined in the legislation with the food groups or items defined in the food consumption survey. Secondly, some potentially important food groups, for example liquid eggs used in industrial processes, to which benzoic acid is commonly added as a preservative, were neglected in this study. It was not possible to disaggregate food consumption data up to this level. Further, benzoic acid in seasonings and condiments and in grape juice fermented for sacramental juice was neglected because of an assumed low consumption and a difficult match with food consumption data.

In conclusion, the intake of benzoic acid in the Belgian adult population does not imply an immediate threat to consumers. Nevertheless, the intake is close to the accepted daily intake level and this maximum level may be exceeded if dietary habits change for example when soft drink consumption increases or when authorisations are extended. No data about children are available at this moment while children are considered the most vulnerable population group. In the future, a food consumption monitoring system should be organised in Belgium to be able to evaluate additive intakes of different population groups on a regular basis.

Acknowledgements

The authors declare not having any conflicts of interest. The authors acknowledge the work of the field work team of the food consumption survey: Stéphanie De Vriese and Michel Moreau and the dietary support from Mia Bellemans, Mieke De Maeyer, Kadija El Moumni and Davy Van Steenkiste. Special thanks go to the respondents and the dieticians during the field work. The members of the steering committee (Leen Rasschaert and Charles Cremer) are greatly acknowledged for their scientific advice. The survey was funded by the Federal Ministry of Health, Food Chain and Environment.

References

1995. European parliament and council directive 95/2/EC of 20 February 1995 on food additives other than colours and sweeteners.

- 403 Beltran-Garcia MJ. 1997. Volatile compounds secreted by the Oyster Mushroom (*Pleurotus*
404 *ostreatus*) and their antimicrobial activities. *Journal of Agricultural and Food Chemistry* 45:
405 4049-4052.
- 406 Bilau M, Matthys C, Vinkx C, de Henauw S. 2008. Intake assessment for benzoates in
407 different subgroups of the Flemish population. *Food Chem Toxicol* 46: 717-723.
- 408 Budavari S, O'Neil MJ, Smith A, Heckelman PE, Kinneary JF. 1996. The Merck index -- an
409 encyclopedia of chemicals, drugs and biologicals. Merck&Co Inc., New York.
- 410 CRIOC - OIVO. 2007. Canaux de distribution des produits alimentaires. Centre de Recherche
411 et d'Information des organisations de consommateurs (CRIOC)
- 412 De Vriese S, Debacker G, de Henauw S, Huybrechts I, Kornitzer M, Leveque A, Moreau M,
413 Van Oyen H. 2005. The Belgian food consumption survey: aims, design and methods. *Arch*
414 *Public Health* 63: 1-16.
- 415 European Commission. 2001. Report from the Commission on dietary food additive intake in
416 the European Union. European Commission, Brussels.
- 417 Food Standards Australia New Zealand. 2005. The 21st Australian Total Diet Study. A total
418 diet study of sulphites, benzoates and sorbates. Food Standards Australia New Zealand
- 419 Guyot C, Bousseta A, Scheirman V, V, Collin S. 1998. Floral Origin Markers of Chestnut and
420 Lime Tree Honeys. *J Agric Food Chem* 46: 625-633.
- 421 Hegnauer R. 1966. *Chemotaxonomie der Pflanzen*.
- 422 Hegnauer R. 1992. *Benzoësäure*.
- 423 International Organisation for Standardisation. 2007. ISO/FDIS 22855:2007 (E) Fruit and
424 vegetable products -- Determination of benzoic acid and sorbic acid concentrations -- High
425 performance liquid chromatography method.
- 426 Iowa State University. 1996. A user's guide to C-SIDE. Software for Intake Distribution
427 Estimation. Department of Statistics and Center for Agricultural and Rural Development;
428 Iowa State University
- 429 Küppers F, Jans J. 1988. Reverse-phase liquid chromatographic determination of benzoic and
430 sorbic acid acids in fresh cheese. *Journal of AOAC* 71: 1068-1071.
- 431 Matejcek D, Klejdus B, Mikes O, Sterbova D, Kuban V. 2003. Application of solid-phase
432 extraction for determination of phenolic compounds in barrique wines. *Anal Bioanal Chem*
433 377: 340-345.
- 434 Nusser SM, Carriquiry AL, Dodd KW, Fuller WA. 1996. A semiparametric transformation
435 approach to estimating usual daily intake distributions. *J Am Stat Assoc* 91: 1440-1449.
- 436 Nyman PJ, Diachenko GW, Perfetti GA, McNeal TP, Hiatt MH, Morehouse KM. 2008.
437 Survey results of benzene in soft drinks and other beverages by headspace gas
438 chromatography/mass spectrometry. *J Agric Food Chem* 56: 571-576.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

439 Pentilla L, Salminen S, Niemi E. 1988. Estimates on the intake of food additives in Finland. *Z*
440 *Lebensm Unters Forsch* 186: 11-15.

441 Sauh MA, Sandine WE, Ayres JW. 1990. Inhibitory effects of microgard on yoghurt and
442 cottage cheese spoilage organisms. *Journal of Dairy Science* 73: 387-393.

443 Sieber R, Bütikofer U, Bosset JO. 1995. Benzoic acid as a natural compound in cultured dairy
444 products and cheese. *International Dairy Journal* 5: 227-246.

445 Slimani N, Valsta L. 2002. Perspectives of using the EPIC-SOFT programme in the context
446 of pan- European nutritional monitoring surveys: methodological and practical implications.
447 *Eur J Clin Nutr* 56 Suppl 2: S63-S74.

448 Steeg E, Montag A. 1987. Nachweis aromatischer Carbonsäuren in Honig. *Zeitschrift für*
449 *Lebensmittel-Untersuchung und -Forschung* 184: 17-19.

450 Suhr KI, Nielsen PV. 2004. Effect of weak acid preservatives on growth of bakery product
451 spoilage fungi at different water activities and pH values. *Int J Food Microbiol* 95: 67-78.

452 Thierry A, Maillard M. 2002. Production of cheese flavour compounds derived from amino
453 acid catabolism by *Propionibacterium freudenreichii*. *Lait* 82: 17-32.

454 Van Renterghem R. 1982. Het natuurlijk benzoëzuurgehalte van yoghurt. *Archives Belges* 40:
455 530-538.

456 Vermeiren L, Devlieghere F, Debevere J. 2002. Effectiveness of some recent antimicrobial
457 packaging concepts. *Food additives and contaminants* 19: 163-171.

458
459
459

Table 1 Maximum permitted level (mg/kg or mg/l) and average concentration level (SD) (mg/kg or mg/l) of benzoic acid in different food groups

FOOD GROUP	Maximum permitted level	Average concentration level	SD	number of samples
<i>use of benzoic acid as an additive</i>				
non-alcoholic flavoured drinks	150	120.5	17.0	10
coca cola	150	128.3	21.7	3
lemonade	150	112.1	10.3	5
ice-tea	150	114.1		1
aromatised water	150	145.9		1
prepared salads	1500	705.0	205.5	10
low-sugar marmalades	500	*		0
confectionary	1500	**		0
chewing gum	1500	**		0
non heat-treated dairy-based desserts	300	**		0
emulsified sauces (>60% fat)	500	**		0
emulsified sauces (<60% fat)	1000	819.0	185.2	5
light sauces	1000	844.6	7.0	2
light mayonnaise	1000	498.6	99.2	2
dressings	1000	768.6		1
non-emulsified sauces	1000	594.3	228.6	3
salted dried fish	200	*		0
semi-preserved fish products	2000	*		0
shrimps, crustaceans and molluscs	1000 [#]	*		0
vegetables in vinegar/oil/brine	2000	*		0
olives and olive-based preparations	500	*		0
liquid soups and broth (not canned)	500	*		0
cooked red beet	2000	*		0
candied fruit	1000	*		0
mustard	1000	*		0
<i>presence of benzoic acid as a natural substance</i>				
wines		2.5 ^{&}		3
fruit juices		3.8 [§]		3
yoghurts total				10
yoghurts with berries		25.7	6.5	7
yoghurts nature		24.1	0.2	3
butter		7.5 [§]		3
syrups		3.8 [§]		6
hard (Gouda) cheeses		13.4	0.5	2
cottage cheese		31.9 [°]		3

SD standard deviation

*No analyses were performed because the Tier 2 showed that the contribution of these products to the total benzoic acid intake was low

** No benzoic acid was found on the labels of these products and consequently no analyses were performed

[#] 6000 mg/kg for grey shrimps

& lower bound method used (3 samples below LOD)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

§ middle bound method used (for fruit juices: 2 samples below LOD and 1 sample below LOQ, for syrups 3 samples below LOD and 3 samples below LOQ)
\$ Upper bound method used (3 samples below LOQ)
° Only one sample above LOQ

For Peer Review Only

Table 2 The mean (SD; P50; P97.5; P99) usual consumption (g/day or ml/day) and the number of consumers (n) of the food groups in which addition of benzoic acid is permitted or in which benzoic acid is present as a natural substance (Belgian food consumption survey, 2004)

FOOD GROUP	Intake (g/day or ml/day)						
	mean	SD	P50	P95	P97.5	P99	n
<i>use of benzoic acid as an additive</i>							
non-alcoholic flavoured drinks	228	251	164	719	884	1107	1540
coca cola	140	201	23	545	673	846	975
lemonade	51	97	0	242	324	446	603
ice-tea	14	45	0	119	169	223	203
aromatised water	23	82	0	174	277	417	211
prepared salads	6	9	0	25	32	41	373
low-sugar marmalades	0.6	3.1	0.0	0.9	8.9	16.8	121
confectionary	1.2	3.6	0.0	7.7	11.5	17.4	256
chewing gum*	0.03	0.87	0.00	0.00	0.00	0.00	15
non heat-treated dairy-based desserts	16	17	11	49	60	77	166
emulsified sauces (>60% fat)	5.8	4.6	4.8	14.8	17.6	21.4	1139
emulsified sauces (<60% fat)	3.9	3.6	3.1	10.9	13.3	16.8	924
light sauces	1.4	2.8	0.0	7.0	9.4	12.8	273
light mayonnaise	0.6	1.1	0.0	2.8	3.6	4.7	254
dressings	1.9	2.3	1.2	6.4	7.6	9.3	503
non-emulsified sauces	17.2	11.1	15.2	38.4	44.5	52.5	1317
salted dried fish*	0.1	3.1	0.0	0.0	0.0	0.0	4
semi-preserved fish products	3.8	3.1	3.1	9.9	11.8	14.4	411
shrimps, crustaceans and molluscs	5.3	5.6	3.9	16.5	20.0	24.7	404
vegetables in vinegar/oil/brine	7.6	8.1	5.0	23.9	30.2	39.0	579
olives and olive-based preparations	0.7	1.8	0.0	4.4	6.3	8.9	99
liquid soups and broth (not canned)*	3.1	35.0	0.0	0.0	0.0	0.0	54
cooked red beet*	0.3	5.8	0.0	0.0	0.0	0.0	29
candied fruit*	0.001	0.038	0.000	0.000	0.000	0.000	1
mustard	0.3	0.2	0.2	0.7	0.9	1.2	289
<i>Presence of benzoic acid as a natural substance</i>							
wines	67	95	20	260	320	402	885
fruit juices	59	79	29	215	269	344	1072
yoghurts: total	39	49	23	129	162	213	943
yoghurts: with fruit	25	39	0	100	125	168	588
yoghurt: plain	14	30	0	78	101	131	440
butter	7	12	1	33	43	58	1164
syrups	3	17	0	20	41	77	124
hard (gouda) cheeses	14	11	12	36	42	51	1568
cottage cheeses	11	26	0	67	92	124	429

The average usual intakes are weighted and adjusted for the age and sex distribution of the Belgian population 2004 and adjusted for interview day and season

The total number of consumers (n) is 3083

* Average intake of the two interview days in stead of usual intake (Nusser method could not be applied due to low number of observations), not adjusted for the age and sex distribution of the Belgian population, neither for interview day and season

Table 3. The mean estimated usual intake (mg/kg bw/day) (SD, % contribution of food groups to the benzoic acid intake, P50, P95, P97.5,P99, % contribution of food groups to the ADI) of benzoic acid in the Belgian population via intake of different food groups, the Tier 2 approach

FOOD GROUP	mean	SD	%	P50	P95	P97.5	P99	% from ADI
non-alcoholic flavoured drinks	0.50	0.31	31.1	0.35	1.59	1.96	2.47	10.0
coca cola	0.30	0.44	18.6	0.05	1.18	1.46	1.85	6.0
lemonade	0.11	0.22	6.8	0.00	0.54	0.74	1.03	2.2
ice-tea	0.03	0.10	1.9	0.00	0.27	0.38	0.50	0.6
aromatised water	0.05	0.19	3.1	0.00	0.40	0.64	0.97	1.0
prepared salads	0.13	0.20	8.1	0.00	0.54	0.68	0.88	2.6
low-sugar marmalades	0.004	0.023	0.3	0.000	0.007	0.063	0.121	0.1
confectionary	0.07	0.08	4.4	0.00	0.17	0.26	0.39	1.4
chewing gum*	0.00073	0.02321	0.05	0.00000	0.00000	0.00000	0.00000	0.02
non heat-treated dairy-based desserts	0.0690	0.0760	4.3	0.0450	0.2190	0.2750	0.3530	1.4
emulsified sauces (>60% fat)	0.0420	0.0340	2.6	0.0340	0.1080	0.1300	0.1600	0.8
emulsified sauces (<60% fat)	0.0570	0.0530	3.5	0.0440	0.1590	0.1950	0.2440	1.1
light sauces	0.0190	0.0370	1.2	0.0000	0.0950	0.1250	0.1680	0.4
light mayonnaise	0.0089	0.0143	0.6	0.0000	0.0380	0.0485	0.0630	0.2
dressings	0.0280	0.0340	1.7	0.0170	0.0950	0.1140	0.1390	0.6
non-emulsified sauces	0.2600	0.1700	16.2	0.2240	0.5940	0.6970	0.8330	5.2
salted dried fish*	0.00018	0.00710	0.01	0.00000	0.00000	0.00000	0.00000	0.004
semi-preserved fish products	0.111	0.096	6.9	0.086	0.302	0.365	0.449	2.2
shrimps, crustaceans and molluscs	0.110	0.150	6.8	0.066	0.404	0.521	0.689	2.2
vegetables in vinegar/oil/brine	0.213	0.220	13.2	0.147	0.657	0.823	1.050	4.3
olives and olive-based preparations	0.0050	0.0122	0.3	0.0000	0.0304	0.0423	0.0585	0.1
liquid soups and broth (not canned)*	0.023	0.280	1.4	0.000	0.000	0.000	0.000	0.5
cooked red beet*	0.010	0.170	0.6	0.000	0.000	0.000	0.000	0.2
candied fruit*	5.449E-06	0.00043	0.0	0.00000	0.00000	0.00000	0.00000	0.0
mustard	0.0039	0.0037	0.2	0.0029	0.0104	0.0137	0.0191	0.1
TOTAL (% from ADI)	1.58 (31.6)	0.95	-	1.39 (27.8)	3.42 (64.8)	3.97 (79.4)	4.72 (94.4)	-

Deleted: ng
Deleted: estimated usual intake (mg/kg bw/day)

The average usual intakes are weighted and adjusted for the age and sex distribution of the Belgian population 2004 and adjusted for interview day and season

The total number of consumers (n) is 3083

*Average in stead of usual intake (due to low number of observations), not adjusted for the age and sex distribution of the Belgian population, neither for interview day and season

Deleted: consumption days (N) is 6166 and the total number of

For Peer Review Only

Table 4. The mean estimated usual benzoic acid intake (mg/kg bw/day) (SD, P50, % contribution of food groups to benzoic acid intake, P95, P97.5, P99, % contribution of food groups to ADI) as an additive (97.3%) and as a natural substance (2.7%) in the Belgian population via intake of different food groups, the Tier 3 approach

FOOD GROUP	mean	SD	%	P50	P95	P97.5	P99	% from ADI
<i>Benzoic acid as additive</i>								
non-alcoholic flavoured drinks	0.40	0.53	34.3	0.28	1.28	1.58	1.99	8.0
coca cola	0.240	0.350	20.6	0.040	0.950	1.180	1.490	4.8
lemonade	0.090	0.180	7.7	0.000	0.433	0.591	0.828	1.8
ice-tea	0.025	0.081	2.1	0.000	0.213	0.303	0.404	0.5
aromatised water	0.043	0.150	3.7	0.000	0.320	0.510	0.780	0.9
prepared salads	0.059	0.095	5.1	0.000	0.252	0.320	0.414	1.2
low-sugar marmalades	0.004	0.023	0.3	0.000	0.007	0.063	0.121	0.1
emulsified sauces (>60% fat)*	0.006	0.015	0.5	0.000	0.038	0.054	0.075	0.1
emulsified sauces (<60% fat)	0.044	0.042	3.8	0.034	0.125	0.154	0.194	0.9
-light sauces	0.0150	0.0300	1.3	0.0000	0.0780	0.1030	0.1370	0.3
-light mayonnaise	0.0044	0.0071	0.4	0.0000	0.0190	0.0242	0.0314	0.1
-dressings	0.0230	0.0270	2.0	0.0140	0.0770	0.0930	0.1140	0.5
non-emulsified sauces	0.1540	0.1000	13.2	0.1330	0.3530	0.4140	0.4950	3.1
salted dried fish**	0.00018	0.00710	0.02	0.00000	0.00000	0.00000	0.00000	0.0
semi-preserved fish products	0.111	0.096	9.5	0.086	0.302	0.365	0.449	2.2
Shrimps, crustaceans, molluscs	0.110	0.150	9.4	0.066	0.404	0.521	0.689	2.2
vegetables in vinegar/oil/brine	0.213	0.220	18.3	0.147	0.657	0.823	1.050	4.3
olives and olive-based preparations	0.0050	0.0122	0.4	0.0000	0.0304	0.0423	0.0585	0.1
soups and broth (not canned)**	0.023	0.280	2.0	0.000	0.000	0.000	0.000	0.5
cooked red beet**	0.0103	0.1700	0.9	0.0000	0.0000	0.0000	0.0000	0.2
candied fruit**	0.00001	0.00043	0.0	0.00000	0.00000	0.00000	0.00000	0.0
mustard	0.0039	0.0037	0.3	0.0029	0.0104	0.0137	0.0191	0.1

Deleted: ng

Deleted: ng

Deleted: estimated usual intake (mg/kg bw/day) of benzoic acid

Benzoic acid as natural substance

wines	0.0023	0.0033	0.2	0.0007	0.0091	0.0113	0.0141	0.05
fruit juices	0.0033	0.0045	0.3	0.0015	0.0121	0.0155	0.0201	0.07
yoghurts total	0.0146	0.0191	1.3	0.0085	0.0490	0.0620	0.0833	0.3
-yoghurts with fruit	0.0095	0.0161	0.8	0.0000	0.0380	0.0487	0.0679	0.2
-yoghurts plane	0.0051	0.0108	0.4	0.0000	0.0283	0.0367	0.0483	0.1
butter	0.00081	0.00140	0.1	0.00012	0.00359	0.00481	0.00656	0.02
syrups	0.000048	0.000190	0.0	0.000000	0.000360	0.000610	0.000960	0.0
hard (gouda) cheeses	0.0028	0.0021	0.2	0.0023	0.0068	0.0080	0.0096	0.06
cottage cheese	0.0049	0.0119	0.4	0.0000	0.0306	0.0415	0.0559	0.01
TOTAL (% from ADI)	1.25 (25.)	0.83	-	1.05 (21)	2.87 (57)	3.4 (68)	4.11(82)	-

* The average usual intakes are weighted and adjusted for the age and sex distribution of the Belgian population 2004 and adjusted for interview day and season

The total number of consumers (n) is 3083

** Average in stead of usual intake (The Nusser method could not be applied due to low number of observations), not adjusted for the age and sex distribution of the Belgian population, neither for interview day and season;

*without mayonnaise because no benzoic acid was found on the labels

Confectionary, chewing gum and non heat-treated dairy-based desserts were set at 0 mg/kg because no benzoic acid was found on the labels

Deleted: ¶

Deleted: ¶
wines ... [1]

Formatted: Font: 10 pt, Bold

Formatted Table

Deleted: number of consumption days (N) is 6166 and the total number

Deleted: as

Deleted: 0

Deleted: there was not found any benzoic acid on the labels

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Page 2: [1] Deleted		Stéfanie Vandevijvere		1/24/2009 3:22:00 PM			
wines	0.0023	0.0033	0.20	0.0000	0.0007	0.0037	0
fruit juices	0.0033	0.0045	0.28	0.0000	0.0015	0.0051	0
yoghurts total	0.0146	0.0191	1.25	0.0000	0.0085	0.0232	0
-yoghurts with fruit	0.0095	0.0161	0.81	0.0000	0.0000	0.0153	0
-yoghurts plane	0.0051	0.0108	0.44	0.0000	0.0000	0.0051	0
butter	0.00081	0.00140	0.07	0.00000	0.00012	0.00111	0
syrups	0.000048	0.000190	0.00	0.000000	0.000000	0.000000	0
hard (gouda) cheeses	0.0028	0.0021	0.24	0.0012	0.0023	0.0039	0
cottage cheese	0.0049	0.0119	0.42	0.0000	0.0000	0.0000	0
% as natural contaminant= 2.7%							
				0.65	1.05	1.63	
TOTAL	1.25 (0.25)	0.83		(0.13)	(0.21)	(0.33)

Figure 1. Cumulative distribution functions of the estimated benzoic acid intake (corrected for age) by the Belgian population (mg/kg bw/day), for men and women separately, Tier 2 approach

Figure 2. Cumulative distribution functions of the estimated benzoic acid intake (corrected for age) by the Belgian population (mg/kg bw/day) for men and women separately, Tier 3 approach

For Peer Review Only