Dietary intake estimate for perfluorooctanesulphonic acid (PFOS) and other perfluorocompounds (PFCs) in English retail foods following determination using standard addition LC-MS/MS

Don Brian Clarke, Antony Lloyd, Simon Hird, David Mortimer, Martin Gem, Victoria Adele Bailey, Anne Routledge

► To cite this version:

Don Brian Clarke, Antony Lloyd, Simon Hird, David Mortimer, Martin Gem, et al.. Dietary intake estimate for perfluorooctanesulphonic acid (PFOS) and other perfluorocompounds (PFCs) in English retail foods following determination using standard addition LC-MS/MS. Food Additives and Contaminants, 2010, 27 (04), pp.530-545. 10.1080/19440040903476590 . hal-00577002

HAL Id: hal-00577002

<https://hal.science/hal-00577002>

Submitted on 16 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dietary intake estimate for perfluorooctanesulphonic acid (PFOS) and other perfluorocompounds (PFCs) in English retail foods following determination using standard addition LC-MS/MS

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2009-334.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	04-Nov-2009
Complete List of Authors:	Clarke, Don; Food and environment research agency, Contaminants and authenticity programme Lloyd, Antony; Food and environment research agency, Contaminants and authenticity programme Hird, Simon; Food and environment research agency, Chemical residues programme Mortimer, David; Food Standards Agency, Food safety: contaminants division Gem, Martin; Food Standards Agency, Food safety: contaminants division Bailey, Victoria; Food Standards Agency, Food safety: contaminants division; University of York, Department of Chemistry Routledge, Anne; University of York, Department of Chemistry
Methods/Techniques:	Chromatography - LC/MS, Survey
Additives/Contaminants:	Environmental contaminants
Food Types:	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review Only

Dietary intake estimate for perfluorooctanesulphonic acid (PFOS) and other perfluorocompounds (PFCs) in English retail foods following determination using standard addition LC-MS/MS

D. B. Clarke* V. A. Bailey[†] A. Routledge[†], A. S. Lloyd, S. Hird, D. N. Mortimer⁺, M. Gem⁺

Food and Environment Research Agency, Department for Environment, Food and Rural Affairs, Sand Hutton, York, YO41 1LZ, UK; [†]The Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK; ⁺Food Standards Agency, 125 Kingsway, London WC2B 6NH

Abstract

The analysis of 252 food samples (UK-produced and imported) purchased from a variety of retail outlets in England was undertaken for the presence of PFOS, PFOA and nine other PFCs. A limit of quantification (LOQ) of 1 µg/kg was achieved for all target analytes, in all samples. Standard addition was used for quantification of PFC levels. All eleven of the targeted PFCs were detected, in 75 of the individual foods. In 70% of the samples, including all meat other than offal, none of the analytes were present at above the LOD. The highest levels found were 59 µg/kg of perfluorooctanesulfonic acid (PFOS) and 63 µg/kg total PFCs (ΣPFCs) in an eel sample, and 40 µg/kg PFOS (62 µg/kg ΣPFCs) in a whitebait sample. The highest level in an offal sample was 10 µg/kg, in a wild roe deer liver. There were 6 samples with ΣPFCs >15 µg/kg (fish, shellfish, crustaceans), a further 7 samples with ΣPFCs 11-15 µg/kg (including a liver), nine with ΣPFCs 6-10 µg/kg (fish and livers), 31 with ΣPFCs 2-5 µg/kg (including kidneys, popcorn and processed peas) and a further 22 with ΣPFCs close to the LOD of 1 µg/kg (including eggs and potatoes). These concentrations indicate that English consumers are being exposed to a low level of PFC contamination from food. The estimated *upper bound* dietary intake of 10 ng/kg bodyweight/day of PFOS for average adult consumers is well below the 0.15 µg (150 ng)/kg bw Tolerable Daily Intake (TDI) set by the European Food Safety Authority.

The *lower bound* adult dietary intake estimate of 1 ng/kg bodyweight/day is similar to estimates undertaken and reported in some other countries, such as Canada, Germany and Spain.

Keywords: perfluorooctanesulphonic acid, PFOS, perfluorocompounds, PFCs, food, fish, liver, TDI, dietary intake

*To whom correspondence should be addressed. e-mail don.clarke@fera.gsi.gov.uk

Introduction

PFOS is now the generally accepted term referring to the individual chemical perfluorooctanesulphonic acid and any closely related compounds that contain the PFOS moiety (C₈F₁₇SO₂), including those that may potentially degrade to PFOS in the environment (Brooke 2004). This term is often used interchangeably with PerFluoro Compounds (PFCs), or PerFluoroAlkylated Substances (PFAS). The chemicals under consideration are best described as an homologous series of fully-fluorinated sulphonic acids, and these homologues can also be present as sulphonamide or carboxylic acid derivatives. The high-energy carbon-fluorine covalent bond makes the core perfluoroalkyl group resistant to hydrolysis, photolysis and microbial or mammalian metabolism. Standard metabolic processes can remove other constituents such as hydrocarbon groups. There are believed to be important introversions of larger to smaller molecular weight PFCs occurring in the environment. PFOS is therefore not readily degradable and can persist and bioaccumulate in the environment, probably by means of covalent protein binding. The US Environmental Protection Agency (EPA) reported in May 2000 that PFOS combines persistence, bioaccumulation and toxicological properties to an extraordinary degree. PFOS exhibits hepatotoxicity, neurotoxicity, carcinogenicity and reproductive toxicity. One of the most disturbing findings was its unusual ability to influence second-generation rodent reproduction at 0.4 mg/kg body weight per day (bw/day) (repeated daily dosing). PFCs are also known to be endocrine disruptors (Jensen 2008).

PFOS has global biospheric distribution, bioaccumulation and biomagnification patterns similar to other persistent organic pollutants (POPs). A recent review provides an up-to-date reference for sources, fate and transport of perfluorocarboxylates (Prevedouros 2006). The increases in global exposure levels, have led to elevated circulatory concentrations of PFOS in mammalian plasma. Retrospectively, this upward trend has been apparent for some decades. Aside from industrial exposure, it is not yet clear as to the exact mechanisms of exposure to specific fluorochemicals and their precursors, or their relative significance, e.g. via fire fighting foam, household and consumer products vs food and water, vs air and dust.

Historically the uses of PFOS have been widespread and include uses as; a metal surface treatment, as a fabric protector (stain repellents - carpets), impregnated in paper (grease proofing - pizza boxes), and leather, in waxes, polishes, paints, varnishes and cleaning products. In Europe the suggested scale of use is 80 tons of PFOS used between 1980 and 2005 by the North Sea oil industry (fire fighting) and an equivalent usage/emission as total European terrestrial pollution.

3M, the major PFC producer, had reduced PFOS production by 98% by the end of 2000 (equivalent to an 88% reduction in world-wide production), with total cessation in 2002 (Olsen 2008, Spliethoff 2008). The global regulation of PFOS/PFOA, together with the voluntary change to alternative products that is already well underway in Westernised countries, should result in a decline in future environmental contamination and hence human exposure. The first declines in human exposure in localised areas have already been reported. The high human exposure to PFCs in North America led to this population and the blood samples collected by the American Red Cross, being the most highly contaminated in the world. However, their PFOS levels appear to have peaked, and are now declining in line with the previously reported half-life of 4-5 years in adult humans (Olsen 2008). A similar decline has very recently been observed in infants (Spliethoff 2008). While declines in US exposure to PFCs appear to have followed the fall in US production, this is not the case globally. PFOS levels appear to be continuing to increase in blood samples taken from Chinese citizens (Jin 2007).

As the estimated half-life of PFOS in the environment is 41 years, the legacy of environmental contamination cannot be expected to be solved rapidly. One of the clearest examples of a temporal trend in the food chain was seen in guillemot eggs obtained from a Baltic island, where the guillemots feed on fish from the Baltic Sea, thus reflecting the general level of contamination in the region. The concentration in eggs rose at an almost linear rate of 18 µg/kg/per year between 1968-2003, increasing from 17 µg/kg in 1968 to 623 µg/kg in 2003 (Holmstrom 2005). As these eggs are a human food source, the trend seen in these eggs can be directly correlated with the human food chain.

Various regulatory limits for acceptable levels of PFOS in the environment are being discussed. The European Food Safety Authority (EFSA) identified 0.03 mg/kg bw per day of PFOS as the lowest no-observed-adverse-effect level (NOAEL, liver pathology) for monkeys and considered this suitable for deriving a human Tolerable Daily Intake (TDI) for PFOS of 0.15 µg/kg bw. Similarly a ten-fold higher human TDI for PFOA of 1.5 µg/kg bw has also been agreed (EFSA 2008).

To date there have been very few studies that have determined PFCs in the human diet. The prevalence of PFOS and its impact on the wider human food chain only became apparent when the 3M Multi-City Study reported that four of the twelve positive food concentrations were from 'control' cities, i.e., those without a fluorochemical industry (3M Company, 2001). The emerging data on the concentrations of PFCs in food throughout the world are forming a picture of a generally low-level contamination in the wider diet, but with individual "hot spots" occurring within the fish and offal food groups. A survey of Swedish Total Diet Study (TDS) food composites (Berger 2007) focused on the four most likely food groups to be contaminated; meat products, dairy products, eggs and seafood. This survey did not positively identify any PFCs and reported less than the LOD (2.2, 3.2 µg/kg PFOS and PFOA, respectively) for all food composites surveyed. However, individual fish samples were found to contain up to 23 µg/kg PFOS (Berger 2007). A Spanish survey of composite foodstuffs found numerous trace levels of PFOS with a maximum of 0.8 µg/kg (Ericson 2008). A larger Canadian TDS study reported a maximum level of 5 µg/kg perfluorononanoic acid (PFNA) in beefsteak. PFOS was measured at 2.7 µg/kg in beefsteak, 2.6 µg/kg in marine fish, 2.1 µg/kg in ground beef and 2.0 µg/kg in

freshwater fish (Tittlemier 2007). A large 7-day duplicate diet study in Germany reported average values from 214 separate daily diets collected from 31 adults as 0.06 and 0.69 $\mu\text{g/kg}$ for PFOS and PFOA, respectively (Fromme 2007). For adults consuming 2.5 kg of food (solid and liquid) this represented mean daily dietary intakes of 123 and 270 ng of PFOS and PFOA, respectively, equivalent to 1.8 and 3.9 ng/kg bw/day. The duplicate diet collection model was therefore not dissimilar in outcome to the TDS model with regard to the intake estimates. Of interest in this study was the detailed analysis of each day's food in which just two PFOA values were unusually high. The higher of these was a concentration of 118 $\mu\text{g/kg}$ in the daily diet. This one-off high value was undoubtedly from smaller portions of a single discrete foodstuff containing at least an order of magnitude higher concentration. This again is consistent with the emerging knowledge on upper-end mg/kg contamination levels in some European freshwater fish from heavily polluted industrialized areas.

In order to put European and UK data into context, recent work in Japan (Guruge 2008) reported cattle, pig and chicken livers to contain mean PFOS concentrations of 34, 54 and 67 $\mu\text{g/kg}$, respectively, with the highest individual PFOS value at 92 $\mu\text{g/kg}$ in a chicken liver. Chinese chicken eggs were all found to contain PFOS in the range of 45-87 $\mu\text{g/kg}$ (Wang 2008). Oysters from Tokyo bay contained 3 $\mu\text{g/kg}$ of PFOS (So 2006).

The aim of the survey was to develop more robust analytical methods, in particular to lower the limit of detection, to enable improved estimates of dietary intakes of the chemicals and to provide data to inform negotiations regarding possible future limits for the chemicals in foods.

Materials and methods

Collection and preparation of food samples

Priority was given to foods in which the chemicals had been detected in the previous survey (potatoes and vegetables (FSA 2006), and foods in which it was anticipated that the chemicals would be present on the basis of literature reports (fish and offal).

A total of 252 samples of various foods (fish, liver, kidney, meat, cheese, milk, eggs, potatoes, vegetables, cereals and jam) were obtained by Ventress Technical Limited. Ventress Technical Limited is an independent consultancy with specialist skills in the practice of food technology, food examinations and offers varied services to science and technology. Using trained sampling personnel and comprehensive field management information systems, it has many years experience of sampling a variety of food products, from bulk consignments on site to individual retail packs on retail sale. The samples for this survey were purchased from retail establishments in ten regions of the UK (covering all four countries but weighted towards England as the most populated of the countries), in two phases, the first in 2007. The second phase, used to follow up on foods in which higher concentrations of the chemicals had been found in phase 1, took place in 2008.

The retail establishments included major and smaller national supermarkets and independent retailers (including convenience stores, butchers, fishmongers, health food shops, farm shops, ethnic stores, licensed market stalls and mail order/internet sales). In selecting the retail establishments and numbers of samples obtained from each, a balance was struck between the relative market shares of supermarkets vs independent retailers and the need to cover a large number of establishments and locate some foods not widely available. All sampling was random. All samples were stored and transported to the Food and Environment Research Agency Laboratory, which carried out the analyses, mostly under similar conditions to those that prevailed when they were on sale. Approximately 400 g of each individual retail food was homogenised and extracted without any further processing. However certain meat and fish samples were frozen immediately after purchase and held in a frozen state throughout further storage and transfer stages to prevent microbial deterioration. The types of foods are described briefly in Table 1. All other food samples and reference materials referred to in this report were obtained by the authors and were used for method development, method validation and quality assurance (QA) purposes only.

Chemicals

All solvents were of HPLC grade. Ammonium acetate (99.99%), acetic acid (glacial), potassium hydroxide pellets were supplied by Sigma (Poole, UK). All other reagents,

solvents and chemicals were of analytical grade, specific LC-MS grade or equivalent. Strata XAW weak anion exchange SPE cartridges (6 cc/200 mg, 33 μ m) were supplied by Phenomenex (Macclesfield, UK). All glassware was either washed three times with methanol or heated to 600°C, and silyanised before use.

Standards

The individual PFCs and internal standards are listed in Table 2. PFOS (perfluorooctanesulphonic acid), PFHxSK (potassium perfluorohexanesulphonic acid) and PFHxA (perfluorohexanoic acid) were supplied by Fluka (Gillingham, UK), PFHpA (perfluoroheptanoic acid), PFOA (perfluorooctanoic acid), PFNA (perfluorononanoic acid), PFDeA (perfluorodecanoic acid), PFUnA (perfluoroundecanoic acid), PFDoA (perfluorododecanoic acid) and PFBSH (perfluorobutanesulphonic acid) and 3 α ,12 α -dihydroxy-5 β -cholic acid-*N*-(2-sulfoethyl)-amide (taurodeoxycholic acid, TDCA) were supplied by Sigma (Poole, UK). PFOSA (perfluorooctanesulphonamide) and TH-PFOS (tetrahydro-PFOS) were supplied by ABCR GmbH (Karlsruhe, Germany), ¹³C₄-PFOA, ¹³C₂-PFDeA, D₉-*n*-EtFOSE (2-(*N*-D₅-ethylperfluoro-1-octanesulfonamido)-D₄-ethanol) and D₃-*n*-MeFOSA (*N*-D₃-methylperfluoro-1-octanesulphonamide) were supplied by Greyhound Chromatography (Birkenhead, UK). Stock standards of those not already in solution were prepared at 1 mg/ml in methanol and stored at +4°C. Stock solution volumes were monitored by recording the weight before and after each aliquot was removed, to check for possible losses of solvent due to evaporation. Working solutions and mixtures were prepared fortnightly and also stored at +4°C. All fluorinated chemicals and reagents were assessed for purity and possible PFC cross contamination. For PFC analytical standards, the area responses of all multiple reaction monitoring (MRM) transitions in each separate 0.5 μ g/ml solution in methanol were measured and assessed as a percentage of the total ion response [assuming equal ionization]. Chemical purities were all >90%, so no corrections were considered to be necessary.

Sample extractions

Quadruple 10 g portions of each homogenised food were weighed out into Falcon tubes (50 ml). The appropriate volumes of internal standard (IS) and standard addition

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

mixtures were added to prepare two unspiked portions; one overspiked at the reporting level (1 µg/kg) and one at 10-times the reporting level (10 µg/kg). The food portions were homogenised for 1-3 mins as required in 20 ml of methanol with an Ultra Turrax (T25 basic with S25N blade). Once homogenised, more methanol was added (*ca.* 40 ml in total) and mixed, while withdrawing the Ultra Turrax blade. Samples were shaken vigorously overnight (16 h), then centrifuged (15 min, 5000 rpm) (Tittlemier 2007). Eggs and milk samples required heating at 70°C for 2 hr to induce protein precipitation. The supernatant methanol extracts were evaporated under a nitrogen stream (80°C, in silanised glass vials) just to dryness, and the residues were re-dissolved in aqueous KOH (25 ml, 0.01 M, sonication 10 min). The aqueous solutions were then re-centrifuged (15 min, 5000 rpm). When required, the supernatants were poured in one continuous gentle movement, without breaking up the floating materials [fat] or disturbing the sediment, into a funnel connected onto the top of a preconditioned SPE cartridge (weak anion exchange)(Taniyasu 2005). Where clear solutions with no floating matter were obtained, these were simply added in repeated *ca.* 5 ml portions to the cartridges as the eluent went through (to *ca.* 1 ml remaining). The cartridges were loaded at a constant drip rate, by increasing from gravity feed to full vacuum as required. After loading, the cartridges were washed with ammonium acetate (2 x 6 ml, 25 mM, pH 4.5) and eluted with basic methanol (4 ml, 0.1% ammonia). The eluates were reduced under a stream of nitrogen gas (60°C), just to dryness and the residues taken up in methanol (400 µl, sonication 10 min). Extracts were transferred into silanised glass microvials (300 µl) for LC-MS/MS determination.

The analysis of cheeses was particularly problematic, due to the excessive amount of material (*ca.* 50 mg) left after the standard clean-up. The clean-up method was therefore extended. The residue was halved by elution of neutral amides off the SPE in methanol (4 ml, neutral pH), and then treating this fraction as a separate extract. Acids, both sulphonic and carboxylic were then eluted as before, using basic methanol (4 ml, 0.1% ammonia). Each fraction still contained *ca.* 25 mg of coextractives. A fluoruous solvent mixture was then used to partition fluorinated from non-fluorinated compounds by liquid-liquid extraction (F-LLE). SPE eluates (neutral or basic) were dried and then redissolved in dichloromethane (10 ml, DCM saturated with water) by sonicating for 10 min. Trifluoroethanol (7.5 ml) and perfluorohexane (FC-72, 2.5 ml)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

were added, the mixture was shaken (1 min), and then sonicated for 10 min. When the residues were in solution, the resulting triphasic system was heated to 80°C for one hour to achieve a monophasic mixture. The mixtures were then cooled to -70°C for one hour and the upper 10 ml of DCM was pipetted out and discarded as the non-fluorinated fraction. The lower two fluorinated layers were pipetted out together and dried under a stream of nitrogen gas (30°C) until just dry, and the residues taken up in methanol (400 µl, sonicated for 10 min). Extracts were transferred into silanised glass microvials (300 µl) for determination by LC-MS/MS.

Chromatography and quantification

Analysis was undertaken by LC-MS/MS. A CTC Pal autosampler (Presearch, UK) and an HP1100 HPLC system with column oven (Agilent, UK) were coupled to an API4000 triple quadrupole mass spectrometer (MDS Sciex Instruments, UK). The guard cartridge was C₈. The HPLC column (5 µm, 60 Å, 2.1 x 150 mm) was Fluorosep RP Octyl phase, thermostatically held at 30°C in the column oven. The injection volume was generally 10 µl. The gradient programme (methanol: aqueous ammonium formate, 5 mM, pH 4) was: 10% methanol increasing to 30% at 0.1 min (linear gradient), to 75% at 7 min and 100% methanol at 10 min, this was held for 5 min (column washing), then decreased to 10% methanol at 15.1 min, this was held 4.9 min at 10% methanol (column re-conditioning). The eluate was diverted to the mass spectrometer between 7 and 19.5 min, whereas from 0-7 and 19.5-20 min it was discarded by valve switching to waste, in order to protect the ion source. A summary of MRM transitions, retention times and solvent-based LODs of the PFC analytical standards are given in Table 2. Analyst 1.4.2 software was used for instrument control, file acquisition and peak integration. The MS detector in MRM mode with a Turbo Ion Spray source was used for quantitative analysis. Data acquisition was conducted in one simultaneous acquisition schedule. Instrumental parameters were optimised by infusion of standard solutions directly into the MS detector (1 µg/ml in 1:1 methanol: aqueous ammonium formate (5 mM, pH 4). The Turbo Ion Spray (TIS) conditions were; turbo-gas 50 psi, curtain-gas 12 psi, nebuliser-gas 50 psi, desolvation temperature 450°C. An Excel spreadsheet was used to calculate PFC concentrations from the standard additions. The standard addition calculation approach is illustrated

in Figure 1, with representative chromatograms demonstrating the response of the less sensitive but highly selective secondary qualifying ion for PFOS in Figure 2.

Analytical quality assurance

Stock standard solutions were prepared at 1 mg/ml in methanol and stored at 4°C. Dilutions of these were prepared bi-weekly to minimise surface binding or evaporation losses. Dispensing of stock solutions was monitored for any possible losses, due to evaporation, by weighing. Working solutions of mixed standards were prepared at 5 and 0.5 µg/ml in methanol for spiking. Foods (10 g) were overspiked at 1 and 10 µg/kg by addition of 20 µl @ 0.5 or 5 µg/ml of a mixed standards solution. Tetrahydro-PFOS (TH-PFOS), ¹³C₄-PFOA, ¹³C₄-PFOS, ¹³C₂-PFDeA, D₉-*n*-Et-FOSE and D₃-*n*-Me-FOSEA were used as internal standards (IS) by addition of 200 µl @ 0.5 µg/ml of a mixed solution, giving a 10 µg/kg IS addition level. Each food sample was analysed in duplicate throughout the entire extraction method to ensure that advantageous point contamination was not mistaken for the presence of any native PFC. Each sample was also overspiked at a minimum of two concentrations (1 & 10 µg/kg). For a specific analyte to be considered present in a sample extract the following criteria had to be met: i) the relative retention times of the analyte had to be comparable to those of a retention time marker, an internal standard, and to authentic analytical standards of each analyte; ii) the peak had to have the correct mass transition, maximising at the correct retention time; iii) the signal to noise ratio of any peak had to be greater than 3:1. In order to prove the absence of a given PFC, the internal standard had to be present in all extracts, the blank extract had to show no signal at the retention time of the target PFC, whilst the overspiked extracts had to show a peak for the target PFC at the required retention time. As many of the foods behave differently with respect to extraction, recovery of analytes through the SPE stages and matrix induced suppression of ionisation, a single LOD (reporting level) for all foods or all target PFCs was not possible. In the absence of any suitable certified reference material (CRM), a number of samples from previous surveys that had been found to contain measurable levels of specific PFCs were re-analysed to demonstrate inter-batch reproducibility (Table 3).

Dietary intakes

Calculations were performed by the UK Food Standards Agency using the Intake2 programme. Intake2 is the FSA in-house system supplied by Dietica for assessing exposure to chemicals in the food we eat and conducting nutrition and consumption assessments using NDNS (National Diet and Nutrition Survey) data. Intake2 is comprised of a bespoke VB front end and a Sagent back end database containing the NDNS consumption surveys, Nutrient information on all foods, and recipe composition data. Users set up a food group containing foods of interest and, if required, the system identifies recipes containing these foods and the percentage within the recipe. For intake assessments, users apply levels of chemicals to the foods selected. Intake2 essentially identifies which consumers have eaten the foods of interest, the amount eaten, and the level of or contaminants within the foods, and provides summary distributional analyses across consumers or the survey population.

All calculations are consumer-based, i.e. where only those who eat a particular food item are included in the calculations, rather than the data being averaged over the whole population. Average and individual age group estimates are calculated based on the most recent UK consumption data for the elderly (65 years and over) (Finch 1998) adults (19- 64 years) (Henderson 2002), school children (4-18 years) (Gregory 2000) and toddlers (1½ - 4½ years)(Gregory 1995)) surveys. (Table 7). High level consumer based estimates are taken as the 97.5% percentile (Table 8). Other calculations are illustrative and based on multiple consumption of single analytes contained in individual samples from Table 4 (Table 9).

Results

In almost three quarters of the samples, including all those of meat other than offal, none of the chemicals was observed above the limit of detection. PFOS was the chemical detected most often, especially in fish, shellfish, liver and kidney, and usually at the highest concentrations. 75 of the food samples contained trace levels of PFOS and other PFCs just above the reporting level of 1 µg/kg. All of the listed eleven perfluoro analytes were found to be present across these food samples. All of the foods found to contain concentrations exceeding the limit of detection have been reported individually, by analyte in descending order of the summed concentration of the individual PFCs (ΣPFCs) of 1-63 µg/kg (Tables 4-6). The highest concentrations

were found in samples of smoked eel and whitebait. PFOSA was the next most frequently detected chemical, but only in certain fish species and crab. PFOA was detected at low concentrations in some samples of whitebait, crab and liver. The other chemicals analysed were detected only occasionally and at low concentrations, but no chemical was absent from and totally undetected in all of the samples. A summary table of the results for all 252 samples, arranged by sample type, illustrates the frequency, range and mean levels of contamination (Table 1). The highest individual concentrations were of 59 µg/kg of perfluorooctanesulphonic acid (PFOS) and 63 µg/kg total PFCs (ΣPFCs) in an eel sample, and 40 µg/kg PFOS (62 ΣPFCs) in a whitebait sample. There were 6 samples with ΣPFCs >15 µg/kg (fish & crustaceans), a further 7 samples with ΣPFCs in the range 11-15 µg/kg (including a liver), nine with ΣPFCs in the range 6-10 µg/kg (fish and livers), 31 with ΣPFCs in the range 2-5 µg/kg (including kidneys, popcorn and processed peas) and 22 with ΣPFCs at the reporting level of 1 µg/kg (including eggs and potatoes).

A set of representative chromatograms for one of the PFOS determinations is given (Figure 2). It was noted in separate LC-TOF experiments that PFOS $[M-H]^-$ m/z 498.927 can be readily distinguished from the isobaric interferences. These have a profile m/z 498.289, 499.289 and 500.289 which, within the constraint of containing a sulphonate grouping (MRM 499>80), gave a single fit $[M-H]^-$ formulae of $C_{26}H_{44}NO_6S$, taurodeoxycholic acid, with the first natural abundance isotope peak $[M+1, -H]^-$ m/z 499.289 being the actual interfering ion. As there is a set of largely indistinguishable di-hydroxycholate epimers about the C_3 , C_6 , C_7 , C_{12} positions, the actual interferences in the various foods differ between samples as do bile salts between the various mammalian, fish and avian sources. As the di-hydroxycholate epimers in foods are secondary metabolites, there is as yet no clear relationship between the primary bile salts in a species and the observed interferences.

The highest positive results were all in the fish and offal food groups, as anticipated. Other primary produce foods, such as shellfish, meat, milk, butter, cheese, cereals and vegetables, were essentially free from contamination (Table 1). There was no observable trend in farming and husbandry practices; free range, organic, farmed and wild foods were all indistinguishable by PFC content.

Discussion

Analytical methods

The analytical method has been validated previously (Lloyd 2009) and performed satisfactorily, with all samples being analysed to meet the required reporting level (RL) (Tables 1-2). The method proved reproducible and accurate over the course of this survey as determined by repeat analysis of in-house reference food samples (Table 3).

Intake estimates

Making the assumptions that these retail food samples reflect the general diet in the UK, it can be assumed that the latter is low in PFC. Dietary intakes of PFOS, PFOA and PFOSA were estimated from the results of this survey using food consumption data from various dietary surveys, in which the food eaten by each of up to 1,724 individual consumers was recorded over a 4- or 7-day period (Gregory 1995, Gregory 2000, Finch 1998, Henderson 2002). Average concentrations of the chemicals found in the oily fish, white fish, shellfish, liver, kidney, meat, milk and milk products, cereals, popcorn, jams, potatoes and vegetables samples were applied to all samples of the relevant food type. These dietary intakes do not take account of food types that were not included in the survey, nor of carp for which there were no recorded consumers in any of the dietary surveys. Adults were assumed to have a body weight of 60 kg. For the other age groups the individual bodyweights of each recorded consumer were used.

The dietary intakes so estimated are presented in Table 7 for consumers in different age groups. These intakes are consumer rather than population intakes, as non-consumers of a given food type are ignored in calculating the average figures. The numbers of recorded consumers of the oily fish species containing the highest concentrations, and of offal are relatively low in all age groups which, when combined with the relatively high concentrations in these food types, inflate the consumer intakes from those food types. However, because foods bread, milk and potatoes, all of which are consumed by nearly all the UK population were also

included, the overall intakes shown in Table 7 also apply to the whole population. The estimated *upper bound* average adult dietary intakes of PFOS and PFOA 0.01 and 0.01 µg/kg bodyweight/day respectively (Table 7) are within the Tolerable Daily Intakes for these chemicals (0.15 and 1.5 µg/kg bodyweight/day respectively) set by the European Food Safety Authority (European Food Safety Authority, 2008), as are the corresponding high level (97.5 percentile) adult intakes of 0.02 and 0.02 µg/kg body weight/day respectively (Table 8).

An alternative method of estimating the dietary intakes of consumers of oily fish is to add the dietary intakes from typical portions of this food to that for the rest of the diet. This method allows for more robust intake estimates from oily fish species for which there were very few or no recorded consumers, e.g. eel, whitebait and carp. The Agency advises that all consumers should eat two portions of fish, one of which should be oily, as part of a balanced diet; that girls and women who may have children should not eat more than two portions of oily fish per week; and that boys, men and women who will not have children should not eat more than four portions of oily fish per week (Scientific Advisory Committee on Nutrition and Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment 2004, Food Standards Agency 2004). A man who ate four 70 g portions of consistently the most contaminated oily fish in the survey (i.e. smoked eel, 59 µg/kg) per week combined with average consumption of the rest of the diet (i.e. excluding the normal contribution from fish) would have a PFOS intake of 0.050 µg/kg bodyweight/day, which is again within the TDI (Table 9).

The concentrations of PFOS and PFOA in fish were consistent with those reported by other countries (Senthilkumar 2007, de Voogt 2008, Nania 2009). The predominance of PFOS accords with evidence that pefluoroalkyl sulphonates such as PFOS are more bioaccumulative than perfluoroalkanoic acids, such as PFOA, with the same fluorinated carbon chain length (Condor 2008). The concentrations found suggest that, if regulatory limits were to be considered, these might be appropriate only for PFOS in fish and animal liver, Limits are not likely to be needed in other foods, or at all for PFOA. However, the results of this survey do not indicate a need for increased consumer protection.

Accurate quantification of PFCs at concentrations below 1 µg/kg in food, while feasible, is complex and costly (Lloyd 2009). It is clear from the results and the intake estimates, that it was possible to demonstrate that exposure to PFOS and PFOA would be considerably below the respective TDIs even for the highest consumers working to an LOD of 1 µg/kg. On this basis, the analytical method described and used in this paper to generate the intake estimates was fit-for-purpose. The *lower bound* estimate for PFOS dietary intake in the UK of 1 ng/kg/day as estimated from the results of this study, was calculated as a consumer rather than population based estimate, which gave a higher result, but this is still comparable to estimates calculated for Spain 1 ng/kg/day (Ericson 2008), Ireland 1 ng/kg/day (Clarke in press), Germany 2 ng/kg bw/day (Fromme), and Canada (4 ng/kg bw/day Tittlemier 2007).

Conclusions

The results from this study indicate an extremely low level of contamination in the English food chain, and they are directly comparable to the findings in other countries that have conducted similar studies. It would therefore be unlikely for any consumer, even an extreme consumer of the more contaminated foods, to exceed the TDI for PFOS or PFOA.

Acknowledgments

The authors would like to thank Ventress Technical Ltd for their assistance in preparing the sampling plan and for obtaining the samples. This work was funded by the UK Food Standards Agency.

References

- 3M Company, 2003. Analysis of PFOS, FOSA and PFOA From Various Food Matrices Using HPLC Electrospray Mass Spectrometry. Analytical Report, June 21, 2001.
- Berger, U., Holmström, K., Glynn, A., Berglund, M., Ankarberg, E., and Törnkvist, A. 2007. Perfluorinated alkyl substances in market basket food samples and fish from Lake Vättern and the Baltic Sea. Rapport till Naturvårdsverket Programområde Miljögiftssamordning Överenskommelse nr 219 0641 Dnr: 721-5953-06Mm. Stockholm/Uppsala 2007-04-03.
- Brooke, D., Footitt, T. A., and Nwaogu, T.A. 2004, Environmental risk evaluation report: Perfluorooctanesulphonate (PFOS) Draft report prepared for the UK Environment Agency. By Building Research Establishment, Watford, and Risk and Policy Analysts, Loddon, Norfolk. http://www.environment-agency.gov.uk/commondata/105385/pfos_rer_sept04_864557.pdf
- Clarke, D. B., A., Lloyd, A. S., and Hird, S., 2008. Dietary intake estimate for perfluorooctanesulphonic acid (PFOS) and perfluorocompounds (PFCs) in Irish retail foods following determinations using standard addition LC-MS/MS. *Food Additives and Contaminants*, In preparation.
- Conder, J. M., Hoke, R. A., De Wolf, W., Russell, M. H., and Buck R. C. 2008. Are PFCAs bioaccumulative? A critical review and comparison with regulatory criteria and persistent lipophilic compounds. *Environ. Sci. Technol.*; **42**, 995-1003.
- EFSA, 2008. Opinion of the Scientific Panel on Contaminants in the Food chain on Perfluorooctane sulphonate (PFOS), perfluorooctanoic acid (PFOA) and their salts. *The EFSA Journal*, **653**, 1-131.
- Ericson, I. Marti-Cid, R., Nadal, M., Van Bavel, B., Lindstrom, G., and Domingo, J. L., 2008. Human exposure to perfluorinated chemicals through the diet; intake of perfluorinated compounds in foods from the Catalan (Spain) market. *J. Ag. Food Chem.*, **56**, 1787-1794.
- European Food Safety Authority, 2008. Perfluorooctane sulphonate (PFOS), perfluorooctanoic acid (PFOA) and their salts. Scientific Opinion of the Panel on Contaminants in the Food Chain, available at http://www.efsa.europa.eu/cs/BlobServer/Scientific_Opinion/contam_ej_653_PFOA_en.pdf?ssbinary=true.

- Food Standards Agency Fluorinated chemicals. UK dietary intakes. *Food Survey Information Sheet*, 2006; 11/06 available at www.food.gov.uk/multimedia/pdfs/fsis1106.pdf.
- Finch, S., Doyle, W., Lowe, C., Bates, C. J., Prentice, A., Smithers, G. and Clarke, P. C., 1998. National Diet and Nutrition Survey: People aged 65 years and over. Volume 1: Report of the diet and nutrition survey. The Stationery Office.
- Food Standards Agency, 2004. Fish and shellfish, available at <http://www.eatwell.gov.uk/healthydiet/nutritionessentials/fishandshellfish/?lang=en>.
- Fromme, H., Schlummer, M., Möller, A., Gruber, L., Wolz, G., Ungewiss, J., Böhmer, S., Dekant, W., Mayer, R., Liebl, B., and Twardella, D., 2007. Exposure of an Adult Population to Perfluorinated Substances Using Duplicate Diet Portions and Biomonitoring Data. *Environ. Sci. & Technol.*, **41** (22), 7928–7933.
- Gregory, J., Lowe, S., Bates, C. J., Prentice, A., Jackson, L. V., Smithers, G., Wenlock, R., and Farron, M. National Diet and Nutrition Survey: young people aged 4-18 years. Volume 1: report of the diet and nutrition survey 2000. The Stationery Office, London.
- Gregory, J., Collins, D. L., Davies, P. S. W., Hughes, J. M., and Clarke, P. C. National Diet and Nutrition Survey: children aged 1½ to 4½ years. Volume 1: report of the diet and nutrition survey 1995. HMSO, London.
- Guruge, K. S., Manage, P. M., Yamanaka, N., Miyazaki, S., Taniyasu, S., and Yamashita, N., 2008. Species-specific concentration of perfluoroalkyl contaminants in farm and pet animals in Japan. *Chemosphere*, **73**, S210-S215.
- Henderson, L., Gregory, J., and Swan, G. The National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 1: Types and quantities of foods consumed 2002. The Stationery Office, London.
- Holmstrom, K. E., Jarnberg, U., and Bignert, A., 2005. Temporal trends of PFOS and PFOA in guillemot eggs from the Baltic sea, 1968-2003. *Environ. Sci. & Tech.*, **39**(1), 80-84.
- Jensen, A. A., and Leffers, H., 2008. Emerging endocrine disruptors. *International Journal of Andrology*, **31**, 161-169.
- Jin, Y., Saito, N., Harada, K., Inoue, K., and Koizumi, A., 2007. Historical Trends in Human Serum Levels of Perfluorooctanoate and Perfluorooctane Sulfonate in Shenyang, China. *Tohoku J. Exp Med.*, **212**, 63-70.

- Lloyd, A. S., Bailey, V. A., Hird, S., Routledge, A., Clarke, D. B., 2009. Mass spectral studies towards more reliable measurement of perfluorooctanesulfonic acid and other perfluorinated chemicals (PFCs) in food matrices using liquid chromatography/tandem mass spectrometry. *Rapid Commun. Mass Spectrom.*, **23**, 2923-2938.
- Nania, V., Pellegrini, G.E., Fabrizi, L., Sesta, G., de Sanctis, P., Luchetti, D., Pasquale, M.D., and Coni, E., 2009. Monitoring of perfluorinated compounds in edible fish from the Mediterranean sea. *Food Chem.*, 115, 951-957.
- Olsen, G. W., Mair, D. C., Church, T. R., Ellefson, M. E., Reagen, W. K., Boyd, T. M., Herron, R. M., Medhdizadehkashi, Z., Nobiletti, J. B., Rios, J. A., Butenhoff, J. L., and Zobel L. R., 2008. Decline in Perfluorooctanesulfonate and Other Polyfluoroalkyl Chemicals in American Red Cross Adult Blood Donors, 2000–2006. *Environ. Sci. & Technol.*, **42(13)**, 4989–4995.
- Prevedouros, K., Cousins, I. T., Buck, R. C., and Korzeniowski, S. H., 2006. Sources, fate and transport of perfluorocarboxylates. *Environ. Sci. & Tech.*, **240(1)**, 32-44.
- Senthilkumar, K., Ohi, E., Sajwan, K., Takasuga, T., and Kannan K. 2007. Perfluorinated compounds in river water, river sediment, market fish, and wildlife samples from Japan. *Bull. Environ. Contam. Toxicol.*, **79**, 427-431.
- Scientific Advisory Committee on Nutrition and Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment, 2004 Advice on fish consumption: benefits & risks. The Stationery Office, available at <http://cot.food.gov.uk/pdfs/fishreport2004full.pdf>.
- So, M. K., Taniyasu, S., Lam, P. K. S., Zheng, G. J., Giesy, J. P., and Yamashita, N. A, 2006. Alkaline digestion and solid phase extraction methods for perfluorinated compounds in mussels and oysters from South China and Japan. *Arch. Environ. Contam. Toxicol.*, **50**, 240-248.
- Splithoff, H. M., Tao, L., Shaver, S. M., Aldous, K. M., Pass, K. A., Kannan, K., and Eadon, G. A., 2008. Use of Newborn Screening Program Blood Spots for Exposure Assessment: Declining Levels of Perfluorinated Compounds in New York State Infants. *Environ. Sci. & Technol.*, **42(14)**, 5361–5367.
- Taniyasu, S., Kannan, K., So, M. K., Gulkowska, A., Sinclair, E., Okazawa, T., and Yamashita, N. A, 2005. Analysis of fluorotelomer alcohols, fluorotelomer acids, and short- and long-chain perfluorinated acids in water and biota. *J Chrom A.*, **1093**, 89-97.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Tittlemier, S. A., Pepper, K., Seymour, C., Moisey, J., Bronson, R., Cao, X.-L., and Dabeka, R. W., 2007. Dietary exposure of Canadians to perfluorinated carboxylates and perfluorooctane sulfonate via consumption of meat, fish, fast foods, and food items prepared in their packaging. *J. Ag. Food Chem.*, **55**, 3203-3210.
- de Voogt, P., van der Wielen, F. W. M., Westerveld, J., D'Hollander, W. and Bervoets, L., 2008. Determination of perfluorinated organic compounds in food and dust. *Organohal. Comp.*, **70**, 714-717.
- Wang, Y., Yeung, L. W., Yamashita, N., Taniyasu, S., So, M. K., Murphy, M. B., and Lam P. K., 2008. Perfluorooctane sulfonate (PFOS) and related fluorochemicals in chicken egg in China. *Chinese Science Bulletin*, **53**(4), 501-507.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Summary of the levels of PFCs by food type.

Class	Type	N	Range ΣPFCs µg/kg	N ≥ LOQ	Mean	% Positive
Fish	Whitebait	6	8-62	6	28	100
	Eel	6	<1-63	3	11	50
	Carp	6	<1-8	5	5	83
	Sprats	3	3-8	3	5	100
	Sardines	6	1-7	6	4	100
	Cod	4	<0-4	2	2	50
	Mackerel	4	<1-3	2	2	50
	Haddock	4	<1-3	3	1	75
	Trout	4	<1-1	2	<1	50
	Herring	4	<1-2	1	<1	25
	Plaice	2	<1-1	1	<1	50
	Salmon	8	<1	0	<1	0
	Sole	2	<1	0	<1	0
Shellfish	Oysters	2	1-1	1	<1	50
Crustaceans	Crab	6	11-20	6	16	100
	Crayfish	1	2	1	2	100
	Prawns	2	<1-1	1	1	50
	Langoustine	1	<1	0	<1	0
Offal	Liver	25	<1-14	19	2	76
	Kidney	12	<1-3	8	1	75
	Other offal	4	<1	0	<1	0
Meat	Carcass meat	16	<1	0	<1	0
Dairy	Milk	11	<1	0	<1	0
	Cheese	10	<1	0	<1	0
Eggs	Chicken	10	<1-1	1	<1	10
	Duck	2	<1	0	<1	0
Oils	Fish	4	<1	0	<1	0
	Vegetable	6	<1	0	<1	0
Plant produce	Grains/bread	12	<1-1	2	<1	17
	Vegetables	63	<1-2	2	<1	3
	Fruit/jam	6	<1	0	<1	0

Range ΣPFCs = the lowest and highest values of the combined concentration of all 11 Individual fluorinated analytes. Mean = lower bound, <LOD =0
N = The number of samples analysed in each group.
N≥LOQ = The number of samples where residues were measured.
Meat = pork, lamb, beef, venison, chicken, turkey.

Table 2. Summary of MRM transitions, retention time, solvent based and matrix based LODs

Test analyte	Primary Ions		Qualifier Ions		MeOH LODs		Matrix based LODs $\mu\text{g/kg}^b$			
	Transition	Product ion	Transition	Product ion	Rt	$\mu\text{g/kg}^a$	Milk	Jam	Veg Oil	Salmon
PFHxA	313.1>269.1	$[\text{C}_5\text{F}_{11}]^-$	313.1>119.1	$[\text{C}_2\text{F}_5]^-$	11.25	0.01	0.025	0.1	0.05	0.3
PFHpA	363.1>319.1	$[\text{C}_6\text{F}_{13}]^-$	363.1>169.1	$[\text{C}_3\text{F}_7]^-$	11.92	0.005	0.01	0.05	0.02	0.01
PFOA	413.1>369.1	$[\text{C}_7\text{F}_{15}]^-$	413.1>169.1	$[\text{C}_3\text{F}_7]^-$	12.43	0.005	0.025	0.01	0.05	0.01
PFNA	463.1>419.1	$[\text{C}_8\text{F}_{17}]^-$	463.1>219.1	$[\text{C}_4\text{F}_9]^-$	12.86	0.001	0.05	0.02	0.05	0.02
PFDeA	513.1>469.1	$[\text{C}_9\text{F}_{19}]^-$	513.1>219.1	$[\text{C}_4\text{F}_9]^-$	13.26	0.001	0.025	0.02	0.02	0.05
PFUnA	563.1>519.1	$[\text{C}_{10}\text{F}_{21}]^-$	563.1>319.1	$[\text{C}_6\text{F}_{13}]^-$	13.66	0.001	0.025	0.02	0.01	0.02
PFDoA	613.1>569.1	$[\text{C}_{12}\text{F}_{23}]^-$	613.1>269.1	$[\text{C}_5\text{F}_{11}]^-$	14.02	0.001	0.025	0.02	0.02	0.05
PFBSH	299.1>80.1	$[\text{SO}_3]^-$	299.1>99.1	$[\text{SO}_3\text{F}]^-$	10.05	0.01	0.05	0.05	0.01	0.05
PFHxS	399.1>80.1	$[\text{SO}_3]^-$	399.1>99.1	$[\text{SO}_3\text{F}]^-$	11.64	0.01	0.05	0.02	0.05	0.1
PFOS	499.1>80.1	$[\text{SO}_3]^-$	499.1>99.1	$[\text{SO}_3\text{F}]^-$	12.59	0.01	0.1	0.1	0.05	0.1
PFOSA	498.1>78.0	$[\text{SNO}_2]^-$	498.1>169.1	$[\text{C}_3\text{F}_7]^-$	13.96	0.002	0.025	0.02	0.1	1
THPFOS	427.1>407.1	$[\text{M}-\text{HF}]^-$	-	-	12.34	-	-	-	-	-
$^{13}\text{C}_2$ -PFDeA	515.1>470.1	$[\text{C}_8^{13}\text{C}_2\text{F}_{19}]^-$	-	-	13.26	-	-	-	-	-
$^{13}\text{C}_4$ -PFOA	417.1>372.1	$[\text{C}_4^{13}\text{C}_3\text{F}_{15}]^-$	-	-	12.42	-	-	-	-	-
$^{13}\text{C}_4$ -PFOS	503.1>80.1	$[\text{SO}_3]^-$	-	-	12.57	-	-	-	-	-
D_9 - <i>n</i> -Et-FOSE	639.1>45.0	$[\text{HCO}_2]^-$	-	-	14.81	-	-	-	-	-
D_3 - <i>n</i> -Me-FOSA	515.1>169.1	$[\text{C}_3\text{F}_7]^-$	-	-	14.68	-	-	-	-	-

Summary of MRM transitions, retention times and solvent based Limits of Determination LODs of PFC analytical standards.

^aMethanol based LODs (3x S/N) were all <10 ng/kg (0.25 ng/ml).

^bConcentrations expressed as $\mu\text{g/kg}$ food equivalent, matrix matched. 10 g food matrix concentrated in the extract 25-fold.

 D_9 -*n*-Et-FOSE 2-(*N*- D_5 -ethylperfluoro-1-octanesulfonamido)- D_4 -ethanol D_3 -*n*-Me-FOSA *N*- D_3 -methylperfluoro-1-octanesulfonamide

Table 3. Interbatch reproducibility data for repeat analysis of food sample contaminated with PFCs.

Sample	Measured PFC concentration µg/kg										
	PFBS	PFHxS	PFOS	PFHxA	PFHpA	PFOA	PFNA	PFDeA	PFUnA	PFDoA	PFOSA
Mackerel			1[2]								1[1]
Dressed crab			6[4]			9[8]	1[2]				1[1]
Whitebait			11[15]								10[27]
Whitebait			12[9]								6[4]
Whitebait			48[40]			4[5]			1[2]		
Cromer crab		1[1]	14[12]			4[4]	2[1]	1[1]			1[2]
Spider crab		2[1]	13[9]			5[6]					
Wild roe deer liver			10[11]			3[6]	1[2]				
Carp roe			765[757]								

Interbatch reproducibility data for repeat analysis of naturally incurred PFC containing food sample.
Results for the second determination are presented in parenthesis.

1 *Table 4. Individual concentrations of PFCs in each food sample.*

Sample	Measured PFC concentration µg/kg											ΣPFCs
	PFHxA	PFHpA	PFOA	PFNA	PFDeA	PFUnA	PFDoA	PFBS	PFHxS	PFOS	PFOSA	
1 Smoked Eel	< 1	< 1	< 1	< 1	2	2	< 1	< 1	< 1	59	< 1	63
2 Whitebait	< 1	< 1	5	< 1	< 1	2	< 1	< 1	1	40	14	62
3 Whitebait	7	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	15	27	49
4 Cromer crab (dressed)	< 1	< 1	4	1	< 1	1	< 1	< 1	2	12	< 1	20
5 Spider crab (whole)	< 1	< 1	6	1	< 1	< 1	< 1	< 1	< 1	13	< 1	20
6 Whitebait	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	9	7	16
7 Whitebait	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	9	6	15
8 Dressed Crab	< 1	< 1	8	2	< 1	< 1	< 1	< 1	< 1	4	1	15
9 Brown crab (whole)	< 1	< 1	6	3	< 1	< 1	< 1	< 1	< 1	3	2	14
10 Wild roe deer liver	< 1	< 1	3	1	< 1	< 1	< 1	< 1	< 1	10	< 1	14
11 Whitebait	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	9	4	13
12 Crab (whole)	< 1	< 1	5	1	< 1	1	1	< 1	< 1	2	3	13
13 Dressed Crab	< 1	< 1	4	1	< 1	< 1	< 1	< 1	< 1	4	2	11
14 Mirror carp	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	8	< 1	8
15 Frozen sprats	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	4	4	8
16 Whitebait	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	5	3	8
17 Carp	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	8	< 1	8
18 Sardines	< 1	< 1	< 1	< 1	< 1	2	< 1	< 1	< 1	3	2	7
19 Venison Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	7	< 1	7
20 Fresh carp	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	6	< 1	6
21 Common carp	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	6	< 1	6
22 Pigs liver	< 1	< 1	2	< 1	< 1	< 1	< 1	< 1	< 1	4	< 1	6
23 Sliced lamb liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	5	< 1	5
24 Pigs kidney	< 1	< 1	1	< 1	< 1	< 1	< 1	< 1	< 1	4	< 1	5
25 Sprats	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	4	5
26 Cornish Sardines	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	3	5
27 Ox Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	5	< 1	5
28 Carp	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	4	< 1	4
29 Bideford cod	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	2	4
30 Sardines	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	3	1	4
31 Pork Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	4	< 1	4
32 Lambs Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	4	< 1	4
33 Sardines	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	1	3
34 Beef liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	3	< 1	3
35 Halal beef liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	3	< 1	3
36 Lambs liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	3	< 1	3

Food Additives and Contaminants

37	English Sprats	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	2	3
38	Whole Herring	3	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	3
39	Mackerel	2	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	3
40	Whole Mackerel	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	1	3
41	Cornish Sardines	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	2	3
42	Ox Kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	3	< 1	3
43	Farmed red deer liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
44	Lambs kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
45	Wild English Eels	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
46	Cod	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
47	Crayfish	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	< 1	< 1	< 1	< 1	2
48	Lambs Liver	< 1	< 1	1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	2
49	Lambs Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
50	Lambs Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
51	Ox Kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	2
52	Marrowfat Peas	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	< 1	< 1	< 1	2
53	Toffee Popcorn	< 1	< 1	< 1	< 1	< 1	< 1	< 1	2	< 1	< 1	< 1	< 1	2
54	Haddock fillet	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
55	Haddock portions	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
56	Smoked eel	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
57	Herring	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
58	Cornish sardines	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
59	Ox liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
60	Wild venison liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
61	Beef kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
62	Lambs kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
63	R. Trout	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
64	Whole R. Trout	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
65	Plaice fillets	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
66	Haddock Fillets	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
67	N. Atlantic Prawns	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
68	Pacific Oysters	< 1	1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1
69	Pork Sliced Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
70	Ox Liver	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
71	Organic pig kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
72	Pig Kidney	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
73	Organic Free Range Eggs	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	1
74	Baby Potatoes [Maris Peer]	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1	< 1	< 1	< 1	< 1	1
75	Sweet Pop Corn	< 1	1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	< 1	1

1 ΣPFCs = combined total concentration of all individual PFCs = PFAS.
2

Table 5. Concentrations ($\mu\text{g/kg}$ fresh weight) of PFOS, PFOA, PFAS and PFOSA in fish and shellfish

Species (No of samples)	PFOS		PFOA		PFAS (lower bound)		PFAS (upper bound)		PFOSA	
	Mean	Range	Mean	Range	Mean	Range	Mean	Range	Mean	Range
Eel (fresh) (3)	1.3	<1-2	<1	-	0.7	0-2	11	11-12	<1	-
Eel (smoked) (3)	20	<1-59	<1	-	21	0-63	31	11-71	<1	-
Herring (4)	1.0	<1-1	<1	-	1.0	0-3	12	11-13	<1	-
Mackerel (4)	1.2	<1-2	<1	-	1.5	0-3	11	11-12	<1	-
Atlantic salmon (farmed) (5)	<1	-	<1	-	0	-	11	-	<1	-
Atlantic salmon (wild) (2)	<1	-	<1	-	0	-	11	-	<1	-
Alaskan salmon (wild) (1)	<1	-	<1	-	0	-	11	-	<1	-
Sardines (6)	2.0	1-3	<1	-	5.5	1-7	13	11-15	2.2	1-3
Sprats (3)	3.0	1-4	<1	-	5.3	3-8	14	12-17	3.3	2-4
Trout (farmed) (4)	1.0	<1-1	<1	-	0.50	0-1	11	-	<1	-
Whitebait (6)	15	<1-40	1.3	<1-5	28	8-62	36	17-68	10	1-27
Carp (6)	5.5	<1-8	<1	-	5.3	0-8	16	11-18	<1	-
All oily fish (47)	4.8	<1-59	1.1	-	6.7	0-63	17	11-71	2.5	1-27
Cod (4)	1.5	<1-2	<1	-	1.5	0-4	12	11-13	<1	-
Haddock (4)	1.0	<1-1	<1	-	1.3	1-2	11	-	<1	-
Plaice(2)	1.0	<1-1	<1	-	0.5	0-1	11	-	<1	-
Sole (2)	<1	<1-1	<1	-	0	-	11	-	<1	-
All whitefish (12)	1.2	<1-2	<1	-	0.8	0-4	11	11-13	1.1	1-2
Crab (6)	6.3	2-13	5.5	4-8	16	11-20	22	18-28	1.7	1-3
Crayfish (1)	<1	-	<1	-	2.0	-	12	12	<1	-
Langoustine (1)	<1	-	<1	-	1.0	-	11	11	<1	-
Pacific oysters (2)	2.5*	1-10*	<1	-	0	-	17*	11-47*	<1	-
Prawns (2)	<1	<1-1	<1	-	0.5	0-1	11	11	<1	-
All shellfish (12)	4.4	1-13	3.3	1-8	8.2	0-20	20	11-47	1.3	1-3

Note: * Means and ranges affected by one or more higher than usual LODs. PFAS = Σ PFCs.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 6. Concentrations (µg/kg fresh weight) of PFOS, PFOA, PFAS and PFOSA in other foods

Food (No of samples)	PFOS		PFOA		PFAS (lower bound)		PFAS (upper bound)		PFOSA	
	Mean†	Range	Mean†	Range	Mean	Range	Mean	Range	Mean†	Range
Chicken liver (2)	<1	-	<1	-	0	-	16	11,20	<1	-
Chicken liver pate (1)	<1	-	<1	-	0	-	11	-	<1	-
Duck liver pate (1)	<1	-	<1	-	0	-			<1	-
Lambs' liver (7)	2.6	<1-5	<1	-	2.6	0-5	13	11-15	<1	-
Ox liver (5)	1.8	<1-5	<1	-	1.4	0-5	12	11-15	<1	-
Pigs' liver (5)	2.2	<1-4	1.2	1-2	2.2	0-6	12	11-15	<1	-
Venison liver (4)	5.0	1-10	1.5	1-3	6.0	1-14	16	11-22	<1	-
All liver (25)	2.5	1-10	1.1	1-3	2.4	0-14	13	11-22	<1	-
Lambs' kidney (4)	1.3	1-2	<1	-	0.8	0-2	11	11-12	<1	-
Ox kidney (4)	1.8	<1-3	<1	-	1.5	0-3	12	11-13	<1	-
Pigs' kidney (4)	1.6	<1-4	<1	-	0.7	0-1	11	-	<1	-
All kidney (12)	1.4	1-3	<1	-	1.3	0-5	12	11-14	<1	-
Lambs' heart (2)	<1	-	<1	-	0	-	11	-	<1	-
Black pudding (2)	<1	-	<1	-	0	-	11	-	<1	-
Bread (4)	<1	-	<1	-	0	-	11	-	<1	-
Cereals (4)	<1	-	<1	-	0	-	11	-	<1	-
Popcorn (4)	<1	-	<1	-	0.8	0-2	11	11-12	<1	-
Cheese (10)	<1.9*	<1-<10*	<1	-	0	-	15*	11-47*	<1	-
Milk (11)	<1	-	<1	-	0	-	11	11-12	<1	-
Eggs (12)	<1	<1-1	<1	-	0.1	0-2	11	-	<1	-
Fish oil supplements (4)	<1	-	<1	-	0	-	11	-	<1	-
Vegetable oils (6)	<1	-	<1	-	0	-	11	-	<1	-
Jams (6)	<1	-	<1	-	0	-	11	-	<1	-
Meat (not offal) (16)	<1	-	<1	-	0	-	11	-	<1	-
Potatoes & products (21)	<1	-	<1	-	0.1	0-1	11	-	<1	-
Vegetables (42)	<1	-	<1	-	0.1	0-2	11	11-12	<1	-

Note: † Upper bound values (where the value of the LOD is used in place of <LOD)
Means and ranges affected by one or more higher than usual LODs. PFAS = ΣPFCs.

1 Table 7. Estimated average UK dietary intakes ($\mu\text{g/kg}$ bodyweight/day) of fluorinated chemicals by consumers of different age ranges
 2 from the diet in 2007-08
 3

Food group	Estimated average dietary exposure ($\mu\text{g/kg}$ bodyweight/day)							
	PFOS		PFOA		PFOSA		PFAS	
	<i>Upper</i>	<i>Lower</i>	<i>Upper</i>	<i>Lower</i>	<i>Upper</i>	<i>Lower</i>	<i>Upper</i>	<i>Lower</i>
<i>Senior citizens</i>								
- living at home	0.009	0.0009	0.008	<0.00005	0.008	0.0003	0.09	0.001
- in old peoples' homes	0.008	0.0004	0.007	<0.00005	0.007	0.0001	0.08	0.0007
<i>Adults</i>	0.01	0.0001	0.01	<0.00005	0.01	0.0004	0.1	0.002
<i>Schoolchildren</i>								
-age 4-6 years	0.02	0.001	0.02	<0.00005	0.02	0.0003	0.3	0.002
-age 7-10 years	0.02	0.0008	0.02	<0.00005	0.02	0.0003	0.2	0.001
-age 11-14 years	0.01	0.0005	0.01	<0.00005	0.01	0.0002	0.1	0.001
-age 15-18 years	0.01	0.0005	0.009	<0.00005	0.009	0.0002	0.1	0.001
<i>Toddlers</i>								
- age 1.5-2.5 years	0.04	0.0008	0.04	<0.00005	0.04	0.0002	0.4	0.001
- age 2.5-3.5 years	0.03	0.001	0.03	<0.00005	0.03	0.0003	0.4	0.002
- age 3.5-4.5 years	0.03	0.0009	0.03	<0.00005	0.03	0.0003	0.3	0.002

4 PFAS = Σ PFCs The estimates for PFAS are both *upper* and *lower bound*. Intake estimates are consumer rather than population based,
 5 where the non-consumers of a given food type are excluded, resulting in higher values.
 6
 7

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 8. Estimated high-level 97.5 percentile UK dietary intakes ($\mu\text{g/kg}$ bodyweight/day) of fluorinated chemicals by consumers of different age ranges from the diet in 2007-08

Food group	Estimated average dietary exposure ($\mu\text{g/kg}$ bodyweight/day)							
	PFOS		PFOA		PFOSA		PFAS	
	Upper	Lower	Upper	Lower	Upper	Lower	Upper	Lower
<i>Senior citizens</i>								
- living at home	0.02	0.004	0.02	0.0002	0.02	0.002	0.2	0.007
- in old peoples' homes	0.02	0.003	0.02	0.0001	0.02	0.001	0.2	0.004
<i>Adults</i>	0.02	0.006	0.02	0.0002	0.02	0.002	0.2	0.009
<i>Schoolchildren</i>								
-age 4-6 years	0.05	0.006	0.04	0.0002	0.05	0.002	0.5	0.01
-age 7-10 years	0.05	0.004	0.03	0.0001	0.03	0.002	0.3	0.006
-age 11-14 years	0.02	0.003	0.02	0.0001	0.02	0.001	0.2	0.004
-age 15-18 years	0.02	0.003	0.02	0.0001	0.02	0.001	0.2	0.006
<i>Toddlers</i>								
- age 1.5-2.5 years	0.08	0.005	0.08	0.0001	0.08	0.001	0.9	0.007
- age 2.5-3.5 years	0.08	0.007	0.08	0.0002	0.08	0.002	0.8	0.01
- age 3.5-4.5 years	0.06	0.005	0.06	0.0002	0.06	0.002	0.6	0.008

PFAS = Σ PFCs

Table 9. Illustrative intakes ($\mu\text{g/kg}$ bodyweight/day) of PFOS and PFOSA from selected fish compared to remainder of diet.

Intake from non-fish part of the diet	PFOS			PFOSA		
	0.010	0.010	0.010	0.010	0.010	0.010
Species of fish	Smoked eel	Whitebait	Smoked eel	Dressed crab	Whole crab	Whitebait
Concentration in fish (mg/kg fresh weight)	0.001	0.04	0.059	0.001	0.003	0.03
Fish portion (gram)	70	70	70	70	70	70
Daily intake from one weekly portion of fish	0.000	0.007	0.010	0.0002	0.0005	0.0045
Total daily intake fish + non-fish intake	0.011	0.017	0.020	0.010	0.011	0.015
Daily intake from two weekly portions of fish	0.000	0.013	0.020	0.0003	0.001	0.009
Total daily intake fish + non-fish intake	0.011	0.024	0.030	0.010	0.011	0.019
Daily intake from three weekly portions of fish	0.001	0.020	0.030	0.0005	0.002	0.014
Total daily intake fish + non-fish intake	0.011	0.030	0.040	0.011	0.012	0.024
Daily intake from four weekly portions of fish	0.001	0.027	0.039	0.0007	0.002	0.018
Total daily intake fish + non-fish intake	0.011	0.037	0.050	0.011	0.012	0.028

Based on representative fish in this survey from Table 4.

Figure 1. Standard addition calculation procedure for an English mackerel fish sample. Both sets of ion responses are first normalised to the $^{13}\text{C}_4\text{-PFOS}$ isotope dilution internal standard. The result is calculated where the line crosses the x-axis. Data are rounded to 1 significant figure, this sample was reported as PFOS contamination of 2 $\mu\text{g/g}$ (primary 1.75, qualifier 1.81). The qualifying ion has a constant 24% of the response of the primary ion response ($0.0897/0.3857$). The average ion ratio for native and overspiked PFOS is equivalent to the precision ratio, 1.81/1.75 4%.

Figure 2. A) Chromatogram of the primary ion $499.1 > 80.1 [M > SO_3]^-$ for PFOS in a sample of whitebait (sample 3 in Table 1, containing 15 µg/kg naturally incurred PFOS) using; A) unspiked sample, B) overspiked at 1 µg/kg, C) overspiked at 5 µg/kg, D) overspiked at 10 µg/kg, E) overspiked at 20 µg/kg and F) overspiked at 50 µg/kg. Other peaks in this MRM are most likely due to epimers of taurodeoxycholic acid, specifically the $^{13}C_1$ native abundance M+1 isotope.

Figure 3. Distribution of individual PFCs (µg/kg) in 75 contaminated foods. The majority of these, 68 foods, contained PFOS, 20 contained PFOSA and 11 contained PFOA.