PCR diagnosis of *P. brevicompactum*, a grape contaminant and mycophenolic acid producer.

Belén Patiño, Angel Medina, Domenech Mirian, González-Jaén M^a Teresa,
Misericordia Jiménez, Vázquez Covadonga

► To cite this version:

Belén Patiño, Angel Medina, Domenech Mirian, González-Jaén M^a Teresa, Misericordia Jiménez, et al.. PCR diagnosis of *P. brevicompactum*, a grape contaminant and mycophenolic acid producer.. Food Additives and Contaminants, 2007, 24 (02), pp.165-172. 10.1080/02652030600967222. hal-00577285

HAL Id: hal-00577285

<https://hal.science/hal-00577285>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PCR diagnosis of *P. brevicompactum*, a grape contaminant and mycophenolic acid producer.

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2006-121.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	10-Jul-2006
Complete List of Authors:	patíño, Belén; University Complutense, microbiology III Medina, Angel; University of Valencia, Microbiology and Ecology Mirian, Domenech; University Complutense, Microbiology III Mª Teresa, gonzález-jaén; University Complutense, Genetics Jiménez, Misericordia; University of Valencia, Microbiology and Ecology Covadonga, Vázquez; University Complutense, Microbiology III
Methods/Techniques:	LC/MS, Molecular biology - PCR, Mycology
Additives/Contaminants:	Mycotoxins
Food Types:	Fruit, Wine

SCHOLARONE™
Manuscripts

PCR identification of *Penicillium brevicompactum*, a grape contaminant and mycophenolic acid producer.

Belén Patiño, ³Ángel Medina, Mirian Doménech, ¹M^a Teresa González-Jaén,
³Misericordia Jiménez and Covadonga Vázquez *

Dpto. Microbiología III, ¹Dpto. Genética, Universidad Complutense de Madrid,
Madrid, Spain and ²Dpto. Microbiología y Ecología, Universidad de Valencia,
Valencia, Spain.

*Covadonga Vázquez Estévez. Departamento de Microbiología III. Facultad de Biología. Universidad Complutense de Madrid. 28040-Madrid, Spain
e-mail: covi@bio.ucm.es Tel.: (34) 913 944 969; Fax: (34) 913 944 964

1

2 **Abstract:**

3 *Penicillium brevicompactum* is a ubiquitous fungal species that contaminates
4 diverse substrates and commodities and produces an array of metabolites toxic to
5 human and animals. In this work, we have obtained evidence, by LC-ion trap-mass
6 spectrometry, of the ability of *P. brevicompactum* strains isolated from grapes to
7 produce mycophenolic acid, a potent immunosuppressor. In order to facilitate early
8 diagnosis of this species on commodities for human and animal consumption, we
9 have developed a rapid, sensitive and specific PCR assay for *P. brevicompactum*.
10 The specific primers were designed on the basis of ITS1-5.8S-ITS2ITS (Internal
11 transcribed Spacers of rRNA genes) multicopy region. This method provides a
12 useful aid to detect the presence of this fungal species in grapes and other
13 commodities in order to prevent the toxins produced entering the food chain.

14

15 **Key words:** *Penicillium brevicompactum*, Mycophenolic acid, PCR-diagnosis,
16 grapes, LC-ion trap-mass spectrometry.

17 **Running title:** Detection of *P. brevicompactum* a mycophenolic acid producer

18

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Introduction

2 Penicillium brevicompactum Dierckx is a ubiquitous fungus which has been
3 isolated from air (Wu *et al.*, 2003), apple production, cheese (Kure *et al.*, 2004;
4 Serra *et al.*, 2003) cashew (Freire *et al.*, 1999), pepper and nuts (Freire *et al.*,
5 2000) and grapes (Barkai-Golan, 1980; Abrunhosa *et al.*, 2001; Serra *et al.*, 2003;
6 Serra *et al.*, 2005). Some *P. brevicompactum* strains from diverse commodities
7 have been described either as mycophenolic acid, brevianamide A, (Paterson *et*
8 *al.*, 1990; Rundberget *et al.*, 2004) or patulin producers (Paterson *et al.*, 2003).
9 Mycophenolic acid-producing strains of *P. brevicompactum* have been isolated
10 from different origins (Vinokurova *et al.*, 2005) such as ginger (Overy & Frisvad,
11 2005), but no reports of mycophenolic acid-producing strains from grapes have
12 been published so far. The mycophenolic acid has been found to affect many
13 physiological processes. It shows antibiotic, anti-tumour, anti-psoriasis and
14 immunosuppressive features, and it is currently used in Medicine (Bentely, 2000).
15 Additionally, it might be relevant in secondary mycotoxicosis (bacterial infection
16 caused by intake of immunosuppressive mycotoxins). Since *P. brevicompactum*
17 produces diverse toxic compounds such as mycophenolic acid, brevianamide A
18 and patulin and it is an ubiquitous fungus, it should be considered as a threat to
19 commodity system (Paterson *et al.*, 2004).

20

21 Penicillium brevicompactum Dierckx (Section Coronata) is included in the
22 important group of terverticillate penicillia which also includes the Sections
23 *Roqueforti*, *Chrysogena*, *Penicillium*, *Digitata* and *Viridicata*. The classification of
24 this group has been in a state of change during the history of the genus.

1
2
3 1 Taxonomic keys to identify *Penicillium* spp. are based primarily on morphological
4
5 2 criteria, such as micromorphology, macromorphology and colours produced in the
6
7 3 mycelium or diffused into the medium (Raper & Thom, 1949; Ramírez, 1982; Pitt,
8
9 4 1991; Samson & Frisvad, 2004). However many authors have shown that some of
10
11 5 these characteristics (i.e. colony texture) are highly variable and therefore do not
12
13 6 always allow unambiguous classification (Frisvad & Filtenborg, 1983). As an
14
15 7 alternative to secondary metabolites profiling (Frisvad & Filtenborg, 1983),
16
17 8 detection of specific enzyme activity and isoenzyme electrophoretic patterns
18
19 9 (Frisvad, 1981; Paterson *et al.*, 1989), have been used. More recently genotypic
20
21 10 characterization (RFLPs, rRNA genes sequence, RAPDs) has proven its utility
22
23 11 (Dupont *et al.*, 1999; Pianzzola *et al.*, 2004). However, although phylogenetic
24
25 12 studies of *P. brevicompactum* and related species have been recently developed
26
27 13 (Peterson, 2000; Peterson *et al.* 2004) a specific method to detect *P.*
28
29 14 *brevicompactum* based on PCR has not been reported.
30
31
32
33
34
35
36
37
38

39 16 The objectives of this work were to analyse the ability to produce mycophenolic
40
41 17 acid by *P. brevicompactum* strains isolated from grapes and to design specific
42
43 18 primers, based on ITS region, and the corresponding PCR assay to detect this
44
45 19 species.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1

2 **Materials and Methods**

3 *Fungal Isolates and culture conditions*

4 All the isolates used in this study, along with their sources, are given in Table I.
5 The isolates were obtained in our laboratory from grapes collected from Spanish
6 vineyards or from culture collections: the Spanish Type Culture Collection (Spain),
7 the Agroindustrial fungi & yeasts collection (MUCL, The Belgian Co-ordinated
8 Collections of Micro-organisms) and the ARS culture collection (NRRL, United
9 States Department of Agriculture).

10

11 [Insert table I about here]

12

13 Cultures were maintained on potato dextrose-agar (PDA) medium (Scharlau
14 Chemie, Barcelona, Spain) at 4° C and stored as spore suspension in 15 %
15 glycerol at –80 °C. The isolates were cultured in 100 mL Erlenmeyer flasks
16 containing 20 mL Sabouraud liquid medium (Scharlau Chemie, Barcelona, Spain).
17 Cultures were inoculated with mycelial disks cut from the plates and incubated at
18 25 °C, 150 rpm. Mycelia from 2-days-old cultures were harvested by filtration
19 through Whatman paper nº 1 and kept at –80 °C for DNA isolation.

20

21 *Fungal Identification*

22 The strains A and U isolated from Spanish grapes (*Tempranillo* variety) were
23 identified by the Spanish Type Culture Collection following the protocol described

1 by Pitt (1999). The strains were inoculated in Czapek yeast extract agar (CYA),
2 incubated for 7 days at 5 °C, 26 °C and 37 °C, in Malt extract agar (MEA) at 26 °C
3 and in 25% glycerol nitrate agar (G25N) at 26 °C. All identifications were confirmed
4 by amplification of 28S rRNA gene (598 bp) and subsequent sequencing. The
5 sequences showed 100% similarity to *P. brevicompactum* strains from the NCBI
6 and MycoBank (CBS) data bases.

7 8 *DNA extraction and PCR amplification.*

9 Genomic DNA of the strains listed in Table I was obtained using either the genomic
10 DNA Extraction Kit (Genomix, Talent, Trieste, Italy), according to the
11 manufacturer's instructions, or the protocol described by Querol *et al.* (1992).

12
13 The ITS1-5.8S-ITS2 region of the nuclear ribosomal DNA from several *Penicillium*
14 strains, including the *P. brevicompactum* strains isolated from grapes, was
15 amplified using primers ITS1 (5' TCCGTAGGTGAACCTGCGG 3') and ITS4 (5'
16 TCCTCCGCTTATTGATATG 3') described elsewhere (White *et al.*, 1990). The
17 PCR reaction was performed in an Eppendorf Mastercycler Gradient (Eppendorf,
18 Hamburg, Germany) using 10 pg-10 ng of genomic DNA. The amplification
19 program used was described by Henry *et al.* (2000) and it is as follows: 1 cycle of 4
20 min 30 s at 95 °C, 40 cycles of 30 s at 95 °C (denaturalization), 30 s at 50 °C
21 (annealing), 60 s at 72 °C (extension) and finally 1 cycle of 3 min at 72 °C. The
22 PCR products generated were purified using the High Pure PCR Product
23 Purification Kit (Roche, Mannheim, Germany) and cloned into the cloning vector
24 pCR4-TOPO with the TOPO TA Cloning Kit for Sequencing (Invitrogen, U.K.).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Plasmids were sequenced in both directions using the ABI PRISM DNA Sequencer
2 (Applied Biosystems, Foster City, USA) according to the manufacturer's
3 instructions in the Genomic Unit of the University Complutense of Madrid (Spain).
4 The ITS1-5.8S-ITS2 sequences of the *P. brevicompactum* strains U, A, and CECT
5 2361 were deposited in the EMBL database under accession numbers AJ877044,
6 AJ877045, and AJ877046, respectively.

7
8 *Development and evaluation of species-specific PCR primers.*

9 The ITS1-5.8S-ITS2 DNA sequences of *P. brevicompactum* and related species
10 obtained in our laboratory and retrieved from data bases (GenBank and EMBL)
11 were analysed and aligned by Clustal method using Dnastar (Lasergene, Madison,
12 Wis., USA). Primers were designed on the basis of these DNA alignments. The
13 primers obtained were named BRE1 (5'GCTGCCGGGGGACATCTGTCC 3') and
14 BRE2 (5'GGCGCCGGCCAGTCCT ACAAAG 3'). The PCR amplification protocol
15 used was as follows: 1 cycle of 4 min 30 s at 95 °C, 25 cycles of 30 s at 95 °C, 20 s
16 at 69.5 °C, 30 s at 72 °C and finally 1 cycle of 5 min at 72 °C. All amplification
17 reactions were carried out in volumes of 25 µL containing 3 µL (10 pg-10 ng) of
18 template DNA, 1.50 µL of each primer (20 µM), 2.5 µL of 10 x PCR buffer, 1 µL of
19 MgCl₂ (50 mM), 0.25 µL of dNTPs (100 mM) and 0.2 µL of Taq DNA polymerase (5
20 U/µL) supplied by the manufacturer (Ecogen, Barcelona, Spain). PCR products
21 were detected in 2% agarose ethidium bromide gels in TAE 1 X buffer (Tris-acetate
22 40 mM and EDTA 1.0 mM). The DNA ladder "Real escala nº2" (Durviz, Valencia,
23 Spain) was used as molecular size marker.

1 Genomic DNA from the strains indicated in Table I was used for PCR amplification
2 using the primers BRE1/BRE2 in the conditions described above. The genomic
3 DNA from all the strains was tested for PCR amplification using primers ITS1 and
4 ITS4 according to Henry *et al.* (2000) using the same master mix used for specific
5 PCR amplification except primers.

7 *Reagents and standards for mycotoxin assays*

8 Mycophenolic acid standards were purchased from Sigma (Sigma-Aldrich,
9 Alcobendas, Spain). Acetonitrile, chloroform, acetic acid, methanol (all LC grade)
10 and phosphoric acid (85%, A.R.) were from J.T. Baker (Deventer, Holland). Pure
11 water was obtained from a Milli-Q Plus apparatus (Millipore, Billerica, Ma., USA)
12 and was used when water was required. Yeast extract and sucrose were
13 purchased from Panreac (Montcada i Reixac, Barcelona, Spain)

15 *Mycotoxin production*

16 One *Penicillium brevicompactum* strain (CECT 2316) and two isolates of this
17 species (A and U) from Spanish grapes (Table I) were tested for mycophenolic
18 acid production using Yeast Extract Sucrose (YES) broth (Singh *et al.*, 1991; Overy
19 & Frisvad, 2005) and grape must:water (20:80 v/v). Grape must:water (20:80 v/v)
20 was used to find out whether this substrate had any effect on the fungal
21 development and/or production of the studied mycotoxins, as two isolates were
22 from grapes grown in Spain.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Liquid medium (50 mL) was poured into 100 mL Erlenmeyer flasks. The flasks
2 were plugged with cotton, covered with aluminium foil and autoclaved at 115 °C for
3 30 min. Sterilised liquid media were inoculated with 1 mL of a spore suspension
4 made in aqueous Tween 80 (0.5%) having 10⁵ spores/mL and then they were
5 incubated at 25 °C for 2 weeks. Three flasks per isolate were assayed for
6 mycotoxin accumulation. Controls consisted of non-inoculated sterilized liquid
7 media incubated under the same conditions.

8

9 *Mycotoxin extraction, clean-up and analysis*

10 Extraction of mycophenolic acid and HPLC analysis was carried out according to
11 the method described by Medina *et al.* (2004) with slight modifications.

12

13 Five ml of culture medium were mixed with 500 µL of a 0.1 M solution of
14 phosphoric acid, extracted with 2 x 5 mL chloroform and, after evaporation of the
15 organic phase, the residue was dissolved in 0.5 mL of mobile phase
16 (acetonitrile:water:acetic acid 47:52:1 v/v/v).

17

18 *Chromatographic equipment*

19 The LC system consisted of a Waters 600 pump equipped with a Waters 717
20 automatic injector and a Waters 474 scanning fluorescence detector (FLD)
21 (Waters). A Waters 996 photodiode array detector (DAD) was usually set after the
22 fluorescence detector. Millennium 32® software ver. 4.0 (Waters) was used to
23 control the chromatograph and to process the signals. Separations were performed
24 on a stainless steel LiChrospher 100 C₁₈ reversed-phase column (250 x 4 mm, 5

1 μm particle size) (Agilent Technologies, Waldbronn, Germany) connected to a
2 guard column (4 x 4 mm, 5 μm particle size) filled with the same phase. The
3 column was kept at 40 °C. Alternatively, separation was performed on a
4 Phenomenex Gemini[®] C₁₈ column (150 x 4.6 mm, 5 μm particle size)
5 (Phenomenex, Torrance, Ca., USA) connected to a Phenomenex Gemini[®] C₁₈
6 SecurityGuard cartridge (4 x 3 mm, 5 μm particle size). The column was kept at 35
7 °C. The flow rate was 1 mL/min and the mobile phase was acetonitrile:water:acetic
8 acid (47:52:1 v/v/v). Mycophenolic acid was quantified by DAD at 254 nm.

9 10 *Toxin confirmation by LC-MS*

11 LC-ion trap-mass spectrometry was used to unambiguously confirm the possible
12 presence of mycophenolic acid. The analysis was carried out on an Agilent 1100
13 liquid chromatograph (Agilent Technologies), equipped with a Zorbax SB-C₁₈
14 column (150 x 4.6 mm, 5 μm particle size) (Agilent Technologies) and coupled to a
15 Bruker Esquire 3000 Plus ion trap mass spectrometer (Bruker Instruments,
16 Billerica, Ma., USA).

17
18 For the analysis of mycophenolic acid the mobile phase was
19 acetonitrile:water:acetic acid (50:49:1, v/v/v). The ionization method was
20 electrospray ionization (ESI) in positive mode by using the following ionization
21 source parameters: N₂ nebulizer gas at 10 psi, dry gas at 5 L/min, dry temperature
22 at 250 °C, and capillary voltage at 4.5 kV.

23

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Results

2 Two specific primers to detect *P. brevicompactum* were designed on the basis of
3 the alignment of ITS1-5.8S-ITS2 regions performed with a representative sample
4 of *Penicillium* species available in our laboratory, including A and U isolates from
5 grapes and closely related species: *P. oslonii* and *P. biourgeianum* (Peterson *et al.*,
6 2004), the latter considered a synonym of *P. bialowiezense* according to Frisvad &
7 Samson (2004). Additionally, 26 sequences of *P. brevicompactum* from the
8 GenBank were also used. Figure 1 shows ITS1-5.8S-ITS2 sequence of *P.*
9 *brevicompactum* isolate A (AJ877045) indicating the position of primers BRE-1 and
10 BRE-2. Primer BRE1 was located within the ITS1, at the position +77, and the
11 primer BRE2 at the position +445 (within the ITS2).

12
13 [insert figure 1 about here]

14
15 All the *Penicillium* strains listed in Table I were tested for amplification using primer
16 pair BRE1/BRE2. A single fragment of about 390 bp was amplified only when
17 genomic DNA from *P. brevicompactum* strains was used. No product was
18 observed with genomic DNA from other *Penicillium* isolates, nor in the case of
19 species belonging to other *genera* (Figure 2). Control amplifications of the genomic
20 DNA with primers ITS1 and ITS4 were positive for all the strains analysed.

21
22 [Insert figure 2 about here]

23

1 The ability of *P. brevicompactum* strains A, U and CECT 2361 to produce
2 mycophenolic acid was tested by chromatographic analysis. The chromatographic
3 analysis of culture media using the Phenomenex Gemini[®] C₁₈ column showed the
4 presence of a peak with a retention time of 5.4 min that was tentatively identified as
5 mycophenolic acid. The identity of mycophenolic acid was confirmed by the UV
6 spectrum that showed absorption maxima at 226.2 nm, 250.9 nm and 305.2 nm,
7 which matched to the absorption maxima shown by the standard. The
8 mycophenolic acid peak was found in the cultures of the three *P. brevicompactum*
9 strains (Figure 3).

10
11 [Insert figure 3 about here]

12
13 The presence of mycophenolic acid in cultures was confirmed by LC-MS based on
14 the protonated molecule $[M+H]^+$ and the most abundant product ion $[(M+H)^+ +$
15 $H_2O)]$ whose m/z ratios were 321 and 338, respectively. The confirmation was also
16 made using the second fragmentation of ion 321 (LC-MS-MS), obtaining in both
17 standard and sample an ion with m/z 303 corresponding to the fragmentation
18 $[(M+H)^+ - H_2O]$.

19
20 Production of mycophenolic acid was compared in two different media, YES and
21 grape must:water (20:80 v/v). The results obtained did not show significant
22 differences between both media (Table II).

23 [insert table II about here]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Discussion

2 We have developed the first PCR detection assay specific for *P. brevicompactum*.
3 The specificity of this assay has been successfully tested in a wide and diverse
4 sample of *P. brevicompactum* strains associated to different substrates, other
5 *Penicillium* species and other fungal species often associated to grapes and
6 related commodities. This PCR method represents a good alternative to
7 conventional identification methods, especially difficult in the case of terverticillate
8 penicillia (Paterson *et al.*, 2004), and does not require highly trained specialists or
9 expensive equipment (Paterson *et al.*, 2004). The PCR assay described in this
10 work also represents an advantage in terms of time of analysis and specificity in
11 comparison with the conventional methods and will facilitate detection of this
12 species in future surveys of agro-products.
13
14 The assay developed in this study targeted the multicopy ITS1-5.8S-ITS2 region in
15 order to improve the sensitivity of the detection. The detection limit of ITS
16 amplification product, defined as the clearly visible product on agarose gels
17 containing ethidium bromide, has been estimated between 1 to 10 pg of DNA
18 template in *Fusarium* (Bluhm *et al.*, 2002). The sensitivity of our PCR assay based
19 on ITS sequences would be in the same order of magnitude and, therefore, more
20 sensitive than primers based on single copy gene, estimated between 0.1 and 1 ng
21 of DNA template per reaction (Bluhm *et al.*, 2002).
22
23 Those features are very important since *P. brevicompactum* is a ubiquitous and
24 cosmopolitan fungi that has been isolated from widespread substrates.

Furthermore, most *P. brevicompactum* strains isolated from commodities other than grapes were capable to produce mycophenolic acid. Some authors have described production of this metabolite by all the *P. brevicompactum* isolates tested (Overy & Frisvad, 2005) while Vinokurova *et al.* (2005), studying isolates from different origins, pointed out that 60 % of the strains tested were able to produce mycophenolic acid. In the present work, we also report production of mycophenolic acid by *P. brevicompactum* isolated from grapes for the first time. The high incidence of *P. brevicompactum* strains capable to produce mycophenolic acid stresses the relevance of this species in terms of food safety.

The levels produced by our isolates grown under the conditions used in this work were relatively low if compared with levels reported by other authors who found *in vitro* production of this compound around 300 mg/L (Vinokurova *et al.*, 2005). In their work, however, these authors suggested a strong influence of culture media on the mycophenolic acid production. Therefore, more studies are necessary to propose an optimised culture medium to evaluate this metabolite.

The cultivation of *P. brevicompactum* collected from grapes on a grape-like media (grape must: water, 20:80 v/v) did not result in enhanced levels of mycophenolic acid production in comparison with YES medium. However, more studies are needed to confirm that production of mycophenolic acid by *P. brevicompactum* isolates would not be positively influenced by grape substrate.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The occurrence of *P. brevicompactum* strains in grapes and their ability to produce
2 mycophenolic acid reported here increases the number and importance of the
3 dietary foods and derivatives susceptible to be contaminated by this metabolite and,
4 therefore, we suggest that its inclusion in future survey programs should be
5 considered by food safety organisations as well as more exhaustive studies on the
6 effects of mycophenolic acid on the immune response in humans (Overy & Frisvad,
7 2005).

8

9 **Acknowledgements**

10

11 The authors wish to thank financial support from the Spanish Government
12 “Ministerio de Educación y Ciencia” (Project AGL-2001-2974-C05-01 and AGL-
13 2004-07549-C05-02/ALI). The authors also acknowledge the Valencian
14 Government (Conselleria d’Empresa, Universitat i Ciència) for financing by Project
15 GV04B-111 and for a research grant.

16

1 References

- 2 Abrunhosa L, Paterson RRM, Kozakiewicz Z, Lima N, Venâncio A. 2001 Mycotoxin
3 production from fungi isolated from grapes. Letters in Applied Microbiology 32:
4 240-242.
- 5 Barkai-Golan R. 1980 Species of *Penicillium* causing decay of stored fruits and
6 vegetables in Israel. Mycopathologia et Mycologia Applicata 54: 141–145.
- 7 Bau M, Bragulat MR, Abarca ML, Minguez S, Cabañes FJ. 2005 Ochratoxigenic
8 species from Spanish wine grapes. International Journal of Food Microbiology
9 98: 125-130.
- 10 Bentley R. 2000 Mycophenolic acid: A hundred year odyssey from antibiotic to
11 immunosuppressant. Chemical Reviews 100: 3801-3825.
- 12 Bluhm BH, Flaherty JED, Cousin MA, Woloshuk CP. 2002 Multiplex Polymerase
13 Chain Reaction Assay for the Differential Detection of Trichothecene- and
14 Fumonisin-Producing Species of *Fusarium* in Cornmeal. Journal of Food
15 Protection 65: 1955-1961.
- 16 Dupont J, Magnin S, Marti A, Brousse M. 1999 Molecular tools for identification of
17 *Penicillium* starter cultures used in food industry. International Journal of Food
18 Microbiology 49: 109-118.
- 19 Freire FCO, Kozakiewicz Z, Paterson RRM. 1999 Mycoflora and mycotoxins of
20 Brazilian cashew kernels. Mycopathologia 145: 103-199.
- 21 Freire FCO, Kozakiewicz Z, Paterson RRM. 2000 Mycoflora and mycotoxins of
22 Brazilian black pepper, white pepper and Brazil nuts. Mycopathologia 149: 13-
23 19.

- 1
2
3 1 Frisvad JC.1981. Physiological criteria and mycotoxin production as aids in
4
5 2 identification of common asymmetric penicillia. Applied and Environmental
6
7 3 Microbiology 41: 568-579.
8
9
10 4 Frisvad JC, Filtenborg O. 1983. Classification of terverticillate penicillia based on
11
12 5 profiles of mycotoxins and other secondary metabolites. Applied and
13
14 6 Environmental Microbiology 46: 1301-1310.
15
16
17 7 Frisvad JC, Samson RA. 2004. Polyphasic taxonomy of *Penicillium* subgenus
18
19 8 *Penicillium*. A guide to identification of food and air-borne terverticillate
20
21 9 Penicillia and their mycotoxin. In: Samson RA, Frisvad JC, editors. Studies
22
23 10 in Mycology. CBS. Utrecht. The Netherlands. Vol. 49,pp. 1-173
24
25
26 11 Henry PC, Iwen S, Hinrichs H. 2000 Identification of *Aspergillus* species using
27
28 12 internal transcribed spacer regions 1 and 2. Journal of Clinical Microbiology
29
30 13 38: 510-515.
31
32
33 14 Kure CF, Skaar I, Brendehaug J. 2004 Mould contamination in production of semi-
34
35 15 hard cheese. International Journal of Food Microbiology 93: 41-49.
36
37
38 16 Medina A, González G, Sáez JM, Mateo R, Jiménez M. 2004 Bee pollen, a
39
40 17 substrate that stimulates ochratoxin A production by *Aspergillus ochraceus*
41
42 18 Wilh. Systematic and Applied Microbiology 27: 261-267.
43
44
45 19 Overy DP, Frisvad JC. 2005 Mycotoxin production and postharvest storage rot of
46
47 20 ginger (*Zingiber officinale*) by *Penicillium brevicompactum*. Journal of Food
48
49 21 Protection 68: 607-609.
50
51
52 22 Paterson RRM, Venancio A, Lima N. 2004. Solutions to *Penicillium* taxonomy
53
54 23 crucial to mycotoxin research and health. Research in Microbiology 155:
55
56 24 507-513.
57
58
59
60

- 1 Paterson RRM, Bridge PD, Crosswaite MJ, Hawksworth DL. 1989. A reappraisal of
2 terverticillate penicillia using biochemical, physiological and morphological
3 features III. An evaluation of pectinase and amylase isoenzymes for species
4 characterization. *Journal of General Microbiology* 135: 2979-2991.
- 5 Paterson RRM, Simmonds MJ, Kemmelmeier C, Blaney W. 1990. Effects of
6 brevianamide A, and its photolysis product Brevianamide D, and ochratoxin
7 A from two *Penicillium* strains on the insect pests *Spodoptera frugiperda* and
8 *Heliothis verescans*. *Mycological Research* 94: 538-542
- 9 Paterson RRM, Kozakiewicz Z, Locke T, Brayford D, Jones SCB. 2003. Novel use
10 of the isoepoxydon dihydrogenase gene probe of the patulin metabolic
11 pathway and chromatography to test penicillia isolates from apple
12 production systems for the potential to contaminate apple juice with patulin.
13 *Food Microbiology*. 20: 359-364.
- 14 Peterson SW. 2000. Phylogenetic analysis of *Penicillium* species based on ITS
15 and LSU-RNA nucleotide sequences. In Samson RA, Pitt JI, editors.
16 Harwood Academic Publisher, Amsterdam. pp. 163-178.
- 17 Peterson SW. 2004. Multilocus DNA sequence analysis shows that *Penicillium*
18 *biourgeianum* is a distinct species closely related to *P. brevicompactum* and
19 *P. oslonii*. *Mycological Research* 108: 434-440.
- 20 Pianzzola MJ, Moscatelli M, Vero S. 2004. Characterization of *Penicillium* isolates
21 associated with blue mold on apple in Uruguay. *Plant Diseases* 88: 23-28.
- 22 Pitt JI. 1991. *A laboratory guide to common Penicillium species*, In: Food Science
23 Pitt JI editor. North Ryde, Australia.

- 1
2
3 1 Querol A, Barrio E, Huerta T, Ramón D. 1992 Molecular monitoring of wine
4
5 2 fermentations conducted by active dry yeast strains. Applied and
6
7 3 Environmental Microbiology 58: 2948-2953.
8
9
10 4 Ramirez C. 1982. Manual and atlas of the penicillia. Elsevier Biomedical press,
11
12 5 Amsterdam.
13
14 6 Raper KC, Thom CA. 1949. Manual of the penicillia. The Williams & Wilkins Co,
15
16 7 Baltimore.
17
18
19 8 Rundberget T, Skaar I, Flaoyen A. 2004. The presence of *Penicillium* and
20
21 9 *Penicillium* mycotoxins in food wastes. International Journal of Food
22
23 10 Microbiology 90:181-188.
24
25
26 11 Samson RA, Frisvad GC. 2004. *Penicillium* subgenus *Penicillium*: new taxonomy
27
28 12 schemes, mycotoxin and other extrolites. In Samson RA, Frisvad GC,
29
30 13 Editors. Vol. 49. CBS. Utrecht. The Netherlands
31
32
33 14 Serra R, Braga A, Venâncio A. 2005. Mycotoxin-producing and other fungi isolated
34
35 15 from grapes for wine production, with particular emphasis on ochratoxin A.
36
37 16 Research in Microbiology 156: 515-521.
38
39
40 17 Serra R, Abrunhosa L, Kozakiewicz Z, Venancio A. 2003 Black *Aspergillus* species
41
42 18 as ochratoxin A producers in Portuguese wine grapes. International Journal
43
44 19 of Food Microbiology 88: 63– 68.
45
46
47
48 20 Singh K, Frisvad JC, Thrane U, Mathur SB. 1991. An illustrated manual on
49
50 21 identification of some seed-borne *Aspergilli*, *Fusaria*, *Penicillia* and their
51
52 22 mycotoxins. Danish government. Institute of Seed Pathology for Developing
53
54 23 Countries Editors. Hellerup. pp. 8-12
55
56
57
58
59
60

- 1
2
3 1 Vinokurova NG, Ivanushkina NE, Kochkina GA, Arinbasarov MU, Ozerskaya SM.
4
5 2 2005 Production of mycophenolic acid by fungi of the genus *Penicillium*
6
7 3 Link. Applied Biochemistry and Microbiology 41: 83-86.
8
9
10 4 White TJ, Burns T, Lee S, Taylor JW. 1990. Amplification and direct sequencing of
11
12 5 fungal ribosomal RNA genes for phylogenetics. Innis MA, Gelgard DH,
13
14 6 Sninsky JJ, White TJ Editors. Academic Press, New York. pp. 315-322.
15
16
17 7 Wu Z, Tsumura Y, Blomquist G, Wang X. 2003 18S rRNA gene variation among
18
19 8 common airborne fungi, and development of specific oligonucleotide probes
20
21 9 for the detection of fungal isolates. Applied and Environmental Microbiology
22
23
24 10 69: 5389-5397.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Sequence of ITS1-5.8S-ITS2 of *Penicillium brevicompactum* isolate A (AJ877045). Specific Primers BRE1/2 are underlined. The genes 18S, 5.8S and 28S are shadowed.

TTCCGTAGGTGAACCTGCGGAAGGATCATTACGGAGTGAGGGCCCTCTGGGTCCAACCTC 60

CCACCCGTGTTTATTTTACCTTGTGCTTCGGCGAGCCTGCCTTTTGGCTGCCGGGGGAC 120

ATCTGTCCCCGGGTCCGCGCTCGCCGAAGACACCTTAGAACTCTGTCTGAAGATTGTAGT 180

CTGAGATTAAATATAAATTATTTAAAACTTTCAACAACGGATCTCTTGGTTCCGGCATCG 240

ATGAAGAACGCAGCGAAATGCGATACGTAATGTGAATTGCAGAATTCAGTGAATCATCGA 300

GTCTTTGAACGCACATTGCGCCCTCTGGTATTCCGGAGGGCATGCCTGTCCGAGCGTCAT 360

TGCTGCCCTCAAGCACGGCTTGTGTGTTGGGCTCCGTCTCCTTCCGGGGGACGGGCCCG 420

AAAGGCAGCGGGCGCACCGGTCCGGTCCTCAAGCGTATGGGGCTTTGTCACCCGCTTTG 480

TAGGACTGGCCGGCGCCTGCCGATCAACCAAACCTTTTCCAGGTGACCTCGATCAGGAGGG 540

ATACCCGCTGAACTTAAGCATATCAATAA 569

Figure 2. PCR amplification using primers BRE1/2 and DNA from *P. brevicompactum* CECT 2316, MUCL 29153, MUCL 38823, MUCL 39543, A and U lanes (2-7) respectively, *Penicillium chrysogenum* CECT 2306 (lane 8), *P. expansum* CECT 2280 (lane 9), *P. simplicissimum* CECT 2747 (lane 10), *P. aurantogriseum* CECT 2320 (lane 11), *P. commune* MUCL 31780 (lane 12), *P. oslonii* NRRL 13634 (lane 13) and *P. biourgeianum* NRRL 865 (lane 14). Lane 1: Non-template control, M: DNA marker

Figure 3. LC-PDA (254nm) chromatogram of water:grape must (80:20 v/v) culture of *P. brevicompactum* (isolate A).

Mycophenolic acid (µg/ml)				
Isolate	YES medium	RSD %	Grape must:water (20:80 v/v)	RSD %
2316	148.878	7.30	90.910	11.37
A	130.528	7.38	115.309	7.18
U	183.256	1.70	134.350	4.81

Table II: Mycophenolic acid production by strains of *P. brevicompactum* isolated from grapes grown in Spain. RSD: relative standard deviation.

Strains	Species	BRE1-2
A (a)	<i>P. brevicompactum</i>	+
U (a)	<i>P. brevicompactum</i>	+
CECT 2316	<i>P. brevicompactum</i>	+
MUCL 29153	<i>P. brevicompactum</i>	+
MUCL 38823	<i>P. brevicompactum</i>	+
MUCL 39543	<i>P. brevicompactum</i>	+
NRRL 2011	<i>P. brevicompactum</i>	+
NRRL 28120	<i>P. brevicompactum</i>	+
NRRL 865	<i>P. biourgeianum</i>	-
NRRL 28149	<i>P. biourgeianum</i>	-
NRRL 13634	<i>P. oslonii</i>	-
NRRL 31467	<i>P. oslonii</i>	-
CECT 2306	<i>P. chrysogenum</i>	-
CECT 2264	<i>P. chrysogenum</i>	-
CECT 2784	<i>P. chrysogenum</i>	-
CECT 2668	<i>P. chrysogenum</i>	-
CECT 2280	<i>P. expansum</i>	-
CECT 2275	<i>P. expansum</i>	-
CECT 2747	<i>P. simplicisimum</i>	-
CECT 2746	<i>P. simplicisimum</i>	-
CECT 2918	<i>P. aurantiogriseum</i>	-
CECT 2320	<i>P. aurantiogriseum</i>	-
CECT 2270	<i>P. corylophium</i>	-
CECT 2273	<i>P. corylophium</i>	-
CECT 2735	<i>P. raistrickii</i>	-
CECT 2737	<i>P. raistrickii</i>	-
CECT 2736	<i>P. raistrickii</i>	-

MUCL 28653	<i>P. commune</i>	-
MUCL 31780	<i>P. commune</i>	-
CECT 2906	<i>P. verrucosum</i>	-
C.AL.P.1 (a)	<i>P. sclerotiorum</i>	-
L (a)	<i>P. polonicum</i>	-
R.Te.P.4 (a)	<i>P. chrysogenum</i>	-
B.Me.P.13 (a)	<i>P. expansum</i>	-
CL.1 (a)	<i>Cladosporium sp</i>	-
UCO.1 (a)	<i>Alternaria consortiale</i>	-
T.MV.F.1 (a)	<i>Fusarium sp.</i>	-
BO.1 (a)	<i>Botrytis sp</i>	-
CECT 2086	<i>Aspergillus carbonarius</i>	-
Z.M.A.29 (a)	<i>Aspergillus niger</i>	-
T.TT.A5 (a)	<i>Aspergillus tubingensis</i>	-
T.MV.A .21 (a)	<i>Aspergillus fumigatus</i>	-
U-2003 (a)	<i>Aspergillus ochraceus</i>	-

Table I: Strains analysed indicating species and the occurrence of PCR amplification product with the pair of primers BRE1-2. (a): Strains isolated from different grape varieties