Effect of pH on ochratoxin A production by *A. niger* aggregate species

Alexandre Esteban, M Lourdes Abarca, M Rosa Bragulat, F Javier Cabañes

► To cite this version:

Alexandre Esteban, M Lourdes Abarca, M Rosa Bragulat, F Javier Cabañes. Effect of pH on ochratoxin A production by *A. niger* aggregate species. *Food Additives and Contaminants*, 2006, 23 (06), pp.616-622. 10.1080/02652030600599124 . hal-00577290

HAL Id: hal-00577290

<https://hal.science/hal-00577290>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of pH on ochratoxin A production by *A. niger* aggregate species

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2005-277.R2
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	10-Jan-2006
Complete List of Authors:	Esteban, Alexandre; Universitat Autònoma de Barcelona, Sanitat i Anatomia Animals Abarca, M Lourdes; Universitat Autònoma de Barcelona, Sanitat i Anatomia Animals Bragulat, M Rosa; Universitat Autònoma de Barcelona, Sanitat i Anatomia Animals Cabañes, F Javier; Universitat Autònoma de Barcelona, Sanitat i Anatomia Animals
Methods/Techniques:	Chromatography - HPLC
Additives/Contaminants:	Ochratoxin A
Food Types:	Coffee, Wine, Dried fruit

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Effect of pH on ochratoxin A production by *A. niger* aggregate species

A Esteban, ML Abarca, MR Bragulat, FJ Cabañes*

Grup de Micologia Veterinària, Departament de Sanitat i d'Anatomia Animals,
Facultat de Veterinària, Universitat Autònoma de Barcelona. 08193 Bellaterra
(Barcelona), Spain.

***Corresponding author:**

F.Javier Cabañes
Grup de Micologia Veterinària
Departament de Sanitat i d'Anatomia Animals
Facultat de Veterinària, Universitat Autònoma de Barcelona
08193 Bellaterra (Barcelona), Spain.

Tel: 34 93 5811749
Fax: 34 93 5812006
e-mail: Javier.cabanes@uab.es

Abstract

The effect of pH (2-10) on growth and ochratoxin A (OTA) production by twelve *Aspergillus niger* aggregate strains was studied in two culture media: Czapek Yeast Autolysate agar (CYA) and Yeast Extract Sucrose agar (YES), during a period of 30 days. The strains were selected to include different sources, different reported abilities to produce OTA and different ITS-5.8S rDNA RFLP patterns. YES was found to be a better culture medium than CYA for OTA production. In this medium, OTA was produced from pH 2 or pH 3 to pH 10 depending on the strain. Our results show the ability of *A. niger* aggregate strains not only to grow but also to produce OTA over a wide pH range. These results will lead to a better understanding of the role of *A. niger* aggregate strains in the OTA contamination of several food commodities.

Keywords: *Aspergillus niger*, coffee, grapes, raisins, ochratoxin A, pH.

Introduction

Among the different mycotoxins which merit special concern for the hazard they represent in food commodities, ochratoxin A (OTA) deserves particular attention. OTA is a nephrotoxic mycotoxin which also possesses carcinogenic, teratogenic, genotoxic and immunosuppressive potential (O'Brien and Dietrich 2005). The International Agency for Research on Cancer (IARC 1993) has classed OTA as a possible human carcinogen (group 2B). The European Commission has established limits for OTA in cereals, dried vine fruits, wine, coffee and infant foods. These limits will be reviewed in the short term and maximum limits for more products such as cocoa, beer, meat products, spices or liquorice will be included (Commission Regulation 2002, 2004, 2005).

OTA was until recently believed to be produced only by *Aspergillus ochraceus* and by *Penicillium verrucosum*. The description of OTA production by two species belonging to *Aspergillus* section *Nigri*, *Aspergillus niger* var. *niger* (Abarca et al. 1994) and *Aspergillus carbonarius* (Horie 1995), may contribute to explain the widespread OTA contamination reported in some food and feed products. This distribution was unclear only taking into account the incidence and natural occurrence of the previously described OTA-producing species.

Within *Aspergillus* section *Nigri* it is known that isolates included in the *A. niger* aggregate can produce a range of different toxic metabolites (Frisvad and Thrane 2000) but it is the ability of some of their members to produce OTA what deserves special concern. It is considered that in substrates such as grapes, raisins and wine

the source of OTA contamination detected is due to *A. carbonarius* and some species included in the *A. niger* aggregate (Abarca et al. 2004). Furthermore, it is worth noting that *A. niger* products hold the GRAS (Generally Regarded as Safe) status from the United States, Food and Drug Administration and are used as a source of enzymes and organic acids for food processing, despite the reported ability to produce OTA (Bigelis and Lausure 1987).

The taxa included in the *A. niger* aggregate are extremely difficult to distinguish one from each other by morphological means. In an attempt to solve the classification of these fungi, several molecular techniques have been carried out in order to clarify the taxonomy and divide these taxa into two or more species (Abarca et al. 2004). On the basis of their ITS-5.8S rDNA restriction fragment length polymorphism (RFLP), the strains were grouped in two patterns, designated N and T (Accensi et al. 1999). To date the OTA-producing isolates with RFLP pattern known are classified as type N (Accensi et al. 2001; Cabañes et al. 2002; Abarca et al. 2003; Leong et al. 2004; Esteban et al. 2004).

Little is known about the environmental factors which can affect the OTA production by species of the *A. niger* aggregate. In a previous study on the effect of temperature, the *A. niger* aggregate strains showed the highest levels of OTA in Yeast Extract Sucrose (YES) agar at 20-25°C (Esteban et al. 2004).

As little is known about the influence of pH, the aim of this study was to establish the effect of pH on growth and OTA production by different species belonging to the *A. niger* aggregate at 25°C over a period of 30 days. The strains were selected to

include different sources, different reported ochratoxigenic ability and different ITS-5.8S rDNA RFLP patterns.

Material and methods

Fungal strains and inocula preparation

Twelve isolates of the *A. niger* aggregate were used in this study (ten of N RFLP pattern and two of T pattern). The reported OTA production, source of isolation and RFLP pattern is shown in Table I. The strains were grown on malt extract agar (Pitt and Hocking 1997) for 7 days at 25°C. Conidia suspensions of each isolate were prepared in an aqueous solution of 0.05% Tween 80. After filtering through sterile cheesecloth they were adjusted to approximately 10⁶-10⁷ conidia per ml as determined by a counting chamber.

Culture conditions

OTA production was determined on two culture media: yeast extract sucrose (YES) agar and Czapek yeast extract (CYA) agar (Pitt and Hocking 1997). The pH of the culture media was varied from 2 to 10. Adjustment of pH was performed before autoclaving by adding HCl (1 and 10 N) and NaOH (1 N). The pH value was measured using a pHmeter GLP21 (Crison Strumenti S.p.A., Carpi, Italy). The sterilization of the media adjusted at pH 2, 3 and 4 was carried out by autoclaving the agar separately from the rest of components in order to avoid solidification problems. The pH of each media was checked after sterilization. Each plate was point inoculated with 1 µl of the adjusted suspension. Plates were incubated at 25°C. Each assay was performed in duplicate.

OTA production and quantification

OTA production was determined after 5, 10, 15, 20 and 30 days of incubation at each pH value following a previously described high-pressure liquid chromatography (HPLC) screening method (Bragulat et al. 2001), being efficient at the range of pH tested. On each sampling occasion, three agar plugs were removed from different points of the colony and extracted with 0.5 ml of methanol. The extracts were filtered and injected into the HPLC. OTA detection and quantification was made by a Waters LCM1 chromatograph with a fluorescence detector Waters 2475 (excitation wavelength: 330 nm/emission wavelength: 460 nm). Twenty μ l of each extract were applied. The mobile phase, with a flow rate of 1 ml/min, consisted of the following isocratic program: acetonitrile, 57%; water, 41% and acetic acid, 2% (Bauer and Gareis 1987). The extracts with the same retention time as OTA (around 6.8 min), were considered positive. Confirmation was made through derivatization of OTA in its methyl-ester (Hunt et al. 1980). The detection limit of the extraction procedure and the HPLC technique was 0.02 ng OTA and the quantification limit of HPLC technique with the extraction procedure was 0.01 μ g g⁻¹ for this mycotoxin.

Data analysis

Data obtained were analyzed statistically by means of the one-way analysis of variance test and Student's test. All statistical analyses were performed using SPSS software (Version 12.0).

Results and discussion

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The twelve strains grew in CYA and YES media adjusted from pH 2 to 10. According to our results, the pH range for growth in *Aspergillus niger* was previously reported to be from 1.4 to 9.8 (Panassenko 1967).

Six out of the 12 strains tested produced detectable levels of OTA in both media. Tables II and III show the concentration detected at each pH value and incubation time in CYA and YES medium respectively. In CYA medium most of the strains produced OTA at all the pH values tested. The strain CBS 618.78 produced OTA from pH 4 to 8 and A-943 was a weak OTA-producing strain with detected values very close to the detection limit of the technique at some pH values. In YES medium, the strains produced OTA from pH 2 (strains A-942, A-943 and CBS 139.52) or pH 3 (strains A-75 and A-136) to pH 10. However, this range was lower for the strain CBS 618.78 which produced OTA from pH 5 to 8.

Mean OTA concentration produced at each pH value by the six positive strains is shown in Figure 1. The concentration achieved by all the strains in YES was higher than in CYA medium except at pH 3 (Figure 1). At this pH value the strains showed a slight higher OTA production in CYA medium. The total mean OTA concentration produced in YES medium ($4.42 \mu\text{g g}^{-1}$) was significantly higher (Student's t $P < 0.01$) than in CYA medium ($1.38 \mu\text{g g}^{-1}$) which supports that YES is better than CYA medium for OTA production. This result was in accordance with previous studies which reported the suitability of YES medium to test OTA production by *A. niger* aggregate strains (Bragulat et al. 2001; Esteban et al. 2004). On the contrary, CYA medium has been reported to be more suitable than YES medium in *A. carbonarius* (Bragulat et al. 2001; Esteban et al. 2004; Esteban et al. 2005).

1
2
3
4
5
6 In both media most of the strains produced OTA after only 5-10 days of incubation
7
8 and showed different optimal values for OTA production. In CYA medium (Table 2)
9
10 the strains showed the optimal OTA production within the range pH 3-8. In YES
11
12 medium (Table 3) the optimal conditions for OTA production were reached within the
13
14 range pH 5-10 and always after 5-10 days of incubation. In both media, the strain A-
15
16 943, isolated from grapes, showed the lowest OTA production in comparison with the
17
18 rest of strains studied.
19
20
21

22
23
24 In agreement with previous studies (Esteban et al. 2004) the OTA content decreased
25
26 with increasing incubation time. It has been suggested that this may be due to the
27
28 fact that the strains remove and assimilate the phenylalanine moiety from the OTA
29
30 molecule as other nitrogen sources become exhausted (Téren et al. 1996).
31
32
33

34
35
36 It is not known yet why the percentage of OTA-producing strains in the *A. niger*
37
38 aggregate is low (Abarca et al. 2004), nor if this is due to a genetic based feature. To
39
40 clarify this fact it is necessary to study the effect of different environmental conditions
41
42 on OTA production. In this study, the six strains initially considered as OTA negative
43
44 did not produce the mycotoxin at any of the pH values tested for the period of 30
45
46 days. In a previous study, these six strains did not produce OTA at any of the
47
48 temperatures tested (Esteban et al. 2004). The strains with the RFLP pattern of type
49
50 T did not produce OTA in the range of pH values assayed. To date OTA production
51
52 has never been demonstrated in type T strains (Accensi et al. 2001; Cabañes et al.
53
54 2002; Abarca et al. 2003; Leong et al. 2004; Esteban et al. 2004).
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

There is no previous published work regarding the effect of pH on OTA production by *A. niger* aggregate strains. Nevertheless, in the case of *A. carbonarius* it has been recently reported that the strains are able to produce OTA in a broad pH range (pH 2 to 10) (Esteban et al. 2005). The results obtained here with the *A. niger* aggregate strains state that they are also able to grow at a wide range of pH values (pH 2-10). This would explain why *A. niger* is so commonly isolated from a great variety of food commodities and is considered the most common *Aspergillus* species responsible for the post-harvest decay of fresh fruit (Pitt and Hocking 1997). The black aspergilli belonging to the *A. niger* aggregate are the predominant *Aspergillus* species isolated from grapes. They are isolated along the different developmental stages of berries although it is at harvesting when the highest levels are observed (Battilani et al. 2003; Serra et al. 2003; Bau et al. 2005). The black spores provide protection from sunlight and UV light, providing a competitive advantage in such habitats (Pitt and Hocking 1997). It is worth mentioning that the pH of grapes increases during ripening, achieving a final pH range of 3-4 (Splittstoesser 1987; FDA 2003). *Aspergillus niger* has been also reported to be the predominant fungi in dried vine fruits (Heenan et al. 1998; Abarca et al. 2003) which show pH ranges of 3.8 to 4.1 (FDA 2003). *Aspergillus niger* aggregate strains have been also isolated from different stages of coffee beans, before and after harvesting (Taniwaki et al. 2003). The pH of unfermented coffee beans can range from 5.4 to 6.4 (Thompson et al. 2001) and therefore, the time before the arrival of the cherries at the factory may represent a high risk for OTA production (Bucheli and Taniwaki 2002). The remarked conditions could support growth and also the production of OTA on these substrates.

1
2
3 It is worth noting that our results show *A. niger* aggregate strains can grow and
4 produce OTA in a wide pH range. Moreover significant amounts of OTA can be
5 produced in only 5 days. This would lead to confirm their contribution in the OTA
6 contamination of substrates, either in the field or after harvesting, due to the presence
7 of the proper environmental conditions for the production of the mycotoxin.
8
9

10
11
12 Further studies on the effect of environmental factors on growth and OTA production
13 are necessary. A better knowledge of the ecology of these species will provide us
14 with management strategies to reduce the OTA contamination of the substrates
15 where they are usually isolated.
16
17

18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Acknowledgements

This research was supported by the Ministerio de Educación y Ciencia of the Spanish Government (AGL2004-07549-C05-03) and by the grant 2002 SGR 00079 from the DURSI, Generalitat de Catalunya.

References

- Abarca ML, Bragulat MR, Castella G, Cabañes FJ. 1994. Ochratoxin A production by strains of *Aspergillus niger* var. *niger*. *Applied Environmental Microbiology* 60:2650-2652.
- Abarca ML, Accensi F, Bragulat MR, Castella G, Cabañes FJ. 2003. *Aspergillus carbonarius* as the main source of ochratoxin A contamination in dried vine fruits from the Spanish market. *Journal of Food Protection* 66:504-506.

- Abarca ML, Accensi F, Cano J, Cabañes FJ. 2004. Taxonomy and significance of black aspergilli. *Antonie van Leeuwenhoek* 86:33-49.
- Accensi F, Cano J, Figuera L, Abarca ML, Cabañes FJ. 1999. New PCR method to differentiate species in the *Aspergillus niger* aggregate. *FEMS Microbiology Letters* 180:191-196.
- Accensi F, Abarca ML, Cano J, Figuera L, Cabañes FJ. 2001. Distribution of ochratoxin A producing strains in the *A. niger* aggregate. *Antonie van Leeuwenhoek* 79:365-370.
- Battilani P, Pietri A, Bertuzzi T, Languasco L, Giorni P, Kozakiewicz Z. 2003. Occurrence of ochratoxin A-producing fungi in grapes grown in Italy. *Journal of Food Protection* 66:633-636.
- Bau M, Bragulat MR, Abarca ML, Minués S, Cabañes FJ. 2005. Ochratoxigenic species from Spanish wine grapes. *International Journal of Food Microbiology* 98:125-130.
- Bauer J, Gareis M. 1987. Ochratoxin A in der Nahrungsmittelkette. *Journal of Veterinary Medicine B* 34:613-627.
- Bigelis R, Lausure LL. 1987. Fungal enzymes and primary metabolites used in food processing. In: Beuchat LR, editor. *Food and Beverage Mycology*. New York: Van Nostrand Reinhold, pp 472-516.
- Bragulat MR, Abarca ML, Cabañes FJ. 2001. An easy screening method for fungi producing ochratoxin A in pure culture. *International Journal of Food Microbiology* 71:139-144.
- Bucheli P, Taniwaki MH. 2002. Research on the origin, and on the impact of post-harvest handling and manufacturing on the presence of ochratoxin A in coffee. *Food Additives and Contaminants* 19:655-665.

- Cabañes FJ, Accensi F, Bragulat MR, Abarca ML, Castellá G, Mínguez S, Pons A. 2002. What is the source of ochratoxin A in wine? *International Journal of Food Microbiology* 79:213-215.
- Comission of the European Communities (2002) Commission regulation (EC) no 472/2002 of 12 March 2002 amending regulation (EC) no 466/2001 setting maximum levels for certain contaminants in foodstuff. *Official Journal of European Communities* L75:18-20
- Comission of the European Communities (2004) Commission regulation (EC) no 683/2004 of 13 April 2004 amending regulation (EC) no 466/2001 as regards aflatoxins and ochratoxin A in foods for infants and young children. *Official Journal of European Communities* L106:13-15.
- Comission of the European Communities (2005) Commission regulation (EC) no 123/2005 of 26 January 2005 amending regulation (EC) no 466/2001 as ochratoxin A. *Official Journal of the European Union* L25:3-5.
- Esteban A, Abarca ML, Bragulat MR, Cabañes FJ. 2004. Effects of temperature and incubation time on production of ochratoxin A by black aspergilli. *Research in Microbiology* 155:861-866.
- Esteban A, Abarca ML, Bragulat MR, Cabañes FJ. 2005. Influence of pH and incubation time on ochratoxin A production by *Aspergillus carbonarius* in culture media. *Journal of Food Protection* 68:1435-1440.
- Frisvad JC, Thrane U. 2000. Mycotoxin production by common filamentous fungi. In: Samson RA, Hoekstra ES, Frisvad JC, Filtenborg, O, editors. *Introduction to food and airborne fungi*. Utrecht: Centraalbureau voor Schimmelcultures. pp 321-331.

- Heenan CN, Shaw KJ, Pitt JI. 1998. Ochratoxin A production by *Aspergillus carbonarius* and *A. niger* isolates and detection using coconut cream agar. *Journal of Food Mycology* 1:67-72.
- Horie Y. 1995. Productivity of ochratoxin A of *Aspergillus carbonarius* in *Aspergillus* section *Nigri*. *Nippon Kingakukai Kaiho* 36:73-76
- Hunt DC, McConnie BR, Crosby NT. 1980. Confirmation of ochratoxin A by chemical derivatisation and high-performance liquid chromatography. *Analyst* 105:89-90.
- IARC. 1993. Some naturally occurring substances: Food items and constituents, heterocyclic aromatic amines and mycotoxins. IARC monographs on the evaluation of carcinogenic risks of chemicals to humans Lyon, vol. 56.
- Leong SL, Hocking AD, Pitt JI. 2004. Occurrence of fruit rot fungi (*Aspergillus* section *Nigri*) on some drying varieties of irrigated grapes. *Australian Journal of Grape and Wine Research* 10:83-88.
- O'Brien E, Dietrich DR. 2005. Ochratoxin A: the continuing enigma. *Critical Reviews in Toxicology* 35:33-60.
- Ono H, Kataoka M, Koakutsu M, Tanaka K, Kawasugi S, Wakazawa M, Ueno Y, Manabe M. 1995. Ochratoxin A producibility by strains of *Aspergillus niger* group stored in IFO culture collection. *Mycotoxins* 41:47-51.
- Panasenko VT. 1967. Ecology of microfungi. *The Botanical Review* 33:189-215.
- Pitt JI, Hocking AD. 1997. *Fungi and Food Spoilage*. London: Blackie Academic & Professional.
- Serra R, Abrunhosa L, Kozakiewicz Z, Venancio A. 2002. Black *Aspergillus* species as ochratoxin A producers in Portuguese wine grapes. *International Journal of Food Microbiology* 88:63-68.

- 1
2
3 Splittstoesser, DF. 1987. Fruits and fruit products. In: Beuchat, LR, editor. Food and
4
5 beverage mycology, 2nd ed. New York: Van Nostrand Reinhold. pp 101-128.
6
7
8 Taniwaki MH, Pitt JI, Teixeira AA, Iamanaka BT. 2003. The source of ochratoxin A in
9
10 Brazilian coffee and its formation in relation to processing methods. International
11
12 Journal of Food Microbiology 82:173-179.
13
14
15 Téren J, Varga J, Hamari Z, Rinyu E, Kebei F. 1996 Immunochemical detection of
16
17 ochratoxin A in black *Aspergillus* strains. Mycopathologia 134:171-176.
18
19
20 Thompson SS, Miller KB, López AS. 2001. Cocoa and coffee. In: Doyle MP, Beuchat
21
22 LR, Montville TJ, editors. Food Microbiology. Fundamentals and Frontiers, 2nd ed.
23
24 Washington: ASM Press. pp 721-733
25
26
27 U.S. Food and Drug Administration [internet]. Approximate pH of foods and food
28
29 products. Available from: <http://www.cfsan.fda.gov/~comm/lacf-phs.html>. Accessed
30
31 2005 September 20.
32
33
34 Varga J, Rigó K, Téren J. 2000. Degradation of ochratoxin A by *Aspergillus* species.
35
36 International Journal of Food Microbiology 59:1-7.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table I. *Aspergillus niger* aggregate strains used in this study.

Species	Strain (source) ^a	OTA production reported (Ref.)	RFLP pattern (Ref.)
<i>A. niger</i> var. <i>niger</i>	A-75 (feedstuffs, CCFVB)	+ (Abarca et al. 1994; Esteban et al. 2004)	N (Accensi et al. 2001)
<i>A. niger</i> var. <i>niger</i>	A-136 (soy, CCFVB)	+ (Abarca et al. 1994; Esteban et al. 2004)	N (Accensi et al. 2001)
<i>A. niger</i> var. <i>niger</i>	A-942 (raisins, CCFVB)	+ (Abarca et al. 2003; Esteban et al. 2004)	N (Abarca et al. 2003)
<i>A. niger</i> var. <i>niger</i>	A-943 (grapes, CCFVB)	+ (Esteban et al. 2004)	N (Esteban et al. 2004)
<i>A. awamori</i>	CBS 139.52 (kuro-koji)	+ (Ono et al. 1995; Esteban et al. 2004)	N (Accensi et al. 2001)
<i>A. foetidus</i>	CBS 618.78 (unknown)	+ (Téren et al. 1996; Esteban et al. 2004)	N (Accensi et al. 2001)
<i>A. niger</i>	CECT 2088 (unknown)	- (Varga et al. 2000; Esteban et al. 2004)	N (Esteban et al. 2004)
<i>A. niger</i>	CBS 554.65 (tannin-gallic acid fermentation)	- (Accensi et al. 2001; Esteban et al. 2004)	N (Accensi et al. 2001)
<i>A. niger</i> var. <i>niger</i>	A-946 (coffee, CCFVB)	- (Esteban et al. 2004)	N (Esteban et al. 2004)
<i>A. niger</i>	CBS 121.55 (otomycosis)	- (Esteban et al. 2004)	N (Esteban et al. 2004)
<i>A. tubingensis</i>	CBS 134.48 (unknown)	- (Accensi et al. 2001; Esteban et al. 2004)	T (Accensi et al. 2001)
<i>A. niger</i> var. <i>niger</i>	A-947 (grapes, CCFVB)	- (Cabañes et al. 2002; Esteban et al. 2004)	T (Cabañes et al. 2002)

^a Abbreviations: CCFVB, Culture Collection of Veterinary Faculty of Barcelona, Bellaterra (Barcelona), Spain; CBS, Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CECT, Colección Española de Cultivos Tipo, Valencia, Spain.

Table II. OTA concentration (mean and mean standard error) produced by the six OTA-producing strains of *A. niger* aggregate in CYA medium at 25°C at each pH value tested.

	days	OTA concentration($\mu\text{g/g}$)								
		2	3	4	5	6	7	8	9	10
A-75	5	nd ^a	1.78 $\pm$ 1.02 ^a	1.92 $\pm$ 1.12 ^a	7.82 $\pm$ 0.55 ^b	1.47 $\pm$ 0.27 ^a	0.74 $\pm$ 0.07 ^a	1.33 $\pm$ 0.77 ^a	0.93 $\pm$ 0.42 ^a	0.64 $\pm$ 0.16 ^a
	10	1.70 $\pm$ 0.49 ^a	1.57 $\pm$ 0.66 ^a	1.59 $\pm$ 0.49 ^a	1.19 $\pm$ 0.03 ^a	0.82 $\pm$ 0.13 ^a	0.69 $\pm$ 0.08 ^a	0.79 $\pm$ 0.23 ^a	0.39 $\pm$ 0.03 ^a	0.41 $\pm$ 0.06 ^a
	15	0.30 $\pm$ 0.06 ^a	3.09 $\pm$ 0.89 ^b	1.38 $\pm$ 0.08 ^a	1.10 $\pm$ 0.31 ^a	0.89 $\pm$ 0.01 ^a	0.81 $\pm$ 0.01 ^a	0.89 $\pm$ 0.06 ^a	0.35 $\pm$ 0.05 ^a	0.21 $\pm$ 0 ^a
	20	0.57 $\pm$ 0 ^a	1.09 $\pm$ 1.01 ^a	1.44 $\pm$ 0.07 ^a	0.41 $\pm$ 0.06 ^a	0.92 $\pm$ 0.02 ^a	0.62 $\pm$ 0.27 ^a	1.00 $\pm$ 0.30 ^a	0.32 $\pm$ 0.04 ^a	0.31 $\pm$ 0.01 ^a
	30	0.18 $\pm$ 0.05 ^a	1.22 $\pm$ 1.20 ^a	1.55 $\pm$ 0.04 ^a	0.95 $\pm$ 0.60 ^a	1.21 $\pm$ 0.91 ^a	0.61 $\pm$ 0.13 ^a	0.87 $\pm$ 0.13 ^a	0.45 $\pm$ 0.08 ^a	0.21 $\pm$ 0.06 ^a
A-136	5	0.03 $\pm$ 0.04 ^a	4.00 $\pm$ 1.13 ^b	2.79 $\pm$ 0.35 ^{ab}	9.31 $\pm$ 2.12 ^c	2.44 $\pm$ 0.47 ^{ab}	1.92 $\pm$ 0.82 ^{ab}	1.54 $\pm$ 0.20 ^{ab}	1.50 $\pm$ 0.13 ^{ab}	1.67 $\pm$ 0.12 ^{ab}
	10	3.68 $\pm$ 0.71 ^a	2.93 $\pm$ 0.20 ^a	1.98 $\pm$ 0.03 ^b	2.01 $\pm$ 0.57 ^b	0.77 $\pm$ 0.23 ^c	1.17 $\pm$ 0.03 ^{bc}	0.90 $\pm$ 0.21 ^c	0.43 $\pm$ 0.19 ^c	0.58 $\pm$ 0.06 ^c
	15	1.14 $\pm$ 0.28 ^a	4.16 $\pm$ 0.54 ^b	2.00 $\pm$ 0.05 ^c	1.83 $\pm$ 0.28 ^c	1.06 $\pm$ 0.12 ^a	0.67 $\pm$ 0 ^a	0.69 $\pm$ 0.27 ^a	0.54 $\pm$ 0.26 ^a	0.51 $\pm$ 0.11 ^a
	20	1.50 $\pm$ 1.08 ^{ab}	2.86 $\pm$ 0.19 ^c	1.98 $\pm$ 0.09 ^{bc}	0.71 $\pm$ 0.06 ^{ab}	1.09 $\pm$ 0.01 ^{ab}	1.02 $\pm$ 0.01 ^{ab}	0.82 $\pm$ 0.25 ^{ab}	0.50 $\pm$ 0.27 ^a	0.46 $\pm$ 0.17 ^a
	30	1.24 $\pm$ 0.71 ^{ab}	2.80 $\pm$ 0.05 ^c	1.80 $\pm$ 0.20 ^b	1.67 $\pm$ 0.31 ^{ab}	1.41 $\pm$ 0.78 ^{ab}	0.81 $\pm$ 0.18 ^{ab}	0.59 $\pm$ 0.13 ^{ab}	0.33 $\pm$ 0.09 ^a	0.30 $\pm$ 0.06 ^a
A-942	5	0.42 $\pm$ 0.18 ^a	7.24 $\pm$ 1.66 ^b	4.23 $\pm$ 0.43 ^c	3.86 $\pm$ 2.28 ^{ac}	1.21 $\pm$ 0.04 ^{ac}	0.88 $\pm$ 0.09 ^{ac}	1.32 $\pm$ 0.34 ^{ac}	0.99 $\pm$ 0.31 ^{ac}	0.44 $\pm$ 0.14 ^a
	10	3.51 $\pm$ 1.84 ^{ab}	4.90 $\pm$ 0.66 ^b	2.91 $\pm$ 0.64 ^{ac}	1.04 $\pm$ 0.02 ^c	0.86 $\pm$ 0.05 ^c	1.00 $\pm$ 0.17 ^c	1.10 $\pm$ 0.37 ^c	0.84 $\pm$ 0.11 ^c	0.52 $\pm$ 0.10 ^c
	15	0.22 $\pm$ 0.01 ^a	8.53 $\pm$ 0.29 ^b	1.92 $\pm$ 0.21 ^c	0.89 $\pm$ 0.06 ^{de}	1.06 $\pm$ 0.13 ^e	0.92 $\pm$ 0.36 ^{de}	1.07 $\pm$ 0.07 ^e	0.91 $\pm$ 0.15 ^{de}	0.37 $\pm$ 0.04 ^{ad}
	20	0.16 $\pm$ 0.01 ^a	7.00 $\pm$ 0.38 ^b	2.62 $\pm$ 0.61 ^c	1.28 $\pm$ 0.21 ^d	1.07 $\pm$ 0.01 ^{de}	0.87 $\pm$ 0.07 ^{ade}	1.22 $\pm$ 0.20 ^d	0.65 $\pm$ 0.04 ^{ade}	0.31 $\pm$ 0.06 ^{ac}
	30	0.36 $\pm$ 0.01 ^{ab}	4.59 $\pm$ 0.31 ^c	1.37 $\pm$ 0.10 ^d	0.49 $\pm$ 0.06 ^{ab}	0.76 $\pm$ 0.21 ^b	0.81 $\pm$ 0.11 ^b	1.28 $\pm$ 0.05 ^d	0.52 $\pm$ 0.16 ^{ab}	0.25 $\pm$ 0.04 ^a
A-943	5	nd ^a	0.14 $\pm$ 0.01 ^b	nd ^a	nd ^a	0.05 $\pm$ 0 ^c	nd ^a	nd ^a	nd ^a	nd ^a
	10	0.10 $\pm$ 0.01 ^a	0.26 $\pm$ 0.06 ^b	nd ^c	nd ^c	0.03 $\pm$ 0.04 ^c	nd ^c	nd ^c	nd ^c	nd ^c
	15	0.19 $\pm$ 0.04 ^a	0.22 $\pm$ 0 ^a	nd ^b	nd ^b	nd ^b	nd ^b	nd ^b	0.09 $\pm$ 0.05 ^c	nd ^b
	20	nd ^a	0.07 $\pm$ 0.09 ^a	nd ^a	nd ^a	0.03 $\pm$ 0.04 ^a	nd ^a	0.06 $\pm$ 0.01 ^a	0.08 $\pm$ 0.04 ^a	nd ^a
	30	nd ^a	0.08 $\pm$ 0.11 ^a	nd ^a	nd ^a	0.05 $\pm$ 0.06 ^a	nd ^a	nd ^a	nd ^a	nd ^a
CBS 139.52	5	4.87 $\pm$ 1.99 ^{ab}	10.50 $\pm$ 0.21 ^c	1.21 $\pm$ 1.03 ^a	3.85 $\pm$ 0.62 ^{ab}	2.90 $\pm$ 1.02 ^{ab}	2.06 $\pm$ 0.41 ^{ab}	2.68 $\pm$ 1.47 ^{ab}	5.69 $\pm$ 1.4 ^b	3.81 $\pm$ 0.42 ^{ab}
	10	14.58 $\pm$ 2.02 ^a	7.77 $\pm$ 2.90 ^b	0.65 $\pm$ 0.91 ^c	2.35 $\pm$ 0.90 ^c	1.87 $\pm$ 0.48 ^c	1.87 $\pm$ 0.33 ^c	2.02 $\pm$ 0.52 ^c	3.24 $\pm$ 1.34 ^c	3.06 $\pm$ 0.57 ^c
	15	2.07 $\pm$ 0.54 ^{ab}	8.63 $\pm$ 1.22 ^c	0.42 $\pm$ 0.28 ^a	1.55 $\pm$ 0.22 ^{ab}	2.30 $\pm$ 0.86 ^{ab}	1.95 $\pm$ 0.25 ^{ab}	3.14 $\pm$ 0.53 ^b	2.79 $\pm$ 0.78 ^b	1.95 $\pm$ 0.12 ^{ab}
	20	1.80 $\pm$ 1.37 ^a	15.35 $\pm$ 4.59 ^b	0.63 $\pm$ 0.80 ^a	1.41 $\pm$ 0.43 ^a	2.04 $\pm$ 0.37 ^a	2.02 $\pm$ 0.42 ^a	2.57 $\pm$ 1.22 ^a	3.64 $\pm$ 1.71 ^a	3.78 $\pm$ 1.58 ^a
	30	2.14 $\pm$ 0.59 ^a	7.69 $\pm$ 2.23 ^b	0.39 $\pm$ 0.55 ^a	3.01 $\pm$ 0.37 ^a	1.76 $\pm$ 0.02 ^a	1.56 $\pm$ 0.18 ^a	2.56 $\pm$ 0.50 ^a	4.16 $\pm$ 2.79 ^a	2.82 $\pm$ 0.15 ^a
CBS 618.78	5	nd ^a	nd ^a	nd ^a	0.12 $\pm$ 0.01 ^{ab}	0.17 $\pm$ 0.01 ^b	0.28 $\pm$ 0.12 ^c	0.08 $\pm$ 0.01 ^{ab}	nd ^a	nd ^a
	10	nd ^a	nd ^a	nd ^a	1.04 $\pm$ 0.50 ^a	0.25 $\pm$ 0.09 ^a	0.52 $\pm$ 0.39 ^a	1.84 $\pm$ 2.47 ^a	nd ^a	nd ^a
	15	nd ^a	nd ^a	nd ^a	0.68 $\pm$ 0.15 ^b	0.73 $\pm$ 0.14 ^b	0.78 $\pm$ 0.34 ^b	0.46 $\pm$ 0.35 ^{ab}	nd ^a	nd ^a
	20	nd ^a	nd ^a	nd ^a	0.07 $\pm$ 0.02 ^a	0.49 $\pm$ 0.11 ^b	0.55 $\pm$ 0.03 ^b	0.25 $\pm$ 0.20 ^a	nd ^a	nd ^a
	30	nd ^a	nd ^a	0.05 $\pm$ 0.06 ^a	0.96 $\pm$ 0.56 ^b	0.27 $\pm$ 0.06 ^a	0.31 $\pm$ 0.09 ^a	0.04 $\pm$ 0.06 ^a	nd ^a	nd ^a

^{a,b,c,d,e} Values with the same superscript within each strain and incubation time are not significantly different ($p < 0.05$); nd: not detected (limit of detection 0.01 $\mu\text{g g}^{-1}$).

Table III. OTA concentration (mean and mean standard error) produced by the six OTA-producing strains of *A. carbonarius* in YES medium at 25°C at each pH value tested.

		OTA concentration(µg/g)								
	days	2	3	4	5	6	7	8	9	10
A-75	5	nd ^a	0.67±0.11 ^a	7.27±0.24 ^{ab}	16.82±0.39 ^b	18.22±3.01 ^b	14.82±7.45 ^b	12.89±2.45 ^b	15.46±3.79 ^b	8.46±0.52 ^{ab}
	10	nd ^a	0.69±0.01 ^a	4.43±0.04 ^{abc}	4.80±1.84 ^{abc}	9.72±1.34 ^c	9.01±1.18 ^c	15.02±4.14 ^d	7.24±0.02 ^{bc}	2.54±0.35 ^{ab}
	15	nd ^a	0.27±0.12 ^a	2.20±0.87 ^{ab}	4.19±1.03 ^{bc}	4.75±0.45 ^{bc}	8.15±2.36 ^d	6.49±0.55 ^{cd}	3.72±1.25 ^{bc}	0.50±0.11 ^a
	20	nd ^a	0.52±0.15 ^a	1.81±0.37 ^{ab}	0.09±0.12 ^a	3.53±0.34 ^b	5.91±0.46 ^c	3.53±1.33 ^b	2.93±0.86 ^b	0.91±0.23 ^a
	30	nd ^a	0.15±0.02 ^a	0.96±0.13 ^{ab}	1.57±0.54 ^{bc}	2.24±0.62 ^c	2.65±0.26 ^c	3.56±0.78 ^d	1.60±0.21 ^{bc}	0.44±0.15 ^{ab}
A-136	5	nd ^a	5.91±0.71 ^a	26.78±4.51 ^{bc}	37.93±1.22 ^d	34.69±1.18 ^{cd}	34.66±2.08 ^{cd}	20.27±0.90 ^{be}	19.90±1.06 ^{bc}	15.94±6.93 ^e
	10	nd ^a	0.91±0.43 ^{ab}	6.19±0.43 ^{ab}	11.55±3.92 ^{ab}	6.39±0.39 ^{ab}	6.55±0.23 ^{ab}	7.08±0.76 ^{ab}	13.98±9.57 ^b	2.62±0.21 ^{ab}
	15	nd ^a	1.04±0.20 ^a	2.34±0.40 ^a	1.75±1.05 ^a	5.33±1.10 ^a	3.55±2.27 ^a	6.37±5.10 ^a	6.15±0.26 ^a	0.29±0.01 ^a
	20	nd ^a	0.39±0.13 ^a	1.56±0.32 ^{ab}	0.18±0.10 ^a	1.54±0.49 ^{ab}	3.84±1.77 ^b	1.83±0.91 ^{ab}	3.43±0.16 ^b	0.69±0.37 ^a
	30	nd ^a	0.22±0.03 ^{ab}	0.73±0 ^{ab}	0.69±0.27 ^{ab}	2.44±0.71 ^c	1.52±0.35 ^{bc}	1.02±0.28 ^{ab}	2.45±0.69 ^c	0.28±0.06 ^{ab}
A-942	5	nd ^a	2.06±0.47 ^{ab}	10.89±1.87 ^{bcd}	5.38±1.09 ^{abc}	12.90±6.56 ^{cd}	11.66±2.36 ^{bcd}	5.49±0.90 ^{abc}	18.23±1.44 ^d	11.38±3.59 ^{bcd}
	10	3.25±3.00 ^a	0.15±0.03 ^a	2.29±1.39 ^a	1.32±0.38 ^a	2.34±0.08 ^a	2.85±0.73 ^a	1.26±0.07 ^a	4.97±1.54 ^a	5.05±2.62 ^a
	15	1.18±0.44 ^{abc}	0.33±0.18 ^a	0.89±0.38 ^{ab}	0.58±0.02 ^a	1.91±0.42 ^c	1.54±0.13 ^{bc}	2.86±0.05 ^d	0.39±0.02 ^a	1.70±0.19 ^{bc}
	20	1.48±1.57 ^a	nd ^a	0.61±0.04 ^a	0.35±0.18 ^a	1.29±0.02 ^a	1.19±0.50 ^a	1.00±0.18 ^a	1.13±0.39 ^a	0.87±0.19 ^a
	30	0.39±0.26 ^a	0.14±0.01 ^a	0.33±0.02 ^a	0.36±0.11 ^a	1.25±0.54 ^a	0.76±0.08 ^a	1.01±0.04 ^a	0.84±0.20 ^a	1.27±0.64 ^a
A-943	5	nd ^a	nd ^a	0.23±0.09 ^a	0.12±0.01 ^a	0.16±0.05 ^a	0.16±0.01 ^a	nd ^a	0.57±0 ^a	1.39±0.48 ^b
	10	0.11±0.15 ^a	0.18±0.26 ^a	nd ^a	nd ^a	0.15±0.02 ^a	nd ^a	nd ^a	0.20±0.06 ^a	0.13±0 ^a
	15	0.12±0.02 ^a	0.17±0.24 ^a	nd ^a	nd ^a	nd ^a	nd ^a	nd ^a	0.07±0.03 ^a	nd ^a
	20	nd ^a	0.12±0.16 ^a	nd ^a	nd ^a	nd ^a	nd ^a	0.08±0.01 ^a	0.09±0 ^a	nd ^a
	30	nd ^a	0.09±0.12 ^a	nd ^a	nd ^a	nd ^a	nd ^a	nd ^a	nd ^a	nd ^a
CBS 139.52	5	0.03±0.04 ^a	16.82±3.14 ^{bc}	17.37±7.70 ^{bc}	58.00±1.34 ^d	10.25±7.95 ^{abc}	14.74±0.28 ^{bc}	19.65±1.80 ^c	19.03±7.65 ^c	1.88±0.23 ^{ab}
	10	0.12±0.16 ^a	3.58±0.75 ^a	1.11±0.79 ^a	17.12±5.13 ^a	9.54±10.88 ^a	11.97±9.62 ^a	16.96±4.77 ^a	42.27±4.59 ^b	18.74±3.28 ^a
	15	nd ^a	3.43±3.29 ^a	9.60±5.33 ^a	16.38±4.24 ^a	13.69±2.76 ^a	9.05±4.79 ^a	8.92±5.58 ^a	7.87±9.38 ^a	5.73±0.86 ^a
	20	nd ^a	4.61±0.70 ^a	4.14±2.49 ^a	4.88±2.61 ^a	10.01±0.40 ^a	12.56±1.92 ^a	12.43±4.50 ^a	17.98±17.78 ^a	5.88±1.21 ^a
	30	nd ^a	2.70±0.08 ^a	3.29±0.02 ^a	11.66±1.34 ^a	6.46±1.87 ^a	7.66±0.70 ^a	6.03±4.05 ^a	11.59±12.43 ^a	2.32±0.11 ^a
CBS 618.78	5	nd ^a	nd ^a	nd ^a	24.56±15.90 ^b	5.49±7.50 ^a	0.25±0.35 ^a	4.12±2.01 ^a	nd ^a	nd ^a
	10	nd ^a	nd ^a	nd ^a	1.17±0.55 ^a	0.95±0.59 ^a	0.21±0.29 ^a	30.56±18.72 ^b	nd ^a	nd ^a
	15	nd ^a	nd ^a	nd ^a	nd ^a	1.63±2.22 ^a	0.40±0.02 ^a	0.95±0.08 ^a	nd ^a	nd ^a
	20	nd ^a	nd ^a	nd ^a	nd ^a	0.12±0 ^a	0.99±0.059 ^b	1.21±0.04 ^b	nd ^a	nd ^a
	30	nd ^a	nd ^a	nd ^a	nd ^a	0.51±0.06 ^b	0.63±0.03 ^c	0.10±0.07 ^a	nd ^a	nd ^a

^{a,b,c,d,e} Values with the same superscript within each strain and incubation time are not significantly different (p < 0.05); nd: not detected (limit of detection 0.01 µg g⁻¹).

For Peer Review Only

Fig.1. Mean OTA concentration produced by the OTA-producing strains at each pH value and culture media tested.