Comparison of ochratoxin A and deoxynivalenol loads of organically and conventionally produced beers sold on the Belgian market

Emmanuel Kossi Tangni, Luc Pussemier, Jean Claude Motte, François van Hove, Yves Jacques Schneider, Carlos van Peteghem, Yvan Larondelle

► To cite this version:

Emmanuel Kossi Tangni, Luc Pussemier, Jean Claude Motte, François van Hove, Yves Jacques Schneider, et al.. Comparison of ochratoxin A and deoxynivalenol loads of organically and conventionally produced beers sold on the Belgian market. Food Additives and Contaminants, 2006, 23 (09), pp.910-918. 10.1080/02652030600743839 . hal-00577299

HAL Id: hal-00577299

<https://hal.science/hal-00577299>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparison of ochratoxin A and deoxynivalenol loads of organically and conventionally produced beers sold on the Belgian market

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2006-022.R2
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	05-Apr-2006
Complete List of Authors:	TANGNI, Emmanuel; Veterinary and Agrochemical Research Center, Quality and Safety Pussemier, Luc; Veterinary and Agrochemical Research Center, Quality and Safety Motte, Jean Claude; Veterinary and Agrochemical Research Center, Quality and Safety Van Hove, François; Université catholique de Louvain, Mycothèque de l'Université catholique de Louvain (BCCMTM/MUCL) Schneider, Yves Jacques; Université catholique de Louvain, Laboratoire de biochimie cellulaire, Institut des Sciences de la Vie Van Peteghem, Carlos; Ghent University, Laboratory of food analysis, Faculty of Pharmaceutical Sciences Larondelle, Yvan; Université catholique de Louvain, Unité de biochimie de la nutrition
Methods/Techniques:	Chromatography - HPLC, Clean-up - affinity columns, Exposure assessment, In-house validation
Additives/Contaminants:	Mycotoxins - trichothecenes, Mycotoxins - ochratoxin A
Food Types:	Beer, Organic foods, Processed foods

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review Only

Comparison of ochratoxin A and deoxynivalenol in organically and conventionally produced beers sold on the Belgian market

ANSELME M.¹, TANGNI E.K.^{2,*}, PUSSEMIER L.², MOTTE J-C², VAN HOVE F³, SCHNEIDER Y-J⁴, VAN PETEGHEM C⁵ & LARONDELLE Y¹

¹: Unité de biochimie de la nutrition, Institut des Sciences de la Vie, Université catholique de Louvain, Croix du Sud 2/8, 1348 Louvain-la-Neuve, Belgium

²: Department of quality and safety, Veterinary and Agrochemical Research Centre (CODA-CERVA), Leuvensesteenweg, 17, B-3080 Tervuren, Belgium

³: Mycothèque de l'Université catholique de Louvain (BCCMTM/MUCL), Unité de Microbiologie, Croix du Sud 3/6, 1348 Louvain-La-Neuve, Belgium.

⁴: Laboratoire de biochimie cellulaire, Institut des Sciences de la Vie, Université catholique de Louvain, Croix du Sud 5/3, 1348 Louvain-la-Neuve, Belgium.

⁵: Laboratory of food analysis, Faculty of Pharmaceutical Sciences, Ghent University, Harelbekestraat 72, 9000 Ghent

Running Head : Ochratoxin A and deoxynivalenol in organic and conventional beers

*: Correspondence: E. K. Tangni, E-mail: emtan@var.fgov.be

Abstract

Beer was chosen as a cereal-derived and homogeneous product which would allow a comparison to be made of organic and conventional production modes in terms of mycotoxin contamination levels. Ochratoxin A (OTA, a storage mycotoxin) and deoxynivalenol (DON, a field mycotoxin) were assessed by HPLC methods in organically and conventionally produced beers sold in Belgium. Immunoaffinity column (OchraTest® and DONPrep®) purification was used prior to HPLC analysis. For in-house validation, recovery experiments carried out with the spiked beers in the ranges of 50 - 200 ng OTA l⁻¹ and 20 - 100 µg DON l⁻¹ led to the overall averages of 91 % (RSD = 10 %, n =9) and 93 % (RSD = 5 %, n = 27), respectively. Organic beers collected during 2003-2004 tended to be more frequently OTA contaminated (95 %, n = 40) than their conventional counterparts (50 %, n = 40). Conventional beers were OTA contaminated at a mean concentration of 25 ng l⁻¹ (range: 19 - 198 ng l⁻¹) whilst organic beers contained a mean level of 182 ng l⁻¹(range: 18 - 1134 ng l⁻¹). High OTA contamination above the limit of 200 ng l⁻¹ (up to 1134 ng l⁻¹) occasionally occurred in organically produced beers. A complementary survey performed with the same brands in 2005 did not confirm this accidental presence of excessive OTA loads (range: 3 - 67 ng l⁻¹ for 10 conventional beers and 19 - 158 ng l⁻¹ for 10 organic beers). Establishing a maximum of 3 µg OTA kg⁻¹ in malt, the application of the regulation EC 466/2001 (entered in force before the last sampling) may be related to the improvement observed. The overall incidence of DON was 67 and 80 % in conventional and organic beers, respectively. DON concentrations ranged from 2 to 22 µg DON l⁻¹ (mean = 6 µg DON l⁻¹) in conventional beers whilst organic beers were contaminated from 2 to 14 µg DON l⁻¹ (mean = 4 µg DON l⁻¹). DON in beers does thus not appear to be a major matter of concern. From the statistical tests, it could be concluded that the variation between different batches was significant (p < 0.0001) in contrast to that observed between different brands, showing thus a lack of homogeneity in the raw materials. This may occur either in organically or in conventionally produced materials. Considering these results, an optimized frequency of controls according to European Regulations EC No 466/2001 and EC No 856/2005 should be recommended in order to reject the irregular batches.

Keywords : Food safety, mycotoxin, ochratoxin A, deoxynivalenol, beer, organic, conventional, Belgium.

Introduction

Due to food crises in the recent past, organic food products have become more popular in Belgium and elsewhere (Pussemier *et al.* 2006). Indeed, the demand for organic foods is constantly increasing mainly due to the consumers' perception that they are healthier and safer than conventional foods (Magkos *et al.* 2003). The situation may however be more complex. Some authors indicate that organic food often contains relatively high amounts of natural toxic compounds produced by fungi or plants, whereas corresponding conventional food tends to contain more synthetic compounds such as pesticide residues (Finamore *et al.* 2004). By contrast, Pussemier *et al.* (2006) have concluded that the influence of organic cultivation methods on the presence of natural toxins is not yet clear. In terms of mycotoxins, even though the foodstuffs resulting from conventional production are not free of mycotoxins, the organic foodstuffs may present a greater risk of contamination. So, there is a need for information related to food safety to inform the consumers of the health benefits and/or hazards of food products of both origins, in order to optimize the impact on health and minimize the risks.

The mycotoxin ochratoxin A (OTA) has strong nephrotoxic, immunosuppressive and teratogenic properties. Its worldwide occurrence as a contaminant of raw agricultural products has been amply documented in a variety of plant products such as cereals, coffee, dried fruits, cocoa, grapes and spices (MacDonald *et al.* 1999, 2004, Ottener & Majerus 2001, Serra *et al.* 2005, Thirumala-Devi *et al.* 2001). Mainly produced by toxigenic strains of the genera *Aspergillus* and *Penicillium* (Pitt 1987, Moss 1998, Samson *et al.* 2004), high OTA levels mainly occur in stored products. Before storage, cereals are prone to be contaminated in the field by fungi of different *Fusarium* species. These fungi may infect the plant and produce trichothecenes, with deoxynivalenol (DON) being the most abundant representative one in cereal grains (Boeira *et al.* 2000). Thus, malting barley, wheat and corn, which are used in brewing either as main ingredients or as adjuncts, can be contaminated by OTA and/or DON. So far, a handful studies have investigated the fate of these mycotoxins

during malting, brewing and fermentation and they have concluded that OTA and DON can at least partially withstand these complex food processing and, finally, may remain present in beer (Niessen & Donhauser 1993, Baxter 1996, Baxter *et al.* 2001, Papadopoulou-Bouraoui *et al.* 2004). Beer is thus a convenient cereal-derived foodstuff allowing to compare the organic and conventional production modes in terms of mycotoxin contamination levels. To our knowledge, the influence of the organic route of beer production on the presence of natural toxins, such as OTA and DON, has not been yet studied. Therefore, the present investigation was initiated to compare the OTA and DON contaminations in organic and conventional beers produced in Belgium, the European country with the greatest diversity of beers. The study includes the development of a method for the determination of DON in beer using immunoaffinity clean-up coupled with HPLC-UV detection.

Materials and methods

Sample collection and handling

Keeping, as far as possible, a balance between organic and conventional products, forty conventionally produced beers (representing 10 brands) and forty organically produced beers (representing 10 brands) were randomly purchased in the supermarkets and retail shops in Belgium from September 2003 to September 2004. All these brands were manufactured in Belgium. To ensure some variation in batches and brands of the sampled materials, two batches per brand and two samples per batch were tested. A complementary sampling was performed on the same brands (10 conventional and 10 organic beers) in March 2005 to check the temporal fluctuation in OTA contamination, especially for previously high OTA contaminated brands. Here, one bottle (one batch) of each brand was investigated. All samples were stored at 4°C till opening for analysis.

Reagents and materials

Pure crystalline forms of OTA and DON were obtained from Sigma Chemical Co. (St Louis, MO, USA). The stock solution of OTA was made in toluene:acetic acid (99:1) whereas that of DON was

made in acetonitrile far-UV (of HPLC grade). The concentrations were determined by spectrophotometry (Wood *et al.* 1996, Petterson 1998). The stock solutions were stored at -18°C until re-use (maximum storage duration of 12 months).

OTA and DON immunoaffinity columns (OchraTest[®] and DONPrep[®]) were obtained from Vicam (Watertown, MA, USA) and from R-Biopharm (Rhône diagnostics Ltd, Glasgow, Scotland), respectively. Microfilters type Millex[®]-HV, SLHV R04NL (0.45µm) and Durapore[®] membrane filters (type 0.22 µm GV, GVWP04700) were purchased from Millipore (Bedford, MA, USA).

Acetonitrile and toluene of HPLC grade were supplied by Lab Scan (Dublin, Ireland). Methanol of HPLC grade and acetic acid *pro analysis* were obtained from Acros Organics (New Jersey, NY, USA) or from Biosolve (Valkenswaard, The Netherlands) while hydrochloric acid (37 %) was purchased from Riedel-de Haën (Seelze, Germany). Phosphate-buffered saline (PBS) solution (NaCl 120 mM, KCl 2.7 mM, phosphate buffer 10 mM, pH 7.4) was supplied by Sigma. Sodium bicarbonate (NaHCO₃) was supplied by Merck (Darmstadt, Germany). All other chemicals were of analytical grade.

HPLC grade water was obtained using a twin system composed by Elix3 and Milli-Q system, both from Millipore (Molsheim, France).

Helium of 99.996 % purity and nitrogen of 99.995 % purity were provided by Air Liquide (Liège, Belgium).

OchraTest method for OTA determination in beer

OTA determination in beer was performed according to the method described by Tangni *et al.* (2002). First, bottles of beer were vigorously shaken and left for 10 min before opening. A sample of beer was taken and decarbonated by shaking again for 10 min. Degassed beer (40 ml) were then diluted with water (5 ml, containing 4 % (w/v) sodium bicarbonate) and with PBS solution (15 ml). Solid particles were separated by centrifugation of the mixture at 820xg during 10 min at 8 ± 2°C.

The supernatant (45 ml) were used for the next steps. Clean-up of the OTA extract was performed with an OchraTest column fitted with a 20-ml syringe reservoir and placed on a vacuum manifold. Before loading the extract, the OchraTest column was conditioned with 20 ml of PBS solution (pH 7.4). The OTA extract was passed through the column at about 1-2 ml min⁻¹. De-ionized water (20 ml) was used to wash the loaded immunoaffinity column and OTA was eluted with methanol (2 ml) and de-ionized water (2 ml). Atmospheric air (*ca* 20 ml) was passed through the column to collect all the eluate. The solution was stored at -20°C prior to HPLC analysis.

As for HPLC detection and quantification, the chromatographic system consisted of a Perkin-Elmer LC049 isocratic pump (Norwalk, CO, USA) equipped with a Rheodyne model 7125 NS injection valve (50 ml) (Rheodyne, Cotati, CA, USA), a RF-551 fluorescence spectrophotometric detector (Shimadzu, Kyoto, Japan) equipped with a 150W xenon lamp ($\lambda_{\text{excitation}} = 332 \text{ nm}$, $\lambda_{\text{emission}} = 462 \text{ nm}$) and a Spectra- Physics SP4290 integrator (San Jose, CA, USA). The analytical column was a HypersilTM BDS reversed phase C18 (15 cm×4.0 mm i.d., 3 mm particles) (Tracer Analytical, Barcelona, Spain). The column was used at room temperature. The mobile phase was made of a mixture of HPLC grade acetonitrile:water:acetic acid (45:54:1), filtered through a 0.22 µm filter membrane, degassed and eluted at a flow-rate of 1.0 ml min⁻¹. The eluate was filtered through a 0.45 µm microfilter and 50 µl were injected for HPLC analysis.

The identity of OTA was confirmed by methyl ester formation according to the method described by Zimmerli & Dick (1995). Contaminated eluates with OTA concentration > 200 ng l⁻¹ beer were used for this preparation.

DONPrep method for DON determination in beer

Degassed beer (20 ml) were taken and mixed with acetonitrile (110 ml). This mixture was evaporated using a rotary evaporator at 45 ± 5°C in order to minimize the volume of water on the DONprep column. As DON is soluble in water, a bigger volume of water affect the DON recovery.

Water (4 ml) containing sodium bicarbonate (5 % w/v) was added to recover the extract, which was then submitted to the following purification step. Clean-up of DON extract was performed with a DONPrep column fitted with a 5-ml syringe reservoir. Column conditioning was not necessary and the solubilized extract was directly allowed to pass through the column by gravity. The column was washed with water (5 ml) and was then dried with atmospheric air. DON was eluted with methanol (2 ml) by back flush. The eluted extract was evaporated to dryness at $35 \pm 5^{\circ}\text{C}$ with a rotary evaporator, reconstituted in 1 ml of methanol:water (10:90) and stored at -20°C prior to HPLC analysis.

As for HPLC detection and quantification, the chromatographic system consisted of two Waters 515 pumps (Milford, MA, USA), an automatic 717 injector and a 2996 DAD UV detector. The chromatograms were processed using Millenium Empower. The Alltima (100 x 4.6 mm id, 3 μm particles) RP-C₁₈ column (Alltech, Deerfield, IL, USA) preceded by an OptiGuard precolumn (10 x 1 mm i.d., 10 μm particles) obtained from Optimize Technology (Oregon City, OR, USA), was thermostatically maintained at 35°C by a temperature controller Model 7990 (Jones Chromatography, Hengoed, UK) during HPLC analysis.

The mobile phase was made of a mixture of HPLC grade acetonitrile:water (5:95), filtered through a 0.22 μm Durapore[®] membrane filter, degassed and eluted at a flow-rate of 1.0 ml min^{-1} . The eluate was filtered through a 0.45 μm microfilter and 50 μl were injected for HPLC analysis.

Validation

To evaluate the analytical procedures, recovery experiments were performed in triplicate with one type of beer spiked at 50, 150 and 200 ng OTA l^{-1} , while DON was added in triplicate samples of three types of beer at 20, 50 and 100 $\mu\text{g l}^{-1}$. The limit of detection (LOD) and limit of quantification (LOQ) of the chromatographic procedures were calculated by the signal-to-noise ratios of 3:1 and 10:1, respectively. The linearity of the detector responses was estimated in experiments with the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

OTA and DON standards. The calibration curves were plotted and their linearity was assessed based on the correlation coefficient (r).

Data handling and statistical analyses

The calibration curves were calculated using the least-squares method. Experimental results are reported as mean $\pm$ standard deviation (SD). Non-detects were assumed to indicate that the sample contained half the LOD in terms of OTA and DON levels, whereas trace levels were assumed to indicate that the sample contained half the LOD + LOQ (Kroes *et al.* 2002). For both mycotoxins, the median and range values are given.

The statistical analysis attempted to first determine the presence of significant differences in contamination frequency between conventional and organic samples at levels over the limit of quantification (Chi-square test). Variance analyses were then performed on quantifiable concentrations in order to underline the effect of the mode of production, of the brand and of the batch on the level of contamination. Correlation analysis was used to study the consistency of the association between OTA and DON levels in the tested samples. Independent sample T-test was used to compare the mean OTA in DON positive and negative samples across the two groups of beers. Statement of significance was based on probability of $p < 0.05$.

Results and discussion

Analytical methods performances

Chromatograms did not show any impurity interfering with OTA or DON. Regarding OTA analysis, the analytical procedures showed a recovery rate of 91 % (RSD = 10 %, n = 9) and the respective LOD and LOQ values of 5 and 18 ng OTA l⁻¹ of beer, within the tested range of spiked beer (50-200 ng OTA l⁻¹).

As for DON analysis, preliminary results (not published) have shown that the DONPrep column has a maximal capacity of about 1.5 μ g DON. With the above described procedure, this amount

corresponds to a concentration of $75 \mu\text{g DON l}^{-1}$ beer. In order to be able to analyze concentrations above $75 \mu\text{g DON l}^{-1}$ beer, we adapted the first step of the method by decreasing the volume of 20 ml of beer. Thus, only 5 ml of degassed and centrifuged beer were mixed with 30 ml of acetonitrile. Assaying DON recovery experiments in triplicate on three types of beer spiked at 20, 50 and $100 \mu\text{g l}^{-1}$, the overall average recovery was $93 \pm 4 \%$ ($n = 27$). The LOD and LOQ values were 2 and $6 \mu\text{g DON l}^{-1}$ of beer, respectively. Note that the recovery rate at $100 \mu\text{g DON l}^{-1}$ estimated at 94 % showed a good precision ($\text{RSD} = 5 \%$, $n = 9$). Care must then be taken not to exceed the capacity of the immunoaffinity column.

Occurrence of OTA in organic and conventional beers marketed in Belgium

Table I presents the OTA occurrence in organic and conventional beers produced in Belgium during the 2003-2004 period.

[Insert table I about here]

OTA was detected in 78 % of conventional samples at a mean level of $25 \pm 38 \text{ ng l}^{-1}$ (range: 19 - 198 ng l^{-1}). All organic beers were OTA contaminated at a mean level of $182 \pm 275 \text{ ng l}^{-1}$ (range: 18 - 1134 ng l^{-1}). Nevertheless, the magnitude between these means was not statistically significant ($p = 0.1758$). OTA above 200 ng l^{-1} was quantified in 10 organically produced beers and in none of conventional samples. The statistical analysis showed that the frequencies of quantifiable levels of OTA were significantly higher ($p < 0.0001$) in contaminated organically produced beers (95 %) than in their conventional counterparts (50 %).

Twenty different brands of beers were tested and no significant brand effect ($p = 0.2093$) in the OTA content was observed for all samples combined. In contrast, the variation between different batches was significant ($p < 0.0001$), showing thus an heterogeneous quality in brewing raw materials (as far as OTA contamination is concerned) within the production of the same brand. This might indicate that different parts of the same stock of raw materials or stocks of various origins

were used for the preparation of different batches. One must be aware that in stored-grain bulks, "hot spots" of OTA may accidentally occur in a given stock, even if the optimal required conditions for mould development have not been reached. Exceptionally, high concentrations of OTA in a batch of beer may thus occur if highly contaminated raw materials were introduced in the brewing process. This finding corroborates with the temporal fluctuation in OTA contamination in Belgian beers observed by Tangni *et al.* (2002). One must realize that the mycotoxin contamination status covers a limited period and that year-to-year variations may often occur as specified by Scott & Kanhere (1995) and Jorgensen (2005).

Numerous surveys have been conducted on OTA contamination in conventionally produced beers across the world. Nakajima *et al.* (1999) analyzed OTA in 46 European beers and found an average concentration of 11.4 ng l^{-1} (range: $1.7 - 66.2 \text{ ng l}^{-1}$) in 43 contaminated samples ($\text{LOD} = 1 \text{ ng l}^{-1}$). In Italy, Visconti *et al.* (2000) analysed the OTA content in 61 samples of 10 domestic and 51 imported beers and found OTA levels ranging from 10 to 135.2 ng l^{-1} ($\text{LOD} = 10 \text{ ng l}^{-1}$). Beers purchased in Denmark were also investigated for OTA contamination by Jorgensen (1998). All samples ($n = 21$) contained traces of OTA ($>1 \text{ ng l}^{-1}$) with a mean content of 49 ng l^{-1} (highest value = 160 ng l^{-1}). Similarly, Araguas *et al.* (2005) studied 21 Spanish beers and found in 67 % of the tested samples a mean of 38 ng OTA l^{-1} (highest value = 205 ng l^{-1}). One must note that these are in accordance with the current study for the conventional type of beer.

So far, a comparison between conventionally (49 samples) and organically (13 samples) produced beers in Belgium has been performed by Tangni *et al.* (2002). These authors found that the mean OTA content was significantly higher ($p = 0.007$) in organic (81 ng l^{-1}) than in conventional (22 ng l^{-1}) beers sampled in 1998-1999. In contrast, no significant difference ($p = 0.233$) was shown for beers collected in 2000-2001, although conventionally produced beers contained higher levels of OTA (49 ng l^{-1}) than organically produced counterparts (29 ng l^{-1}). The highest OTA content (185 ng l^{-1}) was found in an organically marketed beer. In the present study, attention was paid to the

sampling procedure by keeping a balanced sample size between organic and conventional groups. We observed that bad cases of contamination occasionally occur in organically produced beers, with concentrations that can be much higher (up to 1134 ng l⁻¹) than the limit of 200 ng l⁻¹ (FAO 2004). Independently from the mode of its production, high OTA levels in beer have been seldom presented in the recent literature. In fact, Gumus *et al.* (2004) analysed 135 Turkish beers and 15 imported beers and found that 28 % of the samples were OTA contaminated at levels ranging from 100 to 8100 ng l⁻¹. It was found that OTA levels were above 3000 ng l⁻¹ in 6 tested samples. So, the risk of highly contaminated batches occurring from time to time should not be underestimated.

Evidence of temporal fluctuation in OTA contamination in beers

Taking into account the high contamination levels obtained in the above sample, a complementary sampling was performed in March 2005. It was based on the same brands as those analysed in the 2003-2004 period. One bottle (one batch) of each brand was investigated.

Of 10 conventional beers, OTA was detected in 9 samples at mean concentration of 21 ng l⁻¹ (median = 12; range: 3 - 67 ng l⁻¹) whereas all organically produced beers (10) were OTA contaminated at the mean level of 75 ng l⁻¹ (median = 54, range: 19 - 158 ng l⁻¹). So, none of these beers exceeded the limit of 200 ng OTA l⁻¹, showing thus the temporal fluctuation of OTA occurrence in beers belonging to the same brands. One should, however, pay attention to the fact that the samples included in this complementary survey were probably produced after the regulation EC No 466/2001 entered in force, which established a maximum level of 3 µg OTA kg⁻¹ malt (European Commission Regulation 2001). The use of a good quality (as far as OTA contamination is concerned) malt within this legal limit may further avoid the high OTA levels sporadically found in some batches. So, regarding the legal limits, there is a need to continue the work of setting maximum levels for more mycotoxins in more foodstuffs.

Occurrence of DON in organic and conventional beers marketed in Belgium

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table II shows the results of DON contamination in organically (40) and conventionally (40) produced beers in Belgium.

[Insert table II about here]

The overall incidences of DON were of 67 and 80 % in conventionally and organically produced beers, respectively. Statistical analysis showed that the frequencies of quantifiable levels of DON were comparable ($p = 0.4586$) in organic (25 %) and conventional (32 %) beers. The respective averages were of 6 and 4 $\mu\text{g DON l}^{-1}$ whilst the highest levels were of 22 and 14 $\mu\text{g DON l}^{-1}$. The magnitude between these means was not statistically significant ($p = 0.0768$). Analysis of variance revealed homogeneous DON contamination between different brands whatever the mode of production ($p = 0.8639$). In contrast, a statistically significant batch effect ($p < 0.0001$) was noticed within the contaminated samples belonging to the same brand.

As for all samples combined ($n = 80$), the evaluation of DON contamination showed that 74 % of the tested beers were contaminated at a mean level of 5 $\mu\text{g l}^{-1}$. So far, several reports are available on the extent of DON contamination in beer. Of 54 domestic and imported beers in Korea, Shim *et al.* (1997) reported an incidence of 26 % for DON contamination ranging from 1 to 23 $\mu\text{g l}^{-1}$. Molto *et al.* (2000) have found a mean of 14.3 $\mu\text{g l}^{-1}$ (range: 5-221 $\mu\text{g l}^{-1}$) in 22 out of 50 Argentinean beers. The European evidence of DON occurrence in beer is backed up by a Canadian survey of 50 beers brewed in Canada or imported from Europe (Scott *et al.* 1993). This investigation revealed that 29 samples were DON contaminated with an average of 5.4 $\mu\text{g l}^{-1}$ (range: 0.3 - 50.3 $\mu\text{g l}^{-1}$). More specifically, imported beers from Europe were contaminated at levels ranging from 1.1 to 15.8 $\mu\text{g DON l}^{-1}$. Weddeling *et al.* (1994) reported that all the 18 German beers tested were DON contaminated with an average of 5 $\mu\text{g l}^{-1}$. A much higher contamination level was found in another German study by Niessen *et al.* (1993). These authors found that 190 out of 196 German beers were contaminated at levels ranging from 148 to 569 $\mu\text{g l}^{-1}$ (mean = 205 $\mu\text{g l}^{-1}$). These beers were, however, part of a study on gushing and were therefore not randomly selected. Recently,

contamination data have not shown high levels of DON in beer. Indeed, Schothorst & Jekel (2003) analysed 51 beer samples collected in the Netherlands and found that DON occurs at a concentration above $25 \mu\text{g l}^{-1}$ (LOQ value) in only 3 samples, with quantities ranging from 26 to $41 \mu\text{g l}^{-1}$. Papadopoulou-Bouraoui *et al.* (2004) detected DON in 87 % of the analysed beers ($n = 313$, $\text{LOD} = 3.7 \mu\text{g l}^{-1}$) with concentrations ranging from 4.0 to $56.7 \mu\text{g l}^{-1}$. Most samples (73 %) had contamination levels below $20 \mu\text{g l}^{-1}$. Our results compare well with the low level of DON contamination obtained in these previous studies. We therefore conclude that DON contamination in Belgian beers should not be a major matter of concern. So far, no regulatory guideline for DON contamination in beer is available in terms of safe limit. Indeed, the recent European Commission Regulation No 856/2005 amending the EC regulation No 466/2001 as regarding the *Fusarium* toxins states only on cereals and cereal based product not including beer (European Commission Regulation 2005).

From an analytical point of view, the present HPLC analytical procedure allows to detect DON in beer at very low levels and can be routinely used to assay the concentrations of DON in beers. Previous studies of DON determination in beer commonly involved gas chromatography (Scott *et al.* 1993, Molto *et al.* 2000, Schothorst & Jekel 2003) coupled with ECD, FID, MS or MS/MS detectors after derivatisation. Enzyme-linked immunosorbent assay is also used (Niessen *et al.* 1993, Papadopoulou-Bouraoui *et al.* 2004), but accurate quantification with this method is often limited, owing to cross reactivities (Schneider *et al.* 2004).

Co-occurrence of OTA and DON contents in organically and conventionally produced beers

Based on the DON detection limit, two subgroups of DON-positive (levels $\geq \text{LOD}$, $n = 59$) and DON-negative (level $< \text{LOD}$, $n = 21$) were considered.

The mean DON in the positive group ($7 \pm 4 \mu\text{g l}^{-1}$) was obviously significantly higher ($p < 0001$) than the level of contamination in the negative group. The respective concentrations of 8 ± 6 and 5

$\pm 2 \mu\text{g DON l}^{-1}$ were found in conventional and organic positive sub groups. In contrast, the mean OTA ($107 \pm 230 \text{ ng l}^{-1}$) in the DON-positive group was comparable ($p = 0.085$) to the mean OTA ($48 \pm 71 \text{ ng l}^{-1}$) found in the DON-negative group. The same trend was separately observed either for conventionally (30 ± 44 versus $14 \pm 12 \text{ ng OTA l}^{-1}$) or organically (173 ± 296 versus $105 \pm 92 \text{ ng OTA l}^{-1}$) produced beers, highlighting that a real significant co-occurrence of both toxins was not found for the tested beers, whatever their mode of production was. Moreover, non-significant correlation coefficients were derived between OTA and DON levels for either conventional ($r = +0.125$, $p = 0.441$) or organic ($r = +0.091$, $p = 0.576$) group of beers.

If contaminated raw materials are used during the brewing process, beer can be contaminated by all key cereal mycotoxins, except perhaps citrinin, which does not seem to survive the mashing step (Galvano *et al.* 2005). Zearalenone and OTA or aflatoxin B₁ and OTA coexisted in beers analysed in Japan (Nakajima *et al.* 1999) and in South-Africa (Odhav & Neckair 2002). Co-occurrence of trichothecenes such as DON, nivalenol, diacetoxyscirpenol, 3-acetyl deoxynivalenol, 15-acetyl deoxynivalenol and neosolaniol have been found in Canada (Scott *et al.* 1993) as well as in Argentina (Molto *et al.* 2000). Note that the simultaneous occurrence of two or more toxins in cereals has frequently been documented in the literature, e.g. barley, malt, maize and wheat contaminated with DON, nivalenol and zearalenone (Lee *et al.* 1986, Park *et al.* 1992, Tanaka *et al.* 2000), maize contaminated with fumonisin B₁, DON, T-2 toxin, nivalenol and zearalenone (Ali *et al.* 1998), wheat contaminated with DON and OTA (Birzele *et al.* 2000) and barley contaminated with OTA and DON (Olsson *et al.* 2002). These findings highlight the problems associated with ingestion of multiple toxins that may have an interactive effect on the health of the consumers.

Assessment of the contribution of beer to the OTA and DON exposure for the consumers

In recent years, thorough risk assessments of the trichothecenes and of OTA established a provisional maximum tolerable daily intake of $1 \mu\text{g DON kg}^{-1} \text{ bw}$ and a tolerable daily intake (TDI)

of 5 ng OTA kg⁻¹ bw (FAO/WHO 2001). Considering a daily intake of 0.3 l beer per capita in Belgium (CBB 2004) and the mean levels of DON and OTA found in the present study, beer consumption contributes simultaneously to 3 % and 10 % of the TDI of DON and OTA for an adult of 60 kg. We conclude that moderate daily intakes of DON and OTA are provided through beer consumption in Belgium. Beer consumption should thus not lead to acute toxic reactions linked to DON and OTA. It should however be included in calculations of total exposure to DON and OTA, together with other sources of cereals and other foodstuffs, which undoubtedly provide these mycotoxins as well. So far, the effects of a chronic exposure to low levels of these mycotoxins (long-term exposure) are not yet known (Gudmestad *et al.* 1997) and the combined effects of mycotoxins may increase their toxic effects. As an example, Foster *et al.* (1986) showed that DON in naturally contaminated feed has a higher toxicity in comparison to pure DON, suggesting that co-contamination with some unknown toxic metabolite or co-occurrence with a non-toxic compound enhances the toxicity of DON. Moreover, it is noticed that DON belongs to the group of several trichothecenes with a common basic chemical structure, which are produced by *Fusarium* fungi (e.g. T-2 Toxin, HT-2 toxin, nivalenol). These mycotoxins may share common mechanisms of action, which may lead to significant interactions. For all these reasons, the TDI for DON is determined only temporarily (FAO/WHO 2001). Generally, the problem of multi-contamination should not be underestimated.

In conclusion, the present study has further warranted the occurrence of low levels of OTA in beer. OTA was detected in about 90 % of the samples, but the levels were often far below the limit of 200 ng OTA l⁻¹. A few batches from organic beers were found to be much more contaminated, at levels exceeding this limit. The situation appeared to have improved from 2003-2004 to 2005 since none of the sample collected in 2005 exceeded the limit of 200 ng OTA l⁻¹. This improvement may reflect the newly imposed controls on malt. Occurrence of DON in beers should not be a major cause for concern since this toxin was found at low levels in a limited number of beers.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

This research project has been carried out within the framework of the Scientific support plan for a sustainable development policy (SPSD II) Part I “Sustainable production and consumption patterns” (Project CP/30 Mycotoxin contamination of regular and organic foodstuffs). The financial support of the Belgian Federal Science Policy Office is acknowledged here. F. Van Hove also acknowledges the financial support of the BCCM Research Programme (contract BCCM C3/10/003) from the same Office.

References

Ali N, Sardjono-Yamashita A, Yoshizawa T. 1998. Natural co-occurrence of aflatoxins and *Fusarium* mycotoxins (fumonisins, deoxynivalenol, nivalenol and zearalenone) in corn from Indonesia. Food Additives and Contaminants 15:377-384.

Araguas C, Gonzalez-Penas E, Lopez de Cerain A. 2005. Study on ochratoxin A in cereal-derived products from Spain. Food Chemistry 92:459-464.

Baxter ED. 1996. The fate of ochratoxin A during malting and brewing. Food Additives and Contaminants 13 suppl.:23-24.

Baxter DE, Slaiding IR, Kelly B. 2001. Behavior of ochratoxin A in Brewing. Journal of the American Society of Brewing Chemists 59:98-100.

Birzele B, Prange A, Kramer J. 2000. Deoxynivalenol and ochratoxin A in German wheat and changes of level in relation to storage parameters. Food additives and contaminants 17:1027-1035.

Boeira LS, Bryce JH, Steward GG, Flannigan B. 2000. The effect of combinations of *Fusarium* mycotoxins (deoxynivalenol, zearalenone and fumonisin B1) on growth of brewing yeasts. Journal of Applied Microbiology 88:388-403.

- CBB (Belgian Brewery Confederation). 2004. Aperçu du secteur de la Braserie Belge, Statistiques-l'année 2004. Source Banque Nationale et Brasseurs Belges. Available at: [http://www.beerparadise.be/fr/pdf/tabel%202%20frans.jpg].
- European Union, European Commission Regulation No 466/2001 of 8 March 2001 setting maximum levels for certain contaminants in foodstuffs. Official Journal of the European Communities L 77 (16.3.2001), 13p.
- European Union, European Commission Regulation N° 856/2005 amending EC Regulation No 466/2001 as regards *Fusarium* –toxins setting maximum levels for certain contaminants in foodstuffs Official Journal of the European Communities L143 (6.6.2005), 8p.
- FAO. 2004. Food and Nutrition Paper 81: Worldwide regulations for mycotoxins in food and feed in 2003. 183 p. Available at : ftp://ftp.fao.org/docrep/fao/007/y5499f/y5499f00.pdf.
- FAO/WHO. 2001. Joint Food and Agriculture Organization/World Health Organization. (FAO/WHO) Expert Committee on Food Additives, fifty-sixth meeting. Geneva, 6–15 February 2001, pp1-33. Online available at: http://www.fao.org/es/ESN/Jecfa/jecfa56.pdf or at: http://www.inchem.org/documents/jecfa/jecmono/v47je01.htm.
- Finamore A, Britti MS, Roselli M, Bellovino D, Gaetani S, Mengheri E. 2004. Novel approach for food evaluation. Results of a pilot experiment to evaluate organic and conventional foods. Journal of Agriculture and Food Chemistry 52:7425-7431.
- Foster BC, Trenholm HL, Friend DW, Thompson BK, Hartin KE. 1986. Evaluation of different sources of deoxynivalenol (vomitoxin) fed to swine. Canadian Journal of Animal Science 66:1149-1154.
- Galvano F, Titieni A, Piva G, Pietri A. 2005. Mycotoxins in the human food chain. In: D. Diaz, Editor. The mycotoxin Blue Book. Nottingham (England): Nottingham University Press. pp 187-224.

- Gudmestad N, Taylor R, Schwarz P. 1997. How healthy is your malt ? What you should know about a disease that could affect your beer ? The Brewers' Market Guide Online available at [<http://www.brewingtechniques.com/bmg/gudmestad.html>].
- Gumus T, Arici M, Demirci M. 2004. A survey of barley, malt, and beer contamination with ochratoxin A in Turkey. *Journal of the Institute of Brewing* 110:146-149.
- Jorgensen K. 1998. Survey of pork, poultry, coffee, beer and pulses for ochratoxin A. *Food Additives and Contaminants* 15:550-554.
- Jorgensen K. 2005. Occurrence of ochratoxin A in commodities and processed food: A review of EU occurrence data. *Food Additives and Contaminants* 22 (Supplement 1):26-30.
- Kroes R, Müller D, Lambe J, Löwik MRH, Van Klaveren J, Kleiner J, Massey R, Mayer S, Urieta I, Verger P, Visconti A. 2002. Assessment of intake from the diet. *Food and Chemical Toxicology* 40:327-385.
- Lee U, Jang H, Tanaka T, Hasegawa A, Oh Y, Cho C, Sugiura Y, Ueno Y. 1986. Further survey on the *Fusarium* mycotoxins in Korean cereals. *Food Additives and Contaminants* 3:253-261.
- MacDonald S, Wilson P, Barnes K, Damant A, Massey R, Mortby E, Shepherd MJ. 1999. Ochratoxin A in dried vine fruit: method development and survey. *Food Additives and Contaminants* 16:253-260.
- Macdonald S, Prickett TJ, Wildey KB, Chan D. 2004. Survey of ochratoxin A and deoxynivalenol in stored grains from the 1999 harvest in the UK. *Food Additives and Contaminants* 21:172-181.
- Magkos F, Arvaniti F, Zampelas A. 2003. Putting the safety of organic food into perspective. *Nutrition Research Review* 16:211-222.

- Molto G, Samar MM, Resnik S, Martinez EJ, Pacin A. 2000. Occurrence of trichothecenes in Argentinean beer: a preliminary exposure assessment. *Food Additives and Contaminants* 17: 809-813.
- Moss MO. 1998. Recent studies of mycotoxins. Symposium series (Society for Applied Microbiology) 27:62S-76S.
- Nakajima M, Tsubouchi H, Miyabe M. 1999. A survey of ochratoxin A and aflatoxins in domestic and imported beers in Japan by immunoaffinity and liquid chromatography. *Journal of AOAC International* 82:897-902.
- Niessen L, Donhauser S. 1993. Fate of deoxynivalenol in the process of brewing and its prevalence in commercial beer. In K. Scudamore, editor. *Occurrence and Significance of Mycotoxins*. Central Science Laboratory, Slough, 203-207.
- Niessen L, Boehm-Schraml M, Vogel H, Donhauser S. 1993. Deoxynivalenol in commercial beer - screening for the toxin with an indirect competitive ELISA. *Mycotoxin Research* 9:99-109.
- Odhav B, Neckair V. 2002. Mycotoxins in South African traditionally brewed beers. *Food Additives and Contaminants* 19:55-61.
- Olsson J, Börjesson T, Lundstedt T, Schnürer J. 2002. Detection and quantification of ochratoxin A and deoxynivalenol in barley grains by GC-MS and electronic nose. *International Journal of Food Microbiology* 72:203-214.
- Ottener H, Majerus P. 2001. Ochratoxin A (OTA) in coffee: nation-wide evaluation of data collected by German Food Control 1995-1999. *Food Additives and Contaminants* 18:431-435.
- Papadopoulou-Bouraoui A, Vrabcheva T, Valzacchi S, Stroka J, Anklam E. 2004. Screening survey of deoxynivalenol in beer from the European market by an enzyme-linked immunosorbent assay. *Food Additives and Contaminants* 21:607-617.

- 1
2
3 Park KJ, Park AR, Lee YW. 1992. Natural occurrence of *Fusarium* mycotoxins of the 1990 barley
4 crop in Korea. Food Additives and Contaminants 9:639-645.
5
6
7
8
9 Pettersson H. 1998. Intercomparison of trichothecene analysis and feasibility to produce certified
10 calibrants and reference material. European Commission. BCR information reference
11 materials. Brussels - Luxembourg, 143 p.
12
13
14
15
16 Pitt JI. 1987. *Penicillium viridicatum*, *Penicillium verrucosum*, and production of ochratoxin A.
17 Applied and Environmental Microbiology 53:266-269.
18
19
20
21 Pussemier L, Larondelle Y, Van Peteghem C, Huyghebaert A. 2006. Chemical safety of
22 conventionally and organically produced foodstuffs: a tentative comparison under Belgian
23 conditions. Food Control 17:14-21.
24
25
26
27
28
29 Samson RA, Houbraken JAMP, Kuijpers AFA, Frank, JM, Frisvad JC. 2004. New ochratoxin A or
30 sclerotium producing species in *Aspergillus* section *Nigri*. Studies in Mycology 50:45-61.
31
32
33
34 Schneider E, Curtui V, Seidler C, Dietrich R, Usleber E, Martlbauer E. 2004. Rapid methods for
35 deoxynivalenol and other trichothecenes. Toxicology Letters 153:113-121.
36
37
38
39
40 Schothorst RC, Jekel AA. 2003. Determination of trichothecenes in beer by capillary gas
41 chromatography with flame ionisation detection. Food Chemistry 82:475-479.
42
43
44
45 Scott PM, Kanhere SR, Weber D. 1993. Analysis of Canadian and imported beers for *Fusarium*
46 mycotoxins by gas chromatography-mass spectrometry. Food Additives and Contaminants
47 10:381-389.
48
49
50
51
52 Scott PM, Kanhere SR. 1995. Determination of ochratoxin A in beer. Food Additives and
53 Contaminants 12:591-598.
54
55
56
57
58
59
60

- Serra R, Braga A, Venâncio A. 2005. Mycotoxin-producing and other fungi isolated from grapes for wine production, with particular emphasis on ochratoxin A. *Research in Microbiology* 156:515-521.
- Shim WB, Kim JC, Seo JA, Lee YW. 1997. Natural occurrence of trichothecenes and zearalenone in Korean and imported beers. *Food Additives and Contaminants* 14:1-5.
- Tanaka T, Yoneda A, Inoue S, Sugura Y, Ueno Y. 2000. Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography-mass spectrometry. *Journal of Chromatography A* 882:23-28.
- Tangni EK, Ponchaut S, Maudoux M, Rozenberg R, Larondelle Y. 2002. Ochratoxin A in domestic and imported beers in Belgium: occurrence and exposure assessment. *Food Additives and Contaminants* 19:1169-1179.
- Thirumala-Devi K, Mayo MA, Reddy G, Tangni, EK, Larondelle Y, Reddy DV. 2001. Occurrence of ochratoxin A in black pepper, coriander, ginger and turmeric in India. *Food Additives and Contaminants* 18: 830-835.
- Visconti A, Pascale M, Centonze G. 2000. Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography. *Journal of Chromatography A* 888:321-326.
- Weddeling KH, Bäßler MS, Doerk H, Baron G. 1994. Oriented tests for the application of the enzymimmunological process for evidencing deoxynivalenol, ochratoxin A and zearalenone in brewing barley, malt and beer. *Monatsschrift für Brauwissenschaft* 47:94-98.
- Wood GM, Entwistle C, Farnell PJ, Patel S, Boenke A. 1996. The certification of the ochratoxin A mass fraction of two wheat reference materials (CRMs 471 & 472). European Commission BCR information. Brussels-Luxembourg. 58 p.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Zimmerli B, Dick R. 1995. Determination of ochratoxin A at the ppt level in human blood, serum, milk and some foodstuffs by high-performance liquid chromatography with enhanced fluorescence detection and immunoaffinity column cleanup: methodology and Swiss data. *Journal of Chromatography B: Biomedical Applications* 666:85-99.

For Peer Review Only

List of tables

Table I. Occurrence of OTA in organic and conventional beers sold on the Belgian market (period 2003 – 2004).

Table II. Occurrence of DON in organic and conventional beers sold on the Belgian market (period 2003- 2004).

Table I. Occurrence of OTA in organic and conventional beers sold on the Belgian market (period 2003 - 2004).

Occurrence	Conventional	Organic	Total
Number of samples	40	40	80
n.d. ^a	9 (22 %)	0	9 (11 %)
trace ^b	11 (28 %)	2 (5 %)	13 (16 %)
LOQ - 100 ng l ⁻¹	18 (45 %)	22 (55 %)	40 (50 %)
101 - 200 ng l ⁻¹	2 (5 %)	6 (15 %)	8 (10 %)
200.1 - 1200 ng l ⁻¹	0	10 (25 %)	10 (13 %)
Incidence	78 %	100%	89 %
Levels of contamination (ng l⁻¹)			
Mean ^c ± SD	25 ± 38	182 ± 275	103 ± 162
Median ^c	15	45	29
Range ^d	19 – 198	18 – 1134	18 - 1134

^a: n.d., not detected (OTA concentration in sample < 5 ng l⁻¹)

^b: traces means that the concentration of OTA is between the limit of detection (LOD = 5 ng l⁻¹) and the limit of quantification (LOQ = 18 ng l⁻¹)

^c: mean and median are computed assuming that non-detected samples contain half the LOD while samples with traces levels of OTA contain half the LOD + LOQ

^d: range of samples with OTA contents ≥ LOQ

Table II. Occurrence of DON in organic and conventional beers sold on the Belgian market (period 2003- 2004).

Occurrence	Conventional	Organic	Total
Number of samples	40	40	80
n.d. ^a	13 (33 %)	8 (20 %)	21 (26 %)
trace ^b	14 (35 %)	22 (55 %)	36 (45 %)
LOQ - 10 µg l ⁻¹	5 (12 %)	8 (20 %)	13 (16 %)
10 - 20 µg l ⁻¹	6 (15 %)	2 (5 %)	8 (10 %)
20 - 30 µg l ⁻¹	2 (5 %)	0	2 (3 %)
> 30 µg l ⁻¹	0	0	0
Incidence	67 %	80 %	74 %
Levels of contamination (µg l⁻¹)			
Mean ^c ± SD	6 ± 6	4 ± 3	5 ± 4
Median ^c	4	4	4
Range ^d	6 – 22	6 – 14	6 – 22

^a: n.d., not detected (DON concentration in sample < 2 µg l⁻¹)

^b: traces means that the concentration of DON is between the limit of detection (LOD = 2 µg l⁻¹) and the limit of quantification (LOQ = 6 µg l⁻¹)

^c: mean and median are computed assuming that non-detected samples contain half the LOD while samples with traces levels of DON contain half the LOD + LOQ

^d: range of samples with DON contents ≥ LOQ