Determination of bixin and norbixin in meat using liquid chromatography and photodiode array detection.

Herlinde Noppe, Sonja Abuin Martinez, Karolien Verheyden, Joris van Loco, R. Companyo Beltran, Hubert de Brabander

► To cite this version:

Herlinde Noppe, Sonja Abuin Martinez, Karolien Verheyden, Joris van Loco, R. Companyo Beltran, et al.. Determination of bixin and norbixin in meat using liquid chromatography and photodiode array detection.. Food Additives and Contaminants, 2009, 26 (01), pp.17-24. 10.1080/02652030802322564 . hal-00577300

HAL Id: hal-00577300

<https://hal.science/hal-00577300>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Determination of bixin and norbixin in meat using liquid chromatography and photodiode array detection.

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2008-045.R3
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	17-Jun-2008
Complete List of Authors:	Noppe, Herlinde; Ghent University, Faculty of Veterinary Medicines, Veterinary Public Health and Food Safety Martinez, Sonja; Barcelona University, Faculty of Chemistry, Analytical Chemistry Verheyden, Karolien; Ghent University, Faculty of Veterinary Medicines, Veterinary Public Health and Food Safety Van Loco, Joris; Institute of Public Health, Pharmaco-bromatology, Food Section Beltran, R.; Barcelona University, Faculty of Chemistry, Analytical Chemistry De Brabander, Hubert; Ghent University, Veterinary Public Health and Food Safety
Methods/Techniques:	Chromatography - LC/MS, In-house validation, Method validation
Additives/Contaminants:	Residues
Food Types:	Meat

SCHOLARONE™
Manuscripts

Determination of bixin and norbixin in meat using liquid chromatography and photodiode array detection

¹H. NOPPE, ²S. ABUÍN MARTINEZ, ¹K. VERHEYDEN, ³J. VAN LOCO, ²R. COMPANYÓ BELTRAN, ¹H.F. DE BRABANDER

¹*Ghent University, Faculty of Veterinary Medicine, Research group of Veterinary Public Health and Zoonoses, Laboratory of Chemical Analysis, Salisburylaan 133, B-9820 Merelbeke, Belgium*

²*Barcelona University, Faculty of Chemistry, Department of Analytical Chemistry, Avinguda Diagonal, 647, E-08028 Barcelona, Spain*

³*Institute of Public Health, Department of Pharmaco-bromatology, Food Section, J. Wytsmanstraat, B-1050 Brussels, Belgium*

Correspondence: H.F. De Brabander. E-mail: Hubert.DeBrabander@UGent.be

Abstract

In this study, the development of an analytical method that enables routine analysis of the dyes annatto, bixin and norbixin in meat tissue is described. Liquid solid extraction was carried out using acetonitrile. Analysis was by HPLC with photodiode array detection using 2 fixed wavelengths (458 and 486 nm). Additionally, the possibilities of ion trap mass spectrometry (MS) were assessed. Method performance characteristics according to Commission Decision 2002/657/EC were determined, with recoveries between 99 and 102 % and calibration curves being linear in the 0.5 to 10 mg kg⁻¹ range. The limit of quantification was 0.5 mg kg⁻¹.

Keywords: annatto, bixin, norbixin, meat, liquid chromatography, photodiode array

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

As colour is often a key consumer perception for food preference and acceptability, natural and synthetic colorants have extensively been used to colour food, drugs and cosmetics (Bittencourt et al., 2005). Due to restrictions on the use of these synthetic colorants, there has been a growing interest from the food industry to use natural colorants.

Annatto extracts (E160b) are orange/red natural carotenoid dyes obtained from the seeds of the tropical shrub *Bixa Orellana* L. and are nowadays widely used by the food industry to colour many food products including high fat dairy products such as butter and cheese, cereals, snack foods, seasonings, creamers, ice creams, flour, sugar confectionary, soft drinks and fish (Lancaster and Lawrence, 1996, Scotter et al., 1994 & 1995, 2002). The principal coloring component of annatto seeds is cis-bixin, which is soluble in most polar organic solvents to which it imparts an orange colour, but which is largely insoluble in vegetable oil (Scotter et al., 1994 & 1995, 2002). Cis-bixin is the monomethyl ester of cis-norbixin (figure 1) (Scotter et al., 1994). The amounts of the active pigments bixin and norbixin in annatto can vary from less than 1 % to over 85 %, dependent on the type of annatto extract e.g. water, vegetable oil, solvent (Tennant and O’Callaghan, 2005, WHO, 2007) and are affected by degradation agents, such as light, temperature, air, anti- and pro-oxidants and pH (Najar et al., 1988). The degradation products are of commercial significance, since by controlling the degree of the degradation, the orange/red to yellow colour balance of an Annatto formulation can be adjusted (Scotter, 1995).

Notwithstanding the large use of annatto pigments as food colorants, practically no information exists about their toxicological properties for animals and humans. Toxicity was mostly determined in animals using commercial annatto preparations containing undetermined amounts of bixin and norbixin (Engelberth-Holm and Iversen, 1955, Van Esch et al., 1959). Toxicological data about annatto pigments are limited, possibly because food additives derived from natural sources have been given exemption from certification (Hallagan et al., 1995). The Joint FAO/WHO Expert Committee on

Food Additives (JECFA) estimated the ADI for annatto as 0-2.5 mg/kg body weight / day (for a preparation containing 2,6 % carotenoids expressed as bixin) and 0-0.065 mg/kg body weight/day expressed as the pure pigment (WHO, 2007).

In recent years, there has been a growing concern about the presence of these compounds in edible matrices of animal origin. Consequently the use of colorants in foodstuffs has been regulated by the European Union (Council Directive EC/94/36). Depending on the colorant (e.g. annatto or E160b) and the matrix (e.g. fish tissue and dairy products) Maximum Residue Levels (MRLs) have been proposed. Annatto extracts are listed amongst those colorants that may be used separately or in combination in certain foods up to the maximum levels specified (on a ready-for-consumption basis) (Scotter et al., 2002). Other edible matrices, e.g. molluscs, crustaceans, meat tissue, poultry and game, as well as their preparations may not contain added annatto.

Up to now, most analytical studies for annatto concerned with dairy products and high fat foods where annatto has well-established usage, such as cheese and dairy spreads (Scotter et al., 2002). Very few studies have documented the possibility to detect annatto or bixin and norbixin in meat tissue. For the determination of annatto, and more in particular bixin and norbixin, several detection techniques, such as ultraviolet (UV), photo-diode array (PDA) and ion trap mass spectrometry (MS) in combination with liquid chromatography (LC) were evaluated. Within this perspective, our study aimed at developing a routine residue method for the detection of annatto (bixin and norbixin) in meat using liquid chromatography. Additionally, this method was validated in accordance with the Commission Decision 2002/657/EC.

Experimental

Chemicals

Acetonitrile (AcN), ethanol (EtOH) and acetic acid were of HPLC grade and supplied by VWR (Merck, Darmstadt, Germany). HPLC grade Water was obtained from Across Organics (Fairlawn, NJ, USA).

1
2
3 83
4
5 84 *Standards*
6
7
8 85 Bixin and norbixin powders were provided by LGC Promochem (Molsheim, France) with a purity of
9
10 86 88.5 % and 97.1 % respectively. Annatto was purchased from Chromadex, Inc (Daimler St., Santa
11
12 87 Ana, CA, USA) and had a norbixin content of 14-16 %. Stock standard solutions of 200 ng μl^{-1} were
13
14 88 prepared in EtOH. Individual and composite working standards were prepared by appropriate dilution
15
16 89 of the standard stock solutions in 0.1 % acetic acid in AcN/ H_2O (80:20). According to the instructions
17
18 90 of the supplier all standard and working solutions were stored in the dark at -20 °C to minimize
19
20 91 degradation and isomerization.
21
22
23 92
24
25 93 *Extraction and clean-up*
26
27 94 Extraction of 1 g homogenized meat tissue was extracted with 2ml acetonitrile (AcN). For validation
28
29 95 gammon sausage was used and bixin, norbixin or annatto were spiked in the AcN. After vigorously
30
31 96 vortex mixing and centrifugation (5 min at 5500 rpm, 4 °C, 3726 g) (Sorvall RC 5C plus, Goffin
32
33 97 Meyvis, Belgium) 120 μl of the supernatant with addition of 30 μl 0.1 % aqueous acetic acid was used
34
35 98 for chromatographic analysis. For this, 60 μl was injected.
36
37
38 99
39
40 100 *Chromatographic instrumentation*
41
42 101 Analysis was carried out using a High Performance Liquid Chromatography (HPLC) apparatus
43
44 102 comprised of an 1100 series quaternary gradient pump and autosampler (Hewlett Packard, Palo Alto,
45
46 103 CA, USA). Chromatographic separation was achieved using a SymmetryTM C₁₈ column (5 μm particle
47
48 104 size, 150 x 2.1 mm, Waters, Milford, USA). The mobile phase consisted of a mixture of acetonitrile
49
50 105 (A) and 0.1 % aqueous formic acid (B). A linear gradient was run: 80 % A:20 % B, increasing to 95 %
51
52 106 A:5 % B over 10 min, increasing to 100 % A over 5 min (hold 5 min) and finally back to initial
53
54 107 conditions 80 % A:20 % B (hold 10 min). Total run time is 30 minutes at a mobile phase flow of 0.3
55
56 108 ml min^{-1} .
57
58
59 109
60

When detection was performed using photodiode-array detection, a TSP Spectra System UV6000 LP (ThermoFinnigan, San José, CA, USA) with deuterium lamp was used with a detection range between 300 and 580 nm (bandwidth 1 nm). Additionally, two fixed wavelengths, 458 and 486 nm (bandwidth 9 nm), were tested.

Mass spectrometric (MS) detection was carried out with a ThermoFinnigan LCQ DECA ion trap with electrospray ionisation (ESI) interface using the above described chromatographic parameters. Multiple mass spectrometry (MS²) spectra were acquired. The MS method consisted of segments, the first (0-4 min) in negative ionisation mode, with a full scan MS scan event and an MS² scan event (parent ion 379 MW, collision energy 25). The second segment (4-10 min), in positive ionisation mode, consisted also of a full scan MS scan event and an MS² scan event (parent ion 395 MW, collision energy 30). Isolation width was set at 2.0 and activation Q at 0.250.

Data analysis

Chromatograms and spectra were recorded and processed using Xcalibur 1.2. software (ThermoFinnigan, San José, CA, USA). All data were further analyzed using Microsoft® Excel (Microsoft Corporation, USA).

Quality assurance

Before and after sample analysis, a standard mixture (10 ng on column) of the targeted colour agents was injected to check the operation conditions of the LC-PDA devices. For quantitative analysis, calibration standards were spiked at 3 levels in sample matrix 0.5, 1 and 5 mg kg⁻¹. When using this analytical method for routine analysis, quality control of the method was carried out by analysis of blank and spiked comparable sample matrices together with every series of samples. At regular times, secondary quality control, using blind fortified samples, was carried out within the routine analysis.

Validation procedure

Method validation for the detection of annatto, bixin and norbixin in meat tissue was carried out according to Commission Decision 2002/657/EC concerning detection of residues of veterinary drug substances in matrices of animal origin. The examined validation parameters for the complete analytical method were selectivity/specificity, linearity, precision, trueness/recovery, applicability/ruggedness/stability and decision limit ($CC\alpha$) and detection ($CC\beta$) capability. For this, blank sausage meat samples were fortified with a standard solution of a mixture of the dyes at the level of 0.5, 0.75, 1, 2 and 4 mg kg⁻¹ unless otherwise mentioned.

Results and Discussion

Method performance

Up to now, analytical approaches for the detection of annatto, or bixin and norbixin, are described for different matrices, however, not for meat. Within this perspective, the aim of this study was to develop an analytical approach for the detection of annatto, and more in particular bixin and norbixin in meat tissue. The possibilities of 2 detection techniques, photodiode array (PDA) and ion trap mass spectrometry (MS) were evaluated.

With the developed method, the two compounds of annatto, norbixin and bixin, were separated and simultaneously detected by liquid chromatography (LC) using either photodiode array or ion trap mass spectrometry with electrospray ionisation (ESI). Using PDA detection, the main peaks obtained from the chromatographic separation of bixin and norbixin were scanned between 300 and 580 nm and were similar to those reported by Scotter (1995). More in particular, 2 fixed wavelengths were used, 458 and 468 nm. In figure 2, UV and MS² chromatograms and spectra of 10 ng bixin, norbixin and annatto on column are shown. As can be noticed, the PDA-spectra of norbixin and bixin are identical with absorption maxima on 458 and 486 nm. Moreover, a typical increase in absorption intensity was observed around 430 nm for both compounds. It was also observed that annatto mainly consists of norbixin, though both α - and β -isomers of bixin and norbixin were present, consistent with Lancaster and Lawrence (1996) (See figure 2).

Multiple mass spectrometry (MS²) was used to obtain structural information on the compounds corresponding to each chromatographic peak. Bixin and norbixin produced the [M-H]⁺ ion in full scan MS. This molecular ion, more in specific with m/z 379 and 395 for norbixin and bixin respectively, was used as precursor ion for MS². As can be seen in figure 2, the MS² product ions of norbixin (@379) are 253, 291, 310 and 335 whereas the main product ions of bixin (@ 395) are 317, 335, 345, 363 and 377. Using MS², detection is possible from a concentration level of 1 mg kg⁻¹ as the presence of several ions in the spectrum of bixin makes a clear interpretation difficult concerning matrix interferences. With PDA the limit of quantification is 0.5 mg kg⁻¹. Therefore, PDA detection was preferred for use in routine analysis.

Full in-house method validation for bixin and norbixin in meat

Identification

For method validation and interpretation of the obtained results, criteria as described in the Commission Decision 2002/657/EC were used. Targeted dye compounds in meat samples were identified based on comparison of their relative retention time ($\pm 2.5\%$), which is the ratio of the retention time of the analyte to that of the internal standard with a standard or fortified sample. When detection was carried out with PDA, the absorption maxima (± 2 nm) in the spectrum of the analyte were taken into account. Using mass spectrometric detection the ion ratio of the precursor/product ions in the obtained spectrum was used when the corresponding peak in the chromatogram had a signal-to-noise ratio of at least 3:1. In samples, the presence of bixin, norbixin and annatto was confirmed if the retention time, the maximum absorption wavelength and the shape of the UV spectrum of the unknown are in accordance with a fortified sample or a standard mixture. Comparison of the UV spectra of a blank sample fortified at the level of 10 mg kg⁻¹ for norbixin/bixin and at the level of 67 mg kg⁻¹ for annatto (=10 mg kg⁻¹ norbixin) is shown in figure 3.

Specificity/selectivity

One of the general requirements of EC/2002/657, namely the investigation of interferences of matrix components, specificity/selectivity was evaluated through the analysis of blank samples fortified

separately with the considered compounds at national action level (AL). This AL was based on the acceptable daily intake (ADI) for bixin and norbixin and a concentration level of 0.5 mg kg⁻¹ has been postulated. The specificity of the chromatographic method was proved since no significant interfering peaks with a signal to noise ratio of 3 or more were observed at the retention times of the targeted compounds by analysis of fortified samples. As described before, selectivity was proved since conform 2002/657/EC, analytes are identified on the basis of their relative retention time and the ion ratio of the precursor and product ions in the obtained spectrum.

Linearity

The linearity of the method was evaluated using repeated analysis of 5 calibration points in the concentration range of 0.5, 0.75, 1, 2 and 4 mg kg⁻¹. Correlation coefficients were between 0.93 and 1.

Trueness/precision

Because no Certified Reference Material (CRM) was available, trueness of the measurements was assessed by determining the recovery of additions of known amounts of the analytes to blank matrix. For this purpose, blank meat tissue was fortified with 0.5, 0.75, 1, 2 and 4 mg kg⁻¹ in 6 replicates and on 3 different days. All calculated recoveries fulfilled the criterion that recovery should be between 80 and 110 % for a mass fraction of, or greater than, 10 µg kg⁻¹, except for norbixin at the level of 2 mg kg⁻¹. The recoveries at this concentration level appeared to be higher for both compounds although there was no specific explanation. To evaluate the precision of the method, repeatability and within-laboratory reproducibility were determined. Repeatability was evaluated by calculating the coefficients of variation (CV) using the data of 2 series of 3 replicates of fortified samples of an identical matrix at 0.5 up to 4 mg kg⁻¹. These analyses were carried out on two different occasions by the same analyst and under repeatable conditions. According to the Horwitz (HW) equation, the CV for the repeated analysis of fortified samples at a mass fraction of 0.5 and 4 mg kg⁻¹ should not exceed 18 % and 13 % respectively. All calculated CV values for norbixin and bixin were between 3 and 9 %.

Reproducibility

Because no proficiency tests were available for annatto, and more specific for norbixin or bixin, in matrices of animal origin, to evaluate the reproducibility only the within-reproducibility was considered. To this end, 3 series of 5 replicates of fortified samples at 0.5 up to 4 mg kg⁻¹ were analysed by different operators and on different days. The calculated CVs for both different operators, different days of analysis and different concentrations were all between 3 and 17 % for both compounds considered. These values are of the same order of magnitude in comparison with the CVs calculated using the Horwitz equation.

Ruggedness

In order to determine the ruggedness of the analytical method, experiments were carried out by changing factors of the sample pre-treatment and clean-up like different SPE sorbents, extraction solvents and volumes. In addition, a lot of parameters of the chromatographic analysis, e.g. injection volume, liquid phases, chromatographic columns and solvent gradient programs were tested. For the PDA and MS² analysis different ionization modes and wavelengths were tested respectively. The evaluation of the ruggedness of our method resulted in restraining the parameters as described in the Materials and Methods section. Above this, a wide range of commercial processed pork meat products (e.g. sausages, non-smoked gammon, smoked gammon and sliced cold meat products) were analysed using the above described method. In all investigated matrices it was possible to determine annatto, and more specifically norbixin and bixin at 0.5 mg kg⁻¹ using PDA and 1 mg kg⁻¹ using MS² by spiking into the matrix (see also figure 4). No problems were encountered with co-extractive interferences, thus demonstrating the robustness of the method with respect to the analysis of processed meat products.

Decision limit (CC_α) and detection capability (CC_β)

Finally, also the decision limit, which is the concentration level at and above which the sample is non-compliant with 95 % certainty, was calculated for the considered coloring agents. This value was defined as the mean concentration of the spiked blank samples at AL concentration level plus 1.64 times the corresponding standard deviation at this concentration. The detection capability, which is the

1
2
3 248 smallest concentration that can be detected with 95 % certainty, was calculated as the CC α plus 1.64
4
5 249 times the above mentioned standard deviation. The CC α values for norbixin and bixin were 0.6 mg kg⁻¹
6
7 250 ¹ whereas calculation of the CC β values for these compounds gave 0.7 mg kg⁻¹.
8
9
10 251

11
12 252 In routine analysis, quantification of the obtained residues of norbixin and bixin was based on
13
14 253 calibration curves of 3 spikes (0.5, 1 and 5 mg kg⁻¹) in the same matrix of the sample to compensate
15
16 254 for matrix effects.
17
18 255

19
20 256 **Conclusion**
21

22
23 257 In this study, a routine quantitative approach for the determination of annatto or norbixin and bixin in
24
25 258 meat tissue in the low mg kg⁻¹ concentration range is described. Using this validated method, the dye
26
27 259 annatto, and more in particular its compounds bixin and norbixin, can be detected in meat tissue from
28
29 260 the concentration level of 0.5 mg kg⁻¹, which fits into the objectives of the inspection services. The
30
31 261 developed method has an excellent recovery and a good precision. The use of both photodiode array
32
33 262 and MS showed a large flexibility for the screening and confirmation of norbixin and bixin in routine
34
35 263 residue analysis.
36
37 264

38
39 265 **Acknowledgements**
40

41
42 266 The authors are grateful to Mieke Naessens and Els Laverge for assistance in the experimental work
43
44 267 and the skillful operation of the chromatographic devices.
45
46 268

47
48 269 **References**
49

50
51 270
52
53 271 2002/657/EC. 2002. Commission decision implementing council directive 96/23/EC concerning the
54
55 272 performance of analytical methods and the interpretation of results. Official Journal of the European
56
57 273 Communities L221: 8-36.
58
59 274

- 275 Bittencourt C, Felicissimo MP, Pireaux JJ and Houssiau L. 2005. Study of annatto from *Bixa Orellana*
276 seeds: an application of time-of-flight secondary ion mass spectrometry. *Spectroscopy Europe* 17: 16-
277 22
278
279 EC/94/36. 1994. European Parliament and council directive on colours for use in foodstuffs. *Official*
280 *Journal of the European Communities L* 237: 13-29.
281
282 Engelberth-Holm J, Iversen S. 1955. Is the vegetable annatto butter colour cancerogenic?
283 *Acta Pathologica, Microbiologica et Immunologica Scandinavica* 37: 483-492.
284
285 Gloria MBA, Vale SR, Bobbio PA. 1995. Effect of water activity on the stability of bixin in
286 an annatto extract-microcrystalline cellulose model system. *Food Chemistry* 52: 389-391.
287
288 Hallagan JB, Allen DC, Borelleca JF. 1995. The safety and regulatory status of food, drug and
289 cosmetics colour additives exempt from certification. *Food and Chemical Toxicity* 33: 515-
290 528.
291 Lancaster FE and Lawrence JF. 1996. High performance liquid chromatographic separation of
292 carminic acid, α -bixin and β -bixin, and α -norbixin and β -norbixin, and the determination of carminic
293 acid in foods. *Journal of Chromatography A* 732: 394-398.
294 Najar SV, Bobbio FO, Bobbio PA. 1988. Effects of light, air, anti-oxidants and pro-oxidants on
295 annatto extracts (*Bixa orellana*). *Food Chemistry* 29: 283-289.
296 Scotter MJ. 1995. Characterization of the colored thermal degradation products of bixin from annatto
297 and a revised mechanism for their formation. *Food Chemistry* 53: 177-185.

- 298 Scotter MJ, Wilson LA, Appleton GP, Castle L. 2000. Analysis of Annatto (*Bixa Orellana*) Food
299 Coloring Formulations. 2. Determination of Aromatic Hydrocarbon Thermal Degradation Products by
300 Gas Chromatography. Journal of Agricultural and Food Chemistry 48: 484-488.
- 301 Scotter MJ, Thorpe SA, Reynolds SL, Wilson LA and Strutt PR. 1994. Characterization of the
302 principal coloring components of annatto using high performance liquid chromatography with
303 photodiode-array detection. Food Additives and Contaminants 11: 301-315.
- 304
- 305 Sotter MJ, Castle L, Honeybone A, Nelson C. 2002. Method development and analysis of retail foods
306 for annatto food colouring material. Food Additives and Contaminants 19: 205-222.
- 307
- 308 Tennant DR and O'Callaghan M. 2005. Survey of usage and estimated intakes of annatto extracts.
309 Food Research International 38: 911-917.
- 310
- 311 Van Esch GH, Van Genderen H., Vink HH. 1959. Über die chronische verträglichkeit von
312 Annattofarbstoff. Zeitschrift für Lebensmitteleuntersuchung and forschung 111: 93-108.
- 313 Verger P. 1996. One example of utilization of the 'French approach'. Food Additives and
314 Contaminants 4: 417-419.
- 315
- 316 WHO. 2007. Safety Evaluation of certain food additives and contaminants. Online available:
317 whqlibdoc.who.int/trs/WHO_TRS_922.pdf

Figure 1. Chemical structure of bixin (MW 395) (A) and norbixin (MW 378) (B)

1

2 Figure 2. Chromatograms and MS² spectra of norbixin (A) and bixin (B), PDA spectrum of norbixin (C) and bixin (D), PDA spectrum
3 of annatto (E) and MS² spectrum of annatto (norbixin) (F)

1

2 Figure 3. Comparison of the PDA spectra of norbixin and bixin (fortified sample at 10 mg kg^{-1}) (A) and of annatto (fortified sample at
3 67 mg kg^{-1}) (B).

1

2 Figure 4. Chromatograms and PDA spectra of 1 g commercial pork sausage fortified with 0.5 mg kg⁻¹ norbixin (A) and bixin (B).