Determination of sterigmatocystin in cheese by high performance liquid chromatography – tandem mass spectrometry

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2008-122.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	21-Jun-2008
Complete List of Authors:	Versilovskis, Aleksandrs; Latvia University of Agriculture, Department of Chemistry; Ghent University, Department of Bio-Analysis Van Peteghem, Carlos; Ghent University, Pharmacy Department De Saeger, Sarah; Ghent University, Laboratory of Food Analysis
Methods/Techniques:	Clean-up-SPE, Chromatography - LC/MS
Additives/Contaminants:	Sterigmatocystin, Mycotoxin
Food Types:	Cheese

SCHOLARONE™
Manuscripts

3
4 2
5
6 3 **RESEARCH ARTICLE**

7 4
8
9
10 5 **Determination of sterigmatocystin in cheese by high performance liquid**
11
12 6 **chromatography – tandem mass spectrometry**

13
14
15
16 7
17 8 **A. Veršilovskis^{a,b*}, C. Van Peteghem^a, S. De Saeger^a**

18 9
19
20
21 10 *^aDepartment of Bio-Analysis, Laboratory of Food Analysis, Ghent University, Harelbekestraat 72,*
22 11 *9000 Ghent, Belgium; ^bDepartment of Chemistry, Latvia University of Agriculture, Liela iela 2,*
23 12 *Jelgava LV – 3001, Latvia*

24 13
25 14 **Abstract**

26 15 Different cheese samples, produced in Latvia (8) and Belgium (13), were analysed for their
27 16 sterigmatocystin (STC) content. Only two (9.5 %) of the samples were positive for STC with
28 17 concentration levels of 1.23 and 0.52 µg kg⁻¹ respectively. Five (24%) samples contained STC
29 18 above the limit of detection (0.03 µg kg⁻¹) but below the limit of quantification (0.1 µg kg⁻¹), while
30 19 five samples contained STC above LOD, but below LOQ. A sensitive liquid chromatography –
31 20 electrospray positive ionization – tandem mass spectrometry (LC-MS/MS) method, which was
32 21 previously developed for the analysis of STC in grains, was modified and applied to the analysis of
33 22 STC in cheese. This method involved sample extraction with acetonitrile:water (90:10, v/v),
34 23 defatting with n-hexane, solid phase extraction, separation on a reversed phase C₁₈ column and STC
35 24 detection by LC-MS/MS.

36 25
37 26 **Keywords:** Sterigmatocystin, mycotoxin, cheese, LC-MS/MS, *Aspergillus spp.*, *Aspergillus*
38 27 *versicolor*

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

*Correspondence: Aleksandrs Veršilovskis. E-mail: aleksandrs.versilovskis@yahoo.com

Introduction

Mould growth on cheese and, to a less extent, other fermented dairy products stored at low temperatures is a common and recurrent problem for the cheese manufacturer during aging and for the retailer and consumer during storage. Mould growth on cheese causes economic losses from discoloration, poor appearance, and off-flavours. However, of even more serious concern is that some moulds are capable of producing toxic metabolites known as mycotoxins (van Egmond 1989). Mycotoxins are toxic secondary metabolites produced by various fungal species. The presence of these substances in dairy products can have two origins: a) indirect contamination, which results from lactating animals ingesting contaminated feed and b) direct contamination, which occurs because of intentional or accidental growth of moulds on dairy products (van Egmond 1989).

Some of the fungal starters used in cheese manufacturing have been shown to produce various mycotoxins on culture media (Engel and Teuber 1989). Other moulds besides starters may be present in mould cheeses as contaminants, since various fungi are encountered in the environment of the cheese factory and some of these organisms can be potential mycotoxin producers (Lafont et al. 1990; Lund et al. 1995). Reports indicating the occurrence of mycotoxins due to contaminating fungi in mould cheeses are infrequent, in different cheese samples from Finnish supermarkets only mycophenolic acid and roquefortine C were detected in blue cheese samples (Kokkonen et al. 2005). The presence of mycotoxins in dairy products is suggested to be related more to environmental factors causing mould growth than to ingestion of mouldy feed by cows (Lafont et al. 1990).

Aflatoxin producers – *Aspergillus flavus* (*A. flavus*) and *Aspergillus parasiticus* (*A. parasiticus*) were seldom found in cheese. However, *Aspergillus versicolor* (*A. versicolor*), a species known to be capable of producing sterigmatocystin (STC), seemed to be frequently present. *A. versicolor* isolates from cheese and air in the cheese ripening room were all able to produce STC (Lund et al. 1995). STC is of interest because of its relationship to aflatoxin. It contains the same 7,8-dihydrofuro[2,3-*b*]furan moiety, which for STC is fused to a hydroxyl- and methoxy- substituted xanthone (Rodricks 1969). STC is a metabolite not only of *A. versicolor*, but also of *A. flavus*, *A. Sydowi*, *A. nidulans*, *Bipolaris*, *Chaetomium* and *Emiricella* spp. (Davis 1981; Terao 1983; Scott 1994; Scott 2004; Frisvad and Thrane 2004). Its acute toxicity, mutagenity, cytotoxicity and carcinogenity have been reviewed by Terao (1983) and it is classified as a 2B carcinogen by the International Agency for Research on Cancer (IARC 1987). The structure of the adduct of STC at the N-guanine of oligodeoxynucleotides was determined (Gopalakrishnan et al. 1992). A 'no

significant risk' level for humans of $8 \mu\text{g kg}^{-1}$ of body weight per day was estimated by the State of California (Scudamore et al. 1996). STC has been found in cheese contaminated with *A. versicolor* (Scott 1989). Lactose, glycerol, high humidity and temperature are stimulating factors in STC production by *A. versicolor* on cheese (Veringa et al. 1989). Northolt et al. (1980) reported the occurrence of STC in cheese ripening in warehouses in the Netherlands. However, low temperatures (5°C to 7°C) should prevent the growth of *A. versicolor* and production of STC. At these temperatures, the moulds that develop will be *Penicillium* species, which do not produce aflatoxins or STC (Bullerman 1981).

A number of analytical methods have been published to examine STC in cheese matrix. Thin layer chromatography (TLC) with limits of detection (LODs) around $20 \mu\text{g kg}^{-1}$. UV or postcolumn fluorescence with aluminium chloride was described for liquid chromatography (LC) detection. Aluminium chloride was also used as a spray reagent for two dimensional TLC with densitometric detection (LOD $1 \mu\text{g kg}^{-1}$ in feed) (Domagala et al. 1997). The most sensitive method for STC detection in cheese was a liquid chromatography – atmospheric pressure chemical ionization – mass spectrometry (LC-APCI-MS) method with a LOD of $2.4 \mu\text{g kg}^{-1}$ (Scudamore et al. 1996). Therefore no information is available on lower contamination levels which are also of interest.

In this study, a previously developed LC – electrospray positive ionization (ESI) – tandem mass spectrometry (MS/MS) method for the determination of STC in different grains (Versilovskis et al., 2007) (LOD $0.15 \mu\text{g kg}^{-1}$) was modified and applied to the analysis of STC in different cheese samples. This is the first described LC-ESI-MS/MS method for STC determination in cheese with solid phase extraction (SPE) and also, the first time cheese samples produced in Latvia and Belgium were checked for STC contamination.

Materials and Methods

Cheese samples

Cheese samples produced in Latvia (8) and Belgium (13) were purchased during first two month of 2008 year from different local supermarkets in Latvia and Belgium. Samples were homogenized by a food processor (Valentin, SEB, Selongey Cedex, France), and kept frozen (-20°C) before analysis.

Chemicals and reagents

Methanol (HPLC-grade), acetonitrile (HPLC-grade), n-hexane (HPLC-grade) and chloroform (HPLC-grade) were purchased from Merck (Darmstadt, Germany). Formic acid was from Fluka

(Buchs, Switzerland). Deionized water was purified with a Millipore Milli-Q Plus system (Millipore, Brussels, Belgium). Argon (Air Liquide, Belgium) was used as collision gas in mass spectrometry. STC standard was purchased from Sigma (St Louis, MO, USA). A stock solution of $500 \mu\text{g ml}^{-1}$ was prepared by dissolving 5 mg of STC in 10 ml of chloroform. An aliquot (500 μl) of the stock solution was evaporated to dryness under oxygen-free nitrogen at ambient temperature and immediately redissolved in acetonitrile (5 ml) getting a standard solution of $50 \mu\text{g ml}^{-1}$. This was further used to prepare a standard solution of $5 \mu\text{g ml}^{-1}$ of STC in acetonitrile:water (60:40, v/v). This solution was used to spike samples for recovery experiments and to prepare working standards $1.6 \mu\text{g ml}^{-1}$, $0.8 \mu\text{g ml}^{-1}$, $0.2 \mu\text{g ml}^{-1}$, $0.08 \mu\text{g ml}^{-1}$, $0.04 \mu\text{g ml}^{-1}$, $0.008 \mu\text{g ml}^{-1}$ and $0.004 \mu\text{g ml}^{-1}$ as equivalents to $200 \mu\text{g kg}^{-1}$, $100 \mu\text{g kg}^{-1}$, $25 \mu\text{g kg}^{-1}$, $10 \mu\text{g kg}^{-1}$, $5 \mu\text{g kg}^{-1}$, $1 \mu\text{g kg}^{-1}$ and $0.5 \mu\text{g kg}^{-1}$ in cheese matrix.

Sample preparation

The sample preparation was a slight modification of the method of Veršilovskis et al. (2007) for the determination of STC in grains. Ten grams of cheese were weighed and 50 ml of acetonitrile:water (90:10, v/v) and 30 ml of n-hexane were added. The sample was homogenized using a shaker (Agitelec, J. Toulemonde & Cie, Paris, France) for 30 minutes and then centrifuged (3500 rpm, 10 min, + 15°C). The acetonitrile phase was filtered through filter paper 589 (Schleicher & Schuell, Dassel, Germany). Filtered extract (10 ml) was diluted with 30 ml of water and purified using a Strata X (60 mg) SPE column (Phenomenex, Torrance, CA, USA). The SPE column was conditioned with 3.5 ml methanol followed by 3.5 ml of water prior to use. Then, 40 ml of diluted extract was loaded on to the column (using loading reservoirs of 75 ml (UCT, Bristol, PA, UK) and manifold (Waters, Milford, MA, USA)). The column was twice washed with 3.5 ml of acetonitrile:water (30:70, v/v) and STC was eluted with 3 ml of 100 % acetonitrile. The resulting eluate was evaporated to dryness under nitrogen at 37°C and re-dissolved in 250 μl mobile phase (0.02 % formic acid in acetonitrile and 0.02 % formic acid in water (60:40 v/v)). The calibrants were prepared by spiking blank matrix with the standard and prepared in the same way as the samples. Spiked cheese samples were prepared by adding 400 μl , 200 μl , 50 μl , 20 μl and 10 μl of the $5 \mu\text{g ml}^{-1}$ STC standard solution using a digital pipette to 10 g of sample, which was left for 1 h at ambient temperature with occasional agitation to allow the acetonitrile to evaporate. These volumes of standard were equivalent to levels of $200 \mu\text{g kg}^{-1}$, $100 \mu\text{g kg}^{-1}$, $25 \mu\text{g kg}^{-1}$, $10 \mu\text{g kg}^{-1}$ and $5 \mu\text{g kg}^{-1}$ of STC in cheese respectively. Eight replicates at each concentration level were prepared for recovery experiments.

LC-MS/MS analysis

The LC – MS/MS system consisted of a Waters Alliance 2695 separation module with a 100- μ l injection loop (Waters, Milford, MA, USA) coupled to a Quattro LC triple quadrupole mass spectrometer (Waters-Micromass, Manchester, UK) equipped with an electrospray interface. An electrospray ionization (ESI) probe in the positive mode was used in the analysis of STC. The mobile phase was used in isocratic regime. A Phenomenex Luna C₁₈(2) (5 μ m), 150 x 3.0 mm (Phenomenex, Torrance, CA, USA) column was used. The flow rate was 0.3 ml min⁻¹, column temperature was 30°C and the injection volume was 50 μ l. The parameters of the mass spectrometer were optimized using the STC standard. The best response was recorded with the following parameters: cone voltage 30 V, capillary voltage 3.5 kV, extractor 2 V, radio frequency (RF) lens 0.2 V, source temperature 120°C and desolvation temperature 350°C, cone gas flow 108 l h⁻¹, desolvation gas flow 660 l h⁻¹, collision energy 30 eV, dwell 0.2 sec.

For the structural identification in selected reaction monitoring (SRM) mode, the molecular ion [M+H]⁺ (m/z = 325) was fragmented within the MS to its product ions (325>310 and 325>281). Argon at a pressure of 3.5 bar was used as collision gas. A calibration curve was constructed using external standardization in matrix. The product ion with m/z = 281 was used for the quantification of STC. The ratio between peaks of STC obtained on two MRM channels (Peak area (325 > 310) / Peak area (325 > 281)) was used for confirmation of the analyte. This ratio should be 0.7 ± 0.1 for the compound to be confirmed. The MRM chromatograms of blank cheese sample (a), spiked cheese sample (0.5 μ g kg⁻¹) (b), naturally contaminated cheese sample (c) and sample containing traces of STC (d) are shown in **Figure 1**. The following validation parameters were determined: linearity, LOD, limit of quantification (LOQ), recovery and repeatability. The LOD and LOQ of STC were determined by using standards prepared in the blank matrix and calculated using “Masslynx” software integrated calculation (LOD = 3.3 x S/N-ratio; LOQ = 10 x S/N-ratio).

Results and Discussion

Method performance and validation parameters

In LC-MS methods the food matrix often causes changes in responses because matrix components can interfere with the ionization of the analytes (Tang and Kebarle 1993; Spanjer et al. 2007). Because of this matrix effect, the calibrants were prepared in blank matrix (Gouda cheese produced in Belgium), in two linear ranges from 0.5 μ g kg⁻¹ to 25 μ g kg⁻¹ and from 10 μ g kg⁻¹ to 200 μ g kg⁻¹. The method was linear for STC from 0.5 μ g kg⁻¹ to 25 μ g kg⁻¹ ($y = 2921x + 82$, $R^2 > 0.991$) and from 10 to 200 μ g kg⁻¹ ($y = 2656x + 617$, $R^2 > 0.990$). A tolerance of $\pm 10\%$ was accepted for the different calibration points for good linearity. The lowest validated level for STC in cheese was 0.5

$\mu\text{g kg}^{-1}$ (S/N = 25). The LOD and LOQ were $0.03 \mu\text{g kg}^{-1}$ and $0.1 \mu\text{g kg}^{-1}$, respectively. A good recovery was achieved for STC and the relative standard deviation (RSD, %) demonstrated acceptable repeatability (**Table 1**).

Presence of STC in cheese samples

The results on the presence of STC in cheese are shown in **Table 2**. STC above LOQ was only detected in two (15%) of the 13 Belgian cheese samples. In four (50%) of the 8 Latvian cheese samples STC was detected above LOD, but below LOQ. Five (24%) of the 21 analysed samples contained STC above LOD, but below LOQ and five (24%) samples contained traces of STC. In the past only limited studies on the occurrence and stability of STC in cheese were carried out. STC was usually detected in Dutch hard cheeses like Gouda or Edam (van Egmond et al. 1982). Northolt et al. (1980) found that nine of 39 cheeses were positive for STC. These positive samples were obtained from all different cheese categories with levels ranging from 5 to $600 \mu\text{g kg}^{-1}$. In other studies the stability of STC in cheese was checked and it was found to be stable for 3 months at various temperatures (van Egmond 1986). The occurrence, distribution and stability of STC were also studied in Ras cheese in Egypt and 35 % was found positive with a mean concentration of $22 \mu\text{g kg}^{-1}$. In Ras cheese contaminated with spores of *A. versicolor*, toxin production started after 45 days of ripening, reaching a maximum after 90 days. Aged cheese (more than 6 month) inhibited toxin production (Abd Alla et al. 1996). Migration studies of STC in Wilstermarsh cheese after inoculation with *A. nidulans* and incubation at 15°C or room temperature for 40 days was limited to 0.5 cm from the cheese surface (Engel and Teuber 1980). Similar results were obtained with six naturally contaminated Gouda cheeses on which *A. versicolor* was growing (van Egmond et al. 1982).

In our study cheese samples were different from soft (like Kamambert (camembert type) and Bouquet Des Moines) to hard (like Parito, Montenegro and Breugel). Most of hard cheeses do not contain any typical moulds producing STC, so contamination can only be possible during cheese making or cheese storage if they are stored at temperatures above 7°C . Also contamination of camembert type cheese is only possible during cheese making and storage, because *Penicillium camemberti* does not produce neither spores nor STC. But in one Latvian camembert type cheese sample (Kamambert 1) STC was detected, probably due to contamination during cheese making or during storage. Also other cheese samples contained STC traces or STC above LOD.

Our results show that a risk of contamination of cheese with STC exists. These very low concentrations will not acutely affect consumers' health. Only long-term intake of small

1
2 1 concentrations of toxins might have possible adverse effects or synergistic effects with other toxins.
3
4 2

5
6 3 **Conclusions**

7 4 This is the first simple and sensitive LC-ESI-MS/MS method for the determination of STC in
8
9 5 cheese that can be applied for routine and confirmation analyses. This is the first study of STC in
10
11 6 Latvian and Belgium cheese that indicates a low exposure rate to STC.
12
13 7

14
15 8 **Acknowledgements**

16 9
17
18 10 The study was financially supported by the ESF, National programme Name: Support in doctoral
19
20 11 and postdoctoral programme realization and investigation. Contract number:
21
22 12 VPD1/ESF/PIAA/04/NP/3.2.3.1./0001/0005/0067 and European Commision in ERASMUS
23
24 13 lifelong learning agreement within the framework of ERASMUS subprogramm and by Latvian
25
26 14 government for ERASMUS mobility programm realization:contract number: 29 – 07/08.
27
28 15
29 16

30 17 **References**

31
32 18
33 19 Abd Alla EA, Metwally MM, Mehriz AM, Abu Sree YH. 1996. Sterigmatocystin: incidence, fate
34
35 20 and production by *Aspergillus versicolor* in Ras cheese. Nahrung 40(6):310–313.
36
37 21 Bullerman LE. 1981. Public health significance of moulds and mycotoxins in fermented dairy
38
39 22 products. Journal of Dairy Science 64:2439–2452.
40
41 23 Davis ND. 1981. Sterigmatocystin and other mycotoxins produced by *Aspergillus* species. Journal of
42
43 24 Food Protection 44: 711–714.
44
45 25 Engel G, Teuber M. 1980. Formation and distribution of sterigmatocystin in cheese after
46
47 26 inoculation with *A. versicolor* and *A. nidulans*. Milchwissenschaft 35:750-756.
48
49 27 Engel G, Teuber M. 1989. Toxic metabolites from fungal cheese starter cultures (*Penicillium*
50
51 28 *camemberti* and *Penicillium roqueforti*). In: van Egmond HP, editor. Mycotoxins in dairy
52
53 29 products. Elsevier applied science: London. pp 163–192.
54
55 30 Frisvad JC, Thrane U. 2004. Mycotoxin production by common filamentous fungi.
56
57 31 Sterigmatocystin. In: Samson A, Hoekstra S, Frisvad C, editors. Introduction to food- and
58
59 32 airborne fungi. Ponsen & Looyen, Wageningen: The Netherlands. pp. 326.
60
33 Gopalakrishnan S, Liu X, Patel DJ. 1992. Solution structure of the covalent sterigmatocystin-DNA
34
35 34 adduct. Biochemistry 31(44):10790–10801.
35
36 International Agency for Research on Cancer (IARC), 1987. IARC Monographs on the evaluation

- of carcinogenic risks to humans. Summaries and Evaluations, Sterigmatocystin. Suppl. 7:72.
- Kokkonen M, Jestoi M, Rizzo A. 2005. Determination of selected mycotoxins in mould cheeses with liquid chromatography coupled to tandem with mass spectrometry. Food Additives and Contaminants 22(5):449–456.
- Lafont P, Siriwardana MG, DeBoer E. 1990. Contamination of dairy products by fungal metabolites. Journal of Environmental Pathology, Toxicology and Oncology 10(3):99–102.
- Lund F, Filtenborg O, Frisvad JC. 1995. Associated microflora of cheese. Food Microbiology 12: 173–180.
- Northolt MD, van Egmond HP, Soentoro R, Deijl E. 1980. Fungal growth and the presence of sterigmatocystin in hard cheese. Journal of Association of the Official Analytical Chemists 63:115–119.
- Rodricks JV. 1969. Fungal metabolites which contain substituted 7,8-dihydrofuro[2,3-*b*]furan (DHFF) and 2,3,7,8-tetrahydro[2,3-*b*]furans (THFF). Journal of Agricultural and Food Chemistry 17:457–61.
- Scott PM. 1989. Mycotoxigenic fungal contaminants of cheese and other dairy products. In: van Egmond HP, editor. Mycotoxins in dairy products. Elsevier Applied Science: London. pp 193–259.
- Scott PM. 1994. *Penicillium* and *Aspergillus* toxins. In: Muller JD and Trenholm HL, editors. Mycotoxins in grain. Compounds other than aflatoxins. Eaton Press, St. Paul, pp 261–285.
- Scott PM. 2004. Other mycotoxins. 17.4.1 Sterigmatocystin. In: Magan N and Olsen M, editors. Mycotoxins in food. Detection and Control. Woodhead publishing: Cornwall, England. pp 417.
- Scudamore KA, Hetmanski MT, Clarke PA, Barnes KA, Startin JR. 1996. Analytical methods for determination of sterigmatocystin in cheese, bread and corn products using HPLC with atmospheric pressure ionization mass spectrometric detection. Food Additives and Contaminants 13:343–358.
- Spanjer MC, Rensen PM, Scholten, JM. 2007. LC-MS/MS multi-method for mycotoxins after single extraction, with validation data for peanut, pistachio, wheat, maize, cornflakes, raisins and figs. Food Additives and Contaminants 1–18.
- Tang L, Kebarle P. 1993. Dependence of ion intensity in electrospray mass spectrometry on the concentration of the analytes in the electrosprayed solution. Analytical Chemistry 65:3654–3668.
- Terao K. 1983. Sterigmatocystin – a masked potent carcinogenic mycotoxins. Journal of toxicology. Toxin reviews 2:77–110.
- Van Egmond HP, Northolt MD, Paulsh WE. 1982. Distribution and stability of sterigmatocystin in hard cheese. Proceedings 4th meeting on mycotoxins in animal disease, ADAS, Weybridge,

1
2 1 Great Britain, 1 – 3 April, 1981, pp 87–89.
3
4 2 Van Egmond HP. 1989. Introduction. In: Van Egmond HP, editor. Mycotoxins in dairy products.
5 3 Elsevier applied science: London. pp 1–9.
6
7 4 Van Egmond, H.P. and Paulsch, W.E., 1986. Mycotoxins in milk and milk products. Netherlands
8 5 Milk Dairy Journal 40:175–188.
9
10 6 Veringa HA, van den Berg G, Daamen CBG. 1989. Factors affecting the growth of *Aspergillus*
11 7 *versicolor* and the production of sterigmatocystin on cheese. Netherlands Milk Dairy Journal
12 8 43:311–326.
13
14 9 Veršilovskis A, Bartkevičs V, Miķelsone V. 2007. Analytical method for the determination of
15 10 sterigmatocystin in grains using High Performance Liquid Chromatography – Tandem
16 11 Massspectrometry with Electrospray Positive Ionization. Journal of Chromatography A 1157:
17 12 467– 471.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. SRM chromatograms of blank cheese sample (a), spiked cheese sample (0.5 $\mu\text{g kg}^{-1}$) (b), naturally contaminated cheese sample (Chevre Affine – contaminated with 1.23 $\mu\text{g kg}^{-1}$) (c) and cheese sample containing traces of STC (Krievijas) (d).

Deleted: M

Table 1

Average recovery of STC from spiked cheese samples obtained during two days

Deleted: R

Added concentration, $\mu\text{g kg}^{-1}$	Found concentration (Mean $\pm$ S.D.), $\mu\text{g kg}^{-1}$	Mean recovery (n=8), %	Precision RSD, %
5	5 ± 1	97	23
10	10 ± 3	96	27
25	26 ± 7	103	25
100	104 ± 25	104	24
200	205 ± 36	103	17

Table 2

STC presence in different Latvian and Belgian cheese samples

Sample №	Sample Name	Cheese type	Produced in	Found STC concentration, $\mu\text{g kg}^{-1}$	Final decision
1	Zelta	Semi hard	Latvia	0.04	<LOQ
2	Krievijas	Semi hard	Latvia	<LOD	<LOD
3	Hollandes	Hard	Latvia	0.04	<LOQ
4	Parito	Hard	Latvia	0.07	<LOQ
5	Montenegro	Hard	Latvia	<LOD	<LOD
6	Kamambers 1	Soft	Latvia	0.03	<LOQ
7	Kamambers 2	Soft	Latvia	N.D.	<LOD
8	Trikatas Blue	Soft	Latvia	N.D.	<LOD
9	Breugel	Hard	Belgium	<LOD	<LOD
10	Brugs Broodje	Semi soft	Belgium	N.D.	<LOD
11	Brugse Blomme	Soft	Belgium	N.D.	<LOD
12	Bouquet des Moines	Soft	Belgium	N.D.	<LOD
13	Chevre Affine	Soft	Belgium	1.23	Positive
14	Chimay	Semi hard	Belgium	<LOD	<LOD
15	Orval	Semi soft	Belgium	0.03	<LOQ
16	Oud Brugge	Semi hard	Belgium	N.D.	<LOD
17	Passendale	Semi soft	Belgium	N.D.	<LOD
18	St.Paulin	Semi soft	Belgium	<LOD	<LOD
19	Nazareth	Semi hard	Belgium	N.D.	<LOD
20	Watou Special	Soft	Belgium	0.52	Positive
21	Gouda	Hard	Belgium	N.D.	<LOD

LOD = 0.03 $\mu\text{g kg}^{-1}$, LOQ = 0.1 $\mu\text{g kg}^{-1}$, N.D. – not detected

Deleted: (traces)

Deleted: (traces)

Deleted: (traces)

Deleted: (traces)

Deleted: (traces)

Deleted: 27