Residues of rotenone, azadirachtin, pyrethrins, and copper used to control *Bactrocera oleae* (Gmel.) in organic olives and oil

Vito Simeone, Nuray Baser, Donato Perrelli, Gianluigi Cesari, Hamid El Bilali, Patrizia Natale

► To cite this version:

Vito Simeone, Nuray Baser, Donato Perrelli, Gianluigi Cesari, Hamid El Bilali, et al.. Residues of rotenone, azadirachtin, pyrethrins, and copper used to control *Bactrocera oleae* (Gmel.) in organic olives and oil. Food Additives and Contaminants, 2009, 26 (04), pp.475-481. 10.1080/02652030802562938 . hal-00577348

HAL Id: hal-00577348

<https://hal.science/hal-00577348>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Residues of rotenone, azadirachtin, pyrethrins, and copper used to control *Bactrocera oleae* (Gmel.) in organic olives and oil

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2008-243.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	08-Oct-2008
Complete List of Authors:	Simeone, Vito; Agronomic Mediterranean Institute, Organic Agriculture Baser, Nuray; Agronomic Mediterranean Institute, Organic Agriculture Perrelli, Donato; Agricultural Research and Experimentation Centre "Basile Caramia", Agri-Food Cesari, Gianluigi; Agronomic Mediterranean Institute, Organic Agriculture El Bilali, Hamid; Agronomic Mediterranean Institute, Organic Agriculture Natale, Patrizia; Agricultural Research and Experimentation Centre "Basile Caramia", Agri-Food
Methods/Techniques:	Extraction, HPLC, Stripping Voltammetry
Additives/Contaminants:	Copper, Pesticide residues
Food Types:	Olive oil, Organic foods

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

**Residues of rotenone, azadirachtin, pyrethrins, and copper
used to control *Bactrocera oleae* (Gmel.) in organic olives and
oil**

V. SIMEONE¹, N. BASER¹, D. PERRELLI², G. CESARI¹, H. EL BILALI¹,
& P. NATALE²

¹*Mediterranean Agronomic Institute of Bari (C.I.H.E.A.M.-MAI.B.) – Organic
Agriculture Division - Via Ceglie, 9, 70010 Valenzano (BA);* ²*Agricultural
Research and Experimentation Centre “Basile Caramia” – Agri-food and
agro-environmental testing laboratory - via Cisterino, 281, 70010
Locorotondo (BA) simeone@iamb.it*

Abstract

Rotenone, azadirachtin, pyrethrins and copper fungicide decay curves were determined in olives and olive oil samples after experimental trials consisting of one, two, and three applications of each active ingredient were carried out twice in the years 2005 and 2006. Rotenone, azadirachtin and pyrethrins were analysed by extraction with acetonitrile and determined by liquid chromatography; copper was extracted into aqueous HCl and determined by chemical stripping. Pyrethrins were always found but in amounts below the acceptable limits, whereas rotenone and copper residues always exceeded the limits, after the pre-harvest interval, the maximum threshold allowed for olives. As regards the residues in the olive oil, rotenone was found in concentrations higher than those detected on the olives. Copper tended to accumulate in the olives according to the number of applications, whereas in the oil the residue always proved to be lower than the maximum residue limit. The sensitivity of the method applied did not allow the detection of azadirachtin.

Keywords: *organic olive growing, rotenone, azadirachtin, pyrethrins, copper, residues, olives, oil.*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Given the increasing concern about food safety, it is necessary to develop a control system of the production chain, which still displays some weaknesses, particularly for organic products. A knowledge of the post-treatment decay process for the active ingredients applied in organic agriculture on different substrates and in the various growing conditions to control pathogens is an important starting point to define proper application methods.

In the Mediterranean basin, olive growing and olive oil production play a major role in organic agriculture. However, the fungal infection by *Bactrocera oleae* (Gmelin) can seriously affect the quality of olives and therefore the quality of olive oil. Given the high incidence of infection, especially in some areas and on some varieties, appropriate control strategies are required, based also on the use of natural predators and/or products of plant origin (botanicals). The latter generally display a rapid degradation and a low toxicity risk for non-target organisms and mammals. However, their insecticidal action is sometimes of short duration (from a few hours to a few days), their cost is high and their availability on the market is limited. Furthermore, as it occurs with any plant protection product, they can entail risks for the operators and the environment. Among these natural substances, pyrethrins, rotenone, azadirachtin, quassine, nicotine ryanodine, sabadilla and some essential oils are the most used (Bagnoli and Petacchi 2004). As regards azadirachtin, pyrethrum and rotenone, little is known about their residual activity in olives and olive oil.

Azadirachtin A and its isomers act on the feeding behaviour or as growth regulators (Schmutterer 1990). Natural pyrethrins act, mainly by contact or ingestion, on the central and peripheral nervous system, thus causing immediately the insect paralysis (Casida and Quistad 1994). Pyrethrins are used against *B. oleae* as adulticide with a two-day pre-harvest interval.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rotenone is a broad-spectrum bio-insecticide (Copping 1998) which acts by contact and ingestion, inhibiting the respiratory processes. Recent studies have included rotenone among the causes of Parkinson's disease (Sherer et al. 2004).

Copper derivatives can be used in organic agriculture in accordance with the limitations set by Commission Regulation (EC) No 473/2002. The activity is exerted by the copper ion, which is mainly a fungicide, but also a repellent, an anti-oviposition elicitor and is active against the first larval stages of *B. oleae* (Girolami and Cavalloro 1972; Nuzzaci and Pizza 1995; Tzanakakis 1985).

In the present work, the post-treatment decay curves of pyrethrins, azadirachtin, rotenone and copper were studied as these substances were applied on olives in different control programmes against *B. oleae*, to assess their persistence and their transfer into the olive oil.

Materials and Methods

Trials in fields and preparation of samples

All the procedures applied to field studies were carried out under Good Laboratory Practice (GLP). The trials were carried out, in 2005 and in 2006, on two different organically cropped olive tree fields: in 2005, the field, cropped with cv. Nocellara del Belice was at the bio-farm "La Pizzica" in the area of Crispiano (Taranto, Italy); in 2006, the field cropped with cv. Coratina, was located inside the Mediterranean Agronomic Institute, in the area of Valenzano (Bari, Italy). Trees from the two cultivars were about 10 years old, and cropped, vase-trained, with a 6 x 6 (cv. Nocellara) or 8 x 8 (cv. Coratina) m planting distance. Twelve plots were identified per cultivar, each including 6 plants, which were submitted to one to three treatments with azadirachtin (Trial A), rotenone (Trial B), pyrethrum (Trial C) and copper (Trial D). Treatments were carried out with a knapsack sprayer, applying 6 or 10 L per tree, respectively in 2005 and in 2006. The product characteristics,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

dosage and pre-harvest interval are reported in Table I. The first sampling was carried out before the applications and subsequent samplings after the applications, at intervals reported in Table II.

In order to evaluate the residues on the olives, each sample consisted of about 1 kg of olives collected from the whole canopy; in contrast, to assess the residues in the oil, about 20 kg of olives, harvested following the post-harvest interval for each active ingredient, were milled on the harvest day with an electric grinder and the oil was extracted from the paste by centrifugation (15-16% average oil content) at room temperature (12-16 °C).

Extraction of the chemical residues from olives and oil

Protocols by Caboni et al. (2002) and Cabras et al. (2002) were applied to extract azadirachtin, rotenone and pyrethrins from the olives: 25 g of olives and 10 g of Na₂SO₄ were added to 25 mL of acetonitrile, stirred for 10 min, and then sonicated for 15 min. The particulate was removed by filtering. Extracts were kept at 4°C until testing. To detect the same substances in the oil, the extraction protocol by Cabras et al. (2002) was applied. Acetonitrile (10 mL) and 100 µL of oxalic acid (10% aqueous solution) were added to 1 g of oil aliquots. The mix, stirred for 30 min, was then allowed to stand for 15 min for the phase separation; the organic fraction was filtered onto regenerated cellulose microfilters, before being tested by HPLC.

To determine copper on the olives, olive aliquots (20 g) were placed in 100 mL of a 10% HCl aqueous solution. After 30 min stirring, the aqueous phase was recovered and then filtered prior to testing. To assess the presence of copper in the oil, 50 mL of n-hexane and 15 mL of 37% HCL were added to 20 g of oil. The mix was brought to light boil for 30 min and then allowed to cool to room temperature before the quantitative recovery of the lower hydrochloric acid phase. The extraction flask was washed 3 times with 10 mL of distilled water, making the mix boil lightly for 5 min; the aqueous phase was recovered and added to the hydrochloric acid phase collected earlier. The

solution, adjusted to 50 mL with distilled water, was eluted through a Supelclean NVI-CARB SP 500 mg column.

At least three different extractions were carried out on each olive and oil sample.

HPLC analysis of chemical residues

The analytical determination of azadirachtin, pyrethrins and rotenone was carried out by a Thermoquest SpectraSystem[®] Mod. 1000 (Thermo Finnigan, USA) liquid chromatograph, provided with a UV-DAD detector, model UV6000LP (TSP UV 6000 diode array detector) and a Waters X-Terra RP18 (250 x 4.6 mm, 5 μ m) column. The gradients for the pesticide separation were 80 H₂O:20 acetonitrile up to 18 min, 50 H₂O:50 acetonitrile up to 22 min, 40 H₂O: 60 acetonitrile, up to 27 min, 20 H₂O: 80 acetonitrile until the end of the run. Samples of 20 μ L were injected, elution set at 1 mL min⁻¹ flow rate and the column kept at 25°C.

LC analysis for azadirachtin, rotenone and pyrethrins was carried out at 215, 295 and 230 nm, respectively, according to the absorbance peak of the relative UV spectra. Calibration curves were plotted by the External Standard method, reporting the concentrations of increasing standard solutions as a function of the area of the relative chromatographic peaks. High correlation coefficients (0.995-1.000) were obtained between 0.08 and 5 mg kg⁻¹ for azadirachtin, 0.005 and 5.0 mg kg⁻¹ for rotenone, 0.02 and 5.0 mg kg⁻¹ for pyrethrin II and pyrethrin I. The limit of quantification (LOQ) was 0.08 mg kg⁻¹ for azadirachtin, 0.005 mg kg⁻¹ for rotenone and 0.02 mg kg⁻¹ for pyrethrins. The recovery of the active ingredient added to untreated matrices (samples were fortified with 0.5-5 mg kg⁻¹ azadirachtin, 0.02-2 mg kg⁻¹ rotenone, 0.1-2 mg kg⁻¹ pyrethrins) was between 93 and 102%. Retention times were: 18.6 min for azadirachtin, 24.6 min for rotenone, 25.7 min for pyrethrin II and 27.3 min for pyrethrin I.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Copper was determined by the standard addition method by chemical stripping by the equipment ION3 (Steroglass s.r.l, San Martino in Campo, Italy). Copper determination was carried out at a potential range of -800/-100 mV, setting the electro deposition potential at -950 mV and introducing 20 ml of 1 N HCl into the chamber, including 1 ml of Hg(II) and 1 g l⁻¹ of the extract to test. The integration interval was 370/-125 mV. For the quantitative analysis 100 µL of 5 mg kg⁻¹ standard copper were added twice and three addition measurement cycles were performed. The linearity interval was determined between 0 and 1000 µg L⁻¹ (R²>0.995). When olives or oil were fortified with 0.1-1.0 mg/kg copper, the recovery ranged between 90 and 100%.

Data management

Each evaluation of residue concentration is given as the average of the values coming from three different sample extractions ± standard deviation. Such values were converted into their natural logarithms and plotted as a function of time (in days) after applications, by first-order rate equations. Half-time decay values (DT₅₀) were obtained by these plots as ln2/k, where k represents the rate constant, and reported in the tables together with correlation coefficients (R²). .

Results

In both testing years and in the experimental conditions applied, azadirachtin was not detected in any olive or extracted olive oil, even when three applications were carried out and sampling was performed immediately after olive drying. Probably, the levels of azadirachtin residues on the olives were so low as not to be detected by HPLC analysis, as a result of the low sensitivity of the method (0.08 mg kg⁻¹) and the high decay rate of this chemical.

1
2
3
4
5 Rotenone decay and half-life in olives given one to three treatments are
6 indicated in Tables III-V. For olive samples collected for each trial 12 days
7 after the last application, the residue proved to always be much higher than the
8 admitted maximum residue level (MRL 0.05 mg kg^{-1} according to SO G.U..
9 292, issued on Dec. 14 2004). The residue of rotenone found in the olives 12
10 days after a single treatment was 0.40 mg kg^{-1} (2005) and 0.37 mg kg^{-1} (2006),
11 whereas, after three consecutive treatments, it reached 0.52 mg kg^{-1} (2005)
12 and 0.46 mg kg^{-1} (2006). In the 2006 trial including three consecutive
13 treatments, samples were collected up to 30 days after the last application;
14 even at a such long time after application the residue concentration was still
15 high (0.26 mg kg^{-1}). The analysis of the decay rates indicates that, within 5
16 days after the application, rotenone degrades quite rapidly, and then the decay
17 process goes on very slowly. In our experimental conditions, rotenone half-
18 life on the olives ($DT_{50} = 4.5\text{-}5.5$ days; $R^2 = 0.9367\text{-}0.9951$) proved to be
19 slightly higher than that found by Cabras et al. (2002). The rotenone
20 concentration found in the oil samples was always higher than that estimated
21 on the olives (Table VI). Generally, a proportionality was observed between
22 the number of treatments and the residue amount in the oil; indeed, even 15
23 days after the last treatment, rotenone residues in the oil were 16 (0.82 mg
24 kg^{-1}) to 54 (2.69 mg kg^{-1}) times higher than the MRL (0.05 mg kg^{-1}). Residues
25 in the oil were higher than those on the olives by a concentration factor
26 between 2.0 and 5.5.

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45 As regards pyrethrins, the results were expressed as the sum of pyrethrin I and
46 II concentrations. Pyrethrins decay and half -life on olive samples are
47 illustrated in Tables VII-IX. In general, a rapid decay ($DT_{50} = 1.0\text{-}1.5$ days, R^2
48 $= 0.94\text{-}0.9971$) was observed and, from the second day after each application,
49 it resulted into a residue below the admitted MRL on the olives (1 mg kg^{-1}).
50 No accumulation effect was reported on the olives and no residue was found in
51 the oil.
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The extraction technique applied quantified copper on the olive surface, regardless to the oxidation state of the metal. Results concerning copper on the olives are reported in Tables X-XII. In any trial and number of treatments tested, no decay of the copper ion was observed, even 50 days after applications. The accumulation effect was proportional to the number of applications. At the second application, the copper residue already exceeds its MRL on olives (30 mg/kg). In contrast, in both 2006 and 2005 trials, copper was detected in the oil at much lower concentrations than those detected on the olives: concentration ranges in the oil were 0.09-0.13 mg kg⁻¹ in 2005 trial and 0.03-0.17 mg kg⁻¹ in 2006 trial (Table XIII).

Discussion

The study reported here has allowed monitoring of residue rates for pyrethrins, rotenone and copper in olives and olive oil samples after different treatment procedures. Although azadirachtin was one of the chemicals used in the experiments, it was not possible to detect it on the treated olives, even when sampling was performed within 24 hours after application, probably because of the inadequate sensitivity of the method and the rapid decay of this chemical. However, the actual concentrations of azadirachtin on the treated olives must have been below the detection limit of the method (0.08 mg kg⁻¹) and the MRL for this chemical is in fact 0.5 mg kg⁻¹ (Commission Regulation (EC) No 149/2008). Thus, the quantities of azadirachtin used with the described treatments applied to olive trees would result in residue levels of no concern at any time after application. Also pyrethrins proved to be readily degradable and the residue, regardless of the matrix, was always lower than the accepted maximum limit (1 mg kg⁻¹). Therefore, the use of pyrethrins and azadirachtin, within the tested conditions, may be feasible with respect to the current needs of consumers and demands for environmental safety: given the rapid decay observed, these substances can be applied also close to harvest, if necessary.

Conversely, copper was the chemical which showed the slowest decay on the olives. Twenty days after the last application, as indicated in the label the residue level found was similar to that detected immediately after the application; moreover, the active ingredient increased on the olives according to the number of applications. Considering that copper-based products are generally applied to control plant pathogenic bacteria and fungi, and taking into account the strong accumulation effect on the olives, copper should only be used as an insecticide in the framework of an integrated management programme, especially when olives are to be marketed for fresh consumption. Indeed, in the oil, the metal was always found at concentrations lower than the accepted maximum limits. Since copper is concentrated in the olive aqueous fraction, it might be removed from the paste in the extracted water. It is necessary to validate this hypothesis above all for the appropriate management of vegetation water, which might contribute to copper ion accumulation when it is disposed in the soil.

Rotenone was found to be a very stable compound on olives. Its decay, was quite rapid in the early days after its application, decreased when the concentration on the olives reached $0.4\text{--}0.5\text{ mg kg}^{-1}$ (values 10-fold over the limit) and continued slowly, without ever going below the maximum residue limit, not even 12 days after the treatment (two days more than the post-harvest interval). For rotenone, as well, a slight accumulation effect was found in the olives and olive oil, when two or three applications were consecutively performed. The highly lipophilic molecule (Cabras et al. 2002) markedly concentrated in the oil, where it reached an amount even five times higher than that detected on the olives ($1.2\text{--}2.7\text{ mg kg}^{-1}$). The effective biological activity of rotenone makes it a key active ingredient in the control of *Bactrocera oleae*, mainly in the case of heavy infection. However, a possible involvement of rotenone in Parkinson's disease was recently reported (Sherer et al. 2004). Thus, because of the possible hazard to human health, the slow degradation of rotenone on olives has caused concern about its use in the organic cropping of olive trees, as the residue concentration never falls below the legal limits, even

when a single treatment is carried out. Thus, from April 10, 2008, the EU Commission recently removed rotenone from Annex I of the Directive 91-414-CEE, although, its use is still permitted until October 10 2009. Therefore, finding alternatives to the use of rotenone is urgent for bio-farms producing olives for the fresh market and/or olive oil. The much more rapid decay observed with pyrethrins and that it was not possible to even detect azadirachtin residues provided to olive trees in this study, taking into account the positive data on the efficacy against the olive fly, suggest that these active ingredients may represent more ecologically-sound control strategies, which may be beneficial to the consumer, the environment, the operators and the host-pest system.

Acknowledgements

The present work was carried out within the framework of the Interreg III A Project Italy-Albania 2000-2006 “Progetto finalizzato integrato per la diffusione e assistenza tecnica all’implementazione di metodologie per la produzione di prodotti biologici”; Acronym: (PAB). The authors wish to thank Prof. P. Cabras from the Toxicology Dept. of Cagliari University, and Dr S. Pollastro from the Plant Protection Dept. of Bari University, for the scientific review of this paper.

References

- Bagnoli B, Petacchi R, 2004. Problematiche relative all'utilizzo di prodotti di origine vegetale per il controllo della mosca delle olive: i prodotti di origine vegetale (*botanicals*) nella lotta ai fitofagi. http://ento.sssup.it/eventi/olivoBio/relazioni/prodotti_vegetali.htm
- Caboni P, Cabras M, Angioni A, Russo M, Cabras P. 2002. Persistence of azadirachtin residues on olives after field treatment. *Journal of Agriculture Food and Chemistry* 50: 3491-3494.

- 1
2
3
4 Caboni P, Sarais G, Angioni A, Garcia AJ, Lai F, DEdola F, Cabras P. 2006.
5 Residues and persistence of neem formulations on strawberry after field
6 treatment. Journal of Agriculture Food and Chemistry 54: 10026-10032.
7
8
9 Cabras P, Caboni P, Cabras M, Angioni A, Russo M. 2002. Rotenone residues
10 on olives and in olive oil. Journal of Agriculture Food and Chemistry 50:
11 2576-2580.
12
13
14 Casida JE, Quistad GB. 1994. Pyrethrum; a benefit to human welfare. In:
15 Casida JE, Quistad GB, editors. Pyrethrum flowers, production, chemistry,
16 toxicology and uses. Oxford, UK: Oxford University Press. pp. 345-350.
17
18
19 Copping LG. 1998. The BioPesticide Manual. British Crop Protection
20 Council, First Edition. Fernham, UK. pp. 35-36.
21
22
23 Girolami V, Cavalloro R. 1972. Aspetti della simbiosi batterica di *Dacus olaee*
24 Gmelin in natura e negli allevamenti di laboratorio. Annales de la Société
25 Entomologique de France 8: 561-571.
26
27
28 Nuzzaci G, Pizza M. 1995. Nuove metodologie di lotta antidacica.
29 Proceedings of the Symposium "Tecniche, norme e qualità in olivicoltura".
30 Potenza, Italy, December 15-17 1993. pp. 527-535.
31
32
33 Schmutterer H. 1990. Properties and potential of natural pesticides from the
34 neem tree, *Azadirachta indica*. Annual Review of Entomology 35: 271-297.
35
36
37 Sherer TB, Betarbet R, Testa CM, Seo BB, Richardson JR, Kim JH, Miller
38 GW, Yagi T, Matsuno-Yagi A, Greenamyre JT. 2004. Mechanism of
39 toxicity in rotenone models of Parkinson's disease. Journal of Neuroscience
40 23: 10756-10764.
41
42
43 Tzanakakis ME. 1985. Considerations on the possible usefulness of olive fruit
44 fly symbionticides in integrated control in olive groves. In: Cavallaro R,
45 Crovetto A, editors. Integrated Pest Control in olive-groves. Proc.
46 CEC/FAO/IOBC International Joint Meeting, Pisa, April 3-6 1984. pp 386-
47 393.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

For Peer Review Only

Table I. Characteristics of the chemicals applied to the olive trees

Trial	Active Ingredient	Commercial product			Product dosage	Pre-harvest interval (days)
		Company	Name	a.i. (%)		
A	Azadirachtin	Serbios	Diractin®	3	150 ml/hl	3
B	Rotenone	Serbios	Rotena®	6	300 ml/hl	10
C	Pyrethrins	BioIntrachem Italia	BioPirenPlus®	2	150 ml/hl	2
D	Copper	Caffaro	Cuprocaffaro®	50	400 g/hl	20

For Peer Review Only

Table II. Treatments, sampling and harvesting times in 2005 and 2006 trials

Active ingredient	Trial	Treatment schedule ^a		Sampling after each treatment (days)		Harvesting time ^a
2005						
Azadirachtin	A-I	-	-	03/12	0, 1, 2, 5, 9	07/12
	A-II	-	17/11	03/12	0, 1, 2, 5	
	A-III	01/11	17/11	03/12	0, 1, 2	
Rotenone	B-I	-	-	28/11	0, 4, 8, 12, 16	12/12
	B-II	-	12/11	28/11	0, 4, 8, 12, 16	
	B-III	27/10	12/11	28/11	0, 4, 8, 12, 16	
Pyrethrins	C-I	-	-	03/12	0, 1, 2, 3, 9	06/12
	C-II	-	17/11	03/12	0,1, 2, 3, 9	
	C-III	01/11	17/11	03/12	0,1, 2, 3	
Copper	D-I	-	-	14/11	0, 5, 21	06/12
	D-II	-	29/10	14/11	0, 5,10,15	
	D-III	13/10	29/10	14/11	0, 10	
2006						
Azadirachtin	A-I	-	-	03/11	0,1,2,3	08/11
	A-II	-	18/10	03/11	0,1, 2, 3	
	A-III	02/10	18/10	03/11	0,1, 2, 3	
Rotenone	B-I	-	-	31/10	0, 4 , 8, 12, 16, 30	15/11-05/12
	B-II	-	15/10	31/10	0, 4, 8, 12, 16, 30	
	B-III	29/09	15/10	31/10	0, 4, 8, 12, 16, 30	
Pyrethrins	C-I	-	-	03/11	0, 1, 2, 3	08/11
	C-II	-	18/10	03/11	0, 1, 2, 3	
	C-III	02/10	18/10	03/11	0, 1, 2, 3	
Copper	D-I	-	-	15/10	0, 5, 10, 15, 21	07/11
	D-II	-	29/09	15/10	0, 5, 10, 15	
	D-III	13/09	29/09	15/10	0, 5, 10, 15, 20, 29	

^adays/month

Table III. Rotenone decay on olives after one single application in 2006 and 2005 trials

Experimental Time (days)	2006 Trial	2005 Trial
0	$2,2 \pm 0,15^a$	$2,09 \pm 0,37$
4	$1,07 \pm 0,12$	$1,04 \pm 0,25$
8	$0,76 \pm 0,14$	$0,81 \pm 0,2$
12	$0,37 \pm 0,05$	$0,40 \pm 0,26$
16	$0,29 \pm 0,1$	nd
DT ₅₀	5,4	5,3
R ²	0,9767	0,9713

^aconcentration expressed as mg Kg⁻¹ ± standard deviation

nd = not determined

Table IV. Rotenone decay on olives after two consecutive applications in 2006 and 2005 trials

Experimental Time (days)	First application 2006	First application 2005	Experimental Time (days)	Second application 2006	Second application 2005
0	2,42± 0,33 ^a	2,20± 0,4	16	2,73±0,45	2,5±0,4
4	1,09±0,22	0,98±0,13	20	1,02 ±0,23	1,01 ±0,22
8	0,734± 0,12	0,71± 0,14	24	nd	0,733±0,05
12	0,39± 0,09	0,42± 0,13	28	0,42±0,07	0,52±0,07
16	0,26± 0,08	0,24± 0,11	32	0,30±0,04	nd
DT ₅₀	5,0	5,2	DT ₅₀	5,2	4,8
R ²	0,9865	0,9852	R ²	0,9595	0,9636

^aconcentration expressed as mg Kg⁻¹ ± standard deviation

nd = not determined

Table V. Rotenone decay on olives after three consecutive applications in 2006 and 2005 trials

Experimental Time (days)	First application 2006	First application 2005	Experimental Time (days)	Second application 2006	Second application 2005	Experimental Time (days)	Third application 2006	Third application 2005
0	2,56± 0,32 ^a	2,05± 0,3	16	2,74±0,21	2,40±0,3	32	3,03±0,23	2,60±0,3
4	1,0± 0,25	nd	20	1,54±0,29	1,19±0,3	36	1,83±0,32	1,33±0,32
8	0,72± 0,17	0,71± 0,2	24	0,67±0,11	0,67±0,12	40	0,87±0,14	nd
12	0,35± 0,13	0,44± 0,12	28	0,42±0,18	nd	44	0,46±0,08	0,52±0,18
16	0,23± 0,15	0,21± 0,14	32	0,29±0,12	0,32±0,14	48	0,28±0,13	0,31±0,09
DT ₅₀	4,7	5,0	DT ₅₀	4,7	5,3	DT ₅₀	4,5	5,3
R ²	0,9773	0,9934	R ²	0,9815	0,9809	R ²	0,9952	0,9956

^a concentration expressed as mg Kg⁻¹ ± standard deviation

nd = not determined

Table VI. Rotenone residues on olives at harvest and in the extracted oil in 2006 and 2005 trials

Number of applications	2005 Trial			2006 Trial			CF	Olives (30 dala)	Oil (30 dala)	CF
	Olives (12 dala)	Oil (15 dala)	CF	Olives (12 dala)	Oil (15 dala)	CF				
1	0,37±0,05 ^a	1,18±0,06	3,2	0,41±0,01	0,82±0,05	2,0	0,13±0,03	0,27±0,02	2,1	
2	0,41±0,06	2,25±0,05	5,5	0,42±0,06	2,14±0,06	5,1	0,13±0,01	0,68±0,01	5,2	
3	0,56±0,08	2,69±0,08	4,8	0,46±0,02	2,50±0,05	5,4	0,26±0,05	1,04±0,01	4,0	

^aconcentration expressed as mg Kg⁻¹ ± standard deviation
dala = days after last application; CF = concentration factor

Table VII. Pyrethrin decay on olives after one single application in 2006 and 2005 trials

Experimental Time (days)	2006 Trial	2005 Trial
0	0,75± 0,09 ^a	0,77± 0,15
1	0,33±0,04	0,38±0,13
2	0,27± 0,1	0,22± 0,1
3	0,18± 0,1	0,17± 0,08
DT ₅₀	1,7	1,5
R ²	0,9479	0,9684

^aconcentration expressed as mg Kg⁻¹ ± standard deviation

Table VIII. Pyrethrin decay on olives after two consecutive applications in 2006 and 2005 trials

Experimental Time (days)	First application 2006	First application 2005	Experimental Time (days)	Second application 2006	Second application 2005
0	1,2± 0,1 ^a	1,03± 0,11	16	0,88±0,11	0,8±0,11
1	0,62±0,1	0,57±0,09	17	0,54 ±0,05	0,65 ±0,05
2	0,46± 0,08	0,34± 0,08	18	0,36±0,03	0,29±0,03
3	0,22± 0,2	0,14± 0,02	19	0,16±0,1	0,16±0,08
DT ₅₀	1,25	1,06	DT ₅₀	1,26	1,07
R ²	0,9767	0,9854	R ²	0,9756	0,943

^aconcentration expressed as mg Kg⁻¹ ± standard deviation

Table IX. Pyrethrin decay on olives after three consecutive applications in 2006 and 2005 trials

Experimental Time (days)	First application 2006	First application 2005	Experimental Time (days)	Second application 2006	Second application 2005	Experimental Time (days)	Third application 2006	Third application 2005
0	1,2± 0,12 ^{aa}	1,3± 0,11	16	1,3±0,2	1,03±0,11	32	0,74±0,13	0,83±0,1
1	0,65± 0,13	0,63± 0,09	17	0,69±0,1	0,52±0,07	33	0,33±0,12	0,42±0,1
2	0,49± 0,11	0,45± 0,08	18	0,45±0,13	0,30±0,1	34	0,21±0,1	0,25±0,1
3	0,21± 0,11	0,26± 0,07	19	0,27±0,08	0,24±0,1	35	0,13±0,04	0,11±0,08
DT ₅₀	1,26	1,3	DT ₅₀	1,35	1,41	DT ₅₀	1,22	1,05
R ²	0,9658	0,9871	R ²	0,994	0,9586	R ²	0,9791	0,9928

^a concentration expressed as mg Kg⁻¹ ± standard deviation

Table X. Copper concentration on olives after three consecutive applications in 2006 and 2005 trials

Experimental Time (days)	First application 2006	First application 2005	Experimental Time (days)	Second application 2006	Second application 2005	Experimental Time (days)	Third application 2006	Third application 2005
0	18,4± 3,4 ^a	16,4± 3,7	15		32,3±3,4	30		51,0±3,5
5	18,5± 2,8	nd	20	31,9±3,1	30,9±3,1	35	54,9±2,3	50,7±5,7
10	18,6± 3,0	15,5± 2,4	25	30,4±2,1	32,5±4,5	40	nd	51,2±4,1
15	18,6± 2,2	nd	30	30,7±1,5	31,6±3,4	41	53,2±3,7	nd
20	18,0±2,8	nd	35	29,1±3,3	33,0±4,4	45	nd	52,6±3,4
						54	nd	49,3±2,2
						56	52,3±4,4	nd

^a concentration expressed as mg Kg⁻¹ ± standard deviation
nd = not determined

Table XI. Copper concentration on olives after two consecutive applications in 2006 and 2005 trials

Experimental Time (days)	First application 2006	First application 2005	Experimental Time (days)	Second application 2006	Second application 2005
0	17,9± 3,6 ^a	16,3± 3,2	15		33,2±2,6
5	17,4±4,3	14,9±2,4	20	34,1±4,1	33,1±4,0
10	16,6± 3,2	14,7± 3,2	25	nd	30,4±2,7
15	16,5± 2,8	14,3± 3,1	30	nd	30,9±2,1
			34	33,8±3,7	nd
			39	nd	29,7±3,0

^a concentration expressed as mg Kg⁻¹ ± standard deviation

nd = not determined

Table XII. Copper concentration on olives after one single application in 2006 and 2005 trials

Experimental Time (days)	2006 Trial	2005 Trial
0	14,6± 1,6 ^a	15,0±1,8
5	13,9±3,3	14,0±2,3
10	nd	14,4±1,1
15	nd	14,9±2,2
21	14,0± 2,5	14,4± 2,3

^aconcentration expressed as mg Kg⁻¹ ± standard deviation
nd = not determined

Table XIII. Copper concentration in the oil in 2005 and 2006 trials

Number of applications	2005 Trial	2006 Trial
3	0,075± 0,002 ^a	0,029±0,0025
2	0,029± 0,003	0,17±0,0045
1	0,011± 0,0013	0,077±0,0031

^a concentration expressed as mg Kg⁻¹ ± standard deviation