Assessment of Dietary Exposure of Nitrate and Nitrite in France

Céline Menard, Fanny Heraud, Jean-Luc J.L. Volatier, Jean-Charles Leblanc

► To cite this version:

Céline Menard, Fanny Heraud, Jean-Luc J.L. Volatier, Jean-Charles Leblanc. Assessment of Dietary Exposure of Nitrate and Nitrite in France. Food Additives and Contaminants, 2009, 25 (08), pp.971-988. 10.1080/02652030801946561 . hal-00577381

HAL Id: hal-00577381

<https://hal.science/hal-00577381>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assessment of Dietary Exposure of Nitrate and Nitrite in France

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2007-409.R1
Manuscript Type:	Review
Date Submitted by the Author:	21-Jan-2008
Complete List of Authors:	MENARD, Céline; French Food Safety Agency, Quantitative Risk Assessment Team Heraud, Fanny; French Food Safety Agency, Quantitative Risk Assessment Team Volatier, Jean-Luc; French Food Safety Agency, Quantitative Risk Assessment Team Leblanc, Jean-Charles; French Food Safety Agency, Quantitative Risk Assessment Team
Methods/Techniques:	Exposure assessment
Additives/Contaminants:	Nitrate, Nitrite
Food Types:	

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Assessment of dietary exposure of nitrate and nitrite in France

Céline MENARD *, Fanny HERAUD, Jean–Luc VOLATIER, Jean–Charles LEBLANC.

Risk Assessment Scientific Support Office, French Food Safety Agency (AFSSA), 27 – 31
avenue du Général Leclerc, 94701 MAISONS-ALFORT Cedex, FRANCE.

* Corresponding author: Céline MENARD

Phone: + 33 1 49 77 11 74

Fax: + 33 1 49 77 38 92

E-mail: c.menard@afssa.fr

Disclaimer:

The findings and conclusions in this paper are the authors' own and do not necessarily represent the views of the French Food Safety Agency. The authors declare they have no conflict of interests.

Abstract

The aim of this study was to assess the dietary exposure of nitrate and nitrite in France. 13 657 concentration levels of nitrate and nitrite measured in food, representing 138 and 109 food items respectively and coming from French monitoring programs between 2000-2006, were used. Depending on the non-detected and non-quantified analysis treatment, lower and upper concentration mean estimates were calculated for each food item. These were combined with consumption data of 1474 adults and 1018 children from the French national individual consumption survey (INCA1), conducted in 1999 and based on a seven-day food record diary. 18% of spinaches, 6% of salads, 10% of cheeses, 8% of meat products and 6% of industrial meat products exceeded the European nitrate maximum level or maximum residual level. 0.4% of industrial meat products and 0.2% of meat products exceeded their European nitrite maximum level or maximum residual level. Nitrate dietary exposure averaged 40% of the Acceptable Daily Intake (ADI, $3.7 \text{ mg kg}^{-1} \text{ bw day}^{-1}$) for adults and 51 - 54% of the ADI for children with the major contributors being, for adults and children respectively, vegetables (24 and 27% ADI), potatoes (5 and 11% ADI) and water (5 and 5% ADI). The individual nitrate dietary intake of 1.4% (Confidence Interval ($\text{CI}_{95\text{th}}$) [0.8; 2.0]) to 1.5% ($\text{CI}_{95\text{th}}$ [0.9; 2.1]) of adults and 7.9% ($\text{CI}_{95\text{th}}$ [6.2; 9.6]) to 8.4% ($\text{CI}_{95\text{th}}$ [6.7; 10.1]) of children were higher than the ADI. Nitrite dietary exposure averaged 33 - 67% of ADI ($0.06 \text{ mg kg}^{-1} \text{ bw day}^{-1}$) for adults and 67 - 133% of ADI for children, with contributions of additive food-vectors at 33% ADI for adults and 50 - 67% ADI for children. The individual nitrite dietary intake of 0.7% ($\text{CI}_{95\text{th}}$ [0.3; 1.1]) to 16.4% ($\text{CI}_{95\text{th}}$ [14.5; 18.3]) of adults and 10.5% ($\text{CI}_{95\text{th}}$ [8.6; 12.4]) to 66.2% ($\text{CI}_{95\text{th}}$ [63.3; 69.1]) of children were higher than the ADI.

Keywords: Nitrate, Nitrite, Dietary exposure, foods additives, Acceptable Daily Intake (ADI), exposure assessment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Nitrate and nitrite are ions naturally present in the environment. They are used increasingly as food additives to stabilize some meat products and cheeses. But nitrate is mainly used in agriculture as fertilizer. There are many published papers characterizing the presence of nitrate in plant foods such as lettuce and spinach for example or in water. Because of all these sources of nitrate (vegetables, water, meat products) and nitrite (food additives), humans can be exposed to these compounds, although it has been shown that nitrate is also produced endogenously. The human body absorbs the nitrate ingested very easily. However, it is not the nitrate itself that presents a health risk, but its reduction into nitrite (Hartman 1983; FAO/WHO 1998) for which a lot of studies have already shown the effects on human health. The best known health effect of nitrite or nitrate reduced into nitrite is methemoglobinemia in newborn babies, due to the fact that nitrite can prevent hemoglobin from being able to transport oxygen (Comly 1954; FAO/WHO 2002). Moreover, in 1987 Fan et al. showed that high doses of nitrite could have a toxic effect on reproduction (Fan et al. 1987). These observations have since been clearly confirmed (Fan and Steinberg 1996; Manassaram et al. 2006). Nitrite or its degradation compounds have also been shown to be mutagenic. A potential link with the emergence of chromosomal aberrations was demonstrated, as sodium nitrite would lead to an increase in the number of micronucleated polychromatic erythrocytes in vivo in mice (Luca et al. 1987). Finally, nitrite or nitrate reduced to nitrite, could interact with the amide compounds of the stomach to create N-nitroso compounds, which are very carcinogenic for animals in several target organs (Fraser 1985) and even for humans (Broitman et al. 1981). But according to the last conclusions of the Joint FAO/WHO Expert Committee on Food Additives (JECFA) (FAO/WHO 2003), nitrite does not appear to be carcinogenic and the association between the nitrate ingested and an increased risk of cancer is still debated (Van Grinsven et al. 2006; Ward et al. 2005).

Because of the effects of nitrate and nitrite on health, European standards have been established. Nitrate and nitrite could be present in food items as a natural constituent or due to an intentional addition as additive. So, European standards and this study distinguished between additive foods-vectors contributions, food items for which nitrate and nitrite uses are

1
2
3 authorized and other food items, food items for which nitrate and nitrite are natural constituents.
4 Besides, French monitoring Program distinguished 'unprepared' from 'pre-prepared' meat
5 products like chicken wings coated with bread-crumbs or ready-to-eat curried chicken, which
6 are food items obtained by specific food industry processes and called industrial prepared meat
7 products in this study. Concerning agricultural uses of nitrate, the European Commission has
8 established Maximal Levels (MLs) for lettuces (2500 - 3500 mg kg⁻¹ for summer salad and 4000
9 - 4500 mg kg⁻¹ for winter salad) and spinaches (2500 mg kg⁻¹ for summer spinach, 3000 mg kg⁻¹
10 for winter spinach, 2000 mg kg⁻¹ for frozen spinach) (Commission Regulation (EC) No
11 1881/2006). Concerning the use of nitrate as an additive, Maximal residual Level (MrLs) have
12 been established for meat products (250 mg kg⁻¹), cheeses and canned foie gras (50 mg kg⁻¹)
13 (Directive 2006/52/EC). Concerning nitrite uses, the European Commission has established
14 several MrLs (Directive 2006/52/EC) for dry cured meat products as bacon (175 mg kg⁻¹), for
15 dry non-heat-treated meat products (50 mg kg⁻¹) and for others dry cured meat products as dry
16 cured ham (100 mg kg⁻¹). For water, Directive 2003/40/EC also lays down a Maximum Limit for
17 nitrate (50 mg L⁻¹) and nitrite (0.1 mg L⁻¹).
18
19

20
21 Some studies have already estimated dietary exposure of nitrate and nitrite, which
22 generally does not exceed the ADI in the Netherlands, United Kingdom, Korea or New Zealand
23 (Lee et al. 2006; Thomson et al. 2007; Vaessen and Schothorst 1999; Ysart et al. 1999). The
24 main contributors are often potatoes, green vegetables and other vegetables at 33%, 21% and
25 15% respectively of total exposure (Ysart et al. 1999). None of these studies are comparable,
26 however, because some are conducted on the general population and others on adults only.
27 Moreover, some studies are focused on additive food-vectors only (Lee et al. 2006; Verger et al.
28 1998) and do not take account of the whole diet or the actual levels of nitrate and nitrite found in
29 food items.
30
31

32
33 Some studies are available in France. A French national duplicate study was first
34 conducted in 1990 (DGS 1992) to estimate nitrate dietary exposure. Meals eaten outside of the
35 home were analyzed and nitrate intake was in average 147.8 and 84.4 mg/day respectively for
36 adults and children. Water contributed to 22% of the total exposure, some vegetables either
37 consumed in large quantity or highly contaminated contributed to 48% of the total intake.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Concerning nitrite exposure in France, it was shown in 1998 that the Theoretical Maximum Daily Intake of nitrite was on average $0.31 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ (Verger et al. 1998). Another study was conducted in 2005 (Bemrah-Aouachria et al. 2007) to integrate nitrite uses in meat products declared by industries. For high consumers, nitrite dietary exposure was between 0.093 and $0.134 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ for adults and between 0.165 and $0.250 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ for children. These French studies are either theoretical or based on old data and present contrasting results. A new assessment of dietary exposure of nitrate and nitrite was therefore required for the French population.

This study is the first one to estimate dietary intake of nitrate and nitrite in French adults and children, based on individual consumption data and on concentrations measured in several food items and water. This paper reports the levels of nitrate and nitrite present in several French food items, available on the market between 2000 and 2006, and the level of dietary exposure of nitrate and nitrite with a lower and upper estimate, depending on the non-detected and non-quantified analyses treatment. As recommended by the Joint FAO/WHO Expert Committee on Food Additives (FAO/WHO 2003), all nitrate and nitrite food sources and not only just additive food-vectors were considered in this study. As a result, food groups and food items contributing to dietary exposure were also identified.

Methodology

The chronic dietary exposure assessment of nitrate and nitrite in France was conducted by comparing the Acceptable Daily Intake (ADI, $\text{mg kg}^{-1} \text{ bw day}^{-1}$) with the dietary exposure calculated as the sum of all concentration levels (mg kg^{-1}), multiplied by the food consumption (g day^{-1}) and divided by the individual body weight (kg bw).

Individual consumption data

French individual consumption data came from the *Enquête Individuelle et Nationale de Consommation Alimentaire* (INCA 1) conducted between August 1998 and June 1999. This French study was based on a sample of 1474 healthy adults (aged 15 and over) and 1018 children (between 3 and 14 years old), representative of the French population through

stratification (region of residence, size of urban area) and use of the quota method (age, gender, household size, individual occupation and socio-professional status) (Volatier 2000). All subjects were asked to complete a seven-day food record diary (consecutive days) as well as other questionnaires on anthropometrical and socio-economical factors. Participants could estimate portion sizes by comparing the amount they were eating with photographs compiled in a manual adapted from the SuViMax picture booklet (Deheeger 1994). Food records were subsequently coded into 895 food items (INCA classification). Adults identified as under-reporters according to the ratio of energy intake to estimated basal metabolic rate were excluded, as were those for whom no individual body weight was given. Briefly, according to Schofield (Schofield 1985; Schofield et al. 1985) and Goldberg et al (Goldberg et al. 1991), identification of the individuals' reporting status was based on fundamental principles of energy physiology. Basal metabolic rate (BMR) was first estimated using gender-age-weight specific equations computed by Schofield (1985). Dietary reporting status was then assessed as suggested by Goldberg et al. (1991) by computing a ratio of reported energy intake (EI) to estimated basal metabolic rate (BMR). Tabulated cut-off limits for this EI/BMR ratio have been derived by Goldberg et al. (1991) from whole-body calorimeter and doubly labelled water measurements in a wide range of healthy adults. Among our study sample, ratios falling below the 1.1 limit were recognized as being incompatible with long-term maintenance of energy balance and therefore with long-term survival.

Nitrate and nitrite concentration data

Food items. Concentration levels of nitrate and nitrite in foods came from the French national monitoring programs carried out between 2000 and 2006. Monitoring programs are essentially performed on individual composite samples according to sampling guidelines for official control (Commission Regulation (EC) No 1882/2006). These samples are collected on the market and often concern items which are supposed to contribute to dietary exposure. 13 657 samples representing 138 commodities for nitrate and 109 commodities for nitrite were analyzed. All concentration levels and mean estimates were compared to the Maximum Level (ML) or the Maximum residual Level (MrL) currently in force for nitrates (Commission Regulation (EC) No

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1881/2006, Directive 2006/52/EC) and for nitrites (Directive 2006/52/EC). All analytical methods were validated according to standards NF EN 12014-2 for plants and drinks, NF EN 12014-4 for nitrates in meat products, NF V 04-409 for nitrites in meat products, and the French decree of February 5, 1980 for cheese. Analytical limits of the eight French laboratories in charge of the French National Monitoring Program analysis were also obtained, Limits Of Detection (LOD) were between 0.2 and 20 mg kg⁻¹ for nitrate and between 0.03 and 5 mg kg⁻¹ for nitrite and Limits Of Quantification (LOQ) were between 0.5 and 70 mg kg⁻¹ for nitrate and 0.1 and 10 mg kg⁻¹ for nitrite.

Water. For tap water, a distribution of nitrate concentrations was obtained from the French database Sise-eaux 2005 (French Ministry of Health – DDASS-SISE-EAUX). To estimate a nitrate concentration mean in tap water, a distribution was recreated using the “Best Fit” function of @Risk, and the estimated mean was obtained. Because there is no nitrite residue in tap water due to the presence of chlorate, two scenarios were conducted: the first considering the nitrite concentration in tap water to be null and the second considering it to be the Maximum Limit (ML) fixed at 0.1 mg L⁻¹, (Directive 2003/40/EC). The nitrate and nitrite concentration means obtained for tap water were also attributed to boiled water (coffee, tea, soup). For water of source origin, data (2732 and 2721 analyses for nitrate and nitrite respectively) from the French database Sise-eaux 2004-2006 (French Ministry of Health – DDASS-SISE-EAUX) was used. For mineral and sparkling mineral water, data (957 and 460 analyses for nitrate and nitrite respectively) from the French database of the AFSSA-LERH-EMN 2004-2007 (National Mineral Water Database of the Hydrology Laboratory of French Food Safety Agency) was used.

Toxicological Reference Values

ADIs, toxicological reference values established over an entire lifetime, for nitrate and nitrite have been discussed for years. The Joint FAO/WHO Expert Committee on Food Additives (JECFA) (FAO/WHO 2003) and the Scientific Committee on Food (SCF 1995) of the European Commission conducted several assessments. The nitrate ADI used in this study was 3.7 mg kg⁻¹ bw day⁻¹ (SCF 1995, FAO/WHO 2003). For nitrite, because the difference between the

European ($0.06 \text{ mg kg}^{-1} \text{ bw day}^{-1}$) (SCF 1995) and international ($0.07 \text{ mg kg}^{-1} \text{ bw day}^{-1}$) (FAO/WHO 2003) ADI was very low, the European ADI was used in this study to be consistent with the European MRLs and MLs.

Assessment of dietary exposure

As recommended by the international guidelines (FAO/WHO 1985; FAO/WHO 2000), the use of a residue level mean when calculating dietary exposure is a realistic and appropriate estimation of chronic exposure to be compared to the Acceptable Daily Intake (ADI), a toxicological reference value established over an entire lifetime. Nitrate and nitrite concentration means were therefore calculated for each food item. When less 60% of the results were censored, i.e. below the LOD or the LOQ, means per food were calculated by setting all non-detected results to the LOD/2 and all non-quantified results to the $(\text{LOD} + \text{LOQ})/2$. When more than 60% of the results were censored, two mean estimates were calculated, as recommended by the international guidelines (FAO/WHO 1995):

- A lower bound one calculated by setting all non-detected results to 0 and all non-quantified results to the LOD,
- An upper bound one calculated by setting all non-detected results to the LOD and all non-quantified results to the LOQ.

So when more 60% of results were censored, concentration means were obtained by lower and upper bound estimates. When less 60% of results were censored, concentrations means were obtained by a middle bound estimate.

Contamination and consumption data were linked. Consumption of some specific foods analysed in the nitrite national monitoring program was not declared in the INCA survey. In fact, the INCA survey does not distinguish unprepared meat products from industrial prepared meat products obtained by specific industrial processes (coated with bread-crumbs or curried): industrial prepared meat products based on beef, boar, chicken, duck, hare, pork, poultry, quail, rabbit, turkey, veal or lamb, which are new food items less consumed than the corresponding unprepared meat product. In order to be protective and to take account of the potential contribution of these specific food items in the total intake, these industrial prepared meat

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

products were assumed to be consumed at equivalent amounts to the corresponding meat. For example, consumption of the industrial prepared poultry-based meat products was supposed to be equal to the consumption of poultry. So, estimation of nitrite dietary exposure distinguished meat products from industrial prepared meat products contributions and separated additive food-vectors from the other food items contributions..

Dietary exposure of nitrate and nitrite was expressed as a percentage of the ADI established at $3.7 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ and $0.06 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ for nitrate and nitrite respectively (SCF 1995). A risk may occur when the dietary exposure exceeds the ADI. Dietary exposure means were calculated for adults and children. The number and percentage of individual dietary exposure levels exceeding the ADI were also calculated with the respective 95% Confidence Interval (CI_{95th}), considering the normal approximation to the binomial distribution of the probability of an exposure exceeding the ADI. For nitrite, dietary exposure due to additive food-vectors (meat products and industrial meat products) and other foods (the rest of the diet) was calculated separately. Foods were identified as major contributors when the food intake exceeded 5-10% of the ADI (Codex Alimentarius Commission 2005).

Results

Mean of nitrate concentrations

Food items. Nitrate monitoring data concerns 138 foods (Table 1). Only some of these have MLs (Commission Regulation (EC) No 1881/2006), or MrLs. (Directive 2006/52/EC). Concerning salads, 6.2% of salad concentrations exceeded their ML with 23 winter lettuces (4%) and 10 other winter salads (2%) above 4000 mg kg^{-1} and 16 summer lettuces (3%), 4 summer green salads (14%) and 10 other summer salads (2%) above 2500 mg kg^{-1} . But the mean concentration was 1974 mg kg^{-1} for lettuces, 1975 mg kg^{-1} for corn salads and 1783 mg kg^{-1} for other salads. Concerning spinach, 18% of spinach concentrations exceeded their ML with 27 winter spinaches (10%) above 3000 mg kg^{-1} and 21 summer spinaches (8%) above 2500 mg kg^{-1} . But the mean concentration in spinach was 1682 mg kg^{-1} . For additive food-vectors, 6.3% of industrial prepared meat product concentrations exceeded 250 mg kg^{-1} with 2 turkey-based products (4%), 1 rabbit-based product (2%), 25 pork-based products (16%), 1

boar-based product (3%) and 1 veal-based product (5%). But the mean concentration in industrial prepared meat products was below 250 mg kg^{-1} . Concerning unprepared meat products, 311 concentrations (8%) exceeded 250 mg kg^{-1} but the mean concentration was below 250 mg kg^{-1} except in Coppa (mean = 394.2 mg kg^{-1}). The mean concentration in canned foie gras exceeded 50 mg kg^{-1} with 12% of the analysis above 50 mg kg^{-1} . Concerning cheese, 5 concentrations (10%) exceeded 50 mg kg^{-1} but the mean concentration was below 50 mg kg^{-1} , except for processed cheeses and other types of cheese (120 and 60 mg kg^{-1} respectively).

Water. Mean concentrations of nitrate obtained in water (Table 1), were, for the lower and upper bound respectively, 17.4 mg L^{-1} for tap water and boiled water, 1.6 and 2.2 mg L^{-1} for water of source origin, 2.7 and 3.0 mg L^{-1} for mineral and sparkling mineral water.

Mean nitrite concentrations

Food items. Nitrite monitoring data concerns 109 food items (Table 2). Only some of these are concerned by the food additive uses regulation (Directive 2006/52/EC). 9 meat product concentrations (0.2%) and 2 industrial prepared meat product concentrations (0.4%) exceeded 100 mg kg^{-1} but mean concentrations in these food items were below 50 mg kg^{-1} , except in Rosette (94 mg kg^{-1}).

Water. Mean concentrations of nitrite obtained in water (Table 2) were, for the lower and upper bound estimates respectively, 0.0 and 0.1 mg L^{-1} for tap water and boiled water, 0.0 and 0.03 mg L^{-1} for water of source origin and 0.00 and 0.04 mg L^{-1} for mineral and sparkling mineral water.

Nitrate dietary exposure assessment

Nitrate dietary exposure levels were almost the same with the lower and the upper bound estimates (Table 3). For adults, dietary exposure was on average $1.5 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ (40% ADI) and the 97.5th percentile was $3.3 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ (89% ADI). For children, dietary exposure was on average between 1.9 and $2.0 \text{ mg kg}^{-1} \text{ bw day}^{-1}$ (51 and 54% ADI) and the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

97.5th percentile was between 4.8 and 4.9 mg kg⁻¹ bw day⁻¹ (130 and 132% ADI). Nitrate dietary exposure levels for adults and children were on average below the nitrate ADI (3.7 mg kg⁻¹ bw day⁻¹). However, Table 4 shows that 1.4% (CI_{95th} [0.8; 2.0]) to 1.5% (CI_{95th} [0.9; 2.1]) of adults and 7.9% (CI_{95th} [6.2; 9.6]) to 8.4% (CI_{95th} [6.7; 10.1]) of children have an individual dietary exposure exceeding the ADI.

With dietary intakes exceeding 5% of the ADI for adults and children respectively (Table 3), vegetables other than potatoes (24 and 27% ADI), potatoes (5 and 11% ADI) and water including boiled water (5 and 5% ADI) were major contributors to total nitrate intake. Food items contributing to more than 5% of the ADI and considered as major contributors to nitrate dietary intakes were, for adults and children respectively (Table 5), green salads other than lettuce and corn salad (5 and 5% ADI), spinach (3 and 5% ADI) and potatoes (5 and 11% ADI). Other vegetables usually considered to be major contributors (DGS 1992), such as lettuce ,corn salad , radish , green beans or celery , were estimated to contribute to less than 5% of the ADI. Water including boiled water was a major contributor but no individual item contributed to more than 5% of the ADI.

Nitrite dietary exposure assessment

Nitrite dietary exposure levels depended largely on the lower or upper bound estimates chosen (Table 6). For adults, dietary exposure was on average between 0.02 and 0.04 mg kg⁻¹ bw day⁻¹ (33 and 67% of the ADI) and the 97.5th percentile was between 0.05 and 0.08 mg kg⁻¹ bw day⁻¹ (83 and 133% ADI). For children, dietary exposure was on average between 0.04 and 0.08 mg kg⁻¹ bw day⁻¹ (67 and 133% ADI) and the 97.5th percentile was between 0.09 and 0.17 mg kg⁻¹ bw day⁻¹ (150 and 283% ADI). Additive food-vectors contributed on average 0.02 mg kg⁻¹ bw day⁻¹ (33% ADI) for adults and between 0.03 and 0.04 mg kg⁻¹ bw day⁻¹ (50 - 67% ADI) for children. The nitrite dietary intake means for adults and children were almost the same as or higher than the ADI (0.06 mg kg⁻¹ bw day⁻¹). Table 7 shows that between 0.7% (CI_{95th} [0.3; 1.1]) and 16.4% (CI_{95th} [14.5; 18.3]) of adults and between 10.5% (CI_{95th} [8.6; 12.4]) and 66.2% (CI_{95th} [63.3; 69.1]) of children have an individual nitrite dietary intake exceeding the ADI.

Industrial prepared meat products and meat products with intakes at 17 and 17% of the ADI for adults and 33 and 17% of the ADI for children (Table 6) were major contributors (i.e. > 5% of the ADI). Dietary intakes due to vegetables and potatoes exceeded 5% of the ADI, only for upper bound estimation, with contributions equal to 17 and 33% of the ADI for vegetables and 17 and 33% of the ADI for potatoes for adults and children respectively. Major contributors to dietary intake of nitrite, i.e. food contributing to more than 5% of the ADI were, for adults and children respectively (Table 8), industrial prepared meat products made from chicken, beef, .

Discussion

Nitrate dietary exposure

The nitrate dietary exposure level estimated in this study (40% ADI for adults and 51 - 54% ADI for children) is twice as low as the results obtained in the "Diagonale des nitrates" (1.5 versus 2.5 mg kg⁻¹ bw day⁻¹ and 2.0 vs. 4.2 mg kg⁻¹ bw day⁻¹ for adults and children respectively) but is greater than in the Netherlands Duplicate Diet Survey (Vaessen and Schothorst 1999) and in the New Zealand foods and drinking water survey results (Thomson et al. 2007), where nitrate intakes for adults are 30% and 16 % of the ADI respectively. This could be explained by the fact that the Duplicate Diet Survey and the New Zealand foods and drinking water survey are exposure assessment methods which focus less on contributor items than this study method, which is based on monitoring data more or less targeted on food suspected to represent a risk.

The percentage of individual nitrate exposure exceeding the ADI in this study (1.4% Cl_{95th} [0.8; 2.0] to 1.5 Cl_{95th} [0.9; 2.1] of adults and 7.9% Cl_{95th} [6.2; 9.6] to 8.4% Cl_{95th} [6.7; 10.1] of children) is similar to the New Zealand foods and drinking water survey (Thomson et al. 2007). The levels exceeding the ADI are explained by high consumption of the major contributors to total dietary intake, which are vegetables (green salads other than lettuce and corn salad and spinach) and potatoes, consumed in large quantity by the French population. High consumers of vegetables, or vegetarians, have already been identified as subjects more exposed to nitrate (Ministry of Agriculture 1998a; FAO/WHO 2002). These contributors were already identified in France by the "Diagonale des nitrates" (DGS 1992) and other studies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(Thomson et al. 2007). Although lettuce, corn salad, radish, celery stalk, beetroot, Swiss chard and turnip, due to high contamination, and green beans, carrots, cabbage, courgette and leeks, due to high consumption, were previously identified as major contributors in the “Diagonale des nitrates”, they are not included as such in this study. Identification of green salads other than lettuce and corn salad as major contributors and not corn salad or lettuce, which are already known to contribute to nitrate dietary intake (Thomson et al. 2007), could be explained by the fact that some participants in the French INCA1 survey did not know exactly which kind of salad they ate. That is why this study does not show lettuce or corn salad as major contributors of nitrate exposure. The contributions of water (5.4 % ADI for adults and children) were obtained with an estimation of the data mean from several French monitoring databases. This nitrate concentration mean in water is considered to be representative of the expected concentration at the national level but not at the regional level. In fact, in each French region, nitrate concentrations in soil and water depend on the scale of agriculture. In French regions, some nitrate concentrations in tap water even exceed the ML. As a result, these water contributions do not take account of the local consumption of water that could contain a higher concentration of nitrate than the means used in this study. This survey may therefore give an underestimated dietary exposure for the local populations consuming and cooking with this tap water.

Nitrite dietary exposure

The nitrite dietary exposure level estimated in this study (33 - 67% ADI for adults and 67 - 133% ADI for children) is higher than in the Netherlands Duplicate Diet Survey results (Vaessen and Schothorst 1999), where the majority of studied subjects had an intake of less than 10% of the ADI, and in the New Zealand foods and drinking water survey ones (Thomson et al. 2007), where nitrite intakes were approximately 13% of the ADI.

The percentage of individuals with a nitrite exposure exceeding the ADI in this study (0.7% CI_{95th} [0.3; 1.1] to 16.4% CI_{95th} [14.5; 18.3] of adults and 10.5% CI_{95th} [8.6; 12.4] to 66.2% CI_{95th} [63.3; 69.1] of children) is also higher than the results of the New Zealand study where only 1% of the exposure scenarios exceeded the ADI, but this New Zealand study included the major contributors only and did not take account of the whole diet. Several points could explain

these exceedances of the ADI. Firstly, additive food-vectors, for which the use of nitrites as additives is allowed, are the major contributors (33% ADI for adults and 50 - 67% ADI for children). These additive food-vectors contributed on average 1 to 19 times more than the other food items. So, subjects whose nitrite intakes exceed the ADI are high consumers of meat products and industrial meat products. But the additive food-vector contributions, calculated in this study and based on concentrations measured in food, are lower than the results obtained in the first AFSSA study (Bemrah-Aouachria et al. 2007) based on nitrite uses declared by industries, where adult results were 1.2 to 2 times higher (mean at 0.034 - 0.050 mg kg⁻¹ bw day⁻¹ and P97.5th at 0.093 - 0.134 mg kg⁻¹ bw day⁻¹) and children results were 2 to 3 times higher (mean 0.058 - 0.088 mg kg⁻¹ bw day⁻¹ and P97.5th at 0.165 - 0.250 mg kg⁻¹ bw day⁻¹). The higher contributions of additive food-vectors compared to the rest of the diet could be explained by the fact that industrial meat products, items not well identified in the INCA survey and for which no consumption data was available, were assumed to be consumed in the amount equivalent to the consumption data of the corresponding meat, which was consumed in greater quantity. This assumption can overestimate the nitrite dietary exposure.

Moreover, the differences between results of nitrite exposure obtained with the lower and the upper bound estimates are considerable, in particular for vegetables and potatoes. This is due to the fact that analytical limits of the different French laboratories for non-additive food-vectors were higher than for food regulated for the use of nitrites as additives. As a result, although vegetables and potatoes were not quantified, their concentrations, depending on the lower and the upper bound estimates, have an impact on total dietary intake. Indeed vegetables without potatoes and potatoes contributions were 33.4 and 66.6% ADI for adults and children respectively with the upper bound estimate. But only, 4% of lettuce, 23% of others green salads and 8% of spinaches had detected concentration values. These food items, for which at least one concentration value was detected, contributed to nitrate dietary intake at 4.8 and 5.2% of the ADI for adults and children respectively with the high scenario. So, except lettuce, others green salads and spinach, the others vegetables and potatoes, for which 100% of concentrations values are not detected and assumed to be equal to the corresponding LOD, contributed at 28.5 and 61.4% ADI for adults and children respectively. So, the upper bound

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

estimate, using LODs for vegetables and potatoes concentration values, generally not detected, could overestimate the nitrite the dietary intake. This significant difference between analytical limits for some foods has already been identified in the Netherlands Duplicate Diet Survey (Vaessen and Schothorst 1999). Furthermore, vegetables and potatoes are consumed in large quantities by the French population (110 and 63 g day⁻¹ and 67 and 59 g day⁻¹ for adults and children respectively). So the more a food item is consumed, the greater the impact of high analytical limits for this food item is. This can explain the higher exposure of nitrite in this current assessment compared to those in the New Zealand and the Netherlands surveys, where the LOQ can be 10 times smaller than the LOQ reported in this study (Vaessen and Schothorst 1999, Thomson et al. 2007).

Study limits

Contamination data from the national monitoring programs might have been focused on food items which are known to be major contributors to dietary intakes of nitrate and nitrite. This explains why it might not cover the whole diet, because some foods are not taken into account such as cereals products, milk, butter and eggs. Therefore, according to some studies (Kammerer et al. 1992), nitrate concentrations in milk can reach 2 mg L⁻¹, with a milk consumption at 0.4 L day⁻¹ (with a body weight at 60 kg for adults and 30 kg for children), the contribution of milk can reach 0.3 and 0.7% ADI, for adults and children respectively.

Besides, the food nomenclature correspondences between contamination and consumption data were not perfect due to some foods which were analyzed but not declared to be consumed, the INCA 1 survey being not precise enough for the some foods. Moreover, monitoring programs are essentially based on food items for which nitrate and nitrite uses are allowed. So the number of samples varies considerably from one food item to another and the concentration obtained depends on the food items. This could be because some food items like additive food-vectors for nitrite and vegetables for nitrate are analyzed more frequently than other items in the diet. This important heterogeneity in the analysis number can also have an impact on the results of the dietary exposure levels obtained in this study. It is therefore

necessary to make the national monitoring program more homogeneous in terms of number of analyses and more representative in terms of food items consumed at the national level.

Another point to mention is the fact that concentrations measured are reported if the concentration is above the LOQ. But if no result is quantified it does not imply that no nitrate and / or nitrite are present in the food items. This is why two mean estimates are calculated, an upper and a lower one, using analytical limits values. But as it has been shown in this study, there were great differences between analytical limit values. Analytical limits were generally higher for food items for which nitrate or nitrite uses are not allowed. The level of the analytical limits may have a major impact on the exposure estimates that is why upper estimates could be very higher than the lower ones. Improving the sensitivity of these analytical limits, in decreasing analytical limits could refine these upper estimates obtained in this study and decrease the uncertainties on the dietary exposure assessment.

Another point to emphasize is the fact that this study does not take account of the variability of food contamination across regions and seasons. For example, some studies (Ministry of Agriculture 1998a; Vaessen and Schothorst 1999) showed dietary exposure variations according to the season, the nitrate intake mean was 73 mg day⁻¹ in spring vs. 87 mg day⁻¹ in autumn and the nitrite intake mean was 0.6 mg day⁻¹ in spring and below 0.2 mg day⁻¹ in autumn. Cooking vegetables has also been shown to reduce nitrate concentrations by up to 75% (Ministry of Agriculture 1998b). Taking seasons and cooking habits into account would consequently improve this dietary exposure assessment.

Taking account of the endogenous synthesis of nitrate and the proportion of nitrate converted to nitrite in humans, being taken as 5% (mol mol⁻¹) for normally responding individuals and 20% for those showing a high level of conversion, the endogenous synthesis of nitrite would also improve the nitrate and nitrite exposure assessment. But before this, knowledge of the nitrate and nitrite metabolisms must be improved. It would be foreseen to create a new exposure model of dietary exposure, integrating nitrate with a transformed into nitrite part and the excretion urinal rate, and nitrite with the endogenous synthesis part. Nevertheless, based on present toxicological studies, the nitrate and nitrite effects on human health are still not agreed upon, it is therefore important to keep conducting toxicological studies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

to establish more accurate ADIs and because nitrite causes methemoglobinemia, it would also be appropriate to establish an acute reference dose (FAO/WHO 2002).

Lastly, with nitrate and nitrite intakes exceeding the ADIs, the lists and the levels of the respective MLs and MrLs, essentially lower than the residue levels used in this study, might be reconsidered, as recommended by the Joint FAO/WHO Expert Committee on Food Additives (FAO/WHO 2002) and as it was confirmed by the last intake assessment of the WHO (FAO/WHO 2003).

Conclusion

This study is the first French estimate of the dietary exposure of nitrate and nitrite based on concentrations analysed in foods (13 657 analyses in foods and 6870 analyses in water). Nitrate dietary exposure is on average lower than the ADI (40% ADI for adults and 51 - 54% ADI for children) but this exposure is not only due to foods for which MLs and MrLs are fixed (lettuce and spinach). Nitrite dietary exposure is on average fairly similar to the ADI (33 - 67% ADI for adults and 67 - 133% ADI for children) and largely depends on the lower or the upper estimate of the exposure chosen to treat the non-detected and non-quantified analysis results. The major contributors to nitrite dietary exposure are only foods for which nitrite use is authorized. Nitrite dietary exposure, obtained with this approach, is 1 to 3 times lower than the theoretical nitrite dietary exposure previously obtained in France. This is why work on the methodology of dietary exposure assessment must be continued. Analytical limits for non-additive food-vectors would need to be improved in order to refine the exposure assessment. Consumption data must also be refined to take account of all contaminated food items, even those that are rarely consumed, as has been done in the second French individual consumption survey (INCA2), not yet available. Knowledge of the nitrate and nitrite metabolisms and toxicological effects must also be improved to better characterize the risk of dietary exposure of nitrate and nitrite present in food

Acknowledgements

Contamination data has been provided by the French Ministry of consumer Affairs « Direction Générale de la concurrence, de la consommation et de la Répression des Fraudes ».

References

AFSSA-LERH-EMN 2004-2007 (National database of mineral water of the hydrology Laboratory of the French Food Safety Agency).

Bemrah-Aouachria N, Leblanc JC, Volatier JL. 2007. Assessment of dietary exposure to thirteen selected food colors, preservatives, antioxidants, stabilizers, emulsifiers and sweeteners in the French population. In presse.

Broitman SA, Velez H, Vitale JJ. 1981. A possible role of iron deficiency in gastric cancer in Colombia. *Advances in Cirrhosis, hyperammonia, and hepatic encephalopathy* 135: 155-81.

Comly HH. 1954. Cyanosis in Infants caused by Nitrates in Well Water. *Journal of the American Medical Association* 129(2): 112-17.

Codex Alimentarius Commission (CAC). 2005. Procedural Manual. CCFAC Policy for Exposure Assessment of Contaminants and Toxins in food or Food Groups, fifteenth edition, pp. 114-117. Available from <http://www.fao.org/docrep/009/a0247e/a0247e04.htm>

Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs

Commission Regulation (EC) No 1882/2006 of 19 December 2006 Laying Down Methods of Sampling and Analysis for the Official Control of the Levels of Nitrates in Certain Foodstuffs.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Deheeger M, Le Moullec N, Montero P, Potier De Courcy G, Cheroubrier F, Mouget AL, Galan P, Chantrel AM, Preziosi P, Valeix P, LiGotte C, Christides JP, Rolland-Cachera MF, Conti K, Hercberg S. 1994. Portions Alimentaires: manuel photos pour l'Estimation des Quantités Suvimax. Candia Polytechnica, Paris, France.

French Ministry on Health (DGS). 1992. Diagonale des nitrates .Ed ADHEB. Not available.

European Directive 2003/40/EC of 16 May 2003 establishing the list, concentration limits and labelling requirements for the constituents of natural mineral waters and the conditions for using ozone-enriched air for the treatment of natural mineral waters and spring waters.

European Directive 2006/52/EC of 5 July 2006 amending Directive 95/2/EC on food additives other than colours and sweeteners and Directive 94/35/EC on sweeteners for use in foodstuffs.

Fan AM, Steinberg VE. 1996. Health implications of nitrate and nitrite in drinking water: an update on methemoglobinemia occurrence and reproductive and developmental toxicity. Regulatory Toxicology and Pharmacology 23 (1 Pt 1): 35-43.

Fan AM, Willhite CC, Book SA. 1987. Evaluation of the nitrate drinking water standard with reference to infant methemoglobinemia and potential reproductive toxicity. Regulatory Toxicology and Pharmacology 7(2): 135-48.

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 1985. Guidelines for the study of dietary intake of chemicals contaminants. Geneva. Available from: http://www.who.int/foodsafety/publications/chem/Guidelines_chem_contam.pdf

Formatted: English U.S.

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 1995. GEMS/Food-EURO. Reliable Evaluation of low-Level contamination of Food. Kulmbach. Federal republic Germany, p 8. Available from http://www.who.int/foodsafety/publications/chem/en/lowlevel_may1995.pdf

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 1996. WHO Food Additives series 35: Toxicological Evaluation of certain Food additives and contaminants, Forty-fourth meeting of the Joint FAO/WHO. Expert Committee on Food Additives (JECFA). Geneva. Available from <http://www.inchem.org/documents/jecfa/jecmono/v35je01.htm>

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 1998. Guidelines for Drinking-water Quality. Second edition. Geneva. pp.56-67. Available from: http://www.who.int/water_sanitation_health/dwq/2edaddvol2a.pdf

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 2000. Joint FAO/WHO Workshop on Methodology for exposure Assessment of contaminants and Toxins in Food. Geneva. Document WHO/SDE/PHE/FOS/00.5, 20p. Available from: http://www.who.int/foodsafety/publications/chem/en/exposure_june2000.pdf

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 2002. Technical Report Series. Evaluation of certain Food Additives. Fifty-ninth meeting of the joint FAO/WHO. Expert Committee on Food Additives (JECFA). Geneva.

Food and Agricultural Organization/World Health Organisation (FAO/WHO). 2003. WHO Food Additives series 50: Safety evaluation of certain food additives. Fifty-ninth meeting of the joint FAO/WHO. Expert Committee on Food Additives (JECFA). Geneva. Available from <http://www.inchem.org/documents/jecfa/jecmono/v50je01.htm>

Fraser P. 1985. Nitrates: epidemiological evidence. IARC Scientific Publications (65): 183-94.

Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, Prentice AM. 1991.

Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. European Journal of Clinical Nutrition 45(12): 569-81.

Hartman PE. 1983. Review: putative mutagens and carcinogens in foods. I. Nitrate/nitrite ingestion and gastric cancer mortality. Environmental Mutagenesis 5(1): 111-21.

Kammerer M, Pinault L, Pouliquen H. 1992. [Content of nitrate in milk. Relationship with its concentration in the water supply for livestock]. Annales des Recherches Vétérinaires 23(2): 131-8.

Lee CH, Cho YH, Park KH. 2006. Assessment of estimated daily intake of nitrite by average consumption of processed foods in Korea. Food Control 17(12): 950-956.

Formatted: German Germany

Luca D, Luca V, Cotor F, Raileanu L. 1987. In vivo and in vitro cytogenetic damage induced by sodium nitrite. Mutation Research 189(3):333-9.

Manassaram DM, Backer LC, Moll DM. 2006. A review of nitrates in drinking water: maternal exposure and adverse reproductive and developmental outcomes. Environmental Health Perspectives 114(3): 320-7.

Ministry of Agriculture, Fisheries and Food. 1998a. Duplicate diet study of vegetables_ nitrate analysis. Food Surveillance Information Sheet No.165. (London: ministry of agriculture Fisheries and Food). Available from:
<http://archive.food.gov.uk/maff/archive/food/infosheet/1998/no165/165dds.htm>

Ministry of Agriculture, Fisheries and Food. 1998b. Nitrate in vegetables. Food Surveillance Information Sheet N°158. (London: Ministry of Agriculture, Fisheries and Food).
Available from:
<http://archive.food.gov.uk/maff/archive/food/infosheet/1998/no158/158nitra.htm>

European Scientific Committee for Food (SCF) report. 1995. Opinions of the Scientific Committee for food on nitrate and nitrite.

Schofield WN. 1985. Predicting basal metabolic rate, new standards and review of previous work. Human Nutrition Clinical Nutrition 39 Suppl 1: 5-41.

Schofield RF, Kosugi K, Schumann WC, Kumaran K, Landau BR. 1985. Quantitative estimation of the pathways followed in the conversion to glycogen of glucose administered to the fasted rat. Journal of Biological Chemistry 260(15): 8777-82.

Sise-eaux database. 2005. French Ministry of Health – DDASS-SISE-EAUX.

Thomson BM, Nokes CJ, Cressey PJ. 2007. Intake and risk assessment of nitrate and nitrite from New Zealand foods and drinking water. Food Additives and Contaminants 24(2): 113-21.

Vaessen HA, Schothorst RC. 1999. The oral nitrate and nitrite intake in The Netherlands: evaluation of the results obtained by HPIC analysis of duplicate 24-hour diet samples collected in 1994. Food Additives and Contaminants 16(5): 181-8.

Van Grinsven HJ, Ward MH, Benjamin N, de Kok TM. 2006. Does the evidence about health risks associated with nitrate ingestion warrant an increase of the nitrate standard for drinking water? Environmental Health: A global Access Science Source 5: 26.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Verger P, Chambolle M, Babayou P, Le Breton S, Volatier JL. 1998. Estimation of the distribution of the maximum theoretical intake for ten additives in France. Food Additives and Contaminants 15(7): 759-66.

Volatier J-L. 2000. Enquête INCA individuelle et nationale sur les consommations alimentaires. 158 pp TEC et DOC ed, Paris, France.

Ward MH, deKok TM, Levallois P, Brender J, Gulis G, Nolan BT, VanDerslice J. 2005. Workgroup report: Drinking-water nitrate and health--recent findings and research needs. Environmental Health Perspectives 113(11): 1607-14.

Ysart G, Miller P, Barrett G, Farrington D, Lawrance P, Harrison N. 1999. Dietary exposures to nitrate in the UK. Food Additives and Contaminants 16(12): 521-32.

Table 1: Nitrate concentration data

Nitrate concentrations in food expressed in mg kg ⁻¹											
Food item	Number of analyses	% of results < LOD	Estimate of the average concentration		Maximum concentration value	Food item	Number of analyses	% of results < LOD	Estimate of the average concentration		Maximum concentration value
			Lower bound	Upper bound					Lower bound	Upper bound	
Meat products						Vegetables other than potatoes					
Andouille	36	2.8	29.8	29.8	91	Artichoke	1	0.0	16.0	16.0	16
Andouillette	35	11.4	53.3	53.3	294	Asparagus	32	12.5	111.6	111.6	1245
Bacon	10	10.0	114.9	114.9	287	Avocado	2	0.0	1256.0	1256.0	2483
Bacon bits	24	0.0	67.6	67.6	270	Beetroot	3	0.0	644.3	644.3	1141
Black pudding (blood sausage)	41	48.8	17.2	17.2	128	Broccoli	7	14.3	442.4	442.4	1131
Chipolata sausage	231	28.6	69.3	69.3	683	Brussels sprout	4	25.0	24.1	24.1	78
Chorizo dry sausage	116	4.3	164.1	164.1	685	Cabbage, green	16	6.3	498.3	498.3	1855
Cocktail sausage	14	14.3	101.6	101.6	510	Carrot	185	16.2	120.7	120.7	861
Coppa	5	0.0	394.2	394.2	860	Cauliflower	29	10.3	214.2	214.2	910
Dried sausage	142	10.6	85.3	85.3	642	Celeriac	20	10.0	612.7	612.7	1861
Dried sausage (sliced)	438	10.1	95.7	95.7	1248	Celery stalk	17	5.9	1240.8	1240.8	2500
Foie gras, canned	65	13.9	196.6	196.6	11100	Chicory (Belgian endive)	35	2.9	135.1	135.1	414
Frankfurter sausage	45	4.4	67.7	67.7	464	Corn salad	28	0.0	1975.2	1975.2	3600
French saveloy	27	22.2	69.3	69.3	342	Courgette (zucchini)	43	2.3	559.3	559.3	1198
Galantine	66	6.1	65.8	65.8	300	Cucumber	33	3.0	191.6	191.6	1800
Garlic sausage	78	5.1	80.9	80.9	550	Eggplant	23	0.0	284.4	284.4	777
Ham, raw	58	19.0	240.2	240.2	3270	Endive (US chicory)	35	0.0	534.2	534.2	2100
Ham, cooked	346	15.9	71.5	71.5	3000	Fennel	9	0.0	1042.7	1042.7	3345
Head-cheese (Brawn)	39	7.7	133.1	133.1	866	Garlic	4	25.0	120.5	120.5	286
Merguez sausage	382	8.4	160.2	160.2	9500	Green bean	40	2.5	449.0	449.0	922
Montbeliard sausage	6	0.0	199.8	199.8	345	Leek	59	8.5	409.5	409.5	2960
Mortadella	19	21.1	75.0	75.0	341	Lettuce	579	1.9	1973.9	1973.9	5600
Mortaux sausage	22	9.1	52.5	52.5	250	Mushroom	3	33.3	116.5	116.5	341
Pâté	340	16.2	62.3	62.3	2962	Mushroom, canned	6	16.7	24.2	24.2	78
Pâté in pastry.	28	14.3	88.4	88.4	213	Onion	4	75.0	52.5	77.7	210
Pâté, duck	74	25.7	22.0	22.0	412	Other green salads	403	2.7	1783.3	1783.3	6100
Pâté, fish and shellfish	5	20.0	27.0	27.0	42	Parsley	16	6.3	1980.1	1980.1	4300
Pâté, pork liver	45	17.8	91.4	91.4	773	Pea	10	60.0	21.1	21.1	79
Pâté, poultry liver	254	14.6	65.5	65.5	2236	Pepper	34	14.7	158.8	158.8	2870
Rillettes	102	41.2	10.5	10.5	88	Radish	19	0.0	1860.7	1860.7	5000
Rosette dry sausage	24	25.0	101.2	101.2	296	Sorrel	1	100.0	0.0	5.0	
Salami	16	25.0	94.0	94.0	458	Spinach	266	3.4	1681.7	1681.7	8700

<http://mc.manuscriptcentral.com/tfac> Email: fac@tandf.co.uk

Gouda cheese	2	0.0	21.5	21.5	33	Mineral water	957	32.5	2.7	3.0	/
Maasdam® type cheese	1	0.0	12.0	12.0	12	Sparkling mineral water	/		2.7	3.0	/
Mimolette ® type cheese	22	40.9	14.9	14.9	55	Tap water	/	/	17.4	17.4	/
Other cheeses	5	20.0	120.0	120.0	225	Water of source origin	2732	55,6	1.6	2.2	/
Port Salut® type cheeses	2	100.0	0.0	10.0		Boiled Water					
Processed cheese	5	0.0	60.0	60.0	280	Coffee	/	/	17.4	17.4	/
Saint-Paulin® type cheese	1	100.0	0.0	10.0		Tea	/	/	17.4	17.4	/
Appetizer cheese	1	0.0	12.0	12.0	12	Soup	/	/	17.4	17.4	/

When less than 60 % of results < LOD, upper and lower bound estimates of concentration means correspond to middle bound estimates.

Censored results correspond to analyses for which contamination levels were non-detected or non-quantified

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2 : Nitrite concentration data

Nitrite concentrations in food expressed in mg kg ⁻¹											
Food item	Number of analyses	% of results < LOD	Estimate of the average concentrations		Maximum concentration value	Commodities	Number of analyses	% of results < LOD	Estimate of the average concentrations		Maximum concentration value
			Lower bound	Upper bound					Lower bound	Upper bound	
Meat products						Cheese					
Andouille	39	5.1	13.54	13.54	76	Bonbel-Babybel ® type cheese	3	66.7	0.33	1.00	1
Andouillette	48	22.9	11.68	11.68	67	Edam cheese	4	75.0	0.75	1.50	3
Bacon	19	26.3	10.09	10.09	45	Ewe's cheese	2	100.0	0.00	0.10	
Bacon bits	40	5.0	16.65	16.65	59	Gouda cheese	1	100.0	0.00	1.00	
Black pudding (blood sausage)	49	79.6	1.04	2.49	19	Maasdam® type cheese	1	100.0	0.00	1.00	
Chipolata sausage	298	78.9	2.05	3.00	71	Mimolette ® type cheese	17	94.1	0.06	0.84	1
Chorizo dry sausage	139	35.3	3.30	3.30	16	Other cheeses	6	100.0	0.00	1.00	
Cocktail sausage	16	31.3	9.41	9.41	26	Port Salut® type cheese	2	50.0	1.25	1.25	2
Coppa	6	0.0	25.50	25.50	51	Processed cheese	5	60.0	0.70	0.70	1
Dried sausage	176	22.7	8.54	8.54	75	Saint-Paulin® type cheese	1	100.0	0.00	1.00	
Dried sausage (slices)	567	17.1	11.60	11.60	79	Small cheese for aperitif	1	100.0	0.00	1.00	
Foie gras, canned	75	26.7	4.36	4.36	12	Vegetables other than potatoes					
Frankfurter sausage	44	2.3	30.00	30.00	71	Artichoke	1	100.0	0.00	2.00	
French saveloy	45	2.2	34.01	34.01	77	Avocado	1	100.0	0.00	20.00	
Galantine	79	29.1	10.20	10.20	54	Broccoli	2	100.0	0.00	2.00	
Garlic sausage	107	2.8	29.67	29.67	76	Brussels sprout	1	100.0	0.00	2.00	
Ham, raw	73	32.9	8.77	8.77	118	Cabbage, green	7	100.0	0.00	18.57	
Ham, cooked	470	12.1	11.22	11.22	190	Carrot	84	100.0	0.00	4.15	
Head-cheese (Brawn)	51	19.6	12.29	12.29	71	Cauliflower	4	100.0	0.00	2.00	
Merguez sausage	469	72.3	3.04	3.99	220	Celeriac	3	100.0	0.00	2.00	
Montbeliard sausage	6	0.0	19.00	19.00	42	Celery stalk	7	100.0	0.00	2.00	
Mortadella	25	4.0	17.02	17.02	54	Chicory (Belgian endive)	3	100.0	0.00	2.00	
Morteaux sausage	23	4.4	5.89	5.89	40	Corn salad	23	100.0	0.00	2.00	
Pâté	428	53.7	3.44	3.44	36	Courgette (zucchini)	3	100.0	0.00	2.00	
Pâté in pastry	36	19.4	9.93	9.93	34	Cucumber	4	100.0	0.00	2.00	
Pâté, duck	101	64.4	2.70	3.43	43	Endive (US chicory)	10	100.0	0.00	7.40	
Pâté, fish and shellfish	11	90.9	0.09	1.36	1	Garlic	1	100.0	0.00	2.00	
Pâté, pork liver	67	58.2	7.65	7.65	352	Green bean	2	100.0	0.00	15.00	
Pâté, poultry liver	335	49.0	6.01	6.01	500	Leek	15	100.0	0.00	2.53	
Rillettes	136	74.3	1.65	2.60	42	Lettuce	259	99.6	0.10	6.25	25

Rosette dry sausage	28	28.6	93.79	93.79	883	Onion	2	100.0	0.00	11.00	
Salami	22	36.4	4.36	4.36	36	Other green salads	128	97.7	0.65	10.48	55
Sausage meat	66	75.8	3.35	4.53	77	Parsley	7	100.0	0.00	2.00	
Snack	20	15.0	18.23	18.23	164	Pea	2	100.0	0.00	10.00	
Strasbourg sausage	33	9.1	23.53	23.53	68	Pepper	2	100.0	0.00	2.00	
Toulouse sausage	459	74.1	2.45	3.49	76	Radish	13	100.0	0.00	3.39	
Tripes à la mode de Caen	18	77.8	1.17	1.94	14	Sorrel	1	100.0	0.00	2.00	
Vegetable terrines and mousses	1	0.0	4.00	4.00	4	Spinach	112	92.0	10.17	18.88	220
White pudding (white sausage)	72	88.9	0.15	1.10	3	Swiss chard	3	100.0	0.00	2.00	
Other meat products	23	17.4	18.00	18.00	78	Turnip	13	100.0	0.00	2.00	
Offal						Watercress	6	100.0	0.00	11.00	
Liver, poultry	12	75.0	1.08	1.83	6	Potatoes					
Tongue, beef	3	0.0	8.00	8.00	11	Potato	2	100.0	0.00	10.50	
Industrial prepared meat products						Oils					
Hare, stewed	5	60.0	3.30	3.30	11	Rapeseed oil	1	0.0	29.00	29.00	29
Other processed games	28	67.9	3.86	4.68	27	Pastas					
Other processed meat products	4	75.0	1.75	2.50	7	Pasta	4	100.0	0.00	4.00	
Pigeon, roasted	4	75.0	2.25	4.00	9	Quiches					
Processed beef	18	61.1	7.33	8.17	46	Quiche lorraine	1	100.0	0.00	5.00	
Processed boar	44	77.3	2.39	3.25	43	Shellfish and mollusks					
Processed chicken	41	29.3	15.12	15.12	43	Mussels	1	100.0	0.00	1.00	
Processed duck	45	60.0	3.77	3.77	32	Snails	6	66.7	0.67	1.33	3
Processed hare	6	33.3	7.50	7.50	20	Mixed dishes					
Processed lamb	5	20.0	5.70	5.70	15	Tomato	2	100.0	0.00	2.00	
Processed pork	171	32.8	9.75	9.75	155	Water					
Processed quail	7	85.7	0.29	1.71	2	Mineral water	460	92.6	0.00	0.04	/
Processed rabbit	63	46.0	6.15	6.15	44	Sparkling mineral water			0.00	0.04	/
Processed turkey	80	40.0	17.66	17.66	154	Tap water	/	/	0.00	0.10	/
Processed veal	27	81.5	2.22	3.04	23	Water of source origin	2721	95.1	0.00	0.03	/
Roasted pheasant	4	75.0	5.50	6.25	22	Boiled water					
						Coffee	/	/	0.00	0.10	/
						Tea	/	/	0.00	0.10	/
						Soup	/	/	0.00	0.10	/

Wen less 60 % of results < LOD, upper and lower bound estimates of concentration means correspond to middle bound estimates.
Censored results correspond to analyses for which contamination levels were non-detected or non-quantified

Table 3: Consumption and dietary exposure of nitrate

Food groups consumption (g day ⁻¹) and dietary exposure of nitrate (mg kg ⁻¹ bw day ⁻¹)										
Food Groups	Number of consumers	Consumption	Lower bound estimate				Upper bound estimate			
		Mean	Mean	Mean in % ADI	P 97.5	P 97.5 % ADI	Mean	Mean in % ADI	P 97.5	P97.5 % ADI
For Adults > 15 years (n= 1474)										
Alcoholic beverages	338	31.5	<0.05	<0.05	0.2	5.4	<0.05	<0.05	0.2	5.4
Bread	1404	117.7	0.1	2.7	0.2	5.4	0.1	2.7	0.2	5.4
Cheese	494	3.7	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Dried fruit and nuts	6	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Dry vegetables	1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Fruit	813	47.3	<0.05	<0.05	0.1	2.7	<0.05	<0.05	0.1	2.7
Industrial prepared meat products	1450	112.0	0.1	2.7	0.3	8.1	0.1	2.7	0.3	8.1
Meat products	1362	37.0	0.1	2.7	0.2	5.4	0.1	2.7	0.2	5.4
Mixed dishes	1373	11.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Soft drinks	216	7.8	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Offal	104	1.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Oil	1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Pasta	407	35.8	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	0.2	5.4
Potatoes	1132	62.6	0.2	5.4	0.5	13.5	0.2	5.4	0.5	13.5
Quiches	287	8.0	<0.05	<0.05	0.1	2.7	<0.05	<0.05	0.1	2.7
Shellfish and mollusks	17	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Starters	88	1.7	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Vegetables excluding potatoes	1437	110.8	0.9	24.3	2.6	70.3	0.9	24.3	2.6	70.3
Water	1389	556.9	0.1	2.7	0.3	8.1	0.1	2.7	0.3	8.1
Boiled Water	1363	324.6	0.1	2.7	0.3	8.1	0.1	2.7	0.3	8.1
Total		1470.1	1.5	40.5	3.3	89.2	1.5	40.5	3.3	89.2
Children between 3 and 14 years of age (n=1018)										
Alcoholic beverages	20	0.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Bread	921	58.9	0.1	2.7	0.3	8.1	0.1	2.7	0.3	8.1
Cheese	398	3.4	<0.05	<0.05	0.1	2.7	<0.05	<0.05	0.1	2.7
Dried fruit and nuts	1	0	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Dry vegetables		<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Fruits	494	22.3	<0.05	<0.05	0.2	5.4	<0.05	<0.05	0.2	5.4

Industrial prepared meat products	1011	90.9	0.1	2.7	0.5	13.5	0.2	5.4	0.5	13.5
Meat products	932	27.2	0.1	2.7	0.3	8.1	0.1	2.7	0.3	8.1
Mixed dishes	989	9.3	<0.05	<0.05	0.1	2.7	<0.05	<0.05	0.1	2.7
Soft drinks	437	22.7	<0.05	<0.05	0.1	2.7	<0.05	<0.05	0.1	2.7
Offal	44	0.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Oil		<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Pasta	246	36.7	<0.05	<0.05	<0.05	<0.05	0.1	2.7	0.3	8.1
Potatoes	881	59.1	0.4	10.8	1.2	32.4	0.4	10.8	1.2	32.4
Quiches	216	4.8	<0.05	<0.05	0.2	5.4	<0.05	<0.05	0.2	5.4
Shellfish and mollusks	10	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Starters	50	1.0	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Vegetables without potatoes	983	67.1	1.0	27.0	3.7	100.0	1.0	27.0	3.7	100.0
Water	990	477.7	0.2	5.4	0.7	18.9	0.2	5.4	0.7	18.9
Boiled Water	577	45.2	<0.05	<0.05	0.1	2.7	<0.05	<0.05	0.1	2.7
Total		927.3	1.9	51.4	4.8	129.7	2.0	54.1	4.9	132.4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 4 : Individual exposure of nitrate exceeding the ADI

Individual exposure exceeding the ADI				
	Low estimation		High estimation	
	% of people over the ADI	[95% CI]	% of people over the ADI	[95% CI]
Adults > 15 years (n=1474)	1.4	[0.8; 2.0]	1.5	[0.9; 2.1]
Children between 3 and 14 years of age (n=1018)	7.9	[6.2; 9.6]	8.4	[6.7; 10.1]

Table 5: Major food contributors to nitrate exposure

Food consumption (g day ⁻¹) and dietary exposure of nitrate (mg kg ⁻¹ bw day ⁻¹)										
	Consumption		Lower bound estimate				Upper bound estimate			
	Adults > 15 years	Children between 3 and 14 years of age	Adults > 15 years		Children between 3 and 14 years of age		Adults > 15 years		Children between 3 and 14 years of age	
			Mean	Mean % ADI	Mean	Mean % ADI	Mean	Mean % ADI	Mean	Mean % ADI
Vegetables										
Artichoke	4.3	2.0	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Asparagus	2.2	0.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Avocado	2.8	1.6	0.1	2.7	0.1	2.7	0.1	2.7	0.1	2.7
Beetroot	2.4	1.8	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Broccoli	1.9	1.3	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Brussels sprout	1.0	0.6	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Carrot	9.8	8.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Cabbage, green	0.3	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Cauliflower	5.5	3.6	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Celeriac	0.5	0.2	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Celery stalk	1.3	0.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Chicory (Belgian endive)	5.2	1.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Corn salad	1.2	0.3	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Courgette (zucchini)	3.5	1.8	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Cucumber	3.8	3.5	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Eggplant	0.6	0.3	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Endive (US chicory)	0.2	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Fennel	0.4	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Garlic	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Green bean	14.9	9.7	0.1	2.7	0.2	5.4	0.1	2.7	0.2	5.4
Leek	2.0	0.6	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Lettuce	0.7	0.3	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Mushroom	3.5	1.8	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Mushroom, canned	0.2	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Onion	0.4	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Other green salads	8.9	3.0	0.2	5.4	0.2	5.4	0.2	5.4	0.2	5.4

Parsley	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Pea	6.1	7.2	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Pepper	0.5	0.2	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Radish	2.2	1.1	0.1	2.7	0.1	2.7	0.1	2.7	0.1	2.7
Spinach	4.1	3.2	0.1	2.7	0.2	5.4	0.1	2.7	0.2	5.4
Sorrel	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Swiss chard	0.6	0.2	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Tomato	19.1	11.6	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Turnip	0.5	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Watercress	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Total	110.8	67.1	0.9	24.3	1.0	27.0	0.9	24.3	1.0	27.0
Potatoes										
Potato	62.5	59.1	0.2	5.4	0.4	10.8	0.2	5.4	0.4	10.8
Sweet potato	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Total	62.6	59.1	0.2	5.4	0.4	10.8	0.2	5.4	0.4	10.8
Water										
Mineral water	249.8	196.7	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Sparkling mineral water	17.8	15.6	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Tap water	270.6	261.0	0.1	2.7	0.2	5.4	0.1	2.7	0.2	5.4
Water of source origin	18.6	4.3	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Coffee	200.7	10.5	0.1	2.7	<0.05	<0.05	0.1	2.7	<0.05	<0.05
Tea	69.3	8.4	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Soup	54.5	26.3	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Total	881.4	522.9	0.2	5.4	0.2	5.4	0.2	5.4	0.2	5.4

Table 6: Consumption and dietary exposure of nitrite

Food groups consumption (g day ⁻¹) and dietary exposure of nitrite (mg kg ⁻¹ bw day ⁻¹)										
Food Groups	Number of consumers	Consumption	Lower bound estimate				Upper bound estimate			
		Mean	Mean	Mean in % ADI	P 97.5	P 97.5 % ADI	Mean	Mean in % ADI	P 97.5	P97.5 % ADI
For Adults > 15 years (n= 1474)										
Cheese	475	3.51	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Industrial prepared meat products *	1450	99.15	0.01	16.7	0.04	66.7	0.02	33.3	0.04	66.7
Meat products *	1363	37.11	0.01	16.7	0.02	33.3	0.01	16.7	0.02	33.3
Mixed dishes	83	2.36	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Offal	104	1.12	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Oil	1	0.00	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pasta	1132	35.80	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	0.01	16.7
Potatoes	1372	62.52	<0.005	<0.05	<0.005	<0.05	0.01	16.7	0.03	50.0
Quiches	337	6.00	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Shellfish and mollusks	160	0.99	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Vegetables excluding potatoes	1432	99.34	<0.005	<0.05	0.01	16.7	0.01	16.7	0.03	50.0
Water	1389	556.85	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Boiled Water	1363	324.55	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Total		1229.29	0.02	33.3	0.05	83.3	0.04	66.7	0.08	133.3
* Share of additive food-vectors	1474	136.25	0.02	33.3	0.04	66.7	0.02	33.3	0.05	83.3
Share of non-additive food-vectors	1474	1093.04	<0.005	<0.05	0.01	16.7	0.02	33.3	0.05	83.3
Children between 3 and 14 years of age (n=1018)										
Cheese	386	3.31	<0.005	<0.05	<0.005	<0.05	0.00	<0.05	<0.005	<0.05
Industrial prepared meat products *	1006	74.80	0.02	33.3	0.07	116.7	0.03	50.0	0.07	116.7
Meat products *	932	27.21	0.01	16.7	0.03	50.0	0.01	16.7	0.03	50.0
Mixed dishes	30	0.70	<0.005	<0.05	<0.005	<0.05	0.00	<0.05	<0.005	<0.05
Offal	44	0.46	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Oil	/	/	/	/	/	/	/	/	/	/
Pasta	881	36.69	<0.005	<0.05	<0.005	<0.05	0.01	16.7	0.02	33.3
Potatoes	988	59.10	<0.005	<0.05	<0.005	<0.05	0.02	33.3	0.06	100.0
Quiches	206	3.98	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	0.01	16.7
Shellfish and mollusks	72	0.59	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Vegetables excluding potatoes	980	61.90	<0.005	<0.05	0.01	16.7	0.02	33.3	0.05	83.3
Water	990	477.71	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Boiled Water	577	45.23	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05

Total		791.66	0.04	66.7	0.09	150.0	0.08	133.3	0.17	283.3
* Share of additive food-vectors	1018	102.01	0.03	50.0	0.08	133.3	0.04	66.7	0.09	150.0
Share of non-additive food-vectors	1018	689.65	<0.005	<0.05	0.01	16.7	0.05	83.3	0.10	166.7

Table 7: Individual exposure of nitrite exceeding the ADI

Individual exposure exceeding the ADI				
	Low estimation		High estimation	
	% of people over the ADI	[95% CI]	% of people over the ADI	[95% CI]
Adults > 15 years (n=1474)	0.7	[0.3; 1.1]	16.4	[14.5; 18.3]
Children between 3 and 14 years of age (n=1018)	10.5	[8.6; 12.4]	66.2	[63.3; 69.1]

For Peer Review Only

Table 8: Major food contributors to nitrite exposure

Food consumption (g day ⁻¹) and dietary exposure of nitrite (mg kg ⁻¹ bw day ⁻¹)										
	Consumption		Lower bound estimate				Upper bound estimation			
	Adults > 15 years	Children between 3 and 14 years of age	Adults > 15 years	Children between 3 and 14 years of age		Adults > 15 years	Children between 3 and 14 years of age			
				Mean	Mean in % ADI		Mean	Mean in % ADI		
Meat products	Mean	Mean	Mean	Mean in % ADI	Mean	Mean in % ADI	Mean	Mean in % ADI	Mean	Mean in % ADI
Andouille	0.52	0.12	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Andouillette	0.06	0.01	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Bacon	0.02	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Bacon bits	4.57	3.83	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Black pudding (blood sausage)	0.17	0.03	<0.005	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Chipolata sausage	0.33	0.44	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Chorizo dry sausage	0.59	0.12	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Cocktail sausage	0.69	0.72	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Coppa	0.38	0.01	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Dried sausage (slices)	6.30	4.52	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Dried sausage	2.89	1.65	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Foie gras, canned	0.06	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Frankfurter sausage	0.07	0.10	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
French saveloy	1.18	0.82	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Galantine	<0.005	0.01	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Garlic sausage	2.45	1.69	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Ham, cooked.	4.91	2.97	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Ham, raw.	0.39	0.10	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Head-cheese (Brawn)	0.08	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Merguez sausage	2.08	0.95	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Montbeliard sausage	0.09	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Mortadella	0.29	0.26	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Morteaux sausage	0.22	0.21	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Others meat products	0.94	0.42	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pâté	0.09	0.09	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pâté, duck	0.01	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pâté, fish and selffish	0.09	0.02	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05

Pâté in pastry.	0.10	0.01	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pâté, poultry liver	0.02	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pâté, pork liver	0.02	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Pigeon, roasted	0.39	0.03	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Rillettes	0.03	0.01	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Roasted pheasant	1.06	0.39	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Rosette dry sausage	0.10	0.27	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Salami	1.71	1.58	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Sausage meat	0.03	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Snack	0.02	0.06	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Strasbourg sausage	2.54	3.11	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Toulouse sausage	2.89	3.08	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Tripes à la mode de Caen	0.16	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Vegetable terrines and mousses	0.04	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
White pudding (white sausage)	0.00	0.01	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Total	37.11	27.21	0.01	16.7	0.01	16.7	0.01	16.7	0.01	16.7
Industrial prepared meat products										
Processed beef	31.50	29.31	<0.005	<0.05	0.01	16.7	<0.005	<0.05	0.01	16.7
Processed boar	0.07	0.03	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed chicken	22.42	15.52	0.01	16.7	0.01	16.7	0.01	16.7	0.01	16.7
Processed duck	2.35	1.51	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed hare	0.20	0.03	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed lamb	5.28	2.84	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed pork	16.82	10.28	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed quail	0.27	0.25	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed rabbit	4.67	2.24	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed turkey	5.98	5.89	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Processed veal	5.23	3.69	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Other processed game	0.14	<0.005	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Other processed meat products	2.79	2.79	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05	<0.005	<0.05
Total	99.15	74.80	0.01	16.7	0.02	33.3	0.02	33.3	0.03	50.0