Fusarium toxins and total fungal biomass indicators in naturally contaminated wheat samples from north-eastern Poland in 2003

Juliusz Perkowski, Marian Wiwart, Maciej Buśko, Magdalena Laskowska,
Franz Berthiller, Wolfgang Kandler, Rudolf Krska

► To cite this version:

Juliusz Perkowski, Marian Wiwart, Maciej Buśko, Magdalena Laskowska, Franz Berthiller, et al.. Fusarium toxins and total fungal biomass indicators in naturally contaminated wheat samples from north-eastern Poland in 2003. Food Additives and Contaminants, 2007, 24 (11), pp.1292-1298. 10.1080/02652030701416566 . hal-00577463

HAL Id: hal-00577463

<https://hal.science/hal-00577463>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fusarium toxins and total fungal biomass indicators in naturally contaminated wheat samples from north-eastern Poland in 2003

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2006-367.R1
Manuscript Type:	Original Research Paper
Methods/Techniques:	Chromatography - GC/MS, Chromatography - LC/MS
Additives/Contaminants:	Ergosterol, ATP, Mycotoxins - trichothecenes, Volatiles
Food Types:	Wheat, Cereals and grain

SCHOLARONE™
Manuscripts

***Fusarium* toxins and total fungal biomass indicators in naturally contaminated wheat samples from north-eastern Poland in 2003**

Abstract

Concentrations of fungal metabolites were measured in 32 wheat grain samples from north-eastern Poland in 2003. The samples originated from fields cultivated conventionally (but varying in chemical protection level) or cultivated organically. Concentrations of *Fusarium* toxins (HT-2, DON, 3-AcDON, NIV), trichodiene, microbial biomass indicators [fungal ergosterol and general adenosine-5'-triphosphate (ATP)] and seed vigour were assessed. A large variation between samples was observed, depending on their origin. Seed from organic farms contained similar amounts of *Fusarium* toxins but more ergosterol and ATP than conventionally grown and chemically protected seed. The highest levels of toxins and ergosterol were detected in samples from conventional cultivation lacking chemical protection. Intensive agronomic practices (including complete chemical protection) significantly lowered the levels of ergosterol, ATP and trichodiene, as compared with other cultivation systems.

Keywords: wheat, cultivation systems, trichothecenes, ergosterol, trichodiene, ATP

Introduction

In Poland, as in other EU countries, various fungal diseases affect wheat, but special attention is paid to *Fusarium* head blight (FHB). In the literature there is a shortage of convincing data about the influence of cultivation systems on the severity of its disease symptoms (Teich 1989, Champeil et al. 2004, Lemmens et al. 2004, Perkowski et al. 2004). In north-eastern Poland, organic farming is particularly important because of the low level of environmental contamination and the possibility of growing healthy food (Grodzińska et al. 1999, Łukanowski and Sadowski 2002). Literature data reporting contamination by fungal metabolites of organically grown cereals are contradictory (Elmholt and Labouriau 2005, Pussemier et al. 2006). However, some authors point to the significant influence of

environmental conditions (weather in particular) on their concentrations. Malmauret et al. (2002) reported that concentrations of *Fusarium* toxins were higher in wheat and barley seed from organic farms, but their results showed no conclusive evidence whether conventional products were more or less safe than organic ones. Apart from *Fusarium* toxins, also fungal biomass in seed is important. Fungal biomass is usually estimated on the basis of ergosterol or chitin content (Abramson et al. 1998, Ekblad et al. 1998). Also ATP level is measured on the surface of kernels, because it is correlated with levels of some other fungal metabolites (Williams 1989), but it reflects the level of the whole microbial biomass. Seed colonization by fungi often affects its vigour and consequently also sowing value. Infection by pathogens of the genus *Fusarium* often disturbs or even completely inhibits seed germination (Eudes et al. 1997, Packa and Koczowska 1987).

This study was aimed to compare concentrations of fungal metabolites in wheat seed from conventional farms (with more or less intensive use of chemical protection) and organic farms. The comparison involved a complex of fungal metabolites: concentrations of selected *Fusarium* toxins (deoxynivalenol, 3-acetyldeoxynivalenol, the sum of 3-acetyldeoxynivalenol and 15-acetyldeoxynivalenol, nivalenol, fusarenon-X, diacetoxyscirpenol, HT-2 toxin, T-2 toxin and zearalenone), trichodiene (their precursor), and microbial biomass indicators (fungal ergosterol and general microbial ATP). Differences in the studied characters between four cultivation systems and correlations between concentrations of the studied metabolites were analysed.

Material and methods

Samples

The analysis involved 32 grain samples (of 1000 g each) of winter and spring wheat from conventional farms (20 samples) and organic farms (12 samples) located in 7 districts in north-western Poland (Figure 1). Conventional farms were divided into 3 groups: CON I, CON II and CON III. In CON I (9 samples from 6 districts), complete antifungal protection and pesticides were applied. CON II (6 samples from 6 districts) included large fields (over 30 ha) subject to a lower level of chemical protection (fungicidal protection against main foliar diseases, first of all mildew and leaf septorioses), while CON III (5 samples from one district Ostróda) was an experimental farm Bałcyny, where neither fungicides nor pesticides were used. Production in organic farms conformed to the rules applied in this cultivation

system (the seed had the Polish ECO certificate). Production in conventional farms followed the rules of the currently recommended intensive methods of cereal cultivation.

Determination of major A- and B-trichothecenes and zearalenone by LC-MS

Deoxynivalenol (DON), 3-acetyldeoxynivalenol (3-AcDON), the sum of 3-acetyldeoxynivalenol and 15-acetyldeoxynivalenol (AcDONs), nivalenol (NIV), fusarenon-X (FUS-X), diacetoxyscirpenol (DAS), HT-2 toxin, T-2 toxin and zearalenone (ZON) were determined by liquid chromatography – triple quadrupole mass spectrometry as described previously by (Berthiller et al. 2005). 10 g of the ground samples were extracted with 40 mL acetonitrile/water (84/16, v/v) and purified using MycoSep 227 columns (RomerLabs, Tulln, Austria). 4 mL of the purified extracts were evaporated to dryness and dissolved in mobile phase (methanol/water 20/80, v/v containing 5 mM ammonium acetate). An Agilent (Waldbronn, Germany) 1100 HPLC system was used in conjunction to a QTrap MS/MS from Applied Biosystems (Foster City, USA). Atmospheric pressure chemical ionisation was used in negative mode for B-trichothecenes and in positive mode for A-trichothecenes. 25 µl were injected onto a Thermo Aquasil RP-C18 column (4.6 x 100mm, 3µm), thermostated to 30°C. The multiple reaction monitoring mode was used for quantification of the samples. This method was carefully optimized for the quantification of selected Fusarium mycotoxins in cereal. Its limits of quantification (LOQs) are 3 µg/kg DON, 13 µg/kg 3-AcDON, 4 µg/kg AcDONs, 18 µg/kg NIV, 6 µg/kg FUS-X, 1 µg/kg DAS, 4 µg/kg HT-2, 1 µg/kg T-2 and 3 µg/kg ZON. The limits of detection (LODs) were about 3 times lower than the LOQs.

Determination of ergosterol

Samples containing 100 mg of ground grains were placed into 17 mL culture tubes, suspended in 2 mL of methanol, treated with 0.5 ml of 2 M aqueous sodium hydroxide and tightly sealed. Then the culture tubes were placed within 250 mL plastic bottles, tightly sealed and placed inside the microwave oven (Model AVM 401/1WH, Whirlpool, Sweden) operating at 2450 MHz and 900 W maximum output. Samples were irradiated (370 W) for 20 s. and after ca. 5 min. an additional 20 s. After 15 min contents of culture tubes were neutralized with 1 M aqueous hydrochloric acid, 2 mL MeOH were added and extraction with pentane (3 x 4 mL) were carried out within the culture tubes. The combined pentane extracts were evaporated to dryness in a nitrogen stream. Prior to analysis samples were dissolved in 4 mL of MeOH, filtered through 13 mm syringe filters with 0.5 µm pore diameter (Fluoropore

Membrane Filters, Millipore, Ireland) and evaporated to dryness into N₂ stream. The sample extract was dissolved in 1 mL of MeOH and 50 µL was analysed by HPLC technique. Separation was achieved on 150 x 3.9 mm Nova Pak C-18, 4 µm column and eluted with methanol/acetonitrile (90:10) at a flow rate of 0.6 mL /min. Ergosterol was detected with Waters 486 Tunable Absorbance Detector set at 282 nm. Quantification of ERG was made by a comparison of peak areas with those of external standards of ERG. Confirmation of ERG was achieved by a comparison of retention times with the external standard and by co-injection every tenth sample with an ergosterol standard.

Determination of trichodiene

Volatile metabolites were extracted from grain using solid phase microextraction method (SPME). Grain samples (8 g) were placed in 20 mL vials and extracted by means of headspace SPME during 30 min at 40°C with 100 µm polydimethylsiloxane (PDMS) fibre. The analyses were run on a gas chromatograph mass spectrometer (Hewlett Packard GC 6890, MS 5972 A), using an HP – 5MS (0.25 mm x 30 m) capillary column. The injection port temperature was 250°C, the transfer line temperature was 280°C and the analyses were performed with programmed temperature: initial temperature 40°C hold 3 min, from 40°C - 200°C at 10°C/min, 200°C - 280°C at 20°C/min, the final temperature being maintained for 3 min. The helium flow rate was held constant at 0.7 mL/min. Quantitative analysis of trichodiene was performed in selected ion monitoring (SIM) mode, using following ions for trichodiene detection: m/z 67, 93, 108, 109. The amount of trichodiene was estimated comparing the area of its total ion current (TIC) peak with internal standard (tridecane, 25 ng in pentane) and expressed as their ratio (RU). Trichodiene was identified by comparing its mass spectrum with literature data (Jeleń et al. 1995).

Determination of ATP

Adenosine-5'- triphosphate (ATP) measurement was carried out using luciferase enzyme. ATP was extracted from 1.00 g grain samples using 8 mL boiling Tris buffer (0.1 molar solution, pH – 7.75). 100 µL of this extract were mixed with 100 µL reagent containing luciferin and luciferase (LuminATE (QM), Celsis, Netherlands). The emitted light was quantified by Lumac Biocounter M 1500 luminometer and readout as Relative Light Units (RLU). The emitted light correlates with the amount of ATP in the sample.

Seed vigour test

The test was performed in two replicates, according to the guidelines of the International Seed Testing Association (Hampton and Tekrony 1995). Subsamples of 50 surface-sterilized kernels each were placed on strips of filter paper (50 cm × 25 cm). Next the strips were rolled up and deposited in beakers containing a small amount of sterile distilled water and kept in a phytotron in darkness at 22°C. After 7 days, proportions of non-germinated and abnormally germinated grains and of grains with *Fusarium* mycelium were assessed.

Statistical analyses

The significance of differences in mean values was evaluated with the use of Student's *t*-test, and linear correlation analysis was made for all studied parameters.

Results and discussion

The wheat grain samples classified into 4 different groups (CON I, CON II, CON III and ECO) markedly differed in the analysed parameters: HT-2 toxin, DON, 3-AcDON, NIV, sum of the most important group B trichothecenes in grain (DON+NIV together), ERG, TRICH, ATP and seed vigour parameters (non-germinated and abnormally germinated grains as well as grains with *Fusarium* mycelium). The results are presented in Tables 1 and 2 (except of ZEA concentrations because their were below the limit of detection).

The grains were characterized by a low proportion of infestation by fungi of the genus *Fusarium*, ranging from 0 to 4.0%, but differences between the groups were not significant. The proportion of grains with *Fusarium* mycelium was positively correlated with the proportion of abnormally germinated grains, although value of Pearson correlation coefficient was not significant, because during vegetation *Fusarium* is not the only factor influencing abnormal grain germination, but also harvest conditions, kernel damage as well as preinfestation changes. Additionally, since these were productive farms, an accurate determination of the above mentioned factors was difficult. A significant correlation was found between the proportion of non-germinated grains and the proportion of abnormally germinated grains. Mean values of both these parameters were the lowest in CON I, and the highest in ECO.

The mean concentration of trichothecene toxins in CON I and CON II can be regarded as low, and it is comparable with results reported by (Pettersson 1996, Perkowski et al. 1997). In those groups, NIV+DON concentration reached $43.0 \mu\text{g kg}^{-1}$ (CON I) and $30.0 \mu\text{g kg}^{-1}$ (CON II). In organically grown wheat, the NIV+DON level was similar, but the concentration of DON was higher than the concentration of NIV. In comparison with CON I, CON II and ECO, significant differences were found in CON III (conventional but lacking chemical protection), where NIV+DON content was over 14 times higher than the mean for the other groups ($635.2 \mu\text{g kg}^{-1}$ versus $44.3 \mu\text{g kg}^{-1}$), and the major contributor was DON.

Concentrations of trichodiene (TRICH), which is a precursor of trichothecenes (Jeleń et al. 1995), were the highest in samples containing the highest levels of trichothecenes (CON III), while the lowest in organically grown wheat seed. A positive correlation ($r=0.788$, $P=0.05$) was observed between TRICH level and concentration of NIV+DON in seed.

In this study, with the use of so wide-ranging wheat materials, concentrations of ERG (ergosterol, indicator of fungal biomass) and ATP (indicator of microbial biomass in the outer parts of the grain) were measured. To be exact, Reid et al. (1999) examined disease severity, ergosterol content, fungal DNA content and mycotoxin (DON and fumonisin B₁) concentration in silk maize ears channel-inoculated with *F. graminearum* and *F. moniliforme*. The authors reported that although biomass may be the best predictor of disease ratings, the DNA analysis demonstrated that this biomass was not always produced by the expected *Fusarium* sp. However, the above experiments were carried out for maize. Similar conclusions were indicated by Marfleet et al. (1991) as well.

The highest ERG level was 2.39 mg/kg (CON III), but it reached a similar value in ECO. These values were significantly higher than in CON I and CON II, but still were below the threshold of 3 mg/kg reported by Schnürer (Schnürer and Jonsson 1992) for seed showing no symptoms of fungal colonization. In CON I, the lowest ERG level was recorded, and ATP content was also low (2153 RLU). It is interesting that such a low ATP concentration was recorded in all analysed samples of this group. This value was about 8-fold lower than in organically grown wheat seed (16128 RLU). It is probably a result of the lack of chemical protection in organic farming, which is favourable for microbial growth in the outer parts of the grain. The lack of chemical protection and high mineral fertilization combined with all other agronomic practices used in conventional cultivation in CON III could favour fungal

growth at the expense of other microorganisms. This was reflected in a high ERG and mycotoxins concentration and relatively low ATP content. Because of the varied cultivation systems, only a low positive correlation between ERG and ATP content was found when all four groups were analysed jointly. When the groups were analysed separately, the highest correlation between these characters ($r=0.726$, significant at $P<0.01$) was recorded in CON I, which seems to be the most common in that part of Poland.

In conclusion, results of this study show that wheat seed in north-eastern Poland in 2003 was characterized by a low level of colonization by fungi of the genus *Fusarium*, but considerable differences were observed depending on cultivation system. Seed samples from organic farms contained similar amounts of *Fusarium* toxins but higher levels of ERG and ATP than conventionally grown and chemically protected seed samples did. A slightly higher DON content in ECO samples in comparison to CON II samples, and especially CON I samples, and a very low NIV content suggest that while observing the principles of organic farming, especially appropriate crop rotation, the absence of chemical protection does not necessarily have to lead to increased concentrations of these metabolites in grain, which was also emphasized by Champeil et al. (2004). At the conventional cultivation system cereals, especially wheat and maize, account for a very large share of crops in crop rotation. Necrotrophic pathogens from genus *Fusarium* develop dynamically on post-harvest residue of these crops, which promotes the accumulation of FHB symptoms in the successive years, especially in wheat following maize. Dill-Macky and Jones (2000) reported that in case of wheat grown after soy FHB symptoms occur in a much lesser degree. The application of ploughing additionally limits the incidence of this disease, as post-harvest residue is transferred to deeper soil layers. Thus it seems that the application of an appropriate crop rotation scheme in organic farming (with papilionaceous plants), together with comprehensive mechanical soil cultivation reduces the risk of pathogenic species from genus *Fusarium* to such an extent that a lack of chemical protection does not have a significant effect on an increase of *Fusarium* toxin concentrations in wheat kernels. It is possible that a significantly higher dry matter content (including that of nutrients and minerals) in tissues of plants grown in the organic farming system (Kouba 2003) is also accompanied by a higher concentration of phytochemical substances exhibiting fungistatic action connected with resistance to fungal pathogens. It is of interest that conventional farming with no fungicides applied (CON III) resulted in a drastic increase of *Fusarium* toxin concentrations in kernels. This result is especially important in view of the increasingly promoted and implemented sustainable

cultivation system, in which chemical crop protection is reduced, at the simultaneous maintenance of major elements of the conventional farming system (e.g. the application of mineral fertilizers) and at the same time placing more emphasis on crop rotation and mechanical tillage. The most intensive agronomic practices resulted in significantly lower concentrations of ERG, ATP and TRICH, as compared to the other systems of cultivation. The highest levels of ergosterol were detected in samples from conventional cultivation lacking chemical protection. These results form the basis for further research on this subject.

Acknowledgments

This work was realized within a programme of Austrian-Polish scientific and technical cooperation no. 7/2004.

References

- Abramson D, Gan Z, Clear RM, Gilbert J, Marquardt RR. 1998. Relationship among deoxynivalenol, ergosterol and *Fusarium* exoantigens in Canadian hard and soft wheat. International Journal of Food Microbiology 45: 217-224.
- Berthiller F, Schuhmacher R, Buttinger G, Krska R. 2005. Rapid simultaneous determination of major type A- and B-trichothecenes as well as zearalenone in maize by high performance liquid chromatography–tandem mass spectrometry. Journal of Chromatography A 1062: 209–216.
- Champeil A, Fourbet JF, Doré T, Rossignol L. 2004. Influence of cropping system on *Fusarium* head blight and mycotoxin levels in winter wheat. Crop Protection 23: 531-537.
- Dill-Macky R., Jones R. K. 2000. The Effect of Previous Crop Residues and Tillage on *Fusarium* Head Blight of Wheat. Plant Disease 84: 71-76.
- Ekblad A, Wallander H, Näsholm T. 1998. Chitin and ergosterol combined to measure total and living fungal biomass in ectomycorrhizas. New Phytologist 138: 143-149.

- Elmholt S, Labouriau R. 2005. Fungi in Danish soils under organic and conventional farming. *Agriculture, Ecosystems and Environment* 107: 65–73.
- Eudes F, Collin J, Rioux S, Comeau A. 1997. The trichothecenes, a major component of wheat scab pathogenesis. *Cereal Research Communication* 25: 495-496.
- Grodzińska K, Szarek-Łukaszewska G, Godzik B. 1999. Survey of heavy metal deposition in Poland using mosses as indicators. *The Science of the Total Environment* 229: 41-51.
- Hampton JG, Tekrony DM. 1995. *Handbook of vigour test methods*. 3rd Edition Zurich: ISTA.
- Jeleń H, Mirocha C, Wąsowicz E, Kamiński E. 1995. Production of volatile sesquiterpenes by *Fusarium sambucinum* strains with different abilities to synthesize trichothecenes. *Applied and Environmental Microbiology* 61: 3815-3820.
- Kouba M. 2005. Quality of organic animal products. *Livestock Production Science* 80: 33–40
- Lemmens M, Haim K, Lew H, Ruckebauer P. 2004. The effect of nitrogen fertilization on *Fusarium* head blight development and deoxynivalenol contamination in wheat. *Journal of Phytopathology* 152: 1-8.
- Łukanowski A, Sadowski Cz. 2002. Occurrence of *Fusarium* on grain and Hades of Winter wheat cultivated in organic, integrated, conventional systems and monoculture. *Journal of Applied Genetics* 43A: 69-74.
- Malmauret L, Parent-Massin D, Hardy JL. 2002. Contaminants in organic and conventional foodstuffs in France. *Food Additives and Contaminants* 19: 524-532.
- Marfleet L, Magan N, Lacey J. 1991. The relationship between fungal biomass, ergosterol and grain spoilage. In *Proceedings of the 5th International Working Conference on Stored - Product Protection* (ed. F. Fleurat-Lessard & P. Ducom), pp 405-412. INRA: Bordeaux, France.

- 1
2
3 Packa D, Koczowska I. 1987. The effect of *Fusarium culmorum* and *Fusarium nivale*
4
5 metabolites on germination, seedling growth and cytological disturbances of rye, wheat and
6
7 triticale. Mycotoxin Research, Special Edition. European Seminar, "Fusarium-, mycotoxins,
8
9 taxonomy, pathogenicity: 82-85.
10
11
12 Perkowski J, Stachowiak J, Kiecana I, Goliński P, Chełkowski J. 1997. Natural occurrence of
13
14 *Fusarium* mycotoxins in Polish cereals. Cereal Research Communication 25: 379-380.
15
16
17 Perkowski J, Chełkowski J, Goliński P. 2004. Occurrence of mycotoxins in cereals, plants,
18
19 foods and feeds in Poland. In: An overview on toxigenic fungi and mycotoxins in Europe.
20
21 Pettersson H. 1996. Mycotoxins in Scandinavian cereals; 1996; Poland Bydgoszcz:
22
23 Mikotoksyny w żywności, surowcach i paszach przemysłowych: 14-20.
24
25
26 Pussemier L, Larondelle Y, Van Peteghem C, Huyghebaert A. 2006. Chemical safety of
27
28 conventionally and organically produced foodstuffs: a tentative comparison under Belgian
29
30 conditions. Food Control 17: 14–21.
31
32
33 Reid LM, Nicol RW, Ouellet T, Savard M, Miller JD, Young JC, Stewart DW, Schaafsma
34
35 AW. 1999. Interaction of *Fusarium graminearum* and *F. moniliforme* in maize ears: disease
36
37 progress, fungal biomass, and mycotoxin accumulation. Phytopathology Vol. 89, No. 11:
38
39 1028-1037.
40
41
42 Schnürer J, Jonsson A. 1992. Ergosterol levels and mould colony forming units in Swedish
43
44 grain of food and feed grade. Acta Agriculturae Scandinavica Section B-Soil and Plant
45
46 Science 42: 240-245.
47
48
49 Teich AH. 1989. Epidemiology of wheat (*Triticum aestivum* L.) scab caused by *Fusarium*
50
51 spp. In: *Fusarium – Mycotoxins, Taxonomy and Pathogenicity*, Chełkowski J, editor.
52
53 Amsterdam: Elsevier. pp199-225.
54
55
56 Williams AP. 1989. Methodological developments in food mycology. Journal of Applied
57
58 Bacteriology Symposium Supplement: 61-67.
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table 1. Concentration of trichothecenes ($\mu\text{g kg}^{-1}$) and ergosterol (ERG, mg kg^{-1}) and amount of trichodiene (TRICH), adenosine triphosphate (ATP) and vigour test parameters for 32 examined wheat grain samples obtained from different cultivation systems.

Sample	HT-2 ($\mu\text{g kg}^{-1}$)	DON ($\mu\text{g kg}^{-1}$)	3-AcDON ($\mu\text{g kg}^{-1}$)	NIV ($\mu\text{g kg}^{-1}$)	NIV+DON ($\mu\text{g kg}^{-1}$)	ERG (mg kg^{-1})	TRICH (RU) $\times 10^{-3}$	ATP (RLU)	Non-germinated (%)	Abnormal (%)	<i>Fusarium</i> (%)
CON I											
1	13	31	-	-	31	1.08	2.085	3096	0.0	0.0	1.5
2	-	-	-	26	26	0.88	1.579	2844	0.0	0.0	0.0
3	12	-	-	75	75	0.86	5.026	1754	0.0	6.0	2.0
4	53	36	-	39	75	0.79	3.061	2313	1.0	4.0	2.0
5	-	-	-	80	80	0.37	6.630	1874	0.5	2.0	0.0
6	-	-	-	33	33	0.33	0.394	1834	0.5	1.0	0.5
7	-	-	-	-	0	0.64	0.520	1329	0.0	6.0	1.5
8	-	67	-	-	67	0.62	0.467	1110	1.5	2.5	2.5
9	-	-	-	-	0	1.85	0.955	3228	1.5	2.5	4.0
CON II											
10	-	14	-	-	14	2.11	7.768	15821	2.5	4.5	1.0
11	-	11	-	22	33	1.47	7.711	18166	1.5	3.5	0.0
12	-	11	-	(12)	23	0.66	3.116	13582	3.0	3.0	1.5

13	4	26	-	-	26	0.79	3.649	16199	4.5	7.0	1.5
14	6	53	-	(13)	66	0.55	1.319	12706	3.5	4.5	0.5
15	-	17	-	-	17	0.46	2.385	17262	6.0	8.0	3.5
CON III											
16	-	784	18	27	811	1.83	6.293	16642	2.0	4.5	0.0
17	-	480	-	43	523	2.9	3.961	14893	3.0	2.5	2.0
18	66	607	19	(7)	614	2.01	7.866	10944	1.5	3.0	0.5
19	-	190	-	-	190	3.34	8.581	5663	4.0	1.0	1.5
20	-	997	37	41	1038	1.87	3.285	10009	1.5	1.5	0.0
ECO											
21	-	-	-	28	28	3.34	1.528	10010	2.0	6.5	5.0
22	8	-	-	13	16	3.26	1.262	10477	3.5	3.5	2.5
23	-	-	-	-	0	2.01	1.404	12740	4.5	9.5	2.0
24	-	380	-	-	380	2.95	4.843	9772	2.5	10.5	1.5
25	-	12	-	-	12	1.65	0.644	15843	4.5	9.5	2.0
26	-	-	-	-	0	2.44	0.799	14828	3.5	7.5	1.5
27	-	7	-	-	7	0.92	2.009	14287	2.5	7.0	3.5
28	-	60	-	-	60	2.88	0.472	22062	10.5	7.5	1.0
29	-	-	181	-	0	2.84	1.410	31273	3.5	2.0	0.5

30	-	-	-	-	0	2.66	0.501	18701	3.5	2.5	0.0
31	-	-	-	-	0	1.67	0.943	23831	3.5	4.0	3.0
32	-	16	-	-	16	1.39	0.753	9714	0.5	3.0	2.5

T-2, DAS were not detected in the samples (LOQ: 1 µg/kg)

FUS-X was detected only in sample 15 at 12 µg/kg (LOQ: 6 µg/kg)

“-“ means that no peak was found

Results in brackets are below the limit of quantitation

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2. Mean values of analysed characters in wheat seed samples from four cultivation systems

Cultivation System	Mycotoxins ($\mu\text{g kg}^{-1}$)					Proportion of grains (%)					
	HT-2	DON	3-Ac DON	NIV	DON	TRICH (RU) $\times 10^{-3}$	ERG (mg kg $^{-1}$)	ATP (RLU)	Non-germ.	Abnorm. germ.	<i>Fusarium</i> mycelium
CON I	8.7	14.9 ^a	0.0 ^a	28.1 ^b	43.0 ^b	2.302 ^a	0.82 ^a	2153 ^a	0.6 ^a	2.7 ^a	1.6
CON II	1.7	22.0 ^b	0.0 ^a	7.8 ^a	29.8 ^a	4.325 ^{ab}	1.01 ^a	15623 ^b	3.5 ^b	5.1 ^b	1.3
CON III	13.2	611.6 ^c	14.8 ^b	23.6 ^b	635.2 ^c	5.997 ^b	2.39 ^b	11630 ^b	2.4 ^b	2.5 ^a	0.8
ECO	0.7	39.8 ^b	15.3 ^b	3.4 ^a	43.3 ^b	1.381 ^a	2.3 ^b	16128 ^b	3.7 ^b	6.1 ^b	2.1

^{a, b, c} -Values marked with the same letters within columns do not differ significantly ($P>0.05$).

Figure 1. Location of sites of seed samples collection on a map of Poland.