Diarrheic shellfish poisoning due to toxic mussel consumption: first recorded outbreak in Greece

Chrissanthy Vassilios Papadopoulou, Vangelis Economou, Moira Brett, Athina Kansouzidou, George Filioussis, Kostas Charalabopoulos

► To cite this version:

Chrissanthy Vassilios Papadopoulou, Vangelis Economou, Moira Brett, Athina Kansouzidou, George Filioussis, et al.. Diarrheic shellfish poisoning due to toxic mussel consumption: first recorded outbreak in Greece. Food Additives and Contaminants, 2007, 24 (03), pp.297-305. 10.1080/02652030601053139 . hal-00577520

HAL Id: hal-00577520

<https://hal.science/hal-00577520>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Diarrheic shellfish poisoning due to toxic mussel consumption: first recorded outbreak in Greece

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2006-039.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	06-Oct-2006
Complete List of Authors:	Papadopoulou, Chrissanthy; Ioannina University, Medical School, Microbiology; Medical School, Microbiology ECONOMOU, VANGELIS; Medical School, Microbiology Brett, Moira; CPHLS, Food Microbiology KANSOUZIDOU, ATHINA; Hospital for Infectious Diseases, Clinical Microbiology FILIOUSSIS, GEORGE; Medical School, Microbiology CHARALABOPOULOS, KOSTAS; Medical School, Physiology
Methods/Techniques:	Microbiology
Additives/Contaminants:	Environmental contaminants
Food Types:	Fish and fish products

SCHOLARONE™
Manuscripts

Diarrheic shellfish poisoning due to toxic mussel consumption: first recorded outbreak in Greece

V. ECONOMOU¹, C. PAPADOPOULOU¹, M. BRETT², A. KANSOUZIDOU³, K. CHARALABOPOULOS⁴, G. FILIOUSSIS¹, and K. SEFERIADIS⁵

¹Food Microbiology Unit, Microbiology Department, Medical School, University of Ioannina, Ioannina, 45110, Greece

²Marine Biotoxins Unit, Food Safety Microbiology Laboratory, Central Public Health Laboratory Service, 61 Colindale Avenue, London NW9 5HT, UK

³Clinical Microbiology Laboratory, Hospital for Infectious Diseases, G. Labraki Street, Thessaloniki, Greece

⁴Department of Physiology, Clinical Unit, Medical School, University of Ioannina, Ioannina, 45110, Greece

⁵Clinical Chemistry Department, Medical School, University of Ioannina, Ioannina, 45110, Greece

*Corresponding author- Chrissanthy Papadopoulou, e-mail: cpapadop@cc.uoi.gr

Abstract

During the week 14-20th January 2000, 120 people visited the Emergency Departments of Thessaloniki (Northern Greece) hospitals complaining of acute gastrointestinal illness after eating mussels (*Mytilus galloprovincialis*). The symptoms indicated diarrhoeic shellfish poisoning and the toxicity of mussels harvested from Thermaikos Gulf in Thessaloniki during the outbreak, was investigated using mouse bioassays. The bioassays revealed toxicity to mice by the mussel samples, while high numbers of toxic algae *Dinophysis acuminata* were identified in water samples from Thermaikos Gulf. Harvesting of mussels was immediately suspended and a monitoring programme for algal blooms was established from then onwards. During a follow-up of the mussels' toxicity from January 2000 through January 2005, two more mussel samples were found positive for DSP, one harvested in March 2001 from the area of the outbreak (Thermaikos Gulf) and one harvested in January 2001 from North-western Greece (Amvrakikos Gulf). However, no sporadic cases or outbreaks were reported during this period.

Keywords: Diarrheic Shellfish Poisoning, Greece, mussels, algal toxins, Dinophysis, outbreak

1
2
3 1 **Introduction**
4
5
6 2
7

8 3 Diarrheic shellfish poisoning (DSP) is a toxic syndrome caused by the ingestion of
9
10 4 shellfish contaminated with algal toxins produced by marine dinoflagellates belonging to
11
12 5 *Dinophysis* spp. (*D. fortii*, *D. acuta*, *D. acuminata*, *D. caudate*, *D.hastate*, *D. mitra*, *D.*
13
14 6 *rotundata*, *D. tripos*) (Hallegraef 1997; Yasumoto 2000) and *Prorocentrum* spp. (*P. lima*,
15
16 7 *P. maculosum*, *P. concavum* and *P. hoffmianum*) (Van Dolah 2000). The DSP toxins
17
18 8 belong to a group of fat-soluble polyether compounds, including Okadaic acid (OA) and
19
20 9 the closely related Dinophysistoxins (DTX). Gastrointestinal effects have only been
21
22 10 proven for OA, DTX-1 and DTX-3 (Murata *et al.* 1982; Rossini 2000; Vale and Sampayo
23
24 11 2002). DTXs are lipophilic and accumulate in the fatty tissue of shellfish. The toxin profiles
25
26 12 in shellfish display considerable seasonal and geographical variation (Yasumoto *et al.*
27
28 13 1985; Sechet *et al.* 1990). Ordinary cooking processes do not destroy DSP toxins (Gestal
29
30 14 – Otero 2000).
31
32
33
34
35
36
37
38

39 16 DSP symptoms include diarrhoea, nausea, vomiting, abdominal pain and chills, which
40
41 17 manifest within 30 min to a few hours after consumption of contaminated shellfish and
42
43 18 persist for 3 to 4 days (Aune and Yndestad 1993). Although short-term effects are
44
45 19 relatively mild, long-term effects are more severe and chronic exposure may promote
46
47 20 tumour formation in the digestive system (Hallegraef 1997; Rossini 2000). OA and DTX1
48
49 21 are potent inhibitors of protein phosphatases and this mode of action may be linked to the
50
51 22 observed diarrhoea, degenerative changes in the absorptive epithelium of the small
52
53 23 intestine and tumour development (Hallegraef 1997).
54
55
56
57
58
59
60

DSP was first reported in the Netherlands in the 1960s and in Japan in the late 1970s (Kat 1985; Yasumoto *et al.* 1980). Since then it has been recorded in several European countries including France (van Egmond *et al.* 1993), Norway (Rossebo *et al.* 1973; Underdal *et al.* 1985, Torgersen *et al.* 2005), Sweden (Krogh *et al.* 1985), Spain (Campos *et al.* 1982; Fernández *et al.* 1996), Italy (Viviani *et al.* 1995; Ciminiello *et al.* 1997), Portugal (Vale and Sampayo 1999), Belgium (De Schrijver *et al.* 2002) former USSR (Konovalova 1993). There are also reports from Japan (Yasumoto *et al.* 1984; Yasumoto *et al.* 1985; Yasumoto 2000; Kawabata 1989), India (Karunasagar *et al.* 1989), Australia, New Zealand, Thailand, North and South America (Hallegraef 1997; Scoging 1998; Gestal – Otero 2000). From the available literature it is concluded that DSP or the presence of DSP toxins in mussels, is increasingly reported particularly in Europe (Table I). The increased reporting may be the result of increasing knowledge and awareness about toxic shellfish poisonings (DSP, ASP and PSP) and the implementation of surveillance/monitoring programmes. Also, it has to be emphasized that toxin producing algae and toxic mollusks are reported from new areas, but well documented cases in Europe are rare and this may be attributed to under-diagnosis and/or underreporting. The shellfish most frequently implicated in DSP are mussels, but it has also been reported from scallops, clams, cockles, oysters and crabs (Gestal – Otero 2000; Vale and Sampayo 2002; Jørgensen *et al.* 2005; Torgersen *et al.* 2005). In the present paper the first DSP outbreak in Greece is reported, together with the laboratory findings from mussel samples collected during the outbreak and the subsequent four years after the outbreak.

Materials and methods

The DSP outbreak investigation

During the week 14-20th January, 2000, one hundred and twenty people of both sexes, aged from 8 to 70 years old, visited the Emergency Departments of the hospitals in the city of Thessaloniki (Northern Greece), suffering from acute non-febrile gastro-enteritis (Table II). The predominant symptoms reported by the patients were diarrhea, nausea, vomiting, abdominal cramps and chills, which persisted for more than 24 h. All patients reported onset of symptoms within 30 min to 10 h after consuming a mussel meal (deep fried mussels, poached in cheese and tomato hot sauce, mussel pilaf and cooked mussel salads) in various establishments (restaurants, taverns, homes) within the Greater Thessaloniki area. All the hospitalized patients recovered completely within 12 to 36 h after the administration (i.v) of saline solution and were completely asymptomatic upon exit from the hospital. [Insert Table II about here]

Fecal samples from the hospitalised patients were examined for routine bacterial, viral and parasitic agents causing non febrile gastroenteritis, but they were found negative. The common history of the patients (reporting a mussels' meal), based on relative questionnaires, combined with the clinical and laboratory findings were indicative of diarrheic shellfish poisoning, thus the Veterinary Authorities were immediately notified and consumption of mussels was banned until further notice. Mussel dishes are very popular in the coastal areas of Greece, especially in the vicinity of Thessaloniki and accordingly it was estimated that the actual number of people infected was actually much larger than the 120 cases recorded at the hospitals' admissions desks. Hence, the estimates were for a few hundreds of people infected, who either visited privately a

1
2
3 1 general practitioner or asked no medical assistance because their symptoms soon
4
5 2 subsided.
6
7
8 3
9

10 4 The patients who had prepared their meals at home reported purchase of fresh mussels
11
12 5 from seafood retails, supermarkets and open markets in the Thessaloniki greater area
13
14 6 and similar retail sources were reported by the restaurant and tavern owners. The fresh
15
16 7 mussels (*Mytilus galloprovincialis*) sold in the markets of Thessaloniki area and in most
17
18 8 markets in Northern and Northwestern Greece are harvested from aqua cultures located
19
20 9 into the Thermaikos Gulf (Figure 1).
21
22
23
24
25 10
26

27 11 The Veterinary and Sea Biology authorities reported numbers of dinoflagellates ranging
28
29 12 from 3×10^4 to $5,4 \times 10^4$ cells/L of the toxic genus *Dinophysis* spp. in the seawater samples
30
31 13 from Thermaikos Gulf, and specifically of the species *D. acuminata* and *D. sacculus*
32
33 14 (Koukaras and Nikolaidis 2002). No other episodes either of sporadic cases or outbreaks
34
35 15 associated with consumption of mussels exposed to toxin-producing Dinoflagellates of the
36
37 16 DSP type have been reported for the following four years, although the presence of toxic
38
39 17 *Dinophysis acuminata* was reported again in Thermaikos Gulf in March – May 2001
40
41 18 (Koukaras and Nikolaidis 2002). However, suspension of harvesting and withdrawal of
42
43 19 mussels from the market are being applied sporadically in Thermaikos Gulf, whenever
44
45 20 algal blooms are detected during the phytoplankton monitoring program, which has been
46
47 21 established after the outbreak. An algal bloom is a rapid increase in the population of
48
49 22 phytoplankton algae in an aquatic system. The appearance of *Dinophysis*, even at low
50
51 23 densities such as 200 cells per litre, can cause already a toxification of shellfish that is
52
53 24 enough to affect humans (Van Egmond *et al.* 2004). According to the Greek Veterinary
54
55
56
57
58
59
60

and Sea Biology authorities, seafood harvesting is suspended when toxic dinoflagellate counts are higher than 200 cells/L.

The mussel toxicity investigation

Collection of mussel samples

From January 2000 to January 2005, 36 batches (500 g each) of fresh mussels were examined for DSP toxins contamination. The mussel samples (species *Mytilus galloprovincialis*) were randomly collected from local seafood markets. Twenty samples were collected from Thermaikos Gulf (Halastra) in Thessaloniki (Northern Greece) and sixteen samples from Amvrakikos Gulf (Preveza) (Northwestern Greece) (Figure 1). [Insert Figure 1 about here] Two out of the 36 samples were sent to the Food Microbiology Unit of the Microbiology Department; University of Ioannina from colleagues at the Hospital for Infectious Diseases in Thessaloniki during the DSP outbreak. These samples were harvested from the Halastra area of Thermaikos gulf at the 20 January 2000. Samples after April 2001 until January 2005 were collected every 3 months. All fresh mussel samples forwarded to our lab were deep-frozen (-30°C) immediately after collection and were transported by airplane (flight duration 40 min) in well insulated boxes to the Food Microbiology Unit, within less than 24 hours after collection and upon arrival were stored at -30°C until testing. The samples were tested within 7 days time after their arrival in our laboratory. The samples were defrosted at room temperature and were thoroughly cleaned under a stream of tap water after external materials were brushed off the shells. From each sample, 200 g of flesh was homogenized in a blending mixer and a portion of 100 g was weighed using an Ohaus model 1500D balance (Ohaus Co, P.O. Box 2033 19A Chapin Road Pine Brook, NJ 07058) prior to extraction.

Extraction of DSP toxins

The extraction of DSP toxins was performed following the method validated by the Marine Laboratory, Aberdeen, UK, which is the EU designated UK National Reference Laboratory (NRL) for Marine Biotoxins. The method is that of Yasumoto et al (Yasumoto *et al.* 1984) with slight modifications. A portion of 100 g of homogenized mussel flesh was suspended three times and thoroughly mixed with 100 ml of acetone (Scharlau, Analar grade) at room temperature. The aliquots were centrifuged at 3000 rpm for 10 min in a Sorvall GLC-4 desktop centrifuge and the supernatants were collected and pooled. Acetone was removed by rotary evaporation for 1h at 60°C or until all trace of acetone had gone. An Eyela type N rotary evaporator (Eyela Co, 3-3-4 Honcho Nihonbashi, Chuo-ku, Tokyo 103-0023, Japan) was used. The remaining aqueous-lipid extract was separated with diethyl ether (Scharlau, Analar grade), backwashed with a small volume of water and dried by rotary evaporation at 65°C for 2 to 4 h. The remaining material was suspended in 4 ml of 1% Tween 60 (Merck) aqueous solution.

Mouse bioassay

BALB-C strain white mice weighting 19 to 21 g each were used for the mouse bioassay. The mice were weighed prior to intra-peritoneal injection of 1ml of the mussel extract. Three mice were used for each sample. All injected mice and controls were monitored for a period of at least 24 h after injection. Initially the mice were observed constantly for the first 6 h and from then every 1 h for the remaining 18 h. During the 24 h monitoring any behavioral changes or symptoms were recorded in detail. When no typical symptoms of DSP intoxication were observed after 24 h, the mice were monitored for an additional 24 h period. The symptoms that were characterized as typical of DSP intoxication were initial apathy, general weakness, difficulty in movement, spasms, contractions, bluish skin

coloration and respiratory distress and death. According to the Decision 2002/225/EC, death of two out of three mice within 24 hours after inoculation into each of them of an extract equivalent to 5 g of hepatopancreas or 25 g whole body should be considered as a positive result for the presence of one or more of the toxins mentioned, when a mouse bioassay is utilized as a screening method.

Results and discussion

During this investigation positive mouse bioassay results were obtained in 4 out of 36 samples tested. Specifically, 3 samples out of 9 collected from the area of Halastra within a two-year period from the onset of the DSP outbreak were found positive by the mouse bioassay method. The two samples collected during the outbreak were toxic resulting in death of mice within 85-90 min after the intra-peritoneal injection. The third positive sample harvested from the area of Halastra was collected 15 months after the outbreak when the mussel withdrawal was recalled. However, no sporadic cases or outbreak due to DSP or similar gastrointestinal illness in humans were recorded during the period the DSP positive sample was collected.

The fourth positive sample was harvested from the Amvrakikos Gulf in Northwestern Greece, where no positive or suspect DSP cases have ever been recorded. Amvrakikos Gulf is the nearest to Ioannina and like Thermaikos, it is shallow, calm, warm and thought to be the most polluted gulf in the Ionian Sea, which is part of the Adriatic Sea, where there are frequent reports for toxic blooms (Boni *et al.* 1990; Fatorruso *et al.* 1992; Draisci 1995; Viviani *et al.* 1995; Ciminiello *et al.* 1997; 1998; Caroppo *et al.* 2001; Pavela – Vranic *et al.* 2002). Also, it should be noted that the mice injected with the extract of the Amvrakikos sample, in addition to the typical DSP intoxication symptoms, exhibited a

marked hypersensitivity 2 h prior to death, together with convulsions. The symptoms of the mice following intraperitoneal injection are presented in Table III. [Insert Table III about here]

There are a number of studies from different countries concerning DSP outbreaks, sporadic cases or detection of toxins in a variety of seafood (Table I). The DSP outbreak described in this paper is the first recorded in Greece and it was classified as a diarrhetic shellfish poisoning, based on clinical manifestations characteristic for DSP, the common history of all patients reporting consumption of mussels, the absence of any known pathogen in fecal and blood samples from all hospitalized patients, no detection of antibodies to known pathogens, the subsequent analysis of mussels collected during the outbreak and the detection of *Dinophysis acuminata* in the water of Thermaikos Gulf. The actual number of the people affected cannot be precisely estimated, due to lack of sufficient data. However, the number of 120 people needing urgent medical care within a period of less than 7 days indicates a foodborne outbreak, of the group of seafood-borne emerging diseases, the first recorded in Greece. . [Insert Table I about here]

The mouse bioassays performed during the outbreak are indicative of the presence of lipophilic toxins. This finding combined with the increased concentration of *D. acuminata* (Koukaras and Nikolaidis 2002), which is the most common DSP producing species in Europe (Kumagai *et al.* 1996; Morono *et al.* 2002), indicates okadaic acid and / or DTX as the potential cause of this outbreak. It should be noted that the third positive sample from Thermaikos Gulf was collected in March 2001 when, according to other researchers (Koukaras and Nikolaidis 2002) an increased number of *Dinophysis* spp. was observed. Furthermore, the predominant DSP toxins detected in European mussels and especially

1 in the mussels of the Adriatic Sea are okadaic acid and DTX – 2 (Draisci 1995; Pavela –
2 Vranic *et al.* 2002). In a recent publication (Ciminiello *et al.* 2005) concerning
3 investigation of the toxin profile of Greek mussels *M. galloprovincialis* by Liquid
4 Chromatography – Mass Spectrometry, the presence of okadaic acid only (at levels 0.10
5 – 0.20 µg/g) is confirmed. The hypersensitivity and the convulsions observed in mice
6 injected with the extract of the positive sample from Amvrakikos gulf, is indicative of the
7 simultaneous presence of an unknown lipophilic toxin along with okadaic acid and / or
8 DTX – 2. Jumping before death, convulsions and short survival time is observed when
9 mice are injected i.p. with yessotoxin (Ciminiello *et al.* 1997). Although the presence of
10 yessotoxin in mussels has been reported in neighboring areas, it is quite improbable that
11 yessotoxins are the cause of these symptoms. Yessotoxins are poorly soluble in diethyl
12 ether and therefore probably could not co-elute with the other toxins (Ciminiello *et al.*
13 1997). Another possible explanation for the observed symptoms in this sample is that it
14 contained azaspiracid. After intraperitoneal injection of azaspiracid, mice exhibit
15 respiratory difficulties, spasms, paralysis of the limbs and death (Ito *et al.* 2000). Still, the
16 presence of azaspiracid has not been proven by analytical methods in shellfish harvested
17 from the Mediterranean Sea (James *et al.* 2004).

18
19 Detection of DSP is routinely based on mouse bioassays, HPLC methods and
20 immunoassays. It is generally acknowledged that biological tests suffer from considerable
21 variability and lack of sensitivity and because of increasing opposition to animal testing
22 many efforts have been made to detect DSP toxins by HPLC based methods. However,
23 the lack of availability of analytical standards and Certified Reference materials for all of
24 the DSP toxins currently precludes the sole use of chemical methods in routine monitoring
25 programmes. Additionally, the maximum level for DSP toxins set by regulatory authorities

varies considerably from country to country. In general, levels of DSP toxins greater than 200 ng/g in shellfish are considered dangerous for human consumption (Hallegraef, 1997). The potential public health risks and the expanding geographical distribution of the causative dinoflagellates have resulted in the issuing by the Commission of the European Countries of the Decision 2002/225/EC of the 15 May 2002, which is an amendment of the Council directive 91/492/EEC, which states that the maximum level of okadaic acid, dinophysistoxins and pectenotoxins together, in bivalves intended for human consumption (the whole body or any part edible separately) shall be 160 µg of okadaic acid equivalents/kg.

To date, control measures to protect human health against DSP contaminated seafood involve both biological and chemical procedures. At the time of the outbreak in the Food Microbiology Unit we were able to apply only the biological method (mouse bioassay). The accumulation of algal toxins is one of the most critical problems in bivalve aquaculture. Amongst the three most important categories of shellfish poisoning (amnesic, diarrhoeic, paralytic), DSP is the least severe causing only gastrointestinal symptoms. However, the accumulation of toxic algae in many bivalve producing areas has a great economic impact as their appearance leads to the implementation of long term bans on shellfish harvest, which is happening frequently at Thermaikos Gulf aquacultures since the reported outbreak. The factors influencing the growth of the microalgal population include: a) changes in water temperature favoring encysting or excysting of dinoflagellates, b) decreased water salinity, c) increased concentrations of organic substances, d) extended duration of sunlight and e) calm waters (Gestal – Otero 2000; Caroppo *et al.* 2001).

All these factors are present in Thermaikos Gulf (North Aegean Sea), especially in the Halastra site, where the constantly calm and warm waters favor mussel cultures. However, the outfall of three large rivers in the gulf (Gallikos, Loudias and Axios), the waste effluents of Greater Thessaloniki area (second largest city of Greece) and the drainage of the plain of Central Macedonia (a plain with intensive agricultural activities), make Thermaikos Gulf, ideal aquatic environment for the growth of microalgal population and possible toxin production by dinoflagellates. Similar conditions exist at Amvrakikos Gulf (North Ionian Sea), which is part of the South-central Adriatic Sea. Red tide blooms are known to occur in the central Adriatic Sea since the 80's (Boni *et al.* 1990). However, toxic blooms associated with shellfish intoxication are becoming more frequent during the last decade in this geographic area (Fattorusso *et al.* 1992; Viviani *et al.* 1995; Draisci *et al.* 1995; Ciminiello *et al.* 1997; Ciminiello *et al.* 1998; Draisci *et al.* 1999; Caroppo *et al.* 2001; Pavela – Vranic *et al.* 2002).

DSP intoxication is still an under reported illness, which is attracting interest after notification of outbreaks (Gestal – Otero 2000; Yasumoto 1985; Vale and Sampayo 2002). Therefore knowledge of its existence along with the recently increasing reports of red tide blooms spreading from Europe to America to Australia should trigger a constant awareness for this emerging seafood zoonotic disease.

Acknowledgments

The present study was carried out during a bilateral joint research project on Marine Biotoxins, participating the Biotoxins Unit, Food Safety Microbiology Laboratory, CPHLS, London UK and the Food Microbiology Unit, Microbiology Department, Medical School, University of Ioannina, GR; the joint project was financially supported by grants from the

General Secretariat for Research and Technology, The Greek Ministry of Development and The British Council. Also the authors wish to thank the former Director of the Clinical Microbiology Laboratory, Hospital for Infectious Diseases, Dr. V. Daneilides for collecting and sending the two mussel samples during the outbreak. Mouse bioassays comply with the current laws of Greece for laboratory animal experimentations.

References

- Anonymous 1994. Diarrhetic shellfish poisoning associated with mussels. CDR Wkly 4 (22): 101.
- Anonymous 1997. An outbreak of diarrhetic shellfish poisoning. CDR Wkly 7 (28): 247.
- Aune T, Yndestad M. 1993. Diarrhetic shellfish poisoning. Chapter 5. In: Falconer IR, editor. *Algal Toxins in Seafood and Drinking Water*. London, UK, Academic Press. p. 87-104
- Boni L, Mancini L, Milandri A, Poletti R, Pompei M, Viviani R. 1990. First cases of DSP in the Northern Adriatic Sea. International Conference Regione Emilia Romagna; 1990 March 21-24; Bologna. Italy.
- C.E.C. Report of the Scientific Veterinary Committee (Section Public Health) on Diarrhetic Shellfish Poisoning in Europe. 1990. VI/6182/89-EN Rer.2.
- Campos MJ, Fraga S, Marino J, Sanchez FJ. 1982. Red tide monitoring program in NW Spain. Report of 1977-1981, International Council for the Exploration of the Sea, Council Meeting (ICES.CM), 1982/L: 27.
- Carmody EP, James KJ, Kelly SS. 1996. Dynophysistoxin-2 the predominant Diarrhetic Shellfish toxin in Ireland. Toxicon 34:351-359.

- 1
2
3 1 Caroppo C, Congestri R, Bruno M. 2001. Dynamics of *Dynophysis* sensu lato species
4
5 2 (Dinophyceae) in a coastal Mediterranean environment (Adriatic Sea). *Continental*
6
7 3 *Shelf Research* 21: 834 – 1854.
8
9
10 4 Ciminiello P, Fattorusso E, Forino M, Magno S, Poletti, R, Satake M, Viviani R, Yasumoto
11
12 5 T. 1997. Yessotoxin in mussels of the Northern Adriatic Sea. *Toxicon*, 35: 177 – 183.
13
14
15 6 Ciminiello, P., Fattorusso, E., Forino, M., Magno, S., Poletti, R., and Viviani, R. 1998.
16
17 7 Isolation of Adriatoxin, a new analogue of yessotoxin from mussels of the Adriatic
18
19 8 Sea. *Tetrahedron Letters*, 39: 8897 – 8900.
20
21
22 9 Ciminiello P, Dell'Aversano C, Fattorusso E, Forino M, Magno S, Santelia F, Tsoukatou
23
24 10 M. 2005. Investigation of the toxin profile of Greek mussels *Mytilus galloprovincialis*
25
26 11 by liquid chromatography—mass spectrometry. *Toxicon* 47: 174 - 181.
27
28
29 12 De Schrijver K, Maes I, De Man L, Michelet J. 2002. An outbreak of diarrhoeic shellfish
30
31 13 poisoning in Antwerp, Belgium. *Eurosurveillance*, 7:139 – 141.
32
33
34 14 Dickey RW, Fryxell GA, Granade HR, Roelke D. 1992. Detection of the marine toxins
35
36 15 Okadaic acid and Domoic acid in shellfish and phytoplankton in gulf of Mexico.
37
38 16 *Toxicon* 30:355-359.
39
40
41 17 Draisci R, Lucentini L, Giannetti L, Boria P, Stacchini A. 1995. Detection of diarrheic
42
43 18 shellfish toxins in mussels from Italy by ion spray liquid chromatography – mass
44
45 19 spectrometry. *Toxicon*, 33:1591 – 1603.
46
47
48 20 Draisci R, Ferretti E, Palleschi L, Marchiafava C, Poletti R, Milandri A, Ceredi A, Pompei
49
50 21 M. 1999. High levels of yessotoxin in mussels and presence of yessotoxin and
51
52 22 homoyessotoxin in dinoflagellates of the Adriatic Sea. *Toxicon* 37:1187 – 1193.
53
54
55 23 Fattorusso E, Ciminiello P, Constantino V, Magno S, Mangoni A, Poletti R, Pompei M,
56
57 24 Viviani R. 1992. Okadaic acid in mussels of Adriatic Sea. *Marine Pollution Bulletin*
58
59 25 24: 234-237.
60

- 1
2
3 1 Fernández ML, Míguez A, Cacho E, Martínez A. 1996. Detection of Ocadaic acid esters in
4
5 2 the hexane extracts of Spanish mussels. *Toxicon*, 34: 381 – 387.
6
7
8 3 Garcia C, Mardones P, Sfeir A, Lagos N. 2004. Simultaneous presence of Paralytic and
9
10 4 Diarrheic Shellfish Poisoning toxins in *Mytilus chilensis* samples collected in the
11
12 5 Chiloe Island. *Biological Research* 37 suppl. A: 721 – 731.
13
14
15 6 Gestal – Otero JJ. 2000. Nonneurotoxic toxins. In: Botana LM, Editor. *Seafood and*
16
17 7 *freshwater toxins : pharmacology, physiology and detection*. New York. Marcel
18
19 8 Dekker. p. 45 – 64,
20
21
22 9 Hallegraef GM. 1997. Algal toxins in Australian shellfish. In: Hocking AD, Arnold G,
23
24 10 Jenson I, Newton K, Sutherland P, editors. *Foodborne microorganisms of Public*
25
26 11 *Health significance* 5th Edition. Sydney. AIFST (NSW branch). p. 561 – 571.
27
28
29 12 Ito E, Satake M, Ofuji K, Kurita N, McMahon T, James K, Yasumoto T. 2000. Multiple
30
31 13 organ damage caused by a new toxin azaspiracid, isolated from mussels produced
32
33 14 in Ireland *Toxicon*, 38: 917 – 930.
34
35
36 15 James KJ, Fidalgo Sàez MJ, Furey A, Lehane M. 2004. Azaspiracid poisoning, the food-
37
38 16 borne illness associated with shellfish consumption. *Food Additives and*
39
40 17 *Contaminants*, 21 (9): 879 – 892.
41
42
43 18 Jørgensen K, Scanlon S, Jensen LB. 2005. Diarrhetic shellfish poisoning toxin esters in
44
45 19 Danish blue mussels and surf clams. *Food Additives and Contaminants*, 22 (8):743
46
47 20 – 751
48
49
50 21 Karunasagar I, Segar K, Karunasagar I. 1989. Incidence of DSP and DSP in shellfish
51
52 22 along the coast of Karnataka state (India). In: Okaichi T, Anderson DM, Nemoto T,
53
54 23 editors. *Red Tides: Biology, Environmental Science and Toxicology*. Elsevier,
55
56 24 Amsterdam. p. 61 – 64
57
58
59
60

- 1
- 2
- 3 1 Kat M. 1985. *Dinophysis acuminata* blooms, the distinct cause of Dutch mussel poisoning.
- 4
- 5 2 In: Anderson DM, White AW, Baden DG, editors. *Toxic dinoflagellates*. Elsevier,
- 6
- 7 3 Amsterdam. p. 73 – 77,
- 8
- 9
- 10 4 Kawabata T. 1989 Regulatory aspects of marine biotoxins in Japan. In: Natori S,
- 11
- 12 5 Hashimoto K, Ueno Y, editors. *Mycotoxins and Phycotoxins '88*. Elsevier,
- 13
- 14 6 Amsterdam. p. 469 – 476
- 15
- 16
- 17 7 Konovalova GV. 1991. Toxic and potentially toxic dinoflagellates from the far east coastal
- 18
- 19 8 water of the USSR. In: Smayda TJ, Shimizu Y, editors. *Toxic Phytoplankton Blooms*
- 20
- 21 9 *in the Sea; Proceedings 5th International conference on Toxic Marine Phytoplankton*
- 22
- 23 10 *Newport (USA)* . Elsevier, Amsterdam. p. 275 – 279
- 24
- 25
- 26
- 27 11 Koukaras K, Nikolaidis G. 2002. Intense DSP outbreak associated with a *Dinophysis*
- 28
- 29 12 bloom in Thermaikos Gulf (N.W. Aegean Sea, Greece). *Harmful Algae News*, 23: 8.
- 30
- 31
- 32 13 Krogh P, Edler L, Graneli E. 1985. Outbreaks of diarrhetic shellfish poisoning on the west
- 33
- 34 14 coast of Sweden. In: Anderson, DM., White, AW. and Baden, DG, editors. *Toxic*
- 35
- 36 15 *Dinoflagellates*. Elsevier, Amsterdam. p. 501 – 504.
- 37
- 38
- 39 16 Kumagai M, Yanagi T, Murata M, Yasumoto T, Kat M, Lassus R, Rodriguez – Vazquez
- 40
- 41 17 JA. 1986. Okadaic acid as the causative toxin of diarrhetic shellfish poisoning in
- 42
- 43 18 Europe. *Agric. Biol. Chem.*, 50, 2853 – 2857.
- 44
- 45
- 46 19 Madigan TL, Lee KG, Padula DJ, MacNabb P, Pointon AM. 2005. Diarrhetic Shellfish
- 47
- 48 20 Poisoning Toxin in South Australian Shellfish. *Harmful Algae* (article in Press).
- 49
- 50
- 51 21 Mozetic P, Bozic P. 2001. DSP events in Slovenia: A need for legislative regulation.
- 52
- 53 22 *Harmful Algae Management and Mitigation 2001 Abstracts*
- 54
- 55 23 Morono A, Arevalo F, Fernandez ML, Maneiro J, Pazos Y, Salgado C., Blanco J. 2003.
- 56
- 57 24 Accumulation and transformation of DSP toxins in mussels *Mytilus galloprovincialis*
- 58
- 59
- 60

- 1 during a toxic episode caused by *Dinophysis acuminata*. Aquatic Toxicology 62: 269
2 – 280.
- 3 Murata M, Shimatani M, Sugitani H, Oshima Y, Yasumoto T. 1982. Isolation and structural
4 elucidation of the causative toxin of diarrhetic shellfish poisoning. Bulletin of the
5 Japanese Society of Sciences. Fish 48:549 – 552.
- 6 Papadopoulou – Kyriazi A. Public control and supervision of production Shellfish in the
7 Thermaikos gulf and the administrative limits of prefecture Thessaloniki. Proceedings
8 book Vol. A, 3rd Hellenic Symposium on Food Hygiene and Food Technology,
9 Athens, Greece 2004, 463 – 466.
- 10 Pavela-Vrancic M, Vestrovic V, Marasova I, Gillman M, Furey A, James KJ. 2002. DSP
11 toxin profile in the coastal waters of Central Adriatic Sea. Toxicon 40: 1601 – 1607.
- 12 Ramstad H, Hovgaard P, Yasumoto T, Larsen S, Aune T. 2001. Monthly variations in
13 diarrhetic toxins and yessotoxin in shellfish from coast to the inner part of the
14 Sognefjord, Norway. Toxicon 39: 1035 – 1043.
- 15 Rossebo L, Thorson B, Aase R. 1973. Etiologisk ukklar matforgifning etter konsum av
16 blaskjell. Norsk. Vet. Tidsskr, 82: 639 – 642.
- 17 Rossini GP. 2000. Neoplastic activity of DSP toxins: The effects of okadaic acid and
18 related compounds on cell proliferation: Tumor promotion or induction of apoptosis?
19 In: Botana, LM editor. *Seafood and freshwater toxins: pharmacology, physiology and*
20 *detection* Marcel Dekker, New York. p. 257 – 287
- 21 Scoging AC. 1998. Marine biotoxins. Journal of Applied Microbiology, Symposium
22 supplement, 84: 41S – 50S.
- 23 Scoging AC. Bahl M. 1998. Diarrhetic shellfish poisoning in the UK. Lancet 352: 117.
- 24 Sechet V, Safran, P, Hovgaard, P, Yasumoto T. 1990. Causative species of diarrhetic
25 shellfish poisoning (DSP) in Norway. Marine Biology 105: 269 – 274.

- 1
2
3 1 Torgersen T, Aasen J, Aune T. 2005. Diarrhetic shellfish poisoning by okadaic acid
4
5 2 esters from Brown crabs (*Cancer pagurus*) in Norway. *Toxicon* 46:572-578.
6
7
8 3 Underdal B, Yndestad M, Aune T. 1985. DSP intoxications in Norway and Sweden,
9
10 4 Autumn 1984 – Spring 1985. In: Anderson DM, White AW, Baden DG, editors. *Toxic*
11
12 5 *Dinoflagellates*. Elsevier, Amsterdam. p. 489 – 494.
13
14
15 6 Van Egmond HP; Apeldoorn ME van ; Speijers GJA. *Marine biotoxins*. Rome: Food and
16
17 7 Agriculture Organization of the United Nations, 2004 (FAO Food and Nutrition Paper
18
19 8 80)
20
21
22 9 Vale P, Maia AJ, Correia A, Rodrigues SM, Botelho MJ, Casanova G, Silva A, Vilarinho
23
24 10 MG, Silva AD. 2003. An outbreak of diarrhetic shellfish poisoning after ingestion of
25
26 11 wild mussels at the Northern coast in summer 2002. *Electron. Journal Environmental*
27
28 12 *Agricultural and Food Chemistry* 2:449 – 452.
29
30
31 13 Vale P, Sampayo MAM. 1999. Esters of okadaic acid and dinophysitoxin-2 in Portuguese
32
33 14 bivalves related to human poisonings. *Toxicon*, 37:1109 – 1121.
34
35
36 15 Vale P, Sampayo MAM. 2002. First confirmation of human diarrhoeic poisonings by
37
38 16 okadaic acid esters after ingestion of razor clams (*Solen marginatus*) and green
39
40 17 crabs (*Carcinus maenas*) in Aveiro Lagoon, Portugal and detection of okadaic acid
41
42 18 esters in phytoplankton. *Toxicon*, 40: 511 – 517.
43
44
45 19 Van Dolah F. 2000. Diversity of marine and freshwater algal toxins In: Botana, L. M.
46
47 20 editor. *Seafood and freshwater toxins: pharmacology, physiology and detection*.
48
49 21 Marcel Dekker, New York. p. 19 – 43.
50
51
52 22 Van Egmond HP, Aune T, Lassus P, Speijers GJA, Waldock M. 1993. Paralytic and
53
54 23 diarrhoeic shellfish poisons, occurrence in Europe, toxicity, analysis and regulation.
55
56 24 *Journal Natural Toxins* 2: 41-83.
57
58
59
60

- 1
2
3 1 Van Egmond HP, Van Apeldoorn ME, Speijers GJA. 2004. "Marine biotoxins". FAO Food
4
5 2 And Nutrition Paper 80, Rome.
6
7
8 3 Vieytes M, Louzao MC, Alfonso A., Calbado A G, Botana L. 2000. Mechanism of
9
10 4 action and Toxicology. In: Botana, LM, editor. *Seafood and freshwater toxins:*
11
12 5 *pharmacology, physiology and detection*. Marcel Dekker, New York. p. 239 – 256.
13
14
15 6 Viviani R, Boni L, Cattani O, Milandri A, Poletti R, Pompei R, Sansoni G. 1995. ASP,
16
17 7 DSP, NSP and PSP monitoring in "mucilaginous aggregates" and in mussels in a
18
19 8 coastal area of the Northern Adriatic Sea facing Emilia – Romagna in 1988, 1989
20
21 9 and 1991. *The Science of Total Environment*, 165: 203 – 211.
22
23
24 10 Wu JY, Zheng L, Wang JH. 2005. Contamination of shellfish from Shanghai seafood
25
26 11 markets with paralytic shellfish poisoning and diarrhetic shellfish poisoning toxins
27
28 12 determined by mouse bioassay and HPLC. *Food additives and Contaminants* 22:
29
30 13 647-651.
31
32
33 14 Yasumoto T. 2000. Historic considerations regarding seafood safety. In: Botana, LM.
34
35 15 editor. *Seafood and freshwater toxins: pharmacology, physiology and detection*.
36
37 16 Marcel Dekker, New York. p. 1 – 17.
38
39
40 17 Yasumoto, T, Murata M, Oshims Y, Matsumoto GK, Clardy J. 1984. Diarrhetic shellfish
41
42 18 poisoning. In: Ragelis, E.P., editor. *Seafood toxins*. AOAC International. Washington
43
44 19 DC. p. 214-217
45
46
47 20 Yasumoto, T, Murata M, Lee JS, Torigoe K. 1985. Diarrhetic Shellfish toxins.
48
49 21 *Tetrahedron* 41: 1019 – 1025.
50
51
52 22 Yasumoto T, Oshima Y, Sugawara W, Fukuyo Y, Oguri H, Igarishi T, Fujita N. 1980.
53
54 23 Identification of *Dinophysis fortii* as the causative organism of diarrhetic shellfish
55
56 24 poisoning. *Bulletin of the Japanese Society of Sciences Fish.* 46: 1405 – 1411.
57
58
59
60

Table I

Year/month	Country	Number of cases	Seafood	Toxins determined
1984 October	Sweden ^a	Several hundred	Mussels	>170 MU/kg
1984 –1985	Norway ^b	~ 400	Blue mussels	-
1990 August	Canada ^b	16	Mussels	DTX–1: 100µg/100g
1990	France ^c	415	Mussels imported from Denmark	OA: 170 µg/100 g
1990	Mexico Gulf ^d	NR	Oysters (<i>Crassostrea virginica</i>)	OA: 0.162 µg/100g
1991 January	Chile ^b	120	Mussels	OA DTX – 1
1991-1994	Ireland ^e	Several hundred	Mussels (<i>Mytilus edulis</i>)	OA: <0.7 µg/g DTX-2: 6.3 µg/g
1993 Summer- 1994 Summer	Italy ^f	NR	Mussels (<i>M. galloprovincialis</i>)	OA: 32–151 µg/100g DTX–1: 0–0.32 µg/100g
1994 February	Portugal ^g	18	Donax clams (<i>Donax trunculus</i>)	OA: 130 µg /100 g
1994 June	UK ^h	2	Mussels	OA
1997 March	UK ⁱ	49	Mussels	OA: 25.3–36.7 µg/100 g
1997 June	UK ^j	55	Mussels	-

1997 February-June	Croatia ^k	NR	Mussels	OA: 0.133 µg/g DTX-2: 0.1 µg/g
1997 March-November	Norway ^l	NR	Blue mussels	OA/DST-1: 349 µg
1998	Slovenia ^m	Several hundred	Blue mussels	ND
1998 September-November	Spain ⁿ	NR	Mussels	OA: 18–1010 µg /100g
2000 January	Greece ^o	>120	Mussels (<i>M. galloprovincialis</i>)	ND
2001 June-July	Portugal ^p	7	Clams, Razor clams, Green crabs	OA: 32 µg/100 g
2002 February	Belgium ^q	403	Blue mussels imported from Denmark	OA: 22.9 µg/100g
2002 Summer	Norway ^r	~200	Brown crabs (<i>Cancer pagurus</i>)	OA: 29 µg/100g DTX-2: 2 µg/100g
2002 September	Portugal ^s	32	Blue mussels	OA: 1328–1862 µg/100g
2002 August-2004 March	Denmark ^t	NR	Surf clams – Blue mussels	OA: 4.3–163.1 µg/100g
2003 February	Greece ^u	NR	Mussels (<i>M. galloprovincialis</i>)	OA: 0.10-0.20 µg/g
2003 May-October	China ^v	NR	Clams, Scallops Gastropods	OA: 3.2-17.5 µg/100g
2003 December	Australia ^w	NR	Oysters	DSTs total: 0.253

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

				mg/kg
2004 April	Chile ^x	NR	<i>Mytilus chilensis</i>	DTX-1:19 µg/100 g

NR=Not Reported, ND=Not Determined, OA= Okadaic acid, DTX = Dinophysitoxin,
DSTs=Diarrhetic Shellfish Toxins

^a Krogh et al. 1985.

^b Aune and Yndestad 1993.

^c Van Egmond et al. 2004.

^d Dickey et al. 1992.

^e Carmody et al. 1996

^f Draisci et al. 1995.

^g Vale and Sampayo 1999.

^h Anon. 1994.

ⁱ Scoging and Bahl 1998.

^j Anon. 1997.

^k Pavela – Vrančič et al. 2002.

^l Ramstad et al. 2001.

^m Mozetic and Bozic 2001.

ⁿ Moroño et al. 2003.

^o Papadopoulou et al. (present study)

^p Vale and Sampayo 2002.

^q De Schrijver et al. 2002.

^r Torgersen et al. 2005

^s Vale et al. 2003

^t Jorgensen et al. 2005.

^u Ciminiello et al. 2005

^v Wu et al. 2005

^w Madigan et al. 2005

^x Garcia et al. 2004.

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table II.

Characteristic	Values
Demographics	
Male No. (%)	73 (60.8)
Female No. (%)	47 (39.2)
Mean age in years (range)	35 (8-70)
Clinical characteristics	
Mean duration of symptoms in h (range)	5 (0.5-10)
No. (%) of patients with DSP symptoms seeking Hospital assistance	120 (100)
Estimated No. of patients with DSP not seeking medical assistance	>500
Laboratory tests	
No. of patients/no of faecal samples studied	120/120
No. of mussels' samples bought in the same venders as the cases during the onset of the outbreak	2
No. of unconsumed mussels (leftovers) tested	None
No. of mussels' samples bought in the same market as the cases after the end of the outbreak	18

TABLE III.

Sampling Date	Place (one sample/place)	Symptoms observed	Median Mouse survival Time (number of mice dead/number of mice tested)		DSP toxins Result
Outbreak 15/1/2000	Thermaikos Gulf	Apathy, weakness, difficulty in movement, respiratory distress, spastic contractions just before death	1 h 25 min	(3/3)	Positive
Outbreak 19/1/2000	Thermaikos Gulf	Apathy, weakness, difficulty in movement, respiratory distress, spastic contractions just before death	1 h 30 min	(3/3)	Positive
15/4/2000	Thermaikos Gulf	No symptoms	> 48 h	(0/3)	Negative
6/12/2000	Thermaikos Gulf	No symptoms	> 48 h	(0/3)	Negative
19/1/2001	Amvrakikos gulf	Apathy, weakness, difficulty in movement, hypersensitivity, respiratory distress	9 h	(3/3)	Positive
22/2/2001	Thermaikos Gulf Amvrakikos gulf	No symptoms	> 48 h	(0/3)	Negative
23/3/2001	Thermaikos Gulf	Apathy, weakness, difficulty in movement, respiratory distress	5 h	(3/3)	Positive
10/9/2001	Thermaikos Gulf Amvrakikos gulf	No symptoms	> 48 h	(0/3)	Negative
20/12/2001	Thermaikos Gulf	No symptoms	> 48 h	(0/3)	Negative

	Amvrakikos gulf				
15/4/2002	Thermaikos Gulf	No symptoms	> 48 h	(0/3)	Negative
20/7/2002-	Thermaikos Gulf	No symptoms	> 48 h	(0/3)	(0/3)
20/7/2005	Amvrakikos gulf

FIGURE 1.