Allium cepa chromosome aberration and micronucleus tests applied to study genotoxicity of pesticide-treated vegetable extracts

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2006-168.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	03-Nov-2006
Complete List of Authors:	Feretti, Donatella; University of Brescia, Department of Experimental and Applied Medicine Zerbini, Ilaria; University of Brescia, Department of Experimental and Applied Medicine Zani, Claudia; University of Brescia, Department of Experimental and Applied Medicine Ceretti, Elisabetta; University of Brescia, Department of Experimental and Applied Medicine Monarca, Silvano; University of Perugia, Department of Hygiene and Public Health
Methods/Techniques:	Chromatography - HPLC, Health significance, Toxicology
Additives/Contaminants:	Mutagenic compounds, Pesticide residues
Food Types:	Fruit, Vegetables

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

***Allium cepa* chromosome aberration and micronucleus tests applied to study
genotoxicity of extracts from pesticide-treated vegetables and grapes**

Donatella Feretti¹, Ilaria Zerbini¹, Claudia Zani¹, Elisabetta Ceretti¹, Silvano Monarca²

¹Department of Experimental and Applied Medicine, University of Brescia, Viale Europa, 11,
25123 Brescia, Italy (feretti@med.unibs.it);

²Department of Hygiene and Public Health, University of Perugia, Via del Giochetto, 60126
Perugia, Italy (monarca@unipg.it).

Corresponding author:

Prof. Silvano Monarca
Department of Hygiene and Public Health, University of Perugia,
Via del Giochetto, 06126 Perugia ITALY
Tel. +39-075-5857362
Fax +39-075-5857342
E-mail monarca@unipg.it

Field Code Changed

Abstract

The *Allium cepa* assay is an efficient test for chemical screening and *in situ* monitoring for genotoxicity of environmental contaminants. This test has been used widely to study genotoxicity of many pesticides revealing that these compounds can induce chromosomal aberrations in root meristems of *Allium cepa*. Pesticide residues can be present in fruit and vegetables and represent a risk for human health. The mutagenic and carcinogenic action of herbicides, insecticides and fungicides on experimental animals is well known. Several studies have shown that chronic exposure to low levels of pesticides can cause birth defects and that prenatal exposure is associated with carcinogenicity. In this study we evaluated the potential application of plant genotoxicity tests for monitoring mutagens in edible vegetables. The presence of pesticides and genotoxic compounds extracted from twenty-one treated vegetables and eight types of grapes sampled from several markets in Campania, a region in Southern Italy, was monitored concurrently. The extracts were analysed for pesticides by gas chromatography and high-performance liquid chromatography, and for genotoxicity using two plant tests: the micronucleus test and the chromosomal aberration test in *Allium cepa* roots. Thirty-three pesticides were detected, some of which are not approved. Genotoxicity was found in some of the vegetables and grapes tested. *Allium cepa* tests proved to be sensitive in monitoring genotoxicity in food extracts. The micronucleus test in interphase cells gave a much higher mutagenicity than the chromosomal aberration test in anaphase-telophase cells.

Keywords: *Allium cepa* test; genotoxicity; pesticides; vegetables.

1990), in human lymphocytes, in V79 cell line of Chinese hamster and in other organism tests such as fish, crustacea and unicellular algae (Fiskesjö, 1985). Moreover the test can be used to measure also toxicity, studying macroscopic parameters as length of roots, variations in form, colour and consistency of roots, presence of broken root tips, tumors and hooks (Fiskesjö, 1985).

Many mutagenic/carcinogenenic compounds can be present in food and some epidemiological studies have indicated that many dietary factors are related with the development of human cancer (Doll and Peto, 1981). Also pesticide residues can be present in fruit and vegetables and represent a risk for human exposure. Children in particular can be exposed to pesticides by dietary ingestion because they eat more food per body mass than an adult, and their diets differ from those of adults. The children's diet is often rich in foods containing higher levels of pesticide residue, such as juices, fruits and vegetables (Curl et al., 2003). The mutagenic and carcinogenic action of herbicides, insecticides and fungicides on experimental animals is well known. Several studies have shown that chronic exposure to low levels of pesticides can cause birth defects and that prenatal exposure is associated with carcinogenicity (Heeren et al., 2003). Carcinogenicity and neurotoxicity in adults related to pesticides exposure are also reported (Alavanja et al., 2004). Some of these substances have been associated with neuro-developmental toxicity and thyroid hormone imbalance. Many pesticides are recalcitrant and bioaccumulative compounds and they are endocrine disruptors (Vogel, 2005). Toxicological properties, such as mutagenicity, embryotoxicity, teratogenesis and cancerogenesis, are well known for many pesticides (Cueto, 1980; Saleh, 1980; Al-Saleh, 1994; Leiss and Savitz, 1995; De Bertoldi, 1996). Human exposure to agricultural chemicals has been associated with an increase in cancer incidence (Saftlas et al., 1987; Brown et al., 1990; Blair and Zahm, 1991; Cerhan et al., 1998; Settini et al., 2003), even if in some cases only a moderate statistically association is reported (Abdalla et al., 2003). On the other hand studies on the risk assessment for pesticide residues in food suggest a low importance in cancer risk and adverse health effects in humans. The estimate of human risk related to fruit and vegetables

1 consumption must consider that this food is protective for cancer (Block et al., 1992; Swirsky Gold
2 et al., 1997) and concurrently can contain natural substances with genotoxic activity (Ames et al.,
3 1987).
4
5
6
7

8 The extensive use of pesticides can entail risk not only to human health, but also to the environment
9 and non-target organisms. For this reason, according to the European Directive 91/414, the
10 registration of pesticides is regulated in order to avoid harmful effects on human or animal health,
11 and adverse effects on the environment. Moreover pesticides can be move by a diffusive transport
12 which enables their adsorption by soil and a flow transport which cause the movement of
13 compounds from soil surface to groundwater, determining an higher risk for groundwater and
14 drinking water contamination (Guzzella et al., 1996).
15
16
17
18
19
20
21
22

23 The Joint FAO/WHO Meeting on Pesticide Residues (JMPR) has established for food commodities
24 the maximum residue limits (MRLs) of pesticides that can be ingested daily without appreciable
25 human health risk (Environmental Health Criteria, EHC 104, 1990). In Italy, the MRLs are
26 regulated by European Directives 99/65/EC and 99/71/EC. Nevertheless, large quantities of
27 pesticides are released daily into the environment with potential hazards to human genetic material.
28
29
30
31
32
33
34

35 Most pesticides were found capable of inducing mutations in short-term tests (Tezuka et al., 1980;
36 Waters et al., 1982; Plewa et al., 1984; Garrett et al., 1986; Panneerselvam et al., 1995; Ribas et al.,
37 1996; De Marco et al., 2000; Kaya et al., 2000; Zang et al., 2000), although genotoxicity was only
38 present at much higher doses than those utilized in agricultural activities (Rodrigues et al., 1998).
39 140 organophosphate compounds were tested for mutagenic activity in bacteria (*Salmonella*
40 *typhimurium* and *Escherichia coli*) by Hanna and Dyer (1975), who found that 20% of them gave
41 positive mutagenic responses and that this group of chemicals produced base substitutions rather
42 than frame-shift mutations. Out of 228 pesticides under study, 50 were found to be mutagenic in
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

bacterial reversion-assay systems with 5 strains of *Salmonella typhimurium* (TA100, TA98, TA1535, TA1537 and TA1538) and one strain of *Escherichia coli* (WP2 hcr) (Moriya et al., 1983). Hour et al. (1998) carried out a comparative investigation on the mutagenicity of several pesticides using the Ames test (TA100, TA98 and TA1535 strains) and *Salmonella* lactam test (JK947, JK3 and JK1 strains). Six out of 14 pesticides evaluated for genotoxicity using the new strains were highly mutagenic in strain JK3. Dolara et al. (1993, 1994) carried out some studies in Central Italy to determine the toxicological effects of a complex mixture of pesticide residues in common foods. They mixed 15 pesticides, usually detected in food at high concentrations, and then tested this mixture using the *Salmonella*-microsome assay, with and without metabolic activation. No mutagenic activity was observed at concentrations up to 500 mg/plate; however, when they tested the same mixture at concentrations ranging from 0.1 to 20 mg/L on human lymphocytes *in vitro*, they observed a slight but statistically significant increase in sister chromatid exchanges at 1 mg/ml. Bianchi-Santamaria et al. (1997) performed the human lymphocyte micronucleus test on single pesticides and on mixtures of pesticides usually found in the environment and found a weak genotoxicity for benomyl and for three of the four organophosphorous insecticides. However, the various mixtures did not have any synergistic effects.

The cytological effects of individual pesticides have been studied in plants (Ahmed and Grant, 1972) and some studies have been conducted by means of genotoxicity tests: in *Tradescantia* flowers using the *Tradescantia*/stamen hair test and the *Tradescantia*/micronucleus test (Ma et al., 1984; Rodrigues et al., 1998; Mohammed and Ma, 1999), in *Vicia faba* roots (micronuclei and root tip length in *Vicia faba*) (De Kergommeaux et al., 1983; De Marco et al., 2000) and in *Allium cepa* roots (chromosomal aberration in *Allium cepa*) (Fiskesjö et al., 1981; Grant, 1982; Fiskesjö, 1985; Rank and Nielsen, 1993, 1997; Rank et al., 1993; Grover and Satwinderjeet Kaur, 1999). Higher plants provide a valuable genetic assay system for screening and monitoring environmental pollutants (Grant et al., 1994) and comparable results, in term of genetic abnormalities, are obtained

in plant and animal systems (Grant et al., 1981). These studies were performed on single or complex mixtures of pesticides, but a few data on the genotoxicity of residues extracted from fruit and vegetables are yet available.

The aim of this research was to evaluate the potential application of plant genotoxicity test on monitoring of mutagens in food vegetable. The mutagenic activity of extracts of fruit and vegetables used for human consumption was studied using two plant tests in *Allium cepa* roots, the *Allium cepa* micronucleus test, and the *Allium cepa* chromosome aberration test and concurrently the presence of pesticides in these residues by gas-chromatography or high-performance liquid chromatography (HPLC) was analysed.

Materials and methods

We investigated 21 samples of vegetables (beet, cucumber, tomato, cherry tomato, lettuce, round aubergine, pepper) and 8 types of grapes grown in Campania, a region in Southern Italy, and bought from several greengrocer's shops. The samples were extracted for pesticide analyses and for genotoxicity tests.

Fruit and vegetable sampling and pesticide extraction

The samples for the pesticide analyses and for the genotoxicity tests were prepared following two different methods. The method used in the pesticide analyses consisted of the following stages:

1. Sample preparation. Unwashed parts of vegetables and grapes were sliced and whisked to obtain a homogeneous mixture.
2. Residue extraction. A total of 10 g of homogenized sample was mixed with 12 g of diatomaceous earth and placed in an empty 60 ml column. After 10 minutes the column was eluted with 100 ml of dichloromethane.
3. Extract purification or, if chlorophyll was present, on SCX (ion-exchange column). The eluates dried by rotary evaporation at 40°C were re-suspended in 2 ml of

dichloromethane/acetone solution (80:20) and charged on a cartridge of Florisil (60 ml) previously washed with 5 ml of dichloromethane/acetone solution (80:20) for activation, and the eluates were recharged twice on the same type of cartridge (Dagna et al., 1993).

4. Extract purification in presence of chlorophyll. The extracts dried by rotary evaporation and re-suspended in 2 ml of methanol were charged on a cartridge of SCX (60 ml) previously washed with 5 ml of methanol for activation, and the eluate was recharged twice on the same type of cartridge.
5. Gas chromatography and HPLC analysis of the residues. The final eluates were dried by rotary evaporation and re-suspended in 2 ml acetone/hexane solution (1:1) for gas chromatography analysis or in 2 ml water/acetonitrile solution (70:30) for HPLC analysis.

Sample extraction of samples for genotoxicity test was performed as follows:

1. Sample preparation. Unwashed parts of vegetables and grapes were sliced and whisked to obtain a homogeneous mixture.
2. Sample extraction with diatomaceous earth in dichloromethane. 10 g of homogenized sample were mixed with 12 g of diatomaceous earth and then placed in an empty 60 ml column. After 10 minutes the column was eluted with 100 ml of dichloromethane and the eluates were then dried by rotary evaporation at 40°C (Dagna et al., 1993). The extraction was carried out on a total of 50 g of sample.
3. Genotoxicity tests on the extracts. The dried extracts were dissolved in dimethylsulphoxide (DMSO) and diluted in distilled water and analysed for mutagenicity by means of plant tests.

Analysis of the extracts

Pesticide analysis.

Instrumental analysis of the extracts was performed by gas chromatography (Manninen et al., 1987) and by HPLC (Ambrus et al., 1981). Gas chromatography analysis was performed using a

Shimadzu GC17A system (Shimadzu Corporation, Tokyo, Japan) which operates with a bi-channel system: after filtering on a 0.45 µm filter, each extract was injected (1 µl) simultaneously into two columns with different polarities (ZB-5 30 m x 0.32 mm ID x 0.25 µm film and ZB-1701 30 m x 0.32 mm ID x 0.25 µm film) (Phenomenex, Torrance, CA, USA) and connected with electron capture detector (ECD) and nitrogen-phosphorus detector (NPD). This allowed one pesticide to be detected by both columns at the same time, thus avoiding the possibility of overlapping of different pesticides. An ECD detector was used to identify organochloride pesticides, pyrethroids, dicarboximides and thiophthalimides; an NPD detector was used to identify phosphorates, triazoles and acylanines. All these analyses were performed using pesticide standards (Ehrenstorfer GmbH, Augsburg, Germany).

Quantitative gas chromatography was performed using a multiple internal standard (Series M). The temperature programme consisted of an initial temperature of 50°C held for 2 minutes, followed by an increase to 150°C at a rate of 25°C/min and to 270°C at 4°C/min (10 minutes) (Manninen et al., 1987).

High-performance liquid chromatography is the most appropriate method to determine certain types of compounds, such as ureic compounds, growth regulators, benzimidazole and carbamates. Most of these compounds present heat sensitivity and low volatility due to their high molecular weight and/or polarity, leading to long retention on GC columns. Hence, these molecules are not amenable to analysis by gas chromatography. HPLC analysis was performed on a Shimadzu LC10 system (Shimadzu Corporation, Tokyo, Japan) equipped with reverse-phase separation columns packed with 5 µm particles and with pH stability from 2 to 10. The latter parameter is particularly important considering the wide range of molecules used nowadays as pesticides.

Analyses of benzimidazole and other pesticides were carried out with 250 x 4.6 mm ID Luna 5 μ m Phenyl-Hexyl and 5 μ m C₁₈(2) columns (Phenomenex, Torrance, CA, USA) and a DAD detector, whereas for carbamate analysis, separation on 250 x 4.6 mm ID 5 μ m C₈(2) Luna columns (Phenomenex, Torrance, CA, USA) was followed by post-column derivatization with NaOH and o-phthalaldehyde and fluorometric detection. The flow rate was 1 ml/min and quantitative analysis was performed using an external standard. The samples were extracted, purified and dried, and then suspended in a 70:30 water-acetonitrile solution, filtered using a 0.45 μ m cellulose filter and injected (100 μ l) in HPLC.

Toxicity and genotoxicity tests on *Allium cepa* roots.

A phytotoxicity assay was performed prior to the genotoxicity tests on some extracts in order to determine the non-cytotoxic doses. The assay consisted of a 96-hour root toxicity test using *Allium cepa* bulbs grown in extract solutions in distilled water of three kinds of samples (tomato, aubergine, grapes) at concentrations of 225 mg/L, 75 mg/L and 25 mg/L.

Commercial onion bulbs of *Allium cepa* of equal size were carefully cleaned and washed without destroying the primordial roots. For each sample, six onions were exposed to different concentrations of extracts, and natural mineral water known without mutagenic activity was used as a negative control. On three consecutive days the lengths of the roots were measured and macroscopic parameters were considered. The concentrations used caused no visible cytotoxic effects and therefore they were also used for the genotoxicity tests (Fiskesjö, 1985, 1995).

For the plant genotoxicity tests, dried extract equivalent to 50 g of vegetables or grapes was dissolved in 2 ml of DMSO and diluted with distilled water to 225 mg/L, 75 mg/L and 25 mg/L. These solutions were used to analyse micronucleus frequency in interphase and also chromosome aberrations in anaphase-telophase *Allium cepa* root tip cells (Grant, 1982; Rank and Nielsen, 1993,

1997; Rank et al., 1993; Ma et al., 1995; Grover and Satwinderjeet Kaur, 1999). The *Allium cepa* bulbs were put in natural mineral water for pre-germination until the root length was 1.5-2 cm; the roots were then exposed to the extract solutions for 48 hours. Each extract was tested with a series of five onions. After exposure, the root tips were fixed in Carnoy's solution (ethyl alcohol and acetic acid, 3:1) for 24 hours, immediately for chromosome aberrations or after 48 hours' recovery time in mineral water for micronuclei analysis (Ma et al., 1995). After fixation, the roots were transferred to 70% ethyl alcohol and stored in a refrigerator until examined. The root tips were placed in 1N HCl at 60°C for 10 minutes and stained with Feulgen reagent for 1 hour. The microscopic analysis included mitotic index and the recording of micronuclei in interphase cells and of aberrant cells in anaphase-telophase cells. For each extract, the mitotic index was determined by examining 5000 cells from 5 separate slides and calculated as mitotic cells per 1000 cells. The micronuclei were scored in 4000 cells per onion from 4 separate slides and expressed as micronuclei/1000 cells (data were reported as mean $\pm$ standard deviation). Analysis of variance (ANOVA) and Dunnett's *t* test were used to determine the significance of the difference between the samples and negative control at the 5% significance level.

For the *Allium cepa* aberration test, bridges, fragments, laggards, c-mitosis and multipolar anaphases were scored. For each slide, 40 cells in anaphase-telophase were examined, and with 5 of the onions 800 cells per concentration were scored (data were expressed as percentages on 800 cells at least). Statistical analysis was performed by means of a test for the linear trend and the Fisher test.

Each plant test had a negative control with natural mineral water and DMSO 1% and a positive control with potassium dichromate ($K_2Cr_2O_7$, 1.8 μ l/ml) for the *Allium cepa* micronuclei test and sodium azide (NaN_3 , 0.3 μ g/ml) for the aberration test with a 48-hours exposure.

Results

Chemical analyses of the pesticides

The chemical analyses enabled us to identify pesticides with a wide-polarity range in the samples of the vegetables and grapes examined (Table I). The pesticides found in the extracts consisted of various chemical compounds: carbamates (carbaryl, propamocarb), organophosphate compounds (acephate, chlorpyrifos, ethopropos, dimethoate, parathion-methyl, omethoate, azinphos-methyl), benzonitriles (chlorothalonil), amides (iprodione, vinclozolin), acylalanines (methalaxyl), phthalates (folpet) and pyrethroids (fluvalinate, cypermethrin).

[insert Table I here]

Eleven out of twenty-one (52.2%) vegetable samples did not comply with the current law on pesticide residues; eight (38.1%) contained concentrations of pesticides over the MLRs, one (4.8%) contained pesticides not allowed for the type of crop, and one (4.8%) was in breach of the law for both reasons.

Five out of eight grape samples (62.5%) contained concentrations of pesticides over the MLRs.

Only two samples (beet and pepper) contained no pesticide residues.

Allium cepa tests

Table II shows the micronuclei frequencies ($\% \pm SD$) in *Allium cepa* root tip cells after 48 hours of exposure to three doses of vegetable and grape extracts (225 mg/L, 75 mg/L and 25 mg/L) versus a negative control. Only two vegetable samples (9.5%) were negative in the *Allium cepa* micronuclei test and were concurrently not contaminated by pesticides, whereas nineteen vegetable samples (90.5%) were positive in this test. 73.7% of the vegetable samples showed a positive response at all three doses, 15.8% showed a positive response at high and medium doses, and 10.5% were positive only at highest doses.

[insert Table II here]

1 All the grape samples gave positive results in the micronuclei test at all three tested doses at
2
3 the 1% significance level ($p<0.01$).
4

5 Tables IIIa, IIIb and IIIc show the frequency of aberrant cells (%) and the mitotic index in
6
7 *Allium cepa* root tip cells exposed for 48 hours to the vegetable and grape extracts (225 mg/L, 75
8 mg/L, 25 mg/L) versus the negative control.
9
10 [insert Table IIIa here]
11

12 According to the Fisher test, seven vegetable samples (33.3%) were positive using the
13
14 chromosomal aberration test: two were positive at all tested doses, one at high and medium doses,
15
16 and four at the highest dose only (Tables IIIa and IIIb).
17
18 [insert Table IIIb here]
19

20 About the grapes only one was not able to induce chromosomal aberrations in root cells of
21
22 *Allium cepa*. Seven grape samples (87.5%) gave positive results inducing an increase in the
23
24 frequency of aberrations: one at all tested doses, three at high and medium tested doses, and three at
25
26 the highest dose (Table IIIc).
27
28 [insert Table IIIc here]
29

30 In the linear trend analysis, five vegetable samples (sample 8 at $p<0.001$, samples 5 and 7
31
32 at $p<0.01$ and samples 4 and 6 at $p<0.05$) and only one grape sample (sample 2 at $p<0.05$) gave a
33
34 positive response, showing a clear dose-response relation.
35
36
37

38 **Discussion and conclusions**

39 The presence of pesticides and genotoxic compounds extracted from 21 treated vegetables and 8
40
41 grape samples bought in the markets of Campania, a region in Southern Italy, was assessed by
42
43 means of chemical and genotoxicological analysis. Gas chromatography and HPLC analysis
44
45 showed the presence of many pesticides in the vegetable and grape extracts. Sometimes the
46
47 pesticide residues exceeded the maximum concentrations allowed by Italian law. Many vegetable
48
49
50
51
52
53
54
55
56
57
58
59
60

extracts showed chromosomal aberrations and/or micronuclei formation in *Allium cepa* roots, presumably due to the presence of pesticide residue and/or other unknown components.

As previously reported, many pesticides are effective clastogens in plant cells. The *Allium cepa* tests have been successfully carried out in previous research for *in situ* monitoring water samples. This is the first application of the *Allium cepa* tests in food extracts, and the results suggest their applicability for this purpose, especially when using the micronucleus test in interphase cells, which gave much higher mutagenicity than the chromosomal aberration test in anaphase-telophase cells. Micronucleus frequency was much higher compared to the negative control, with a dose-response relation in some cases. The lowest dose of extracts (25 mg/L) induced an increase of micronuclei in most of the vegetable samples and in all the grape samples. Fewer samples showed chromosomal aberrations. Some positive samples (vegetable samples 4, 5, 6, 8 and 17 and grape sample 2) showed a dose-response relationship. The only two vegetable samples resulted without pesticide residues have not induced increase of micronuclei frequency and chromosomal aberrations too (samples 1 and 14).

The higher sensitivity of the *Allium cepa* micronuclei test compared to the chromosomal aberration test in *Allium cepa* can be explained by the fact that chromosomal aberrations are more closely associated with DNA damage (bridges, laggards), whereas the induction of micronuclei in root meristems of *Allium cepa* is the manifestation of chromosome damage and disturbance of the mitotic process. The micronucleus is formed by a new membrane developing around the chromatin matter that failed to move to either pole during the anaphase of mitosis. Such chromatin matter arises either from anomalous disjunction of chromosomes due to spindle abnormalities or from breakage of chromosomes resulting in the formation of acentric fragments, dicentric chromosomes and chromatin bridges. Therefore, the induction of micronuclei suggests that the pesticide residues in the vegetables may have been either spindle inhibitors or clastogens. Recently a research has

1 shown in about 7,000 subjects that an increase of micronucleus frequency in lymphocytes predicts
2
3 the risk of carcinogenesis in human (Bonassi et al., in press).
4
5

6
7 The micronuclei test in *Allium cepa* was positive for both the vegetable and grape extracts
8
9 contaminated by pesticides outlawed in Italy and for samples contaminated by legal pesticides in
10
11 Italy. This agrees with the hypothesis of potential synergistic activity between the components of a
12
13 chemical mixture used in agriculture. The only two negative vegetable samples were not
14
15 contaminated by pesticides. This is in agreement with considerations by Ennever et al. (1988), who
16
17 describe the *Allium* test as one of the most sensitive tests as it produces low percentages of false
18
19 negatives. Like all other tests on plants, it seems to be less specific because it gives a high degree of
20
21 false positives. With reference to the significance of micronuclei test in plants for human health
22
23 hazards it is recognized that if a chemical compound causes damage to chromosomes in plant cells,
24
25 it can be potentially dangerous also for the chromosomes of mammalian cells.
26
27

28 The genotoxic effects detected may be the result of the synergistic activity of the different pesticide
29
30 residues present in the vegetable and grape extracts, even if many natural substances in plant
31
32 material might be genotoxic (Ames et al., 1987). In this study unwashed vegetables and grapes were
33
34 used, but it would be useful to test thoroughly washed vegetables and grapes to evaluate if the
35
36 genotoxicity is really related to pesticide residues in food. In fact washing processes remove a large
37
38 amount of residues, and ensure a large margin of safety (Jayakrishnan et al., 2005; Radwan et al.,
39
40 2005; Lentza-Rizos et al., 2006).
41
42

43 Moreover gas-chromatography and HPLC analyses looked for only some pesticides in vegetables
44
45 and grapes, usually utilized in agricultural practice, and this fact can explain the variability of the
46
47 results because other compounds not considered here could be present in the extracts which were
48
49 tested.
50
51

The *Allium cepa* test appears to be a fast, inexpensive and easy to handle method, and, as other short-term mutagenicity tests, is particularly suitable to analyze complex mixtures of contaminants and to reveal synergistic effects. For these reasons this chemical/biological approach permits to study different but complementary aspects of food contamination and could be helpful to better evaluate cancer hazards related to diet.

Acknowledgements

This paper is dedicated to the memory of the conceivers of the research, Domenica Biscardi and Raffaele De Fusco, memorable friends and researchers of the University of Naples and Salerno. The authors express their deepest appreciation to Dr. Giovanni Petrone of D'Aniello Laboratory of Sant'Egidio Montalbino, Salerno, Italy, for his assistance in this research.

References

Abdalla MH, Gutierrez-Mohamed ML, Farah IO. 2003. Association of pesticide exposure and risk of breast cancer mortality in Mississippi. *Biomedical Sciences Instrumentation* 39: 397-401.

Ahmed M, Grant WF. 1972. Cytological effects of pesticides Phosdrin and Bladex on *Tradescantia* and *Vicia faba*. *Canadian Journal of Genetics and Cytology* 14: 157-165.

Alavanja MC, Hoppin JA, Camel F. 2004. Health effects of chronic pesticide exposure: cancer and neurotoxicity. *Annual Review of Public Health* 25: 155-197.

Al-Saleh IA. 1994. Pesticide: a review article. *Journal of Environmental Pathology, Toxicology and Oncology* 13: 151-161.

Ambrus A, Lantos J, Visi E, Csatos I, Sarvari L. 1981. General method for determination of pesticides residues in samples of plant origin, soil and water. I. Extraction and cleanup. *Journal of Association of Official Analytical Chemists* 64: 733-742.

Ames BN, Magaw R, Swirsky Gold L. 1987. Ranking possible carcinogenic hazards. *Science* 27: 271-280.

Bianchi-Santamaria A, Gobbi M, Cembran M, Arnaboldi A. 1997. Human lymphocyte micronucleus genotoxicity test with mixtures of phytochemicals in environmental concentrations. *Mutation Research* 388: 27-32.

Blair A, Zahm SH. 1991. Cancer among farmers. *Occupational Medicine* 6: 335-354.

Block G, Patterson B, Subar A. 1992. Fruit, vegetables, and cancer prevention: a review of the epidemiological evidence. *Nutrition and Cancer* 17: 1-29.

Brown LM, Blair A, Gibson R, Everett GD, Cantor KP, Schuman LM, Burmeister LF, Van Lier SF, Dick F. 1990. Pesticide exposures and other agricultural risk factors for leukemia among men in Iowa and Minnesota. *Cancer Research* 50: 6585-6591.

Bonassi S, Znaor A, Ceppi M, Lando C, Chang WP, Holland N, Kirsch-Volders M, Zeiger E, Ban S, Barale R, Bigatti MP, Bolognesi C, Cebulska-Wasilewska A, Fabianova E, Fucic A,

Hagmar L, Joksic G, Martelli A, Migliore L, Mirkova E, Scarfi MR, Zijno A, Norppa H, Fenech M. An increase micronucleus frequency in peripheral blood lymphocytes predicts the risk of cancer in humans. *Carcinogenesis*, in press.

Cabrera GL, Rodriguez DMG. 1999. Genotoxicity of soil from farmland irrigated with wastewater using three plant bioassays. *Mutation Research* 426: 211-214.

Cerhan JR, Cantor KP, Williamson K, Lynch CF, Torner JC, Burmeister LF. 1998. Cancer mortality among Iowa farmers: recent results, time trends, and lifestyle factors. *Cancer Causes and Control* 9: 311-319.

Cueto C Jr. 1980. Consideration of the possible carcinogenicity of some pesticides. *Journal of Environmental Science and Health - Part B: Pesticides, Food Contaminants and Agricultural Wastes* 15: 949-975.

Curl CL, Fenske RA, Elgethun K. 2003. Organophosphorus pesticide exposure of urban and suburban preschool children with organic and conventional diets. *Environmental Health Perspectives* 111 (3): 377-382.

Dagna L, Gasparini G, Icardi ML, Sesia E. 1993. Metodo multiresiduo di estrazione su colonna per la determinazione di residui di fitofarmaci in matrici vegetali e liquide. (Multiresidue analysis method based on column extraction of residue of pesticide in vegetables and liquid samples). *Bollettino dei Chimici Igienisti* 44: 383-397.

De Bertoldi M. 1996. Genotoxic effects of pesticides. *European Journal of Cancer Prevention* 5: 397-399.

De Kergommeaux DJ, Grant WF, Sandhu SS. 1983. Clastogenic and physiological response of chromosomes to nine pesticides in the *Vicia faba* in vivo root tip assay system. *Mutation Research* 124: 69-84.

1 De Marco A, De Salvia R, Polani S, Ricordy R, Sorrenti F, Perticone P, Cozzi R, D'Ambrosio C,
2
3 De Simone C, Guidotti M, Albanesi T, Duranti G, Festa F, Gensabella G, Owczarek M.
4
5 2000. Evaluation of genotoxic and cytotoxic properties of pesticides employed in Italian
6
7 agricultural practices. *Environmental Research* 83: 311-321.
8
9 Dolara P, Torricelli F, Antonelli N. 1994. Cytogenetic effects on human lymphocytes of a mixture
10
11 of fifteen pesticides commonly used in Italy. *Mutation Research* 325: 47-51.
12
13 Dolara P, Vezzani A, Caderni G, Coppi C, Torricelli F. 1993. Genetic toxicity of a mixture of
14
15 fifteen pesticides commonly found in the Italian diet. *Cell Biology and Toxicology* 9: 333-
16
17 343.
18
19 Doll R, Peto R. 1981. The causes of cancer: quantitative estimates of available risks of cancer in the
20
21 United States today. *Journal of the National Cancer Institute* 66: 1191-1308.
22
23 Ennever FK, Andreano G, Rosenkranz HS. 1988. The ability of plant assays to predict
24
25 carcinogenicity. *Mutation Research* 205: 99-105.
26
27 Fiskesjö G. 1985. The Allium test as a standard in environmental monitoring. *Hereditas* 102: 98-
28
29 112.
30
31 Fiskesjö G. 1995. Allium test. *Methods in Molecular Biology* 43: 119-127.
32
33 Fiskesjö G, Lassen C, Renberg L. 1981. Chlorinated phenoxyacetic acids and chlorophenols in the
34
35 modified Allium test. *Chemico-Biological Interactions* 34: 333-344.
36
37 Garrett NE, Stack HF, Waters MD. 1986. Evaluation of the genetic activity profiles of 65
38
39 pesticides. *Mutation Research* 168: 301-325.
40
41 Grant WF. 1982. Chromosome aberration assay in Allium. A report of U.S. Environmental
42
43 Protection Agency Gene-Tox Program. *Mutation Research* 99: 273-291.
44
45 Grant WF. 1994. The present status of higher plant bioassay for the detection of environmental
46
47 mutagens. *Mutation Research* 310: 175-185.
48
49
50
51
52
53
54
55
56
57
58
59
60

- Grant WF. 1999. Higher plant assays for the detection of chromosomal aberrations and gene mutations – a brief historical background on their use for screening and monitoring environmental chemicals. *Mutation Research* 426: 107-112.
- Grant WF, Sandhu SS, De Serres FJ, Gopalau HN, Svendsgaard D, Veleminsky J, Becking GC. 1994. Environmental monitoring for genotoxicity with plant system. Results and recommendations. *Mutation Research* 310: 257-263.
- Grant WF, Zinov'eva-Stahevitch AE, Zura KD. 1981. Plant genetic test system for the detection of chemical mutagens. In: Stich HF, San RHC, editors. *Short Term Test for Chemical Carcinogens*, Springer Verlag, New York, pp. 200-216.
- Grover IS, Dhingra AK, Neeta A, Ladhar SS. 1990. Genotoxicity of pesticides and plant systems. In: Mendelsohn LM, Albertini RJ editors. *Mutation and the Environment, Part E*, Wiley-Liss, New York, pp. 91-106.
- Grover IS, Satwinderjeet Kaur. 1999. Genotoxicity of wastewater samples from sewage and industrial effluent detected by the *Allium* root anaphase aberration and micronucleus assays. *Mutation Research* 426: 183-188.
- Guzzella L, De Paolis A, Bartone C, Pozzoni F, Giuliano G. 1996. Migration of pesticides residues from agricultural soil to groundwater. *International Journal of Environmental Analytical Chemistry* 65: 261-275.
- Hanna PJ, Dyer KF. 1975. Mutagenicity of organophosphorus compounds in bacteria and *Drosophila*. *Mutation Research* 28: 405-420.
- Heeren GA, Tyler J, Mandeya A. 2003. Agricultural chemical exposures and birth defects in Cape Province, South Africa. A case-control study. *Environmental Health: a Global Access Science Source* 2: 11-19.
- Hour TC, Chen L, Lin JK. 1998. Comparative investigation on the mutagenicities of organophosphate, phthalimide, pyrethroid and carbamate insecticides by the Ames and lactam tests. *Mutagenesis* 13: 157-166.

Kaya B, Creus A, Yanikoglu A, Cabre O, Marcos R. 2000. Use of the *Drosophila* wing spot test in the genotoxicity testing of different herbicides. *Environmental and Molecular Mutagenesis* 36: 40-46.

Jayakrishnan S, Dikshit AK, Singh JP, Pachauri DC. 2005. Dissipation of Lambda-cyhalothrin on tomato (*Lycopersicon esculentum* Mill.) and removal of its residues by different washing processes and steaming. *Bulletin of Environmental Contamination and Toxicology* 75: 324-328.

Leiss JK, Savitz DA. 1995. Home pesticide use and childhood cancer: a case-control study. *American Journal of Public Health* 85: 249-252.

Lenza-Rizos C, Avramides EJ, Kokkinaki K. Residues of azoxystrobin from grapes to raisins. 2006. *Journal of Agricultural and Food Chemistry* 54: 138-141.

Ma TH, Harris MM, Anderson VA, Ahmed K, Mohammed KB, Bare JL, Lin G. 1984. Tradescantia-micronucleus (Trad-MCN) tests on 140 health-related agents. *Mutation Research* 138: 157-167.

Ma TH, Xu Z, Xu C, McConnell H, Vailterra Rabago E, Arreola AG, Zhang H. 1995. The improved Allium/Vicia root tip micronucleous assay for clastogenicity of environmental pollutants. *Mutation Research* 334: 185-195.

Manninen A, Kuitunen ML, Julin L. 1987. Gas chromatographic properties of the M series of universal retention index standards and their application to pesticide analysis. *Journal of Chromatography* 394: 465-471.

Monarca S, Feretti D, Collivignarelli C, Guzzella L, Zerbini I, Bertanza G, Pedrazzani R. 2000. The influence of different disinfectants on mutagenicity and toxicity of urban wastewater. *Water Research* 34: 4261-4269.

Monarca S, Feretti D, Zerbini I, Alberti A, Zani C, Resola S, Gelatti U, Nardi G. 2002a. Soil contamination detected using bacterial and plant mutagenicity tests and chemical analyses. *Environmental Research* 88: 64-69.

- Monarca S, Richardson SD, Feretti D, Grottolo M, Thruston AD, Zani C, Navazio G, Ragazzo P, Zerbini I, Alberti A. 2002b. Mutagenicity and disinfection by-products in surface drinking water disinfected with peracetic acid. *Environmental Toxicology and Chemistry* 21: 309-318.
- Monarca S, Rizzoni M, Gustavino B, Zani C, Alberti A, Feretti D, Zerbini I. 2003. Genotoxicity of surface water treated with different disinfectants using in situ plant tests. *Environmental and Molecular Mutagenesis* 41: 353-359.
- Mohammed KB, Ma TH. 1999. Tradescantia micronucleus and – stamen hair mutation assays on genotoxicity of the gaseous and liquid forms of pesticides. *Mutation Research* 426: 193-199.
- Moriya M, Ohta T, Watanabe K, Miyazawa T, Kato K, Shirasu Y. 1983. Further mutagenicity studies on pesticides in bacterial reversion assay systems. *Mutation Research* 116: 185-216.
- Panneerselvam N, Sinha S, Shanmugam G. 1995. Genotoxicity of the herbicide fluchloralin on human lymphocytes in vitro: chromosomal aberration and micronucleus tests. *Mutation Research* 344: 69-72.
- Plewa MJ, Wagner ED, Gentile GJ, Gentile JM. 1984. An evaluation of the genotoxic properties of herbicides following plant and animal activation. *Mutation Research* 136: 233-245.
- Radwan MA, Abu-Elamayem MM, Shiboob MH, Abdel-Aal A. 2005. Residual behaviour of profenofos on some field-grown vegetables and its removal using various washing solutions and household processing. *Food and Chemical Toxicology* 43: 553-557.
- Rank J, Nielsen MH. 1993. A modified Allium test as a tool in the screening of genotoxicity of complex mixtures. *Hereditas* 118: 49-53.
- Rank J, Nielsen MH. 1997. Allium cepa anaphase-telophase root tip chromosome aberration assay on N-methyl-N-nitrosourea, maleic hydrazide, sodium azide, and ethyl methanesulfonate. *Mutation Research* 390: 121-127
- Rank J, Jensen AG, Skov B, Pedersen LH, Jensen K. 1993. Genotoxicity testing of the herbicide Roundup and its active ingredient glyphosate isopropylamine using the mouse bone marrow

micronucleus test, Salmonella mutagenicity test, and *Allium* anaphase-telophase test. Mutation Research 300: 29-36.

Ribas G, Surrallés J, Carbonell E, Xamena N, Creus A, Marcos R. 1996. Genotoxicity of the herbicides alachlor and maleic hydrazide in cultured human lymphocytes. Mutagenesis 11: 221-227.

Rodrigues GS, Pimentel D, Weinstein LH. 1998. In situ assessment of pesticide genotoxicity in an integrated pest management program I - *Tradescantia* micronucleus assay. Mutation Research 412: 235-244.

Saftlas AF, Blair A, Cantor KP, Hanrahan L, Anderson HA. 1987. Cancer and other causes of death among Wisconsin farmers. American Journal of Industrial Medicine 11: 119-129.

Saleh MA. 1980. Mutagenic and carcinogenic effects of pesticides. Journal of Environmental Science and Health - Part B: Pesticides, Food Contaminants and Agricultural Wastes 15: 907-927.

Settimi L, Masina A, Andron A, Axelson O. 2003. Prostate cancer and exposure to pesticides in agricultural settings. International Journal of Cancer 104: 458-461.

Smaka-Kincl V, Stegnar P, Lovka M, Toman MJ. 1996. The evaluation of waste, surface and ground water quality using the *Allium* test procedure. Mutation Research 368: 171-179.

Steinkellner H, Mun-Sik K, Helma C, Ecker S, Ma TH, Horak O, Kundi M, Knasmüller S. 1998. Genotoxic effects of heavy metals: Comparative investigation with plant bioassay. Environmental Molecular and Mutagenesis 31: 183-191.

Swirsky Gold L, Stern BR, Slone TH, Brown JP, Manley NB, Ames BN. 1997. Pesticide residues in food: investigation of disparities in cancer risk estimates. Cancer Letters 117: 195-207.

Tezuka H, Ando N, Suzuki R, Terahata M, Moriya M, Shirasu Y. 1980. Sister-chromatid exchanges and chromosomal aberrations in cultured Chinese hamster cells treated with pesticides positive in microbial reversion assays. Mutation Research 78: 177-191.

- 1 Vogel JM. 2005. Perils of paradigm: complexity, policy design, and the endocrine disruptor
2 screening program. *Environmental Health: a Global Access Science Source* 4: 2-13.
- 3
4 Waters MD, Sandhu SS, Simmon VF, Mortelmans KE, Mitchell AD, Jorgenson TA, Jones DC,
5
6 Valencia R, Garrett NE. 1982. Study of pesticide genotoxicity. *Basic Life Sciences* 21: 275-
7
8 326.
- 9
10 Zang Y, Zhong Y, Luo Y, Kong ZM. 2000. Genotoxicity of two novel pesticides for the earthworm,
11
12 *Eisenia fetida*. *Environmental Pollution* 108: 271-278.
- 13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table I. Pesticides found in the vegetable and grape samples.

Sample no	Vegetables	Pesticides (mg kg ⁻¹) ^a
1	Beet	nd
2	Beet	Endosulfan (0.87); Methalaxyl (1.02); Deltamethrin (0.12)
3	Cucumber	Dichloran (0.6); Procymidone (0.02); Folpet (0.07)
4	Lettuce	Dichloran (0.1); Endosulfan (0.03); Propamocarb (3.69); Chlorothalonil (0.12)
5	Lettuce	Dichloran (0.04); Procymidone (0.01); Vinclozolin (0.02); λ-Cyhalothrin (0.04)
6	Lettuce	Dichloran (0.02); Endosulfan (0.04); Propamocarb (12.6)
7	Roman lettuce	Procymidone (0.04); Delthamethrin (0.01); Endosulfan (0.08); Chlorothalonil (0.44)
8	Roman lettuce	Dichloran (0.24); Dichlofluanid (0.03); Iprodione (2.97); Chlorothalonil (0.08); Bifenthrin (1.10)
9	Roman Lettuce	Propamocarb (5.93); Iprodione (4.83); Cymoxanil (5.69)
10	Round aubergine	Dichloran (0.01); Iprodione (0.11); Chlozoline (0.01)
11	Round aubergine	Dichloran (0.12); Procymidone (0.03); Ethoprophos (1.35)
12	Round aubergine	Dichloran (0.02); Iprodione (0.73); Azinphos-methyl (0.04); Acrinathrin (0.05)
13	Pepper	Acephate (0.85); Chlorothalonil (0.09)
14	Pepper	Nd
15	Pepper	Fenarimol (0.01); Fluvalinate (0.02); Chlorothalonil (0.10)
16	Sweet pepper	Procymidone (0.40); Ethoprophos (0.64)
17	Sweet pepper	Dimethoate (0.17); Acephate (15.3); Omethoate (0.71); λ-Cyhalothrin (0.11)
18	Sweet pepper	Endosulfan (0.07); Iprodione (1.16)
19	Cherry tomato	Chlorothalonil (0.04); Dichlofluanid (0.02); Procymidone (0.02)
20	Tomato	Dichloran (0.09); Procymidone (0.03); Vinclozolin (0.09)
21	Tomato	Procymidone (0.16); Iprodione (1.51); Fenarimol (0.10); Fluvalinate (0.41)
Sample No	Grapes	Pesticides (mg kg ⁻¹)
1	Italian grapes	Dimethoate (0.10); Vinclozolin (0.92); Procymidone (1.44); Fenarimol (0.03); Hexaconazole (0.12); Myclobutanil (0.27)
2	Italian grapes	Parathion-ethyl (0.06); Folpet (0.10); Procymidone (0.44); Cypermethrin (0.04)
3	Italian grapes	Procymidone (0.03); Endosulfan (0.07)
4	Italian grapes	Parathion-methyl (0.16); Procymidone (1.58); Endosulfan (0.24); Bromopropylate (0.03); Iprodione (7.55); Pyrimethanil (3.10); Fenithrothion (0.75)
5	Italian grapes	Parathion-ethyl (0.16); Folpet (1.10); Procymidone (0.17); Acephate (0.04); Myclobutanil (0.56)
6	Italian grapes	Parathion-ethyl (0.11); Folpet (1.03); Procymidone (0.14); Acephate (0.05); Myclobutanil (0.49)
7	Italian grapes	Pyrimethanil (0.30); Parathion-ethyl (0.05); Procymidone (0.25); Bromopropylate (0.22); Omethoate (0.26)
8	Palierian grapes	Omethoate (0.19); Carbaryl (0.57); Parathion-ethyl (0.07); Iprodione (0.44); Procymidone (0.92); Chlorpyrifos-ethyl (0.23)

^aThe values in bold are the residues found in concentrations exceeding the maximum values permitted by Italian law or the pesticides not allowed for that food product under the same law (99/65/EC and 99/71/EC)

nd = not detected

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table II. Mean frequency (%±SD) of micronuclei in *Allium cepa* root tips exposed to pesticide-treated vegetables and grapes for 48 hours.

Samples	Negative control	Micronucleus frequency (% ± SD)		
		225 mg/L	75 mg/L	25 mg/L
Vegetables				
1	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
2	0.0 ± 0.0	3.9 ± 1.1**	2.6 ± 0.6**	1.5 ± 0.5**
3	0.2 ± 0.7	2.6 ± 1.2**	2.6 ± 1.2**	0.8 ± 0.7
4	0.1 ± 0.2	3.7 ± 0.9**	3.3 ± 1.2**	2.2 ± 1.4**
5	0.1 ± 0.3	4.4 ± 1.1**	3.5 ± 0.9**	1.9 ± 0.7**
6	0.5 ± 0.7	4.6 ± 3.8**	3.5 ± 1.4**	1.0 ± 0.9
7	0.0 ± 0.0	4.2 ± 2.0**	0.4 ± 0.6	0.3 ± 0.6
8	0.2 ± 0.4	2.6 ± 2.5**	0.7 ± 1.0	0.4 ± 0.7
9	0.2 ± 0.4	4.2 ± 2.2**	2.6 ± 1.4**	1.7 ± 1.4**
10	0.2 ± 0.4	5.4 ± 0.9**	4.2 ± 1.4**	2.3 ± 0.7**
11	0.5 ± 0.6	4.3 ± 1.8**	1.9 ± 0.9**	1.3 ± 0.7
12	0.0 ± 0.0	3.8 ± 0.7**	2.7 ± 0.9**	1.5 ± 0.6**
13	0.6 ± 0.6	4.7 ± 2.1**	4.1 ± 0.7**	2.7 ± 0.9**
14	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
15	0.2 ± 0.7	5.1 ± 1.2**	2.7 ± 0.9**	1.9 ± 0.9**
16	0.2 ± 0.7	3.8 ± 0.9**	3.7 ± 1.2**	2.6 ± 1.2**
17	0.1 ± 0.2	4.6 ± 0.8**	3.8 ± 0.8**	1.5 ± 0.6**
18	0.1 ± 0.2	7.4 ± 1.4**	4.3 ± 0.8**	2.3 ± 1.0**
19	0.1 ± 0.2	4.0 ± 0.7**	4.1 ± 1.0**	4.0 ± 0.7**
20	0.3 ± 0.6	5.8 ± 0.9**	3.6 ± 0.8**	1.8 ± 0.7**
21	0.2 ± 0.4	5.5 ± 0.9**	4.4 ± 1.2**	1.6 ± 0.7**
Grapes				
1	0.0 ± 0.0	4.0 ± 0.8**	3.8 ± 0.6**	1.4 ± 0.5**
2	0.0 ± 0.0	4.0 ± 0.6**	2.3 ± 0.9**	1.5 ± 0.5**
3	0.0 ± 0.0	4.3 ± 0.7**	2.2 ± 0.8**	1.5 ± 0.5**
4	0.0 ± 0.0	4.4 ± 0.7**	3.4 ± 0.5**	2.2 ± 0.8**
5	0.0 ± 0.0	4.7 ± 0.7**	3.5 ± 0.5**	1.1 ± 0.7**
6	0.0 ± 0.0	3.2 ± 0.8**	2.6 ± 0.7**	1.4 ± 0.5**
7	0.0 ± 0.0	3.5 ± 0.6**	3.5 ± 0.7**	1.1 ± 0.8**
8	0.0 ± 0.0	4.2 ± 0.7**	3.1 ± 0.7**	1.5 ± 0.6**

** Statistically significant versus negative control according to Dunnett's *t* test (*p* < 0.01)
Positive control (K₂Cr₂O₇, 1.8 µg/ml): 6.3 ± 1.5%

Table IIIa. Mitotic index and anaphase aberrations in *Allium cepa* roots exposed to the vegetable extracts (samples 1-10) for 48 hours.

Samples	Negative control (%)		Doses (mg/L)	Mitotic index (%)	Aberrant cells ^a (%)	Statistical tests	
	Mitotic index (%)	Aberrant cells (%)				Fisher's exact test	Linear trend
1	21.6	0.0	225	20.2	0.0	ns	ns
			75	20.8	0.0	ns	
			25	20.8	0.0	ns	
2	21.6	0.0	225	20.8	0.6	ns	ns
			75	20.4	0.6	ns	
			25	20.8	0.4	ns	
3	21.6	0.0	225	14.2	0.0	ns	ns
			75	15.6	0.0	ns	
			25	15.2	0.0	ns	
4	21.4	0.0	225	15.4	0.8	*	*
			75	13.8	0.6	ns	
			25	13.3	0.0	ns	
5	21.4	0.0	225	13.0	0.6	ns	**
			75	23.0	0.0	ns	
			25	13.7	0.0	ns	
6	19.7	0.0	225	18.2	2.5	***	*
			75	19.8	1.4	***	
			25	20.6	1.0	**	
7	19.7	0.0	225	14.8	1.1	**	ns
			75	17.0	0.8	*	
			25	14.6	0.8	*	
8	19.7	0.0	225	16.6	1.4	***	***
			75	16.8	0.5	ns	
			25	16.0	0.0	ns	
9	19.7	0.0	225	16.6	0.1	ns	ns
			75	16.9	0.0	ns	
			25	16.4	0.0	ns	
10	20.8	0.0	225	15.8	0.6	ns	ns
			75	16.0	0.5	ns	
			25	15.8	0.3	ns	

*Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.05$)

**Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.01$)

***Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.001$)

ns = not significant

^aPositive control (NaN₃, 0.3 µg/ml): 5.5%

Table IIIb. Mitotic index and anaphase aberrations in *Allium cepa* roots exposed to the vegetable extracts (samples 11-21) for 48 hours.

Samples	Negative control (%)		Doses (mg/L)	Mitotic index (%)	Aberrant cells ^a (%)	Statistical tests	
	Mitotic index (%)	Aberrant cells (%)				Fisher's exact test	Linear trend
11	20.8	0.0	225	15.8	0.0	ns	ns
			75	15.6	0.0	ns	
			25	16.6	0.0	ns	
12	20.8	0.0	225	20.0	0.6	ns	ns
			75	20.0	0.5	ns	
			25	19.8	0.4	ns	
13	20.8	0.0	225	17.2	1.1	**	ns
			75	18.4	0.4	ns	
			25	18.4	0.5	ns	
14	20.8	0.0	225	19.4	0.0	ns	ns
			75	19.6	0.0	ns	
			25	19.4	0.0	ns	
15	20.8	0.0	225	15.8	0.0	ns	ns
			75	14.0	0.0	ns	
			25	15.6	0.0	ns	
16	22.0	0.0	225	18.2	0.0	ns	ns
			75	17.0	0.0	ns	
			25	18.2	0.0	ns	
17	22.0	0.0	225	16.0	0.9	*	**
			75	13.9	0.4	ns	
			25	16.0	0.0	ns	
18	22.0	0.0	225	17.0	0.0	ns	ns
			75	16.2	0.0	ns	
			25	15.6	0.0	ns	
19	22.8	0.0	225	11.3	0.4	ns	ns
			75	13.2	0.3	ns	
			25	11.6	0.0	ns	
20	22.8	0.0	225	16.4	1.3	*	ns
			75	16.3	0.8	*	
			25	13.8	0.5	ns	
21	22.8	0.0	225	20.2	0.5	ns	ns
			75	21.0	0.5	ns	
			25	21.0	0.1	ns	

*Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.05$)

**Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.01$)

***Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.001$)

ns = not significant

^aPositive control (NaN₃, 0.3 µg/ml): 5.5%

Table IIIc. Mitotic index and anaphase aberrations in *Allium cepa* roots exposed to the grape extracts for 48 hours.

Samples	Negative control (%)		Doses (mg/L)	Mitotic index (%)	Aberrant cells ^a (%)	Statistical tests	
	Mitotic index (%)	Aberrant cells (%)				Fisher's exact test	Linear trend
1	20.6	0.0	225	20.4	0.6	ns	ns
			75	20.8	0.6	ns	
			25	20.8	0.5	ns	
2	20.6	0.0	225	21.6	0.8	*	*
			75	20.1	0.8	*	
			25	20.8	0.0	ns	
3	20.6	0.0	225	21.0	0.8	*	ns
			75	21.2	0.4	ns	
			25	20.5	0.3	ns	
4	20.6	0.0	225	20.2	1.0	**	ns
			75	19.6	0.8	*	
			25	19.6	0.8	*	
5	21.0	0.0	225	21.2	1.0	**	ns
			75	21.6	0.5	ns	
			25	19.6	0.5	ns	
6	21.0	0.0	225	21.0	0.8	*	ns
			75	20.8	0.6	ns	
			25	22.0	0.4	ns	
7	21.0	0.0	225	19.3	1.4	***	ns
			75	20.2	1.0	**	
			25	18.4	0.6	ns	
8	21.0	0.0	225	20.0	0.8	*	ns
			75	21.0	0.9	*	
			25	20.2	0.4	ns	

*Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.05$)

**Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.01$)

***Statistically significant versus negative control according to Fisher's exact and linear trend tests ($p < 0.001$)

ns = not significant

^aPositive control (NaN₃, 0.3 µg/ml): 5.5%