Determination of aflatoxin B1 in tiger nut based soft drinks

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2005-170.R1
Manuscript Type:	Original Research Paper
Methods/Techniques:	Chromatography - HPLC
Additives/Contaminants:	Mycotoxins – aflatoxins
Food Types:	Beverages

SCHOLARONE™
Manuscripts

Determination of aflatoxin B₁ in tiger nut based soft drinks

ISABEL ARRANZ ¹, JOERG STROKA ¹ and MICHAEL NEUGEBAUER ²

¹ Food Safety & Quality Unit, Institute for Reference Materials and Measurements, European Commission – Joint Research Centre, Retieseweg, B-2440 Geel (Belgium).

² Pharmazeutisches Institut der Universitaet Bonn, Kreuzbergweg 26, D-53115 Bonn (Germany)

Formatted: Font: 10 pt, Complex Script Font: 10 pt

Formatted: German Germany

Abstract

An analytical method for the determination of Aflatoxin B₁ (AfB₁) in a tiger nut based soft drinks named 'horchata' is described. The method is based on an immunoaffinity clean-up, followed by HPLC separation and fluorescence detection after electrochemical post-column derivatisation (PCD). The detection limit (S/N = 3) and the quantification limit (S/N = 10) were 0.02 µg/kg and 0.06 µg/kg respectively. The mean recovery at a level of 2 µg/L was 88 % (n = 6) and the coefficient of variation (CV) 9 %. The method was applied to conduct a small market survey for a beverage named 'horchata' that is frequently consumed by all kind of population in Southern Europe. A total of 22 samples from Spanish and Belgian supermarkets were analysed. As a result only one sample was found to contain AfB₁ at the LOQ of the method.

Deleted: The described

Deleted: allows

Deleted: kg

Deleted: l

Deleted:

Deleted: 8.

Deleted: children

Deleted: below

Deleted: D

Deleted: 0.06 µg/kg

Keywords

Aflatoxin B₁, tiger nuts, 'chufa', 'horchata', High-performance Liquid-Chromatography.

INTRODUCTION

Tiger nut or 'chufa' is a plant that botanically belongs to the genus *Cyperus esculentus* L. There are two varieties with different applications: *Cyperus esculentus* L. var. *esculentus* and

Deleted: N

Deleted: "

Deleted: "

Cyperus esculentus L. var. *sativus*. The second one is cultivated for human consumption and the rhizomes are used to obtain the tiger nuts. (Fact Sheet HS-583)

In Europe, the cultivation of tiger nuts is widely spread through the eastern part of Spain (Region of Valencia). Tiger nuts are also grown in Central Africa and in the southern part of United States (Florida).

The Spanish annual consumption of tiger nuts is about 5-7 million of kg/year, and around 3 million kg were imported during 2004. Exporting countries were Nigeria, Niger, Burkina Faso, Benin, Mali and Ghana. The main destination of the tiger nuts in Spain is the production of 'horchata' (40-55 millions litres/year), a soft drink consumed by all kind of population including children (with special importance due to their low body weight).

'Horchata' is either available as liquid, hailstorm or in congealed consistency. It is considered mainly as a refreshing drink, but is also as nutritive due to its caloric contents. After collection of the tiger nuts, they are washed and sorted in order to separate the product from soil, insect- or otherwise damaged tiger nuts that can be present at the collection. This is an important step for minimizing possible aflatoxin contamination present in the raw material and it is nowadays performed with the help of industrial machines. In the next step the tiger nuts are dried to reduce the water content from 50 to 11 %, this is the last step in the processing and it should be performed carefully since mycotoxin producing micro organisms can still start to grow on not sufficiently dried tiger nuts.

For the preparation of 1 L of 'horchata' approx. 200 g of tiger nuts and 150 g of sugar are used. This product must be distinguished from another different type of 'horchata', that is based on rice and vanilla, is made in Central and South America. (Mosquera et al. 1996).

Tiger nuts are also used for the production of oil for human consumption and they are an important animal feed ingredient in the southern United States (Kelley and Fredricksin 1991)

Deleted: Tiger nuts are also used ... [2]
Deleted: For its production
Formatted ... [3]
Deleted: , collected tiger nuts
Deleted: are
Deleted: them from rests of sand
Deleted: and
Formatted ... [4]
Formatted ... [5]
Deleted:
Deleted: other materials
Deleted:also mouldy and bit ... [6]
Deleted: reducing
Deleted: the content of
Formatted ... [7]
Deleted: s
Deleted: in
Formatted ... [8]
Deleted: ated
Formatted ... [9]
Deleted: T
Deleted: is the
Deleted: and
Deleted: drying in which the aim is
Deleted: ied
Deleted:
Formatted ... [10]
Deleted: .
Deleted: T
Deleted: slow and
Deleted: is the critical one due to
Deleted: there is a
Deleted: the
Deleted: risk
Deleted: of contamination with
Deleted: that remaining
Deleted:
Deleted: that
Formatted ... [11]
Deleted: occur
Deleted:
Formatted ... [12]
Deleted: should be wash and le ... [13]
Deleted: , next is to blend this ... [14]
Formatted ... [15]
Deleted: The process ends with ... [16]
Formatted ... [17]
Deleted: A
Formatted ... [18]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In 1996 Bankole and Esegbe recognized tiger nut as one of the commodities susceptible to aflatoxin contamination and detected aflatoxins in 35% of tiger nut samples collected from Nigeria with concentrations ranging from 10 to 120 µg/kg. Adebajo (1993) also reported the presence of aflatoxin in tiger nut at toxicological unsafe levels. Aflatoxins are considered one of the most significant mycotoxins, they are natural toxins produced by fungi affecting several food and feed commodities. They are classified as carcinogen to humans by the International Agency for Research on Cancer (IARC 1993; JECFA 1996). In Europe, regulations on aflatoxins exist for various products but not for tiger nuts or products thereof.

EU maximum levels for AfB₁ and total aflatoxins are 2 µg/kg and 4 µg/kg, respectively for most food products (Van Egmond 2004). Generally in the EU, imported food products from third countries are controlled as they enter the Community and in cases of consumer health risks, EU Member States can exchange information on contamination cases in the so called “Rapid Alert System for Food and Feed” (RASFF) (http://europa.eu.int/comm/food/food/rapidalert/index_en.htm). In 2002, three notifications were made about contamination levels of aflatoxin in EU imported tiger nuts. And in April 2004, one notification was made about tiger nuts with a contamination level of 300 µg/kg AfB₁ that were imported from Mali (http://europa.eu.int/comm/food/food/rapidalert/reports/week15_en.pdf).

To our knowledge no method has been reported for the analysis of AfB₁ in ‘horchata’. The here-described method has been developed on the basis of a previously described one by Stroka et al (2000). A small survey on commercially available ‘horchata’ from Spanish and Belgian supermarkets was conducted after method development.

- Deleted: A different type of horchata, based on rice and vanilla, is made in Central and South America.
- Deleted: class I human
- Formatted: Not Highlight
- Deleted: ,
- Deleted: for AfB₁ and
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: L
- Deleted: kg
- Deleted: respectively
- Deleted: ng/g
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: L
- Deleted: total aflatoxins in
- Deleted: The EU strategy for consumer protection (Commission Regulation No. 178/2002)
- Deleted: considers controls throughout the entire food and feed chain.
- Deleted: aims to provide a methodology for the determination of actions to protect health with the use of three components of risk communication (risk assessment, risk management, and risk communication).
- Deleted: (Regulation EC/178/2002).
- Deleted: As a result of this strategy
- Deleted: ,
- Deleted: , products
- Deleted: Generally food products
- Formatted: Not Highlight
- Deleted: import
- Deleted: s that are imported from third countries to the EU are carried out on the product as it enters the Community
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: while
- Deleted: . In cases
- Deleted: within the EU these controls are performed throughout the entire food and feed chain
- Field Code Changed
- Deleted: The determination is performed by HPLC with fluorescent detection and the need of transforming AfB₁ in a nonquenchable derivat(... [19]
- Deleted: ,
- Deleted: 2002).
- Deleted: The market

MATERIALS AND METHOD

Samples

The survey was carried out on 'horchata', from supermarkets in Spain and Belgium (imported product from Spain) with production dates from 2004. The method has been tested for 8 brands and different batches of some of these brands with a total of 22 samples.

Reagents

Solvents and reagents. Solvents used were complying with grade ACS-ISO. Acetonitrile and methanol were HPLC grade from Merck (Darmstadt, Germany). Immunoaffinity columns type EASI-EXTRACT™ aflatoxins were from Rhone-rBiopharm (Glasgow, Scotland).

Water. Purified water was delivered by the Millipore water purification device Milli-Q Gradient A10 (Bedford, USA).

Deleted: a

Deleted: (

Deleted: Bedford,

Phosphate-buffered saline (PBS). PBS was prepared by adding 0.2 g KCl, 0.2 g KH₂PO₄, 1.16 g anhydrous Na₂HPO₄, and 8.0 g NaCl in 1.0 L water. The pH was adjusted to 7.4 with NaOH c=(0.1 mol/L).

Standards. AFB₁ standard solution with a concentration of 10.0 µg/mL dissolved in chloroform was obtained from the Dutch National Institute for Public Health and the Environment, RIVM (Bilthoven, The Netherlands). From this, a dilution with a final concentration of 0.10 µg/mL solution was used for calibration standards purposes. Portions of 125, 250, 500, 750 and 1000 µL from this last standard were evaporated in 250 mL

Deleted:

Deleted: , p

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

volumetric flasks, afterwards re-dissolved with 75 mL of methanol and filled up with water.

The following concentrations were achieved: 0.05, 0.10, 0.20, 0.30, 0.40 in ng/mL.

For recovery experiments, 5 mL of a 0.40 $\mu\text{g/mL}$ AfB₁ standard solution was used to fortify 1 L of blank matrix the day prior to analysis, resulting in a level of fortification of 2 $\mu\text{g/L}$.

Apparatus.

HPLC system. The HPLC equipment consisted of a combination of Gilson HPLC pump type 309 with a solid phase extraction unit ASPEC XL (Milano, Italy), with a 100 μL injection loop. The fluorescence detector was from Waters type 474 (Milano, Italy). The PCD with KOBRA cell was from Rhone -rBiopharm (Glasgow, Scotland). The column used was a Supelcosil LC-18, 25 cm x 4.6 mm, 5 μm .

Sample preparation and clean-up.

25 mL of PBS and 25 mL of the 'horchata' were mixed and filtered over glass micro fibre filter (Whatman GF/A), 25 mL of this filtrate were passed through an immunoaffinity column (IAC) at a flow rate of approximately 3 mL/min. The IAC was washed twice with 10 mL of water at the same speed. For elution of AfB₁, 0.5 mL methanol was applied by gravity on the IAC. The eluate was collected in a volumetric flask of 5.0 mL. After 1 min, a second portion of 0.5 mL methanol was applied. This step was repeated once more after another minute and air was pressed through the IAC to collect most of the applied methanol. The volumetric flask, containing the combined portions of IAC eluate was filled up to approx. 4.5 mL, shaken and filled up further with water to the mark and shaken again. The clear extracts were directly injected into the HPLC system.

HPLC equipment and reverse phase conditions

Deleted: calculations

Deleted: the

Deleted: n

Deleted: the

Deleted: ces

Deleted: extraction

Deleted: . The

Deleted: contamination

Deleted: was

Deleted: l

Deleted: ¶

Deleted: Phosphate-buffered saline (PBS). PBS was prepared by adding 0.2 g KCl, 0.2 g KH₂PO₄, 1.16 g anhydrous Na₂HPO₄, and 8.0 g NaCl in 1.0 L water. The pH was adjusted to 7.4 with NaOH c=(0.1 mol/L).¶

Deleted: Extraction procedure

The detector was set at 360 nm (excitation wavelength) and 420 nm (emission wavelength). The mobile phase was water/acetonitrile/methanol [62 + 22 + 16 (v+v+v)] with 120 mg of KBr and 350 µL of HNO₃ (4 mol/L) per litre of mobile phase. The flow rate was at 1 mL/min. Injection volume was 100 µL.

RESULTS AND DISCUSSION

It could be shown, that the clean-up procedure by immunoaffinity chromatography after dilution of the sample with PBS and glass micro fibre filtration is fast, simple and reliable despite a solid residue content of the product of a required minimum of 12 % and a fat content of around 2 % in 'horchata' drinks. Chromatograms were free of interferences in the region of the AFB₁ peak (see Figure 1).

“[Insert Figure 1 here]”

The mean recovery of the method was 88 % (n = 6). These recovery experiments were performed using blank samples spiked at a level of 2 µg/L of AFB₁ in the product. This shows that the method procedure is suitable to purify and concentrate the analyte by immunoaffinity from the matrix and that the matrix has no significant effect on the recovery during clean up. Additional recovery experiments at lower levels were not carried out as results from collaborative studies and own experience showed that recovery figures for the here used methodology (immunoaffinity clean-up with HPLC-fluorescence detection) and this type of analyte (aflatoxins) are rather stable in terms of fortification levels, while being more influenced by the matrix used for analysis (Arranz et al. 2005, Dragacci et al. 2001 and Stroka et al. 2000, 2001, 2003).

Deleted: applied

Deleted: is

Deleted: are

Deleted: f

Deleted: about

Deleted: kg

Deleted: [CollStudiesAf

Formatted: Highlight

Deleted:].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The resulting relative standard deviation was 9 % under repeatability conditions. On the basis of the assumption that the internal reproducibility is linked by a factor of 3/2 with the internal repeatability, the estimate for the internal reproducibility is 14%.

Based on a “fitness for purpose approach” which is characterised by the determination of an acceptable precision parameter that is derived from the function given below, the calculated reproducibility of 14% from this study was compared with the limit calculated from the function below. This approach has recently been implemented in EU legislation on mycotoxin method performance requirements (Commission Directive No. 38/2005) and can be used for those cases where no fully validated method precision (full collaborative trial) is available.

$$U_f = \sqrt{(LOD / 2)^2 + (\alpha \times C)^2}$$

In this formula U_f is the maximum acceptable standard uncertainty in $\mu\text{g/kg}$, LOD is the Limit of Detection expressed in $\mu\text{g/kg}$, α is a factor (0.1 – 0.2) that depends on the concentration of interest (C in $\mu\text{g/kg}$). For concentration below 50 $\mu\text{g/kg}$, α is 0.2. The resulting maximum acceptable standard uncertainty for a target level of 2 $\mu\text{g/L}$ (assuming a liquid density of 1 g/mL for ‘horchata’) and a LOD of 0.02 $\mu\text{g/L}$ is 0.40 $\mu\text{g/L}$. Expressed as relative standard deviation this figure is 20 %.

As the internal reproducibility is mostly accepted as standard uncertainty, the calculated figure of 14% shows that the method can be considered as “fit for purpose” according to this scheme.

In none of the tested 22 sample materials AfB₁ could be quantified. Only in one case the presence of AfB₁ could be detected at the level of the estimated LOQ (Figure 1).

CONCLUSION

- Deleted: (RSD)
- Deleted: obtained precision
- Deleted: in
- Deleted: this “sing
- Deleted: le laboratory validation”
- Deleted: is currently discussed for use in future
- Deleted: [Directive
- Formatted: Not Highlight
- Deleted:]
- Deleted: to be
- Formatted: Not Highlight
- Deleted: criteria
- Deleted: are
- Deleted: ¶
- Deleted: kg
- Deleted: l
- Deleted: kg
- Deleted: l
- Deleted: kg
- Deleted: l
- Deleted: kg
- Deleted: 0
- Deleted: kg
- Deleted: This is equivalent to a maximum allowed repeatability of
- Deleted: RSD
- Deleted: The experimentally obtained RSD of 9 % is below the calculated minimum repeatability requirement
- Deleted: showing
- Deleted: identified
- Deleted: slightly above the
- Deleted: at a level of 0.07 $\mu\text{g/kg}$
- Deleted: at a contamination level
- Deleted: below
- Deleted: of 0.06 $\mu\text{g/kg L}$
- Deleted:
- Deleted: ¶

A rather simple and fast method for the determination of AfB₁ in 'horchata' was developed and validated under single laboratory conditions. It could be demonstrated that the described method shows good performance characteristics when compared with minimum requirement data calculated from a "fitness for purpose" function that is currently discussed for use in EU legislation on mycotoxin methodology in official food control. The described method therefore is a good candidate for such use.

Based on this small survey, no indication was found for a consumer risk from the 'tiger nut' based soft drink as sold on the Spanish and Belgian market during the period 2004. However, a survey with more samples would be necessary to obtain a full picture, which was partially difficult as there is a limited amount of producers in Spain.

Deleted: carried out with this new method

Acknowledgements

The authors are grateful to the RIVM in Bilthoven (The Netherlands) for providing us with the aflatoxin standard and to the Regulating Council of Denomination of Origin "Chufa de Valencia" (CRDO) in Valencia (Spain) for providing us with some technical information about the 'horchata' consumption in Spain.

Deleted: National Institute for Public Health and the Environment (

Deleted:)

Deleted: "

Deleted: "

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Adebajo LO 1993. Survey of aflatoxins and ochratoxin A in stored tubers of *Cyperus esculentus*. *Mycopathologia* 124, 41-46.

2. Arranz I, Sizoo E, van Egmond E, Kroeger K, Legarda TM, Burdaspal P, Reif K and Stroka J. 2005. Mini-collaborative study for the determination of Aflatoxin B₁ in medical herbs. *Journal of AOAC International*. (Accepted).

3. Bankole SA and Esegbe DA. 1996. Occurrence of mycoflora and aflatoxin in marketed tiger nut in Nigeria. *Crop Research* 11 (2): 219-223.

4. Commission Directive (ECC) No. 38/2005 of 6 June 2005 laying down the sampling methods and the methods of analysis for the official control of the levels of *Fusarium* toxins in foodstuffs. Available from: http://europa.eu.int/eur-lex/pri/en/oj/dat/2002/l_031/l_03120020201en00010024.pdf Via the INTERNET. Accessed 2005 October 04.

5. Dragacci S, Grosso F and Gilbert J. 2001. Immunoaffinity column cleanup with liquid chromatography for determination of aflatoxin M₁ in liquid milk: collaborative study. *Journal of AOAC International*. Vol. 84: 437-43.

6. Fact Sheet HS-583. Horticultural Sciences Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Available from: <http://edis.ifas.ufl.edu/MV050>. Via the INTERNET. Accessed 2005 September 26.

7. International Agency for Research on Cancer, IARC Monographs on the evaluation of carcinogenic risk to humans, World Health Organisation, Lyon, Vol. 56, 1993.

Deleted: <#>Agriculture quality policy for the "Protected Designation of Origin" of "Chufa de Valencia". [internet]
Available from: .¶

Formatted: Bullets and Numbering

Formatted: English U.K.

Formatted: Indent: Before: 0 pt, Hanging: 17.85 pt, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 pt + Tab after: 18 pt + Indent at: 0 pt, Tabs: Not at 262.8 pt + 489.6 pt

Formatted: Not Highlight

Formatted: English U.K., Not Highlight

Formatted: Not Highlight

Formatted: English U.K.

Formatted: Italian Italy

Formatted: Bullets and Numbering

Deleted: ,

Deleted: <#>Code of Practice for the Prevention and Reduction of Mycotoxin Contamination in Cereals, including annexes on Ochratoxin A, Zearalenone, Fumonisin and Tricothecenes, CAC/RCP 51-2003. Available from: http://www.ipfsaph.org/servlet/BinaryDownloadServlet?filename=kopool_data/codex_0/en_cxc_051e.pdf. Via the INTERNET. Accessed 2005 October 04. ¶

Formatted: Highlight

Formatted: Indent: Before: 0 pt, Hanging: 17.85 pt, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 pt + Tab after: 18 pt + Indent at: 0 pt

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic

Formatted: Not Highlight

Deleted: <#>Commission Regulation (ECC) No. 178/2002 of 28 January 2002 laying down the general principle(... [20]

Field Code Changed

Formatted: English U.K.

Formatted: English U.K.

Field Code Changed

Formatted: English U.K.

Formatted: English U.K.

Field Code Changed

Formatted: English U.K.

Formatted: English U.K.

Formatted: English U.K.

Formatted: Bullets and Numbering

8. JECFA. 1996. Summary of evaluations performed by the Joint FAO/WHO Expert Committee on Food Additives (JECFA) 1956-1995, first through forty-fourth meetings. Geneva, FAO/WHO, ILSI Press.

9. Kelley JR and Fredricksin LH. 1991. Chufa biology and management. United States. Department of Interior. Fish and Wildlife Leaflet 13.4.18: 1-6.

10. Legal basis of the Rapid Alert System for Food and Feed" (RASFF). Available from: http://europa.eu.int/comm/food/food/rapidalert/index_en.htm. Via the INTERNET. Accessed 2005 June 07.

11. Mosquera LA, Sims CA, Bates RP and O'Keefe SF 1996. Flavor and Stability of "Horchata De Chufas". Journal of Food Science. Volume 61, No. 4.

12. RASFF on week 2005/14. Available from: http://europa.eu.int/comm/food/food/rapidalert/reports/week15_en.pdf. Via the INTERNET. Accessed 2005 June 07.

13. Stroka J, Ankla E, Joerissen U. and Gilbert J. 2000. Immunoaffinity Column Cleanup with Liquid Chromatography Using Post-Column Bromination for Determination of Aflatoxins in Peanut Butter, Pistachio Paste, Fig Paste, and Paprika Powder: Collaborative Study. Journal of AOAC International. Vol. 83: 320-340.

14. Stroka J, Ankla E, Joerissen U. and Gilbert J. 2001. Determination of Aflatoxin B₁ in Baby Food (Infant Formula) by Immunoaffinity Column Cleanup Liquid Chromatography Using Post-Column Bromination: Collaborative Study. Journal of AOAC International. Vol. 84: 1116-1123.

15. Stroka J, von Holst C, Ankla E, and Reutter M, 2003. Immunoaffinity Column Cleanup with Liquid Chromatography Using Post-Column Bromination for Determination of Aflatoxin B₁ in Cattle Feed: Collaborative Study. Journal of AOAC International. Vol. 86: 1179-86.

Deleted: ,

Deleted: ,

Formatted: English U.K.

Formatted: English U.K., Not Highlight

Formatted: English U.K.

Formatted: English U.K., Not Highlight

Deleted:

Formatted: English U.K.

Deleted: <#>Papadopoulou – Bouraoui A, Stroka J and

Deleted: <#> ,

Deleted: <#>Ankla E. 2002. Comparison of two post column derivatization systems, ultraviolet irradiation and electrochemical determination, for the liquid chromatographic determination of aflatoxins in food. Journal of AOAC International. Vol. 85. No. 2: 411-416.

Deleted: <#>The Regulating Council of Denomination of Origin "Chufa de Valencia [internet]. Available from: <http://www.chufadevalencia.org/>. Accessed 2005 June 07.

Formatted: English U.K.

Formatted: Not Highlight

Deleted: , Ankla E, Joerissen U, Gilbert J.

Deleted: ,

Deleted: ,

Formatted: Bullets and Numbering

Deleted: 1

Formatted: German Germany, Not Highlight

Formatted: Bullets and Numbering

Formatted: German Germany, Not Highlight

Formatted: German Germany

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

16. The Regulating Council of Denomination of Origin "Chufa de Valencia [internet].
Available from: <http://www.chufadevalencia.org/>. Accessed 2005 June 07.

17. Van Egmond H. 2004. Natural toxins: risks, regulations and the analytical situation in
Europe, Review. Analytical and Bioanalytical Chemistry 378: 1152-1160.

Formatted: Indent: Before: 0 pt, Hanging: 17.85 pt, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 pt + Tab after: 18 pt + Indent at: 0 pt, Tabs: 262.8 pt, Centered + 489.6 pt, Right

Deleted: ¶
<#>¶

Deleted: Literature to add for collstudy:¶
Dragacci (M1 in milk)¶
Stroka (animal feed)¶
Stroka (baby food)¶
Arranz (medical herbs)¶

For Peer Review Only

Page 2: [1] Deleted arranis 9/26/2005 5:03:00 PM

chufa

Page 2: [1] Deleted arranis 9/26/2005 5:02:00 PM

Kilos

Page 2: [1] Deleted arranis 9/26/2005 3:16:00 PM

K

Page 2: [1] Deleted arranis 9/26/2005 5:22:00 PM

Importing

Page 2: [1] Deleted arranis 9/26/2005 5:24:00 PM

mainly

Page 2: [2] Deleted arranis 10/5/2005 9:12:00 AM

Tiger nuts are also used for the production of oil for human consumption and they are an important animal feed ingredient in the southern United States (Kelley and Fredricksin 1991)

In Spain, the quality of the tiger nuts must fulfil the specifications of the agriculture quality policy of the European Union and the name of the end product "Chufa de Valencia" is a product defined according to the conditions of a Protected Designation of Origin ((ECC) No. 2081/91)

The quality of tiger nuts depends on the terrain used for cultivation. Therefore, it has been reported that also the quality and flavour of horchata is influenced by the production area (Mosquera et al. 1996)

The end product

Page 2: [2] Deleted arranis 9/26/2005 3:18:00 PM

h

Page 2: [3] Formatted strokjo 10/6/2005 12:09:00 PM

Not Highlight

Page 2: [4] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [4] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [5] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [6] Deleted	strokjo	10/6/2005 11:47:00 AM
,		
Page 2: [6] Deleted	strokjo	10/6/2005 11:47:00 AM
also mouldy and bitten by insects tiger nuts should be eliminated.		
Page 2: [7] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [8] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [9] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [10] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [11] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [12] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [13] Deleted	strokjo	10/6/2005 11:52:00 AM
should be wash and leave to soak during at least 12 hours, afterwards they should get		
wash again and mixed with		
Page 2: [14] Deleted	strokjo	10/6/2005 11:52:00 AM
, next is to blend this mixture and to sieve it		
Page 2: [15] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [16] Deleted	strokjo	10/6/2005 11:53:00 AM
The process ends with the addition of 1 L of water, the beverage should be served cold.		
Page 2: [17] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		
Page 2: [18] Formatted	strokjo	10/6/2005 12:09:00 PM
Not Highlight		

Page 3: [19] Deleted strokjo 10/6/2005 12:44:00 PM

The determination is performed by HPLC with fluorescent detection and the need of transforming AfB₁ in a nonquenchable derivative is solved by PCD involving electrochemical bromination (Kobra cell) (Papadopoulou-Bouraoui et al.

Page 9: [20] Deleted strokjo 10/6/2005 12:03:00 PM

Commission Regulation (ECC) No. 178/2002 of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. Available from: [Regulation EC/178/2002](http://ec.europa.eu/food/food/food_safety_and_hygiene/regulation_178_2002_en.htm). Via the INTERNET. Accessed 2005 June 07.

Figure 1:

Chromatogram from a horchata sample containing aflatoxin B₁ at the estimated LOQ (0.06 µg/L).