Generation of recombinant pestiviruses using a full-genome amplification strategy

Thomas Bruun Rasmussen, Ilona Reimann, Åse Uttenthal, Immanuel Leifer, Klaus Depner, Horst Schirrmeier, Martin Beer

► To cite this version:

Thomas Bruun Rasmussen, Ilona Reimann, Åse Uttenthal, Immanuel Leifer, Klaus Depner, et al.. Generation of recombinant pestiviruses using a full-genome amplification strategy. *Veterinary Microbiology*, 2010, 142 (1-2), pp.13. 10.1016/j.vetmic.2009.09.037 . hal-00578396

HAL Id: hal-00578396

<https://hal.science/hal-00578396>

Submitted on 20 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Generation of recombinant pestiviruses using a full-genome amplification strategy

Authors: Thomas Bruun Rasmussen, Ilona Reimann, Åse Uttenthal, Immanuel Leifer, Klaus Depner, Horst Schirrmeier, Martin Beer

PII: S0378-1135(09)00454-4
DOI: doi:10.1016/j.vetmic.2009.09.037
Reference: VETMIC 4594

To appear in: *VETMIC*

Please cite this article as: Rasmussen, T.B., Reimann, I., Uttenthal, Å., Leifer, I., Depner, K., Schirrmeier, H., Beer, M., Generation of recombinant pestiviruses using a full-genome amplification strategy, *Veterinary Microbiology* (2008), doi:10.1016/j.vetmic.2009.09.037

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Generation of recombinant pestiviruses using a full-genome amplification
strategy**

Thomas Bruun Rasmussen^{1*}, Ilona Reimann², Åse Uttenthal¹, Immanuel Leifer²,
Klaus Depner², Horst Schirrmeier² and Martin Beer²

¹DTU National Veterinary Institute, Technical University of Denmark, Lindholm,
DK-4771 Kalvehave, Denmark

²Institute of Diagnostic Virology, Friedrich-Loeffler-Institut, D-17493 Greifswald-
Insel Riems, Germany

*Corresponding author

Thomas Bruun Rasmussen
DTU National Veterinary Institute (DTU Vet)
Technical University of Denmark
Lindholm
DK-4771 Kalvehave
Phone : +45 35887850
Fax : +45 35887901
Email : tbrur@vet.dtu.dk

22 **Abstract**

23 Complete genome amplification of viral RNA provides a new tool for the
24 generation of modified viruses. We have recently reported a full-genome
25 amplification strategy for recovery of pestiviruses (Rasmussen et al., 2008). A
26 full-length cDNA amplicon corresponding to the Border disease virus-Gifhorn
27 genome was generated by long RT-PCR and then RNA transcripts derived from
28 this amplicon were used to rescue infectious virus. Here, we have now used this
29 full-genome amplification strategy for efficient and robust amplification of three
30 additional pestivirus strains: the vaccine strain C and the virulent Paderborn
31 strain of Classical swine fever virus plus the CP7 strain of Bovine viral diarrhoea
32 virus. The amplicons were cloned directly into a stable single-copy bacterial
33 artificial chromosome generating full-length pestivirus DNAs from which
34 infectious RNA transcripts could be also derived.

35

36 Keywords: *Pestivirus*, *Flaviviridae*, long RT-PCR, full-genome amplicon,
37 infectious clone, pBeloBAC11

38

39 **Introduction**

40 The genus *Pestivirus* is part of the family *Flaviviridae* and contains economically
41 important animal viruses such as Classical swine fever virus (CSFV), Bovine
42 viral diarrhoea virus (BVDV) and Border disease virus (BDV). Pestivirus virions
43 are enveloped and contain a positive-stranded RNA genome of about 12.3 kb
44 which has a single long open reading frame flanked by non-translated regions
45 (5'NTR and 3'NTR), which are important for viral replication. Generation of
46 pestivirus cDNA clones is a general prerequisite for genetic manipulation of
47 their genomes, a process that can be both lengthy and laborious. Several
48 infectious cDNA clones containing complete pestivirus genomes have been
49 described (Meyers et al., 1996; Moormann et al., 1996; Ruggli et al., 1996;
50 Vassilev et al., 1997; Fan and Bird, 2008). These cDNA clones have provided
51 new options for directed genetic manipulations of pestiviruses. However,
52 construction of these infectious cDNA clones has been hampered due to the
53 relatively large size of the viral RNA genome and because of genetic instability
54 of the cloned cDNA in combination with the plasmid vectors in the bacterial
55 host. Therefore, new strategies are needed to facilitate construction of stable
56 infectious cDNA clones from a wider range of strains.

57 Generation of infectious cDNA clones can be facilitated by using
58 long RT-PCR for full-genome amplification. This approach has been applied
59 with success to a number of RNA viruses from the *Flaviviridae* (Tellier et al.,
60 1996; Gritsun and Gould, 1998; Zhang et al., 2001). For pestiviruses, the
61 methodology has been used to obtain the entire open reading frame for
62 sequencing studies (Jones et al., 2006) as well as for virus rescue (Rasmussen

et al., 2008). Full-length genome amplification facilitates, in addition to virus rescue, direct full-length sequence analysis and the amplicons can also be directly inserted into suitable cloning vectors.

In the present study, we developed an optimized, efficient and streamlined method for the robust amplification of full-length cDNAs of four strains (BVDV-CP7, BDV-Gifhorn, CSFV-C and CSFV-Paderborn). This procedure was followed by the insertion of the BDV and CSFV amplicons into the stable single-copy vector pBeloBAC11 (Wang et al., 1997). This bacterial artificial chromosome (BAC) has been shown to successfully maintain the full-length cDNAs of Japanese encephalitis virus (Yun et al., 2003) and BVDV-SD1 (Fan and Bird, 2008). Using this procedure, we have generated new full-length cDNA clones of BDV and CSFV, which enlarges the range of cloned pestivirus genomes that can be manipulated.

Materials & Methods

The different viruses used in this study were; BVDV-CP7 from bovine KOP-R cells (10^6 TCID₅₀/ml); CSFV-Paderborn from porcine PK15 cells (10^7 TCID₅₀/ml; vaccine strain CSFV-C from porcine PK15 cells ($10^{4.75}$ TCID₅₀/ml); and BDV-Gifhorn from ovine SFT cells (10^6 TCID₅₀/ml). Infected cells were grown in Dulbecco's Modified Eagle Medium supplemented with 10% BVDV-free foetal bovine serum at 37°C in a humidified atmosphere containing 5% CO₂. The total RNA was extracted from the cells using a combined Trizol/RNeasy protocol. This protocol was used to obtain high-quality full-length genomic RNA. Briefly, total RNA was extracted from 1 ml cell supernatant or cell lysate with 3 ml Trizol

LS reagent (Invitrogen) and 0.8 ml chloroform. After phase separation the aqueous phase was mixed with an equal volume of 75% ethanol and 0.7 ml of the resulting mix was transferred sequentially to the same RNeasy column (Qiagen). The column was washed according to the manufacturer's protocol. Total RNA was eluted from the column with 4 times 30 µl nuclease free water and stored at -80°C until use. Pestivirus genomes were amplified from total RNA preparations using long RT-PCR. It was found that the third RNA eluate empirically performed the best in long RT-PCR, presumable due to a higher proportion of full-length genomic RNA in this fraction.

The total RNA was reverse transcribed at 50°C for 90 minutes using SuperScript III reverse transcriptase kit (Invitrogen) and specific cDNA primers (Table 1). For improved long RT-PCR efficiency, the resulting cDNA was subsequently treated with RNase H to remove the RNA template. Full-length PCR amplification was performed using primers specific for the 5'NTR and the 3'NTR of each strain (Table 1) with the Accuprime High Fidelity kit (Invitrogen) which consists of a mixture of *Taq* and proofreading *Pyrococcus* *GB-D* DNA polymerases. Reactions containing 2 µl cDNA were then amplified in a final volume of 50 µl using 94°C for 30 seconds followed by 35 cycles of 94°C for 15 seconds, 65°C for 30 seconds and 68°C for 12 minutes. Samples (1-10 µl) of each PCR reaction were analysed on a 1 % agarose gel in TBE buffer alongside a 1kb DNA ladder (Fermentas).

Standard techniques were used for cloning of the full-length amplicons into pBeloBAC11 (Invitrogen). Briefly, pBeloBAC11 (7507 bp) was digested with *NotI* and afterwards treated with Antarctic phosphatase (NEB).

111 The full-length amplicons of about 12.3 kb were also digested with *NotI* and
112 purified using the Nucleotide removal kit (Qiagen). Equal amounts of vector and
113 amplicons were mixed and ligated in 25 µl reactions overnight at 16°C using
114 DNA ligase for long fragments (Takara Bio) according to the manufacturer's
115 recommendations. The products (1 µl) were electroporated into
116 electrocompetent DH10B T1 phage resistant *E.coli* (Invitrogen) and
117 transformants harbouring the cloned amplicons were isolated from selective LB
118 plates containing 12.5 µg/ml chloramphenicol. Small scale preparations of DNA
119 from selected transformants were obtained using the BAC miniprep protocol
120 previously described (Warming et al., 2005) and the presence of the full-length
121 pestivirus cDNA was determined following restriction analysis using *NotI*.
122 Selected BAC DNAs were digested with *NotI*, *in vitro* transcribed using T7 RNA
123 polymerase and the RNA transcripts were tested for infectivity as previously
124 described (Reimann et al., 2004; Rasmussen et al., 2007). Immunostaining of
125 electroporated porcine SK6 cells (CCLV RIE262) was performed with the pan-
126 pestivirus antibody (C16) targeting the non-structural NS3 proteins (Reimann et
127 al., 2004).

128

129 **Results**

130 Full-length cDNAs corresponding to four different pestivirus strains were
131 successfully amplified in single long RT-PCRs using an optimized version of the
132 full-genome amplification strategy described by Rasmussen et al. (2008). The
133 original protocol was modified by substituting the Elongase enzyme mix with the
134 Accuprime High Fidelity kit. This polymerase mix, in combination with the

Superscript III reverse transcriptase, efficiently produced, from each of the different viral RNAs, amplicons of approximately 12.3 kb, corresponding to the full-length genomes (Fig. 1). The Accuprime High Fidelity polymerase mix increased the specificity and efficiency of the long RT-PCR compared to the Elongase enzyme mix and produced distinct products of the expected size without smears and aberrant amplification products (Fig. 1). High yields (several micrograms) of DNA were obtained in each long RT-PCR. For amplification of complete genomes in a single RT-PCR, specific primers were designed based on nucleotide sequences of the terminal 5'NTR- and 3'NTR- genomic ends (Table 1). The primers for amplification of BDV-Gifhorn were described in Rasmussen et al. (2008). The primers for BVDV-CP7 (Meyers et al., 1996, Accession number BVU63479), and for the CSFV-C (Accession number AY259122) were based on published sequences. Similarly, the 5'NTR primer for CSFV-Paderborn was designed from Accession number AY072924 (Oleksiewicz et al., 2003) but this sequence lacks approximately 75 nucleotides from the terminus of the 3'NTR. Therefore, the 3'Paderborn reverse primer was designed using a nucleotide alignment of known full-genome CSFV sequences. Generally, knowledge of less than 50 nucleotides of the terminal sequences from each of the strains was needed for successful amplification of the whole genome.

By incorporating unique *NotI* restriction sites within the primers (Table 1), the amplicons were designed for subsequent insertion into pBeloBAC11. Full-length amplicons of the two strains CSFV-C and CSFV-Paderborn, and also the BDV-Gifhorn were obtained using the respective *NotI*

159 primers (Fig. 2). Amplification and cloning of BVDV-CP7 was not attempted by
160 this strategy due to an internal *NotI* site in the cDNA sequence.

161 The amplified genomes were digested with *NotI* and ligated directly
162 into the *NotI* site in pBeloBAC11. Electroporation of DH10B *E.coli* cells with the
163 ligated products yielded small and large bacterial colonies (from a few to
164 several hundreds) after incubation for 24 hours at 37°C. Both small and large
165 colonies were tested for the presence of full-genome amplicons. Generally, two
166 thirds of the tested transformants carried a BAC containing an insert of the
167 expected size of approximately 12.3 kb (Fig. 3). One third of the transformants
168 either carried BACs with smaller inserts, empty BACs or no visible BAC vector.
169 There was no correlation between the size of the BAC inserts and the size of
170 the bacterial colonies.

171 Testing of the BAC clones for their ability to generate virus
172 infectivity in cells is ongoing in our laboratories. The preliminary results of this
173 evaluation show that infectious pestiviruses can be rescued from the DNAs
174 generated by direct insertion of full-length genomes into the BAC vector. For
175 example, selected BAC clones harbouring the CSFV-Paderborn full-length
176 cDNA were *NotI* digested, *in vitro* transcribed and the RNA transcripts were
177 tested for infectivity by electroporation into porcine cells. RNA derived from one
178 out of two tested BAC clones (pBeloPader10) was shown to efficiently replicate
179 in porcine cells. The cells electroporated with RNA transcripts from
180 pBeloPader10 showed a high number of NS3-positive cells (as judged by
181 immunofluorescence, Fig. 4) 24 hours later and infectious virus could be
182 recovered after 1st and 2nd cell passages. In addition, RNA transcripts derived

183 from three BAC clones harbouring BDV-Gifhorn full-length cDNA have been
184 tested for infectivity in ovine cells. Following electroporation, viral RNA derived
185 from each of these three BAC clones was found to be replicating in the cells
186 (data not shown). However, only rather few cells were scored as positive after
187 the 1st cell passage indicating a possible growth defect in these constructs.

188

189 **Discussion**

190 The use of the full-length genome amplification strategy for amplification of
191 pestivirus genomes has several advantages; (i) the same optimized protocol
192 can be used for amplification of a wide selection of strains; (ii) only limited
193 terminal nucleotide sequence information of a given strain is needed for
194 amplification of the whole genome; (iii) multiple cloning steps can be
195 circumvented since full-length amplicons can be directly *in vitro* transcribed or
196 inserted into plasmid vectors; and (iv) construction of genetically modified
197 viruses can be facilitated using stable cDNA clones.

198 In the present study, four different pestiviruses representing each of
199 three main types (CSFV, BVDV and BDV) were amplified by long RT-PCR. This
200 demonstrates that the recently described full-length genome amplification
201 strategy (Rasmussen et al., 2008) is of a generic nature since it can be utilized
202 for amplification of diverse pestivirus strains. Furthermore, the direct cloning of
203 the amplified genomes into the stable single-copy BAC vector, pBeloBAC11,
204 shows that we have created an efficient, streamlined strategy for direct
205 preparation of new pestivirus cDNA clones. Full-length cDNAs corresponding to
206 other members of the *Flaviviridae*, cloned in pBeloBAC11, have previously been

207 shown to be stable after multiple generations in *E.coli* cells (Yun et al., 2003;
208 Fan and Bird, 2008) and we anticipate that our constructs will display similar
209 stability in the bacterial host but this does need to be determined.

210 Our new strategy allows construction of stable infectious BAC
211 DNAs from a single full-length PCR product, which gives an increased flexibility
212 in the design of new genetically modified pestiviruses. Since long RT-PCR
213 preserves the genetic variants present in a virus population (Chumakov, 1996)
214 many independent full-length cDNA clones can be obtained and tested which
215 increases the probability of finding infectious constructs. For targeted design of
216 genetically modified pestiviruses, the efforts can now be expedited and focused
217 on, in principal, any pestiviral strain and is hence not limited by the availability of
218 existing infectious cDNAs. Furthermore, our full-genome amplification protocol
219 obviates the time-consuming and costly process of construction and screening
220 of infectious clones by traditional methods. Therefore, the full-genome
221 amplification strategy significantly simplifies and streamlines the workflow for
222 the generation of new recombinant pestiviruses and also facilitates direct full-
223 length sequence analysis.

224

225 **Acknowledgements**

226 We are grateful to Gabriela Adam and Doreen Reichelt for technical assistance.
227 We thank Dr. Donata Kalthoff for her valuable advice regarding cloning in BAC
228 vectors. This work was supported by the EU Network of Excellence, EPIZONE
229 (FOOD-CT-2006-016236) and by the Danish Research Council for Technology
230 and Production Sciences (DRCTPS grant 274-07-0198).

231 **Conflict of interest statement**

232 None

233

234 **References**

235 Chumakov, K.M., 1996. PCR engineering of viral quasispecies: a new method
236 to preserve and manipulate genetic diversity of RNA virus populations. *J. Virol.*
237 70, 7331-7334.

238 Fan, Z.C., Bird, R.C., 2008. An improved reverse genetics system for
239 generation of bovine viral diarrhea virus as a BAC cDNA. *J Virol Methods* 149,
240 309-315.

241 Gritsun, T.S., Gould, E.A., 1998. Development and analysis of a tick-borne
242 encephalitis virus infectious clone using a novel and rapid strategy. *J. Virol.*
243 *Methods* 76, 109-120.

244 Jones, L.R., Zandomeni, R.O., Weber, E.L., 2006. A long distance RT-PCR
245 able to amplify the Pestivirus genome. *J. Virol. Methods* 134, 197-204.

246 Meyers, G., Tautz, N., Becher, P., Thiel, H.J., Kummerer, B.M., 1996. Recovery
247 of cytopathogenic and noncytopathogenic bovine viral diarrhea viruses from
248 cDNA constructs. *J. Virol.* 70, 8606-8613.

249 Moormann, R.J., van Gennip, H.G., Miedema, G.K., Hulst, M.M., van Rijn, P.A.,
250 1996. Infectious RNA transcribed from an engineered full-length cDNA template
251 of the genome of a pestivirus. *J. Virol.* 70, 763-770.

- 252 Oleksiewicz, M.B., Rasmussen, T.B., Normann, P., Uttenthal, A., 2003.
253 Determination of the sequence of the complete open reading frame and the
254 5'NTR of the Paderborn isolate of classical swine fever virus. *Vet. Microbiol.* 92,
255 311-325.
- 256 Rasmussen, T.B., Reimann, I., Hoffmann, B., Depner, K., Uttenthal, A., Beer,
257 M., 2008. Direct recovery of infectious pestivirus from a full-length RT-PCR
258 amplicon. *J Virol Methods* 149, 330-333.
- 259 Rasmussen, T.B., Uttenthal, A., Reimann, I., Nielsen, J., Depner, K., Beer, M.,
260 2007. Virulence, immunogenicity and vaccine properties of a novel chimeric
261 pestivirus. *J Gen Virol* 88, 481-486.
- 262 Reimann, I., Depner, K., Trapp, S., Beer, M., 2004. An avirulent chimeric
263 Pestivirus with altered cell tropism protects pigs against lethal infection with
264 classical swine fever virus. *Virology* 322, 143-157.
- 265 Ruggli, N., Tratschin, J.D., Mittelholzer, C., Hofmann, M.A., 1996. Nucleotide
266 sequence of classical swine fever virus strain Alfort/187 and transcription of
267 infectious RNA from stably cloned full-length cDNA. *J. Virol.* 70, 3478-3487.
- 268 Tellier, R., Bukh, J., Emerson, S.U., Miller, R.H., Purcell, R.H., 1996. Long PCR
269 and its application to hepatitis viruses: amplification of hepatitis A, hepatitis B,
270 and hepatitis C virus genomes. *J. Clin. Microbiol.* 34, 3085-3091.
- 271 Vassilev, V.B., Collett, M.S., Donis, R.O., 1997. Authentic and chimeric full-
272 length genomic cDNA clones of bovine viral diarrhea virus that yield infectious
273 transcripts. *J. Virol.* 71, 471-478.

- 274 Wang, K., Boysen, C., Shizuya, H., Simon, M.I., Hood, L., 1997. Complete
275 nucleotide sequence of two generations of a bacterial artificial chromosome
276 cloning vector. *Biotechniques* 23, 992-994.
- 277 Warming, S., Costantino, N., Court DL, Jenkins, N.A., Copeland, N.G., 2005.
278 Simple and highly efficient BAC recombineering using galK selection. *Nucleic
279 Acids Res.* 33, e36.
- 280 Yun, S.I., Kim, S.Y., Rice, C.M., Lee, Y.M., 2003. Development and application
281 of a reverse genetics system for Japanese encephalitis virus. *J Virol* 77, 6450-
282 6465.
- 283 Zhang, F., Huang, Q., Ma, W., Jiang, S., Fan, Y., Zhang, H., 2001. Amplification
284 and cloning of the full-length genome of Japanese encephalitis virus by a novel
285 long RT-PCR protocol in a cosmid vector. *J. Virol. Methods* 96, 171-182.
286
287

Table 1: Primers used for long RT-PCR^a

Primers for CSFV-C and CSFV-Paderborn

5'Paderborn_T7 (forward)	<u>TAATACGACTCACTATAGT</u> ATACGAGGTTAGCTCGTCCTCGTGACAAACATT
5'Paderborn_T7_NotI (forward)	ATATGCGGCCGCT <u>TAATACGACTCACTATAGT</u> ATACGAGGTTAGCTCGTCCTCGTGACAAACATT
3'Paderborn (reverse)	GGGCCGTTAGGAAATTACCTTAGTCCAACCTGTGGA
3'Paderborn-NotI (reverse)	ATATGCGGCCGCGGCGCCGTTAGGAAATTACCTTAGTCCAACCTGTGGA
3'Paderborn_cDNA (reverse)	GGGCCGTTAGGAAATTACCTTAGT
5'C-strain_T7 (forward)	<u>TAATACGACTCACTATAGT</u> ATACGAGGTTAGTTCATTCTCGTATACACGATTGGACAAATC
5'C-strain_T7_NotI (forward)	ATATGCGGCCGCT <u>TAATACGACTCACTATAGT</u> ATACGAGGTTAGTTCATTCTCGTATACACGATTGGACAAATC

Primers for BVDV-CP7

5'BVDV CP7_T7 (forward)	TAATACGACTCACTATAGTATACGAGAATTAGAAAAGGCAC
3'BVDV CP7 (reverse) ^b	GGGTGACGTCGGGTGTACCCCTCATAC

Primers for BDV-Gifhorn

5'T7Gif-NotI (forward)	ATATGCGGCCGCT <u>TAATACGACTCACTATAGT</u> ATACGAGAGTAGTTCAGGCTCGTATGCAAAATTGGGTGTTTC
3'Gif-NotI (forward)	ATATGCGGCCGCGGGGCTGTTAGGGTTTTTCTTAATCCAACCTATGGACTTCAG

^a T7 promoter is underlined, and NotI sites are shown in italics.

^b Same primer was also used as cDNA primer.

307 Fig. 1. Full-genome amplification of pestiviruses by long RT-PCR using T7-
 308 5'NTR and 3'NTR primers. The pictures show agarose gel electrophoresis of
 309 the full-length amplicons of BDV-Gifhorn (A, 1 μ l), BVDV-CP7 (B, 1 μ l), CSFV-C
 310 (C, 10 μ l) and CSFV-Paderborn (D, 10 μ l).

311

312 Fig. 2. Full-genome amplification of pestiviruses by long RT-PCR using T7-
 313 5'NTR and 3'NTR primers containing *NotI* restrictions sites. The picture shows
 314 agarose gel electrophoresis of the full-length amplicons of BDV-Gifhorn (A, 10
 315 μ l), CSFV-C (B, 10 μ l) and CSFV-Paderborn (C, 10 μ l). The non-specific bands
 316 in A and C are most likely caused by the increased complexity of the primers
 317 (extra *NotI* sites).

318

319 Fig. 3. *NotI* digestion of BAC transformants. DNA preparations from BAC
 320 transformants were digested with *NotI* and analysed by agarose gel
 321 electrophoresis. Four have the correct size insert (A-C, E), whereas one (D) has
 322 a smaller insert, and another (F) contained no detectable BAC vector.

323

324 Fig. 4. Recovery of infectious CSFV-Paderborn derived from *in vitro* transcribed
 325 pBeloPader10 RNA electroporated into porcine SK6 cells. The pictures show
 326 immunofluorescence (IF) staining of cells expressing the NS3 proteins after the
 327 1st (Panel A) and 2nd (Panel B) cell culture passage. Panels C and D are control
 328 cells electroporated without any RNA transcripts.

329

330

331

332 Figure 1.

333

334

335

336

337 Figure 2.

338

339

340

341 Figure 3.

342

343

344

345 Figure 4.

346

347

348