Genetic architecture of open-angle glaucoma and related determinants

Wishal D Ramdas, Najaf Amin, Leonieke Me van Koolwijk, Cecile Ajw Janssens, Aisa Demirkan, Paulus Tvm de Jong, Yurii Aulchenko, Roger Cw Wolfs, Albert Hofman, Fernando Rivadeneira, et al.

► To cite this version:

Wishal D Ramdas, Najaf Amin, Leonieke Me van Koolwijk, Cecile Ajw Janssens, Aisa Demirkan, et al.. Genetic architecture of open-angle glaucoma and related determinants. *Journal of Medical Genetics*, 2010, 48 (3), pp.190. 10.1136/jmg.2010.083337 . hal-00583892

HAL Id: hal-00583892

<https://hal.science/hal-00583892>

Submitted on 7 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetic architecture of open-angle glaucoma and related determinants

Wishal D. Ramdas MD MSc,^{1,2} Najaf Amin DSc,¹ Leonieke M.E. van Koolwijk MD MSc,^{1,3} A. Cecile J.W. Janssens PhD,¹ Aisa Demirkan MSc,¹ Paulus T.V.M. de Jong MD PhD,^{4,5} Yuri S Aulchenko PhD,¹ Roger C.W. Wolfs, MD PhD,^{1,2} Albert Hofman MD PhD,¹ Fernando Rivadeneira MD PhD,^{1,6} Andre G. Uitterlinden PhD,^{1,6,7} Ben A. Oostra PhD,⁷ Hans G. Lemij MD PhD,³ Caroline C.W. Klaver MD PhD,^{1,2} Johannes R. Vingerling MD PhD,^{1,2} Nomdo M. Jansonius MD PhD,^{1,8} Cornelia M. van Duijn PhD.¹

¹ Department of Epidemiology, Erasmus Medical Center, Rotterdam, the Netherlands

² Department of Ophthalmology, Erasmus Medical Center, Rotterdam, the Netherlands

³ Glaucoma service, The Rotterdam Eye Hospital, Rotterdam, the Netherlands

⁴ Department of Ophthalmogenetics, The Netherlands Institute for Neuroscience, RNAAS, Amsterdam, the Netherlands

⁵ Department of Ophthalmology, Academic Medical Center, Amsterdam, the Netherlands

⁶ Department of Internal Medicine, Erasmus Medical Center, Rotterdam, the Netherlands

⁷ Department of Clinical Genetics, Erasmus Medical Center, Rotterdam, the Netherlands

⁸ Department of Ophthalmology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

To be submitted to: Journal of Medical Genetics

Word count: Summary: 249 (max. 250); Manuscript: ~4,061

Number of tables/figures: 6 (max. 6); **references:** 28 (max. 50); **Supplementary Material:** Yes

Conflict of interest: none

Corresponding author: Johannes R. Vingerling, MD, PhD, Dept. Ophthalmology and Epidemiology,

Erasmus Medical Center Rotterdam, PO Box 2040, 3000 CA Rotterdam, The Netherlands

Telephone: (+31) 6-50221140

Fax: (+31) 10-2620852

E-mail: j.vingerling@erasmusmc.nl

Keywords: Genetics; Vertical cup-disc ratio; Intraocular pressure; Open-angle laucoma; Polygenomic;

Genetic epidemiology

SUMMARY

Background – Although the vertical cup-disc ratio (VCDR) and intraocular pressure (IOP) are important determinants of open-angle glaucoma (OAG), it is unclear to what extent the genetic origin of these traits overlap with those of OAG. We evaluated whether the same genes that determine VCDR and IOP also predict OAG.

Methods – Genetic risk scores were constructed from single nucleotide polymorphisms (SNPs) using genome-wide association data of 9,326 participants from the Rotterdam Study cohorts (mean $\pm$ standard deviation age: 64.6 ± 9.1). These risk scores were used to calculate the explained variance of VCDR and IOP in an independent cohort (Erasmus Rucphen Family study) consisting of 1,646 participants (mean $\pm$ standard deviation age: 46.8 ± 14.1) and the OAG risk in a subset of the Rotterdam Study cohorts. To evaluate false positive findings, we generated two new variables containing randomly sampled values to serve as a negative control.

Results – The explained variance of VCDR increased when increasing the number of SNPs included in the risk score, suggesting a polygenic model. We found no clear evidence for a similar model for IOP, suggesting that a small number of SNPs determine the susceptibility to IOP. The SNPs related to IOP in terms of p-values contributed little to VCDR. The risk scores associated with VCDR were also associated significantly with OAG. This suggests a common polygenic background for VCDR and OAG.

Conclusions – We found evidence for a polygenic model underlying one of the major traits of OAG, VCDR, and OAG itself. The IOP did not show any evidence for such a model.

INTRODUCTION

Open-angle glaucoma (OAG) is a neurodegenerative disease that leads to progressive damage to the retinal ganglion cells and nerve fibers, resulting in glaucomatous visual field loss. It is one of the leading causes of irreversible blindness in the world, affecting over 60 million persons worldwide.[1] Although the etiology of OAG is still obscure, it has been recognized for long that genetic factors play a role in its pathogenesis. Relatives of patients with OAG have an increased risk of OAG.[2] Several rare genetic variants associated with a high risk of disease have been implicated in familial forms of OAG, but these variants only explain a small percentage of OAG patients.[3] This raises the question whether the genetic etiology of OAG is determined by a large number of rare variants with major effects on the disease risk (rare variant, common disease hypothesis) or whether there are multiple common variants underlying the disease (common variant-common disease hypothesis).

Genome-wide association studies (GWAS) have identified a large number of common variants (single nucleotide polymorphisms [SNPs]) with a low risk for a large number of common diseases (e.g. type 2 diabetes, Alzheimer's disease, and various forms of cancer), but also for diseases that are less common such as multiple sclerosis and Crohn's disease.[4] GWAS have also been powerful in studies on quantitative risk factors of diseases such as blood pressure, glucose levels and height.[4] More recently, GWAS were used to identify polygenic forms of inheritance for schizophrenia, implicating ten thousands of genetic variants covering the whole genome rather than a distinct number of variants.[5] The essence of the latter approach is that a gene discovery sample is used to define risk scores, and the risk is subsequently predicted in an independent sample. In this approach, not only the variants that are genome-wide significant are used but also those who do not reach this threshold. This method can also be applied to test for overlapping of genes between diseases and their underlying liability.

If there is a polygenic model that explains part of the susceptibility to a disease, the classical theory of Fisher predicts that the number of risk alleles is distributed according to the Gaussian distribution in the population.[6] It is easy to show that if there are >30 genetic variants, no person in the population carries zero risk alleles but everyone carries a gradient number of risk alleles.[7] Thus, a polygenic model implies that individuals in a population always carry risk alleles for a disease and that the liability to disease increases with the number risk alleles one carries. It also follows from this model that

underlying a disease that is defined as present or absent, there is probably a quantitative trait. The persons in the tail of the distribution of this trait are considered to be “diseased”. If the quantitative trait (or risk factor) is related to the genotypic distribution and increases with the number of risk alleles, the risk of disease will behave accordingly.

If we translate this to OAG, there are two clinical measures consistently associated with this disease and may thus determine its (polygenic) liability to OAG. First, OAG is characterized by damage to the optic nerve head, which is visible as an (increased) excavation (cupping) upon ophthalmological examination. The extent of the excavation is commonly quantified as the vertical cup-to-disc ratio (VCDR), ranging from 0 to 1.[8] Both an increase in VCDR and an unusually large VCDR at a single observation may indicate glaucomatous changes of the optic nerve head.[9] The second clinical measure that may determine the (polygenic) liability to OAG is the intraocular pressure (IOP). Current treatment for OAG is based on lowering the IOP. VCDR as well as IOP are highly heritable, making these two variables potentially important endophenotypes for OAG.[10] Until now it is unknown to which extent the genes involved in VCDR overlap with those involved in IOP, and to which extent they actually predict OAG.

The aim of this study was to investigate whether there is a polygenic model in which many variants apply to VCDR and IOP and whether these genes also predict the risk of OAG. For this study, we used the Rotterdam Study I, Rotterdam Study II and Rotterdam Study III (RS-I, RS-II and RS-III, respectively) as discovery cohorts to derive genetic risk scores that capture the genes involved in VCDR and IOP. Next, we assessed whether these risk scores predicted VCDR and IOP in an independent target cohort (the Erasmus Rucphen Family [ERF] study). For the prediction of OAG, RS-I served as the target cohort, because no OAG cases were included in ERF.

METHODS

Study Populations

The RS-I is a prospective population-based cohort study of 7,983 residents aged 55 years and older living in Ommoord, a suburb of Rotterdam, the Netherlands. Baseline examination for the ophthalmic part took

place between 1991 and 1993; follow-up examinations were performed from 1997 to 1999 and from 2002 to 2006.[11, 12] RS-I included 188 OAG cases that either already had OAG at baseline or developed OAG during follow-up.

The RS-II and RS-III are two other prospective population-based cohort studies of respectively 3,011 residents aged 55 years and older and 3,392 residents aged 45 years and older. The rationale and study design are similar to that of the RS-I.[11] The baseline examination of RS-II took place between 2000 and 2002; the follow-up examination was performed from 2004 to 2005. Baseline examination of RS-III took place between 2006 and 2009.[13]

The ERF Study is a family-based cohort in a genetically isolated population in the southwest of the Netherlands with over 3,000 participants aged between 18 and 86 years. Cross-sectional examinations took place between 2002 and 2005. The rationale and study design of this study have been described elsewhere.[14, 15] All measurements in these studies were conducted after the Medical Ethics Committee of the Erasmus University had approved the study protocols and all participants had given a written informed consent in accordance with the Declaration of Helsinki.

Ophthalmic examination

The ophthalmic assessment in RS-I and RS-II, both for baseline and follow-up, included a medical history, autorefraction, keratometry, IOP measurement, visual field testing and optic nerve head imaging with simultaneous stereoscopic photography (ImageNet; Topcon Corporation, Tokyo, Japan) of both eyes after mydriasis with tropicamide 0.5% and phenylephrine 2.5% eye drops.

RS-III was similar to RS-I except for optic nerve head imaging with Heidelberg Retina Tomograph 2 (HRT; Heidelberg Engineering, Dossenheim, Germany). The ophthalmic assessment in ERF included a medical history, autorefraction, keratometry, IOP measurement, and optic nerve head imaging with HRT of both eyes after pharmacologic mydriasis.

In all cohorts we measured IOP at baseline and at every follow-up round with Goldmann applanation tonometry (Haag Streit, Bern, Switzerland) after applying oxybuprocaine 0.4% eye drops. The median value of three successive measurements was recorded.[16]

The OAG definition was based on the presence of glaucomatous visual field loss. Details have been described elsewhere. Neither VCDR nor IOP were part of the diagnostic criteria of OAG.[17, 18]

ImageNet, which was used for optic nerve head imaging in RS-I and RS-II, takes simultaneous stereoscopic images of the optic disc at a fixed angle of 20°, using a simultaneous stereoscopic fundus camera (Topcon TRC-SS2; Tokyo Optical Co., Tokyo, Japan). Images were analyzed using the ImageNet retinal nerve fiber layer height module. On each stereoscopic pair of optic disc images four points were marked on the disc margin, defined as the inner border of the peripapillary ring or the outer border of the neural rim, if a scleral ring was visible. Next, the software drew an ellipse using these points to outline the disc margin and to determine the cup. The amount of correspondence between the marked points on the two images of the stereoscopic pair is expressed as a “bad points” percentage, which indicates the percentage of points lacking correspondence. This percentage can be used as an indicator of image quality. Images with 25% or more bad points were excluded.[19]

HRT 2, used for optic nerve head imaging in RS-III and ERF, uses a focused 670-nm diode laser light beam to acquire scans of the optic nerve head region, using the confocal principle. The HRT obtains, during one scan, three series of 16 to 64 confocal frontal slices. From each of these series, a 3-dimensional image of the optic nerve head is reconstructed, from which the software calculates several optic disc parameters. To define the cup, the HRT places a reference plane 50 μm below the peripapillary retinal surface in the region of the papillomacular bundle. Imaging was performed after entering the participant’s keratometry data into the software and after adjusting the settings in accordance with the refractive error. In RS-III all HRT 2 data were converted to HRT 3. As an indicator of image quality we used the topographic standard deviation of the scan, which is a measure of the variability among the three series of a single HRT scan. Scans with a topographic standard deviation exceeding 50 μm were excluded. The inter-observer variability and agreement for both systems have been described elsewhere.[20][*Ramdas, et al. unpublished data*]

Genotyping

In the RS-I, RS-II and RS-III cohorts, DNA was genotyped using the Illumina Infinium II HumanHap arrays according to the manufacturer's protocols. After exclusion of participants for reasons of low-quality DNA, a total of 5,974 participants were available with genotyping data from RS-I (HH550 v3.0®), 2,157 participants were from RS-II and 2,082 from RS-III (HH550Duo and HH610Quad®). Details are described elsewhere.[21]

In ERF, DNA was genotyped on four different platforms (Illumina 6k, Illumina 318K, Illumina 370K and Affymetrix 250K), which were then merged and imputed to 2.5 million SNPs hapmap using build 36 HapMap (release 22) CEU population as a reference cohort. After exclusion of participants for whom genotyping data were unavailable, 2,385 had genotyping data.

Extensive quality control analyses have been performed in each cohort.

Statistical analysis

Statistical analysis within the discovery cohorts

If data of both eyes were available, we randomly chose one. In case of missing or unreliable baseline data of both eyes, we used follow-up data whenever possible. For each of the discovery cohorts (comprising RS-I, RS-II and RS-III) linear regression models were used to examine the associations between the risk scores and VCDR and IOP, adjusted for age and gender. For VCDR we also adjusted for optic disc area. For IOP the mean value of both eyes was included and adjusted for IOP lowering treatment. This adjustment was done by adding 30% upon the measured IOP for medical treatment and by fixing the IOP at 30 mmHg for surgical/laser treatment in the past.

To summarize the results through the three discovery cohorts, we performed a meta-analysis with Metal for Linux (www.sph.umich.edu/csg/abecasis/metal) using the inverse variance method of each effect size estimate.[22]

All statistical analyses were performed using SPSS version 15.0.0 for Windows (SPSS inc., Chicago, IL, USA; 2006), PLINK[23] and R statistical package version 2.10.1 for Mac (www.r-project.org).

Risk scores within the independent cohort

The risk score profiling method tests the association of a genetic score variable that reflects a combined effect of a number of selected SNPs with a trait. Details of this method have been described elsewhere.[5] The results of the SNPs of the meta-analyzed discovery cohorts were categorized according to p-value and examined in an independent cohort (ERF). The p-value thresholds were as follow: $p < 10^{-10}$, $p < 10^{-9}$, $p < 10^{-8}$, $p < 10^{-7}$, $p < 10^{-6}$, $p < 10^{-5}$, $p < 10^{-4}$, $p < 10^{-3}$, $p < 10^{-2}$, $p < 0.1$, $p < 0.2$, $p < 0.3$, $p < 0.4$, $p < 0.5$, $p < 0.6$, $p < 0.7$, $p < 0.8$, $p < 0.9$, $p < 1.0$. For each category the effect (beta) of each SNP was multiplied by the corresponding number of alleles of each participant to calculate a score for that particular SNP. The mean of all these scores for the respective participant was the risk score. In case of missing SNP data from a participant of the replication cohort the allele frequencies were used and multiplied by the betas. Next, we ran linear regression analyses with the VCDR as the outcome measure and the risk scores adjusted for age and gender as the determinant in order to calculate the proportion of explained variance (PEV) of each group of SNPs (i.e. p-value threshold) as determined in the discovery cohort. A p-value of 0.05 or smaller was considered as statistically significantly associated with the trait.

The same analysis was also done with IOP as the outcome. To examine whether a “non-ophthalmic” variable reveals the same results we performed the same analysis for a negative control. For this variable we used a sampling method without replacement. We sampled the VCDR respectively IOP in the target cohort, and thus, the distribution remained the same. In addition, we created a second normal distributed variable by assigning a random number (with mean: 10; standard deviation: 1) to all participants.

For the analysis of the family-based data (ERF) we used SOLAR version 4.1.5 for Linux to adjust for the pedigree structure.[24]

Finally, we applied the same approach for the OAG cases of RS-I by calculating the Nagelkerke R-square of OAG for the VCDR and IOP risk scores. This was done using logistic regression analyses adjusted for age and gender. Since, as a matter of course, individuals with OAG may have an increased VCDR and/or IOP, we performed a secondary analysis in which we excluded RS-I from the meta-analysis of the discovery cohorts to avoid possible biased results. Differences in general characteristics between cases and controls were analyzed with independent t-tests and chi-square statistics.

Only autosomal SNPs that were available in all four cohorts (~520.000 SNPs) were included. To have less overlap of linkage disequilibrium blocks we did not use imputed data. SNPs that deviated significantly

from Hardy-Weinberg equilibrium ($p < 0.0001$) or that had a minor allele frequency less than 0.05 were excluded from the present study.

RESULTS

Study samples

Of the 5,974 participants that were genotyped in the RS-I cohort 5,312 had valid VCDR data. The remaining 662 were excluded because of missing or unreliable data. From the RS-II cohort 2,048 from 2,157 genotyped participants were included in this study. For 109 persons we did not have (reliable) data. From the RS-III cohort a total of 1,966 participants were included. The target cohort for the quantitative analysis (ERF) included 1,646 participants. Table 1 summarizes the general characteristics for the discovery cohorts and the replication cohort. The participants of the target set were significantly younger ($p < 0.001$), and had lower VCDR values ($p < 0.001$). The latter difference can be explained by the use of different measurement techniques for optic disc imaging (since we analyzed the cohorts separately and subsequently meta-analyzed the findings, this difference will not influence our findings). The distributions of VCDR and IOP for the four cohorts are shown in Supplementary Figure S1. Genetic outliers of non-European ancestry were excluded. No institutional heterogeneity between the cohorts or residual population sub-stratification was noticed after inspecting the genotype data.[25] Inflation factors for the included cohorts for VCDR and IOP ranged from 1.024 to 1.061 and from 1.006 to 1.037, respectively.

Risk score analysis

For VCDR, age and gender explained 0.3% of the total variance. Figure 1A presents the PEV attributable to the risk scores when risk scores from the VCDR analysis in the discovery sample were used to predict VCDR in the target sample ERF. Figure 1B shows the PEV when the same risk scores were used to predict a random variable (negative control). For VCDR, the risk score based on the p-value threshold of $p < 10^{-10}$ (first bar) consisted of 9 SNPs (Table 2) and explained only 0.1% (in addition to age and gender; region above the dotted line) of the variance in VCDR (Figure 1A) in the target population. Increasing the number of SNPs included in the risk scores (i.e. increasing the p-value thresholds for the SNPs to be

included) resulted in a gradual increase in the PEV until the $p < 10^{-2}$ threshold was used (consisting of 7,260 SNPs; $p = 8.68 \times 10^{-4}$). The PEV increased up to 1.0% ($p = 1.91 \times 10^{-4}$) at the threshold of $p < 0.2$ for the SNPs to be included in the risk score and then stabilized. This suggests that there is a polygenic model underlying the genetics of VCDR. No similar pattern was observed for the negative controls, i.e., the random variable used to evaluate false positive associations other than due to confounding by admixture (lowest observed $p = 0.204$). For the second negative control, based on sampling without replacement, increasing the p-value thresholds for the SNPs did not have any effect on the PEV, neither for the negative control based on random generated numbers (Figure 1B; Supplementary Figure S2).

Figures 1C and 1D present the corresponding graphs for IOP. For IOP, age and gender explained 2.5% of the total variance, while the first risk score based on a threshold of $p < 10^{-8}$ explained an additional 0.2% (region above the dotted line) of the variation in IOP in the target sample ERF. This score included only one SNP and did not reach nominal significance ($p = 0.093$). However, the addition of more SNPs did not yield any increase in PEV. None of the more extended risk scores were even marginally significant.

To evaluate whether there is a shared genetic component between VCDR and OAG and IOP and OAG, we tested the association of risk scores based on VCDR and IOP in the discovery sample with OAG cases and controls in RS-I. Table 3 shows the descriptive data of the OAG cases and controls in RS-I. A total of 5,304 participants of RS-I had complete data for optic nerve head measurements, IOP and reliable visual fields. Of these, 171 were classified as having OAG. This classification was based on the presence or absence of glaucomatous visual field loss. OAG cases were significantly older and more often men compared to controls. Not surprisingly, OAG cases had a larger VCDR and an increased IOP compared to controls, although these measures were not part of the classification process.

Figure 2A displays the PEV (Nagelkerke R-square) of OAG in RS-I attributable to VCDR genetic risk scores. Age and gender explained 4.0% of the variance of OAG and the first risk score based on the p-value threshold of $p < 10^{-10}$ (first bar) explained only an additional 0.3% of the OAG variance. However, when more SNPs with higher p-values were added to the score, the PEV increased sharply from $p < 10^{-2}$

and kept on increasing up to the inclusion of SNPs in the risk score with $p < 0.3$ in the discovery set. The latter score explained up to 4.7% of the variance of OAG above the attribution of age and gender, which is more than what is explained by age and gender together. The pattern of Figure 2A suggests a polygenic model for OAG as well and also a large genetic component shared by both VCDR and OAG. In a secondary analysis we excluded RS-I from the meta-analysis of the discovery cohorts and found a similar trend (Figure 3A). Nevertheless, as a consequence the PEV was much lower due to a major reduction in power. Results for IOP were similar to VCDR if we included RS-I, in that the risk scores were significantly associated with OAG (Figure 2B). However, in contrast to VCDR the evidence for a polygenic model underlying IOP disappeared in the secondary analyses (i.e. after excluding RS-I from the meta-analysis of the discovery cohorts; Figure 3B).

DISCUSSION

In the present study of GWAS data from Caucasian participants, we found evidence for a polygenic model underlying VCDR, one of the major known measures for OAG. In contrast, IOP did not show significant evidence for a polygenic model in the current study. Of these two traits, VCDR is the most relevant trait related to the diagnosis of OAG, while IOP is the most relevant trait for OAG treatment in patients with elevated IOP. Of interest is the finding that the risk scores based on the VCDR also predicted OAG. Although both VCDR and elevated IOP are putative endophenotypes for OAG, only VCDR showed evidence for a joint polygenic origin with OAG.

In the approach of the present study a large discovery cohort is crucial, while the size of the target cohort is less important. Therefore, we combined data of three independent cohorts from the Rotterdam Study, and used a single target cohort (ERF). We also repeated our analysis in which we used RS-III as a second replication cohort, rather than as one of the discovery cohorts. The results were similar, further supporting the evidence for a polygenic model of VCDR (data not shown).

At first sight, it seems unexpected that we did not find a similar model for IOP as for VCDR, because an

elevated IOP has been implicated as the most important determinants of both glaucomatous optic neuropathy (often referred to as an increased VCDR) and glaucomatous visual field loss. In some patients with OAG the IOP is not elevated. This form of disease is referred to as normal tension glaucoma, which represents about 15-40% of all patients with OAG.[26] In these patients, however, IOP lowering also causes a decrease in the rate of disease progression, suggesting that IOP matters in normal tension glaucoma as well. Due to the relatively low number of cases in our study, we did not separately analyze patients with normal tension glaucoma. For both traits, IOP and VCDR, strong evidence of genetic correlation with OAG susceptibility has been found. These two traits also correlate with each other in our population (Pearson's $p < 0.001$) and in other populations.[27] Nonetheless, when we tested whether the risk scores based on IOP predicted VCDR in the target cohort we found 0.54% as the highest significant PEV for VCDR of IOP-related SNPs (at $p < 10^{-8}$; data not shown). This finding suggests there is little overlap in the genes determining IOP and VCDR. Combined with the current findings this suggests that there are other, possibly rare variants with strong effects or yet unidentified environmental risk factors that determine IOP. However, a drawback of our analysis of IOP is that we did not adjust for the central corneal thickness, while the measurement of IOP may also be affected by central corneal thickness.[28]

All OAG cases were only derived from RS-I. A potential problem in the interpretation of the findings on OAG might be that the OAG cases derived from RS-I were also included in the discovery cohorts for the VCDR genetic risk scores. Although this may create an autocorrelation between the VCDR risk scores defined in the discovery set and OAG prediction conducted in the same set, this problem also occurred for the IOP scores. As the IOP score did not associate with the risk of OAG, it appears unlikely that any autocorrelation bias explained the overlap in genes involved in VCDR and OAG. Furthermore, it is important to realize that the diagnosis for OAG in RS-I was primarily based on glaucomatous visual field loss, thus not on VCDR or IOP. Moreover, the results did not alter when we excluded RS-I from the discovery cohorts.

As far as we know, the current study is the first to describe the phenomenon of a polygenic model for optic disc cupping (expressed in the VCDR). We further showed that the same risk scores predicted

OAG. This finding sheds new light on the etiology of OAG. The amount of variance explained by the VCDR risk scores exceeded that of age and gender. Nevertheless the PEV of both was still too low to allow the prediction of OAG in persons at risk.

Rare variants with large effects have also been implicated in the etiology of OAG. However, the current data suggest that many SNPs (probably with very small effects) may collectively account for a substantial proportion of variation in VCDR. This implies that many common variants have not been identified yet, because of low effect estimates of risks associated with single SNPs. Larger GWAS are needed to detect those variants and to further replicate our findings.

In conclusion, the present study has three major implications. First, in our large epidemiological study there is little overlap between the genes involved in VCDR and IOP. Second, the current study provides evidence for a polygenic model in VCDR involving many common SNPs that are not shared by a random variable, but specific to VCDR itself. This polygenic basis is not shared by the IOP, which may suggest that either the trait IOP has underlying rare variants, or the variants involved in IOP have such small effects that they were not well detected in the discovery set because of lack of power. Third, the genetics of VCDR overlap convincingly with that of OAG.

ACKNOWLEDGEMENTS

Financial support: The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. The generation and management of genome-wide association studies (GWAS) genotype data for the Rotterdam Study is supported by the Netherlands Organisation of Scientific Research NWO Investments (nr. 175.010.2005.011, 911-03-012). This study is funded by the Research Institute for Diseases in the Elderly (014-93-015; RIDE2), the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO)/Netherlands Consortium for Healthy Aging (NCHA) project nr. 050-060-810. The genetic study in the Erasmus Rucphen (ERF) Study was supported by the Center for Medical Systems Biology (CMSB) of NGI. The ophthalmic part of the Rotterdam Study and ERF are supported by the Netherlands Organization for Health Research and Development (ZonMw) grant 2200.0035; Lijf en Leven, Krimpen a/d Lek; MD Fonds, Utrecht. Oogfonds Nederland, Utrecht; Stichting Nederlands Oogheekkundig Onderzoek, Nijmegen/Rotterdam; Swart van Essen, Rotterdam; Netherlands Organisation for Scientific Research (NWO); Bevordering van Volkskracht, Rotterdam; Blindenhulp, The Hague; Landelijke Stichting voor Blinden en Slechtzienden (LSBS), Utrecht; Rotterdamse Vereniging voor Blindenbelangen, Rotterdam; OOG, The Hague; Algemene Nederlandse Vereniging ter Voorkoming van Blindheid (ANVVB), Doorn; The Rotterdam Eye Hospital Research Foundation (Stichting Wetenschappelijk Onderzoek Het Oogziekenhuis [SWOO] Prof. Dr. H.J. Flieringa, Rotterdam); Laméris Ootech BV, Nieuwegein; Topcon Europe BV, Capelle aan de IJssel, all in the Netherlands, and Heidelberg Engineering, Dossenheim, Germany.

The sponsors or funding organizations had no role in the design, conduct, analysis or publication of this research.

We thank Pascal Arp, Mila Jhamai, Dr. Michael Moorhouse, Jeannette Vergeer, Manijn Verkerk, and Sander Bervoets for their help in creating the GWAS database, and Dolinda Pottuit and Raph de Haas for their help in analyzing the optic disc images. The authors are grateful to the study participants, the staff

from the Rotterdam and ERF Study and the participating general practitioners and pharmacists.

REFERENCES

- [1] Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. *Br J Ophthalmol*. 2006 Mar;90(3):262-7.
- [2] Wolfs RC, Klaver CC, Ramrattan RS, van Duijn CM, Hofman A, de Jong PT. Genetic risk of primary open-angle glaucoma. Population-based familial aggregation study. *Archives of ophthalmology*. 1998 Dec;116(12):1640-5.
- [3] Allingham RR, Liu Y, Rhee DJ. The genetics of primary open-angle glaucoma: a review. *Exp Eye Res*. 2009 Apr;88(4):837-44.
- [4] Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, McCarthy MI, Ramos EM, Cardon LR, Chakravarti A, Cho JH, Guttmacher AE, Kong A, Kruglyak L, Mardis E, Rotimi CN, Slatkin M, Valle D, Whittemore AS, Boehnke M, Clark AG, Eichler EE, Gibson G, Haines JL, Mackay TF, McCarroll SA, Visscher PM. Finding the missing heritability of complex diseases. *Nature*. 2009 Oct 8;461(7265):747-53.
- [5] Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, Sklar P. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009 Aug 6;460(7256):748-52.
- [6] Fisher RA. The correlation between relatives on the supposition of Mendelian inheritance. *Philosophical Transactions of the Royal Society of Edinburgh*. 1918;52:399-433.
- [7] Janssens AC, van Duijn CM. Genome-based prediction of common diseases: advances and prospects. *Hum Mol Genet*. 2008 Oct 15;17(R2):R166-73.
- [8] Kwon YH, Fingert JH, Kuehn MH, Alward WL. Primary open-angle glaucoma. *N Engl J Med*. 2009 Mar 12;360(11):1113-24.
- [9] Quigley HA. Open-angle glaucoma. *N Engl J Med*. 1993 Apr 15;328(15):1097-106.
- [10] van Koolwijk LM, Despriet DD, van Duijn CM, Pardo Cortes LM, Vingerling JR, Aulchenko YS, Oostra BA, Klaver CC, Lemij HG. Genetic contributions to glaucoma: heritability of intraocular pressure, retinal nerve fiber layer thickness, and optic disc morphology. *Invest Ophthalmol Vis Sci*. 2007 Aug;48(8):3669-76.

- [11] Hofman A, Breteler MM, van Duijn CM, Krestin GP, Pols HA, Stricker BH, Tiemeier H, Uitterlinden AG, Vingerling JR, Witteman JC. The Rotterdam Study: objectives and design update. *Eur J Epidemiol.* 2007;22(11):819-29.
- [12] Stricker BH, Stijnen T. Analysis of individual drug use as a time-varying determinant of exposure in prospective population-based cohort studies. *Eur J Epidemiol.* 2010 Apr;25(4):245-51.
- [13] Hofman A, Breteler MM, van Duijn CM, Janssen HL, Krestin GP, Kuipers EJ, Stricker BH, Tiemeier H, Uitterlinden AG, Vingerling JR, Witteman JC. The Rotterdam Study: 2010 objectives and design update. *Eur J Epidemiol.* 2009;24(9):553-72.
- [14] Aulchenko YS, Heutink P, Mackay I, Bertoli-Avella AM, Pullen J, Vaessen N, Rademaker TA, Sandkuijl LA, Cardon L, Oostra B, van Duijn CM. Linkage disequilibrium in young genetically isolated Dutch population. *Eur J Hum Genet.* 2004 Jul;12(7):527-34.
- [15] Pardo LM, MacKay I, Oostra B, van Duijn CM, Aulchenko YS. The effect of genetic drift in a young genetically isolated population. *Ann Hum Genet.* 2005 May;69(Pt 3):288-95.
- [16] Dielemans I, Vingerling JR, Hofman A, Grobbee DE, de Jong PT. Reliability of intraocular pressure measurement with the Goldmann applanation tonometer in epidemiological studies. *Graefes Arch Clin Exp Ophthalmol.* 1994 Mar;32(3):141-4.
- [17] Wolfs RC, Borger PH, Ramrattan RS, Klaver CC, Hulsman CA, Hofman A, Vingerling JR, Hitchings RA, de Jong PT. Changing views on open-angle glaucoma: definitions and prevalences--The Rotterdam Study. *Invest Ophthalmol Vis Sci.* 2000 Oct;41(11):3309-21.
- [18] Czudowska MA, Ramdas WD, Wolfs RC, Hofman A, Jong PT, Vingerling JR, Jansonius NM. Incidence of Glaucomatous Visual Field Loss: A Ten-Year Follow-up from the Rotterdam Study. *Ophthalmology.* 2010 Jun 28.
- [19] Rolando M, Iester M, Campagna P, Borgia L, Traverso C, Calabria G. Measurement variability in digital analysis of optic discs. *Doc Ophthalmol.* 1994;85(3):211-22.
- [20] Ikram MK, Borger PH, Assink JJ, Jonas JB, Hofman A, de Jong PT. Comparing ophthalmoscopy, slide viewing, and semiautomated systems in optic disc morphometry. *Ophthalmology.* 2002 Mar;109(3):486-93.

- [21] Rivadeneira F, Styrkarsdottir U, Estrada K, Halldorsson BV, Hsu YH, Richards JB, Zillikens MC, Kavvoura FK, Amin N, Aulchenko YS, Cupples LA, Deloukas P, Demissie S, Grundberg E, Hofman A, Kong A, Karasik D, van Meurs JB, Oostra B, Pastinen T, Pols HA, Sigurdsson G, Soranzo N, Thorleifsson G, Thorsteinsdottir U, Williams FM, Wilson SG, Zhou Y, Ralston SH, van Duijn CM, Spector T, Kiel DP, Stefansson K, Ioannidis JP, Uitterlinden AG. Twenty bone-mineral-density loci identified by large-scale meta-analysis of genome-wide association studies. *Nat Genet.* 2009 Nov;41(11):1199-206.
- [22] Sanchez-Meca J, Marin-Martinez F. Weighting by Inverse Variance or by Sample Size in Meta-Analysis: A Simulation Study. *Educational and Psychological Measurement.* 1998;58(2):211-20.
- [23] Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *American journal of human genetics.* 2007 Sep;81(3):559-75.
- [24] Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *American journal of human genetics.* 1998 May;62(5):1198-211.
- [25] Liu F, Wollstein A, Hysi PG, Ankra-Badu GA, Spector TD, Park D, Zhu G, Larsson M, Duffy DL, Montgomery GW, Mackey DA, Walsh S, Lao O, Hofman A, Rivadeneira F, Vingerling JR, Uitterlinden AG, Martin NG, Hammond CJ, Kayser M. Digital quantification of human eye color highlights genetic association of three new loci. *PLoS genetics.* 2010;6:e1000934.
- [26] Dielemans I, Vingerling JR, Wolfs RC, Hofman A, Grobbee DE, de Jong PT. The prevalence of primary open-angle glaucoma in a population-based study in The Netherlands. The Rotterdam Study. *Ophthalmology.* 1994 Nov;101(11):1851-5.
- [27] Charlesworth J, Kramer PL, Dyer T, Diego V, Samples JR, Craig JE, Mackey DA, Hewitt AW, Blangero J, Wirtz MK. The path to open-angle glaucoma gene discovery: endophenotypic status of intraocular pressure, cup-to-disc ratio, and central corneal thickness. *Invest Ophthalmol Vis Sci.* 2010 Jul;51(7):3509-14.
- [28] Wolfs RC, Klaver CC, Vingerling JR, Grobbee DE, Hofman A, de Jong PT. Distribution of central corneal thickness and its association with intraocular pressure: The Rotterdam Study. *Am J Ophthalmol.* 1997 Jun;123(6):767-72.

LICENCE STATEMENT

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in JMG and any other BMJ PGL products and sublicences such use and exploit all subsidiary rights, as set out in our licence (<http://group.bmj.com/products/journals/instructions-for-authors/licence-forms>)."

FIGURE LEGENDS

Figure 1. The explained variance of risk scores of the vertical cup-disc ratio (VCDR) and intraocular pressure (IOP) for each p-value threshold for their respective trait in the target cohort ERF (A and C, respectively); and the explained variance of the risk scores of VCDR and IOP for each p-value threshold for the negative control (B and D, respectively).

The y1-axis (left; histogram) depicts the explained variance (in %) against different p-value thresholds on the x-axis that were used to construct the risk scores. The y2-axis (right; line) depicts the p-values from the association analysis of the risk scores with the trait. Dotted line represents the attribution of age and gender to the explained variance.

Figure 2. The Nagelkerke R-square of risk scores of the vertical cup-disc ratio (VCDR) and intraocular pressure (IOP), adjusted for age and gender, for each p-value threshold on open-angle glaucoma (OAG) in RS-I (A and B, respectively).

The y1-axis (left; histogram) depicts the explained variance (in %) against different p-value thresholds on the x-axis, which were used to construct risk scores. The y2-axis (right; line) depicts the p-values from the association analysis of the risk scores with the trait.

Figure 3. The Nagelkerke R-square of risk scores of the vertical cup-disc ratio (VCDR) and intraocular pressure (IOP), adjusted for age and gender, for each p-value threshold on open-angle glaucoma (OAG) in RS-I (A and B, respectively) after excluding RS-I from the discovery cohort. To display the pattern more clear the attribution of age and gender has already been subtracted.

The y1-axis (left; histogram) depicts the explained variance (in %) against different p-value thresholds on the x-axis, which were used to construct risk scores. The y2-axis (right; line) depicts the p-values from the association analysis of the risk scores with the trait.

TABLES

Table 1. Characteristics of the four study populations, presented as mean $\pm$ standard deviation (range) unless stated otherwise

	Discovery cohorts			Target cohort
	RS-I	RS-II	RS-III	ERF
Total sample size (N)	5,312	2,048	1,966	1,646
Age (years)	68.0 $\pm$ 8.4 (55 - 99)	64.3 $\pm$ 7.8 (55 - 98)	55.6 $\pm$ 5.5 (45 - 89)	46.8 $\pm$ 14.1 (18 - 84)
Gender, N(%) women	3,099 (58.3)	1,109 (54.2)	1,102 (56.1)	942 (57.2)
Vertical cup-disc ratio*	0.50 $\pm$ 0.14 (0.00 - 0.89)	0.50 $\pm$ 0.14 (0.05 - 0.87)	0.42 $\pm$ 0.17 (0.00 - 1.00)	0.46 $\pm$ 0.15 (0.00 - 0.84)
Intraocular pressure (mmHg)**	14.8 $\pm$ 3.4 (5.0 - 58.5)	14.4 $\pm$ 3.4 (7.0 - 31.5)	13.6 $\pm$ 3.0 (5.0 - 30.0)	14.8 $\pm$ 3.0 (6.0 - 30.0)

* = In RS-I and RS-II measured with ImageNet and in RS-III and ERF with Heidelberg Retina

Tomograph; ** = Sample sizes for intraocular pressure analyses were 5,794 for RS-I, 2,102 for RS-II,

2,041 for RS-III, and 2,035 for ERF

Table 2. Number of single nucleotide polymorphisms (SNPs) included in each p-value category presented as N(%)

p-value threshold	Vertical cup-disc ratio	Intraocular pressure
<10 ⁻¹⁰	9 (0.0)	-
<10 ⁻⁹	10 (0.0)	-
<10 ⁻⁸	12 (0.0)	1 (0.0)
<10 ⁻⁷	18 (0.0)	2 (0.0)
<10 ⁻⁶	34 (0.0)	4 (0.0)
<10 ⁻⁵	61 (0.0)	13 (0.0)
<10 ⁻⁴	179 (0.0)	88 (0.0)
<10 ⁻³	966 (0.2)	755 (0.1)
<10 ⁻²	7,260 (1.4)	6429 (1.2)
<0.1	60029 (11.5)	56,798 (10.9)
<0.2	114,122 (21.8)	110,338 (21.1)
<0.3	166,995 (31.9)	163,261 (31.2)
<0.4	219,096 (41.9)	215,470 (41.2)
<0.5	270,510 (51.7)	267,418 (51.2)
<0.6	321,288 (61.5)	318,625 (60.9)
<0.7	371,823 (71.1)	369,764 (70.7)
<0.8	422,175 (80.8)	420,547 (80.4)
<0.9	472,496 (90.4)	471,532 (90.2)
<1.0	522,762 (100.0)	522,782 (100.0)

Table 3. Descriptive data of open-angle glaucoma cases and controls in RS-I, presented as mean $\pm$ standard deviation (range) unless stated otherwise

	Cases (N=171)	Controls (N=5,133)	p-value	RS-I (Total N=5,304)
Age (years)	77.4 $\pm$ 7.2 (56 – 94)	74.0 $\pm$ 7.5 (55 – 99)	< 0.001*	74.1 $\pm$ 7.5 (55 – 99)
Gender, N(%) women	81 (47.4)	3,013 (58.7)	< 0.001*	3,094 (58.3)
Vertical cup-disc ratio	0.59 $\pm$ 0.15 (0.15 – 0.89)	0.50 $\pm$ 0.14 (0.00 – 0.83)	< 0.001*	0.50 $\pm$ 0.14 (0.00 – 0.89)
Intraocular pressure (mmHg)	16.8 $\pm$ 4.7 (5.5 – 39.7)	14.7 $\pm$ 3.2 (5.0 – 35.8)	<0.001*	14.7 $\pm$ 3.3 (5.0 – 39.7)

* = significant at a p-value of 0.05