



Limiting mycotoxins in stored wheat

Naresh Magan, David Aldred, Kalliopi Mylona, Ronald J.W. Lambert

► To cite this version:

Naresh Magan, David Aldred, Kalliopi Mylona, Ronald J.W. Lambert. Limiting mycotoxins in stored wheat. Food Additives and Contaminants, 2010, 27 (05), pp.644-650. 10.1080/19440040903514523 . hal-00591170

HAL Id: hal-00591170

<https://hal.science/hal-00591170>

Submitted on 7 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Limiting mycotoxins in stored wheat

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2009-333.R1
Manuscript Type:	Special Issue
Date Submitted by the Author:	26-Nov-2009
Complete List of Authors:	Magan, Naresh; Cranfield University, Applied Mycology Aldred, David; Cranfield University, Applied Mycology Group Mylona, Kalliopi; Cranfield University, Applied Mycology Group Lambert, Ronald; Cranfield University, Applied Mycology Group
Methods/Techniques:	Mycology, Quality assurance
Additives/Contaminants:	Mycotoxins - trichothecenes, Mycotoxins - fungi, Antioxidants
Food Types:	Cereals and grain, Animal feed

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Limiting mycotoxins in stored wheat – a review

Naresh Magan, David Aldred, Kalliopi Mylona and Ronald J.W. Lambert

Applied Mycology Group, Cranfield Health, Cranfield University, Bedford MK43 0AL, U.K.

Corresponding author: Prof. N. Magan, Applied Mycology Group, Cranfield Health,
Cranfield University, Bedford MK43 0AL, U.K. Tel: +44 1234 758308; fax: +44 1234
758380; E.mail: n.magan@cranfield.ac.uk

Key words: spoilage fungi, mycotoxins, dry matter losses, boundary models, modified
atmospheres, preservatives, biocontrol, post-harvest management

Abstract

The quality of harvested wheat grain can deteriorate markedly during the post-harvest management stages. The biotic factors such as grain type and ripeness, coupled with the prevailing abiotic factors such as water content and temperature, and also preservative concentration will influence the safe storage life and the level of contamination with mycotoxins such as deoxynivalenol (DON) produced pre-harvest and zearalenone (ZEA) produced post-harvest by *Fusarium graminearum* and *Fusarium poae* respectively, ochratoxin (OTA) produced by *Penicillium verrucosum* post-harvest in cool damp northern European climates, and perhaps T-2 and HT-2 toxins produced by *Fusarium langsethiae*. This review presents recent data on the relationship between dry matter losses caused by *F. graminearum* under different environmental regimes (water activities, temperatures) and the level of contamination with DON. This is important as poor post-harvest drying and storage management may exacerbate DON contamination already present pre-harvest. It is thus critical to relate the environmental factors in stored wheat grain during storage, especially of intergranular relative humidity (RH) and temperature, to safe storage periods without spoilage or risk from increased DON contamination. The growth/no growth and DON/no DON (*F. graminearum*) and OTA/no toxin production (*P. verrucosum*) have been used to build a model with a simple interface to link the temperature and RH values to the potential risk level which may allow growth or toxin production. This paper also considers the use of modified atmospheres, preservatives and biocontrol to minimise DON and OTA in moist wheat grain. These approaches together with clear monitoring criteria and hygiene could contribute to better post-harvest management of stored temperate cereals and ensure that mycotoxin contamination is minimised during this key phase in the food/feed chain.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

In Europe the key approach to minimising natural contaminants in the food chain has been by developing minimisation strategies within a HACCP framework. Thus, the cereal food chain for different end products has been studied in detail (Aldred and Magan, 2004; Aldred et al., 2004; Magan et al., 2008). This has included the monitoring of critical control points (CCPs) both pre- and post-harvest. With regard to mycotoxins as a hazard in temperate cereals (DON, OTA) the main drivers have included EU legislative limits, customer specifications and consumer perceptions. A prevention strategy must aim to minimise the occurrence of these mycotoxins below the legislative limits based on best practice, both pre- and post-harvest. The key CCPs post-harvest have been identified for the wheat food/feed chain. In temperate cereals the main mycotoxins of concern are ochratoxin OTA (*Penicillium verrucosum*), deoxynivalenol (DON; *Fusarium graminearum*, *Fusarium culmorum*), zearalenone (*F. graminearum*, *Fusarium poae*), and T-2 and HT-2 toxins (*Fusarium langsethiae*). There are legislative limits for the first three of these and for the latter two, legislation is imminent.

Cereal grain, whether effectively dried or slightly moist, is alive and respiring and thus has to be treated with care during harvesting, drying, storage and subsequent downstream processing along the food chain. The concept of considering stored grain as an ecosystem was originally developed by Wallace and Sinha (1971). This was an important milestone as they included all the key biotic and abiotic interactions which might occur during storage including grain respiration, insect pests, mycoflora, insect/fungal interactions, pest immigration and emigration, heterogeneity of moisture and moisture migration, temperature and O₂/CO₂, and for animal feed the presence of preservatives. This holistic approach was

instrumental in facilitating some of the advances made during the last decade (Magan & Aldred, 2007).

Respiration, dry matter losses and mycotoxin contamination

Harvested grain must be dried to <14.5% moisture content (m.c., wet weight basis) to ensure that no mould spoilage or any pre-harvest contamination with mycotoxins is exacerbated. Often, grain is harvested at 16-20% m.c. because of the better efficiency of modern combine harvesters. However, this does subsequently require heated or ambient drying to be used to conserve grain quality. Where drying has not been carried out effectively, the temperature will rise quickly and a succession of spoilage moulds may develop resulting in spontaneous heating with temperatures of 60-70°C being reached, ending with thermophilic actinomycetes and complete loss of quality. This is accompanied by a significant increase in the chance for mycotoxin contamination.

Studies have been carried out to examine the relationship between grain respiration, mould activity and quality loss (dry matter) during these critical periods between harvesting, drying and storage. These have shown that in wheat and barley, dry matter losses of only 0.22 – 0.44% at 20-25°C and 0.90 water activity ($a_w = 19-21\%$ m.c.) results in visible moulding already being present (Lacey et al., 1999). Indeed, studies in the USA suggested that in other cereals such as maize, losses of 0.5% dry matter can lead to rejection of the grain for human consumption and that this was accompanied by a significant chance of aflatoxin contamination (Seitz et al., 1982). Fleurat-Lessard (2002) extensively reviewed data on general spoilage and dry matter losses and suggested that the rate of CO₂ production in cereals can be effectively used to model and predict global changes in cereal quality.

1
2
3 100 Recent studies at Cranfield University have tried to relate the activity of specific
4
5 101 mycotoxigenic species under different environmental factors which might contribute to both
6
7
8 102 dry matter losses and the chances of exceeding the EU legislative limits for DON. Thus
9
10 103 experiments were carried out with stored winter wheat inoculated with spores of *F.*
11
12 104 *graminearum* at 0.86, 0.93 and 0.95 a_w and different temperatures (15-30°C) in chambers
13
14 105 with the same equilibrium relative humidity atmosphere and measuring the respiration using
15
16 106 GC analyses over periods of 4-5 days. The final *F. graminearum* populations and DON
17
18 107 contamination after 10 days were also analysed. The effect of temperature x a_w on total
19
20 108 calculated dry matter losses over periods of 4 days storage are shown in Figure 1. This shows
21
22 109 the effect that these two parameters have on colonisation rates and quality loss. DON
23
24 110 concentrations were found to be above the legislative limits (>1250 ppb) at 30°C and 0.95
25
26 111 and 0.93 a_w in only 10 days, while at 15-20°C and all a_w levels the concentrations were less
27
28 112 than the legislative limits over these short periods. Similar data on dry matter losses caused
29
30 113 by *P. verrucosum* over 7-14 days showed that dry matter losses were also between 1-1.5% at
31
32 114 0.95 a_w , and lower at 0.85-0.90 a_w and 25°C (Magan and Aldred, 2007). This approach may
33
34 115 be beneficial for combining dry matter losses in relation to m.c.s during drying and initial
35
36 116 storage with the range of potential risk conditions for OTA and DON, relative to the EU
37
38 117 legislative limits. Previous comprehensive respiration data sets in relation to a_w and
39
40 118 temperature of stored wheat, have been used to develop the possible relationship between the
41
42 119 variables. Thus, the results obtained as $O_2\ d^{-1}\ kg^{-1}$ dry grain (R) was divided by the incubation
43
44 120 temperature ($\square$) to give units of $O_2\ d^{-1}\ kg^{-1}$ dry matter $^{\circ}C^{-1}$. There was a linear relationship
45
46 121 ($r^2=0.9594$) between $R/\square$ and a_w (Figure 2, Lacey et al., 1997). This is based on O_2
47
48 122 consumption, while our new data is based on CO_2 production. Thus some conversion will be
49
50 123 required to relate our new data to the previous data. However, it may be possible to relate our
51
52 124 present data to the risk of exceeding the EU legislative limits for mycotoxigenic fungi. This
53
54
55
56
57
58
59
60

could then be a valuable practical aid if developed for different grains and nuts prone to contamination with different mycotoxins.

Boundary models of growth/no growth and mycotoxin/no mycotoxin and stored grain management

It is essential that grain is effectively dried immediately after harvest. Often heated air drying is used to achieve the target safe moisture content of about 14.5%. However, sometimes in the harvesting season grain is left for short periods of time before drying because of logistical issues. In some European countries ambient air drying is always used. Where heated drying is used the grain is less affected by the prevailing weather conditions, especially in damp autumn seasons in northern Europe. In the UK, where ambient drying is often used, this can result in slow drying with a moist front moving vertically up through the grain bed, especially in flat bed dryers/stores. This can result in grain at the top of the store being rewetted, while grain at the bottom can become over dried. This can lead to layers of mouldy grain with the potential for increased DON or OTA contamination.

Measurements can be made within stored grain silos to measure the changes in temperature and relative humidity in the inter-granular air spaces in the grain during drying and storage. This has been done at the top and in the middle of a flat bed store. Generally, any leaks or even small openings via damage to the structure can lead to an increase in temperature and intergranular m.c. This can often occur during the early periods after drying and storage phase (5-10 days) where both these parameters can increase, especially in the top layers which can increase the risk from spoilage and mycotoxin contamination (Magan et al., 2008).

1
2
3 150 We have previously developed boundary layer contour maps for growth and DON and OTA
4
5
6 151 production by *F. culmorum* and *F. graminearum* and for *P. verrucosum* (Hope and Magan,
7
8 152 2003; Hope et al., 2004; Cairns et al., 2005). This data is useful as a guide to conditions
9
10 153 where there may be a risk of growth and toxin production. A polynomial equation has been
11
12 154 used to model these growth/no growth and toxin/no toxin boundaries. Figure 3 shows an
13
14 155 example of the data for *F.graminearum* in relation to minimal growth rates (0.1 mm day⁻¹)
15
16 156 and for DON production (0.01 ppm).
17
18
19
20 157

21
22 158 It is possible to link the temperature and RH levels monitored in the intergranular atmosphere
23
24 159 to these boundary models. Thus, by inputting the data into the model one can check directly
25
26 160 whether there is a risk of conditions conducive to growth or to DON production. The inset in
27
28 161 Figure 3 shows the simple front end diagram which can be used for this purpose. This would
29
30 162 enable dials related to temperature and RH to be moved to determine a low (green colour) or
31
32 163 high (red colour) risk of growth or mycotoxin (DON, OTA) in store. The model could easily
33
34 164 be modified to include the legislative limits for specific mycotoxins and applied to a range of
35
36 165 mycotoxigenic fungi, and food commodities. We have developed the growth and toxin
37
38 166 models for *A. flavus* (aflatoxin), *F. verticillioides* (fumonisins), *F. culmorum* (DON) and *P.*
39
40 167 *verrucosum* (OTA) and for the new *A.ochraceus* grouping (Abdel-Hadi & Magan, 2009).
41
42
43
44 168

45
46
47
48 169 Other models have also been developed for *P.verrucosum* and OTA production by Lindblad
49
50 170 et al. (2004). They developed very useful models which relate the potential for exceeding the
51
52 171 EU legislative limit for OTA in cereals to the CFUs isolated from grain. They showed that
53
54 172 $>10^3$ CFUS g⁻¹ on a selective medium for *P. verrucosum* would represent a risk of OTA
55
56 173 contamination >5 µg kg⁻¹ stored grain which is the EU legislative limit. They also developed
57
58 174 models of toxin free storage times under different temperature and moisture content
59
60

conditions. These are very useful practically for the cereal industry. Studies by Jonsson et al. (2000) suggest that the maximum storage time without any visible moulding was probably reduced by 50% if the mc at harvest was decreased by 1-3% ($=0.05 a_w$) or if the storage temperature was increased by 5°C, for example in a mild autumn year, in temperate cereals.

Modified atmosphere storage, chemical preservation systems and biocontrol

Moist grain, especially for animal feed use is sometimes stored in sealed silos to enable the natural respiration of the grain to increase the CO₂ concentration and inhibit spoilage moulds and the potential for mycotoxin formation post-harvest. However, if there are any leaks or the conditions remain micro-aerophilic then there is potential for both growth and mycotoxin formation. Studies suggest that increasing concentrations of CO₂ to 25 or 50% has little effect on growth or OTA production by *P. verrucosum* or indeed *Aspergillus ochraceus* (*=Aspergillus westerdijkiae*). Spore germination of these two fungi is unaffected, although germ tube extension and growth can be affected by up to 75% CO₂ (Cairns-Fuller, 2004; Cairns-Fuller et al., 2005). However, this does not mean that mycotoxin production is significantly affected. While inhibition of OTA did occur when a_w and CO₂ were increased, the effect was not synergistic and sometimes inhibition was not to a level below the legislative limits.

The effect of modified atmosphere on *F. culmorum* and other species has been examined and showed that they are micro-aerophilic and thus very tolerant of very low O₂ concentrations (Magan & Lacey, 1984). Paster (2000) has reviewed the effect of modified atmospheres on mycotoxigenic and other moulds and this has shown that there is very little information on mycotoxigenic Fusaria. Work has been carried out with *Fusarium sporotrichioides* and this demonstrated that T-2 toxin was reduced by 80% in the presence of 50% CO₂/20%O₂ but that

mycelial colonisation was not affected by <60% CO₂ (Paster et al., 1986; Paster and Menashero, 1988).

Preservatives are often added to moist grain, especially for grain destined for animal feed. Commercial products are mainly based on mixtures of salts of aliphatic acids such as propionic, sorbic and benzoic acids. However, these are all fungistats and require effective contact with the product. Thus mixing is very important to ensure that inhibition of spoilage is maintained. Work has been carried out to try and identify alternative compounds which may be used to replace these aliphatic acid based products. Since *P. verrucosum* often causes problems post-harvest then the use of natural and novel preservatives may have an impact on OTA contamination. Aldred et al. (2008) screened a number of essential oils and antioxidants for control of both growth and OTA production by both *P. verrucosum* and *A. ochraceus*. They found that while essential oils such as clove, cinnamon and thyme were effective in vitro, in situ on grain they were relatively ineffective. This may perhaps be due to binding of the essential oil components to grain constituents. However, they did find that anti-oxidants, such as butylhydroxyanisole (BHA), propyl paraben, and especially resveratrol, were effective on naturally contaminated and grain artificially inoculated with *P. verrucosum*.

A range of essential oils (23) and 6 antioxidants were screened for control of growth and DON and NIV production by *F. culmorum*, DON production by *F. graminearum* and T-2 and HT-2 by *F. langsethiae* (Hope, 2004). Of these, three essential oils (bay, clove, cinnamon oil) and two antioxidants (propyl paraben and hydroxymethylanisole) were found to be effective in controlling growth of these species in the range 50-200 ppm at 15 and 25°C at 0.995 and 0.955 a_w in vitro on wheat-based media. More detailed studies on stored wheat grain at 15 and 25°C and essential oil/antioxidants treatments (100ppm; 500ppm) were carried out. This

showed that 500 ppm significantly reduced or inhibited growth at all the environmental combinations examined. However, at 100 ppm the essential oils/antioxidants stimulated growth under some experimental condition. Cinnamon and clove essential oils were the most effective inhibitors of growth regardless of temperature, a_w or species. The essential oils and antioxidants had a variable effect on inhibition of mycotoxin production. BHA and clove oil inhibited DON at both 100 and 500 ppm. However, propyl paraben and cinnamon oil enhanced NIV production at 100 ppm and intermediate a_w levels. Table 1 shows the efficacy of 500 ppm of these potential preservatives on control of DON and T-2 and HT-2 by *F. graminearum* and *F. langsethiae*, respectively, when inoculated onto wheat grain and stored for 28 days.

Studies have been carried out to use inoculants such as yeasts or lactic acid bacteria to control stored moist cereals destined for animal feed. The applied inoculants need to be able to become established and dominate the stored grain ecosystem to minimise the contamination with mycotoxins. A significant amount of work has been carried out to use xerotolerant yeasts such as *Pichia anomala* to control spoilage of feed grain, especially in airtight storage. It has been shown that this competitive yeast can prevent spoilage by micro-aerophilic fungi such as *Penicillium roqueforti* (Druverfors et al., 2002). Recently, different fluidised bed dried formulations of this yeast has been examined for the specific control of *P. verrucosum* and OTA contamination of stored wheat (Mokiou and Magan, 2008). These studies showed that formulation had an impact on the efficacy of control of both *P. verrucosum* populations and the relative effectiveness of controlling OTA in stored wheat grain.

Many lactic acid bacteria (e.g. *Lactobacillus acidophilus* strains) have been screened for the ability to either degrade or bind different mycotoxins (Fuchs et al., 2008). They have become

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

attractive inoculants because of the indication that they can give beneficial health effects. Studies have suggested that they can detoxify aflatoxin B1. Others studies suggest that they may bind or degrade zearalenone, trichothecenes and fumonisins (El-Nezami et al., 2002). However, less work has been carried out in situ under realistic storage conditions, especially intermediate moisture and temperature conditions. Potential does exist for this approach to pay dividends in relation to preservation of moist cereals specifically for animal feed purposes.

Conclusions and future strategies

There are a number of key points which are important post-harvest in any management system to minimise mycotoxin contamination especially downstream for processing purposes. These include

- (a) regular and accurate moisture determination,
- (b) minimising time between harvesting and drying; this will relate to holding time/temperature prior to drying as well as the actual drying conditions,
- (c) efficient and prompt drying of wet grain to target moisture contents (temperate cereals <14.5%) by heated drying where possible as ambient drying can result in spoilage problems
- (d) infrastructure for quick response, including provision for segregation and appropriate transportation conditions,
- (e) appropriate storage conditions at all stages in terms of moisture and temperature control, and general maintenance and hygiene of facilities for prevention of pest and water ingress which can lead to pockets of metabolic water and initiate spoilage,
- (f) ability to efficiently identify and reject material below specified standards in terms of both fungal diseases and, at some stages, mycotoxin levels (e.g. when passing onto a third party),

(g) operation of approved supplier systems. This involves setting specifications for acceptance/rejection.

Other important areas which need to be considered are those in relation to better representative sampling and early and rapid detection of mycotoxigenic fungi post-harvest. Problems do still arise in relation to obtaining a representative sample both pre- and post-harvest. Whitaker (2004) and Whitaker et al. (2000) have shown that in stored cereals, nuts and other commodities that the error in obtaining a representative sample can be 25-60% of the total error. There are specific EU sampling plans (e.g. for DON; Commission Directive 2005/38/EC) which are prescribed but have been found cumbersome to use. The question arises as to whether one should use a regular grid type sampling approach or use a random sample approach. Which type of sampling regime should be employed may further be dictated by the type of mycotoxin as different contaminants may be distributed homogeneously or heterogeneously or spatially either in regular pockets or randomly. Recent research has been undertaken to try and use geostatistics to try and unravel this complex issue. This approach is beneficial if you have the exact grid positions of the samples in three dimensions, it can then be used on any scale. Recent studies by Rivas et al. (2009 a, b) suggest that mycotoxins such as DON, have a spatial structure in the store and thus a regular grid may be more precise than using a random sampling approach. This may not be the case for other mycotoxins and in other commodities and thus further work is imperative to enable more accurate representative samples to be taken and improve the accuracy of such sampling.

We believe that there are new technologies which need to be utilised in the future to improve the strategies to minimise mycotoxins in temperate cereals. As has been mentioned earlier the potential for using remote sensing especially in silos linked to post-harvest models of

1
2
3 298 mycotoxigenic moulds integrated with the legislative limits could be very useful. There are
4
5
6 299 now many useful molecular-based techniques which could be applied effectively post-
7
8 300 harvest. For example, Schmidt-Heydt et al. (2000) utilised the expression of the OTA
9
10 301 polyketide synthase (*otapksPV*) to examine temporal colonisation of stored wheat grain of
11
12 302 different moisture contents (14-24%) and this was correlated with OTA production analysed
13
14
15 303 by HPLC and populations by CFUs. Thus the RT-PCR approach could be used as a sensitive
16
17 304 measure of the quality status with regard to OTA contamination in wheat during 3 months
18
19 305 storage. These powerful molecular tools have been applied pre-harvest but not widely post-
20
21 306 harvest. This is supported by the recent work by Suanthe et al. (2009) who have developed
22
23 307 multiplex RT-PCR for detection and quantification of mycotoxigenic *Aspergillus*, *Penicillium*
24
25 308 and *Fusarium* species.
26
27
28
29 309
30
31 310 In the future another important potential impact is that environmental change. This may have
32
33 311 an impact on the relative importance of different mycotoxins in different geographical
34
35 312 regions. This may become important and have a significant impact on which mycotoxins may
36
37 313 become important in the next 5-10 years. There is a lack of data on the impact that twice or
38
39 314 three times the existing CO₂ concentrations combined with 2-3°C elevation in temperature
40
41 315 might have on colonisation and mycotoxin production. A certain amount of this information
42
43 316 can be generated by modelling approaches. However, the potential for stimulation of
44
45 317 mycotoxin production is more difficult to predict. In conclusion, while this paper has dealt
46
47 318 with post-harvest issues in temperate cereals it has to be seen in the context of a more holistic
48
49 319 food chain systems approach with better and more effective management post-harvest being
50
51 320 just one component of this.
52
53
54
55
56 321
57
58
59
60 322

References

- Abdel-Hadi, A. & Magan, N. (2009). Influence of environmental factors on growth, sporulation and ochratoxin A and B production of the new grouping of the *A. ochraceus* group. *World Mycot Journ* In Press.
- Aldred, D. & Magan, N. (2004). Prevention strategies for tricothecenes. *Toxicol Lett* 153: 165-171.
- Aldred D., Olsen, M. & Magan, N. (2004). The use of HACCP in the control of mycotoxins: the case for cereals. Chapter 7 In: *Mycotoxin in food: detection and control*, Eds. Magan, N. & Olsen, M., Woodhead Publishing, Cambridge, U.K. pp. 139-173.
- Aldred, D; Cairns-Fuller, V. & Magan, N. (2008). Environmental factors affect efficacy of some essential oils and resveratrol to control growth and ochratoxin A production by *Penicillium verrucosum* and *A. westerdijkiae* on wheat grain. *J Stored Prod Res* 44: 341-346.
- Cairns-Fuller, V., Aldred, D. & Magan, N. (2005). Water, temperature and gas composition interactions affect growth and ochratoxin A production by isolates of *Penicillium verrucosum* on wheat grain. *J Appl Microbiol* 99: 1215-1221.
- Druverfors U, Jonsson N, Boysen ME, Schnürer J. 2002. Efficacy of the biocontrol yeast *Pichia ananmala* during long term storage of moist feed grain under different oxygen and carbon dioxide regimens. *FEMS Yeast Research* 2: 389-394.
- Fuchs, S., Sontag, G., Stidl, R., Ehrlich, V., Kundi, M., Knasmüller, S. (2008). Detoxification of patulin and ochratoxin A, two abundant mycotoxins by lactic acid bacteria. *Food Chem Toxicol* 46: 1398-1407.
- El-Nezami, H., Chrevatidis, A., Auriola, S., Salminen, S., Mykkanen, H. (2002). Removal of common *Fusarium* toxins in vitro by strains of *Lactobacillus* and *Propionobacterium*. *Food Addit Contam* 19: 680-686.

- 348 Fleurat-Lessard F (2002). Qualitative reasoning and integrated management of the quality of
349 stored grain: a promising new approach. J Stored Prod Res 38: 191-218.
- 350 Hope, R. & Magan, N. (2003). Two dimensional environmental profiles of growth,
351 deoxynivalenol and nivalenol production by *Fusarium culmorum* on a wheat-based
352 substrate. Letts Appl Microbiol 37: 70-74.
- 353 Hope, R., Aldred, D. and Magan, N. (2005). Comparison of the effect of environmental factors
354 on deoxynivalenol production by *F. culmorum* and *F. graminearum* on wheat grain.
355 Letts Appl Microbiol 40: 295-300
- 356 Jonsson N, Petterson H and Schnurer J (2000). Study of the relationship between storage
357 conditions and the growth of moulds and production of Ochratoxin A in grain-
358 preliminary results. Proceedings of European Agricultural Engineering Symposium
359 2000, Warwick University, U.K. pp.
- 360 Lacey J, Hamer A and Magan N (1994). Respiration and losses in stored wheat under
361 different environmental conditions. In: Highley E, Wright EJ, Banks HJ and Champ
362 BR (eds), Proc. Of 6th Int. Working Conference on Stored Product Protection, CAB
363 International, Canberra, Australia, Vol II, pp. 1007-1013.
- 364 Lacey, J., Hamer, A. and Magan, N. (1997). Respiration of wheat grain stored in different
365 environments. In Proc. Int. Conf. Controlled Atmosphere and Fumigation of Stored
366 Products, Eds Donahaye, E.J., Navarro, S. & Varnava, A. pp. 113-122.
- 367 Lindblad, M., Johnsson, P., Jonsson, N., Lindqvist, R. and Olsen, M. (2004) Predicting
368 noncompliant levels of ochratoxin A in cereal grain from *Penicillium verrucosum*
369 counts. J Appl Microbiol 97: 609-616.

- 370 Magan, N. & Lacey, J. (1984). Effects of gas composition and water activity on growth of
371 field and storage fungi. *Trans Br Mycol Soc* 82: 305-314.
- 372 Magan, N. & Aldred, D. (2007). Post-harvest control strategies: minimizing mycotoxins in
373 the food chain. *Int J Food Microbiol* 119: 131–139.
- 374 Magan, N., Olsen, M. & Aldred, D. (2008). Prevention strategies for trichothecenes and
375 ochratoxin in cereals. Chapter 32, In *Mycotoxins: detection methods, management,*
376 *public health and agricultural trade*, eds. J.Leslie, R.Bandyopadhyay, A.Visconti. pp.
377 369-383. CABI BioSciences, Wallingford, U.K.
- 378 Mokiou, S. & Magan, N. (2008). Physiological manipulation and formulation of the
379 biocontrol yeast *Pichia anomala* for control of *Penicillium verrucosum* and
380 ochratoxin contamination of moist grain. *Biocontrol Sci Technol* 18: 1063-1073.
- 381 Paster, N. (1990). Modified atmospheres for preventing moulds and mycotoxins in stored
382 grain. Chapter 4 in *Food preservation and modified atmospheres*. Eds. Calderon, M.
383 & Barkai-Golan, R. (1990) CRC Press, Boca Ratton, FL, USA. Pp. 39-55.
- 384 Petersson S, Wittrup Hansen M, Hult K & Schnurer J (1998). Ochratoxin A accumulation in
385 cultures of *Penicillium verrucosum* with the antagonistic yeast *Pichia anomala* and
386 *Saccharomyces cerevisiae*. *Mycol Res* 102: 1003-1008.
- 387 Paster, N. & Menashero, N. (1988) Inhibition of T-2 toxin production on high moisture corn
388 kernels by modified atmospheres. *Appl Environ Microbiol* 54: 540-543.
- 389 Paster, N., Menashero, N., Lacey, J. & Fanelli, C. (1992) Synergism between methods for
390 inhibiting the spoilage of damp maize during storage. *Postharvest Biol Technol* 2:
391 163-170.

- 392 Rivas Cosado, M., Parsons, D., Weightman, R., Magan, N., Oraggi, S. (2009a). Geostatistical
393 analysis of the spatial distribution of mycotoxin concentration in bulk cereal. Food
394 Addit Contam 26: 867-873.
- 395 Rivas Casado, M., Parsons, D., Weightman, R.M., Magan, N., Oraggi, S. (2009b). Modelling
396 the spatial distribution of mycotoxin concentration in bulk commodities to design
397 effective and efficient sampling strategies. Food Addit Contam 26: 1298-1305.
- 398 Seitz L.M., Sauer D.B. & Mohr H.E. (1982). Storage of high moisture corn: Fungal growth
399 and dry matter loss. Cereal Chem 59: 100-105.
- 400 Sinha, R.N. (1995). The Stored Grain Ecosystems. In Stored Grain Ecosystems, pp. 1-32, eds.
401 D.S.Jayas, N.D.G.White, W.E. Muir, Marcell Dekker, New York.
- 402 Suanthie, Y., Cousin, M.A & Woloshuk, C.P. (2009). Multiplex real-time PCR for detection
403 and quantification of mycotoxigenic *Aspergillus*, *Penicillium* and *Fusarium*. J Stored
404 Prod Res 45: 139-145.
- 405 Schmidt-Heydt, M., Richterz, W., Michulec, M., Butinger, G. & Geisen, R. (2007).
406 Molecular and chemical monitoring of growth and ochratoxin A biosynthesis of *P.*
407 *verrucosum* in wheat stored at different moisture conditions. Mycotoxin Res 23: 138-
408 146
- 409 Whitaker, T.B. (2004). Sampling for mycotoxins. In Mycotoxins in Food: detection and
410 control, eds. N.Magan, M.Olsen. Woodhead Publishing Ltd. P 69.
- 411 Whitaker, T.B., Hagler Jr, W.M., Giesbrecht, F.G. & Johansson, A. (2000). Sampling,
412 Sample Preparation, and Analytical Variability Associated with Testing Wheat for
413 deoxynivalenol. Journal of AOAC International 83: 1285–1292.
- 414
415
416
417

418

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends

Figure 1. Comparison of dry matter losses caused by *Fusarium graminearum* when inoculated onto wheat grain stored at three different water activities and four different temperatures stored for 4 days. Respiration was measured using GC analyses.

Figure 2. Linear regression of the ratio of O₂ consumption rate/temperature at different water activity levels, corrected for temperature for wheat 7 days storage (adapted from Lacey et. al., 1997).

Figure 3. The growth/no growth and deoxinivalenol/no deoxynivalenol boundaries used to develop the polynomial based model to describe the contour diagrams for *F. graminearum*. Insert: shows the simple front end diagram which could be developed and linked to real time temperature and relative humidity data by movement of the dials to indicate low or high risk of growth and mycotoxin production.

Table 1. Effect of 500 ppm of three essential oils (extracts of bay leaf (Bay), cinnamon (Cin) and clove) and two anti-oxidants (butylhydroxyanisole (BHA), propyl paraben (PP) on (a) deoxynivalenol (DON, ppm) production by *F.graminearum* at 15, 25°C and 0.995 and 0.95 water activity, and (b) T-2 and HT-2 (ppm) by *F.langsethiae* on stored wheat grain at 15, 25°C and 0.95 water activity.

(a) <i>F.graminearum</i>		DON (PPM)			
Temperature (°C)		15		25	
Water activity		0.995	0.95	0.995	0.95
Control		10.11	0.25	3.52	44.07
Bay		0.55	0.30	1.72	2.12
BHA		0.43	0.27	0.91	2.15
Cin		0.57	0.30	1.05	55.11
Clove		0.45	0.32	0.85	1.60
PP		0.75	0.40	1.05	3.10

L.S.D. (P=0.05): Temperature x water activity: 0.25 ppm

(b) <i>F.langsethiae</i>					
Temperature (°C)		15		25	
Mycotoxin		T-2	HT-2	T-2	HT-2
Control		0.53	7.25	0.63	5.07
Bay		0.13	0	0.50	4.71
BHA		0	0	0.79	8.93
Cin		0	0	0	0
Clove		0.13	0.55	4.20	6.52
PP		0.22	0.75	0.42	2.20

L.S.D. (P=0.05): 1.25




Figure 1: Magan et al.




Figure 2: Magan et al.




Figure 3: Magan et al.