Effect of haplotypes on the pharmacokinetics and renin-inhibiting effect of aliskiren

Tuija Tapaninen, Pertti J. Neuvonen, Mikko Niemi

► To cite this version:

Tuija Tapaninen, Pertti J. Neuvonen, Mikko Niemi. Effect of haplotypes on the pharmacokinetics and renin-inhibiting effect of aliskiren. *European Journal of Clinical Pharmacology*, 2010, 66 (9), pp.865-870. 10.1007/s00228-010-0836-z . hal-00594969

HAL Id: hal-00594969

<https://hal.science/hal-00594969>

Submitted on 23 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Effect of ABCB1 haplotypes on the pharmacokinetics and
renin-inhibiting effect of aliskiren**

Journal:	<i>European Journal of Clinical Pharmacology</i>
Manuscript ID:	EJCP-2010-0093.R1
Type of submission:	Original
Date Submitted by the Author:	21-Apr-2010
Complete List of Authors:	Tapaninen, Tuija; University of Helsinki, Department of Clinical Pharmacology Neuvonen, Pertti; University of Helsinki, Department of Clinical Pharmacology Niemi, Mikko; University of Helsinki and Helsinki University Central Hospital, Department of Clinical Pharmacology

**Effect of *ABCB1* haplotypes on the pharmacokinetics and
renin-inhibiting effect of aliskiren**

Tuija Tapaninen, Pertti J. Neuvonen and Mikko Niemi

Department of Clinical Pharmacology, University of Helsinki and Helsinki University
Central Hospital, Helsinki, Finland

Correspondence and requests for reprints:

Mikko Niemi, MD
Department of Clinical Pharmacology,
Helsinki University Central Hospital
PO Box 705, FI-00029 HUS, Finland
Fax: +358-9-471 74039
E-mail: mikko.niemi@helsinki.fi

Word count: 2153 words (excluding the title page, summary, references, tables, and
figures)

A revised manuscript (EJCP 2010-0093) for *European Journal of Clinical
Pharmacology*, including 21 pages, 1 table, and 1 figure (April 2010).

ABSTRACT

Purpose

This study aimed to investigate possible effects of *ABCB1* haplotypes on the pharmacokinetics and renin-inhibiting effect of aliskiren.

Methods

Eleven healthy volunteers homozygous for the *ABCB1* c.1236C-c.2677G-c.3435C (CGC) haplotype and eleven homozygous for the c.1236T-c.2677T-c.3435T (TTT) haplotype ingested a single 150-mg dose of aliskiren. Plasma aliskiren concentrations were measured up to 72 h, its excretion into urine up to 12 h, and plasma renin activity up to 24 h.

Results

The *ABCB1* haplotypes had no significant effect on the pharmacokinetics or renin-inhibiting effect of aliskiren. The geometric mean ratio (95% confidence interval) of aliskiren peak plasma concentration and area under the plasma concentration-time curve from 0 h to infinity in participants homozygous for the TTT haplotype to those in participants homozygous for the CGC haplotype were 1.14 (0.66, 1.96; $P=0.631$) and 1.01 (0.58, 1.76; $P=0.960$), respectively.

Conclusions

These data suggest that the *ABCB1* c.1236C-c.2677G-c.3435C and c.1236T-c.2677T-c.3435T haplotypes have no clinically meaningful effect on the pharmacokinetics of aliskiren.

KEYWORDS

Pharmacokinetics, Pharmacogenetics, *ABCB1* haplotypes, Aliskiren

INTRODUCTION

The direct renin inhibitor aliskiren is poorly absorbed and has a low oral bioavailability of 2-3% [1]. Aliskiren is mainly eliminated by biliary excretion [1]. In addition, about 0.4% of the oral dose is excreted unchanged into urine and about 1.4% undergoes oxidative biotransformation via CYP3A4 [1]. Aliskiren is a substrate of P-glycoprotein (also known as ABCB1) [2], an efflux transporter expressed in tissues with an excretory or barrier function [3, 4]. The P-glycoprotein and CYP3A4 inhibitors, itraconazole and cyclosporine [5-9], have raised the area under the plasma concentration-time curve (AUC) of aliskiren by about 6.5-fold and 5-fold, respectively [10, 11], supporting a major role of P-glycoprotein in limiting the intestinal absorption of aliskiren. Moreover, rifampicin, an inducer of P-glycoprotein and CYP3A4 [12, 13], has reduced the AUC of aliskiren by 56% [14].

A common synonymous single nucleotide polymorphism (SNP) in the *ABCB1* gene encoding P-glycoprotein, c.3435C>T, has been variably associated with reduced expression and function of P-glycoprotein *in vitro* and increased plasma concentrations of P-glycoprotein substrate drugs *in vivo* [15-17]. The mechanism underlying these associations is not completely understood, but it has been suggested that c.3435C>T affects *ABCB1* mRNA stability [18]. Alternatively, the associations may be explained by linkage disequilibrium between c.3435C>T and a functional sequence variation, or that c.3435C>T and sequence variations in linkage disequilibrium with it form a haplotype conferring altered P-glycoprotein activity [16, 19]. The *ABCB1* c.3435C>T

SNP is in strong linkage disequilibrium with the c.1236C>T (synonymous) and c.2677G>T/A (p.Ala893Ser/Thr) SNPs within the same gene [17, 19, 20].

The variant alleles c.1236T, c.2677T, and c.3435T are common in Europeans and European Americans, with reported frequencies of occurrence ranging from 39 to 46%, 39 to 47%, and 54 to 60%, respectively [17, 19-22]. In the Finnish population, the two most common haplotypes based on these SNPs are c.1236T-c.2677T-c.3435T (TTT) and c.1236C-c.2677G-c.3435C (CGC), with frequencies of 42.7% and 34.4%, respectively [17]. The *ABCB1* TTT haplotype has been associated, for example, with increased plasma concentrations of and enhanced lipid-lowering response to simvastatin (acid) and atorvastatin, compared with the *ABCB1* CGC haplotype [17, 23, 24].

Considerable interindividual variability exists in the pharmacokinetics of aliskiren [25], but the background of this variability is unknown. Our hypothesis was that genetic variability in *ABCB1* affects the pharmacokinetics of aliskiren. Therefore, we investigated the effects of the *ABCB1* CGC and TTT haplotypes on the pharmacokinetics and pharmacodynamics of aliskiren.

METHODS

Subjects

Twenty-two healthy White Finnish volunteers participated in this study after giving written informed consent. They were recruited from a group of more than 700 subjects genotyped for the *ABCB1* c.1236C>T (rs1128503), c.2677G>T/A (rs2032582), and c.3435C>T (rs1045642) SNPs [17]. To minimize variability in aliskiren pharmacokinetics due to other genetic variants, all subjects were genotyped for the functionally significant *ABCB1* c.1199G>A (p.Ser400Asn; rs2229109) [17] and *CYP3A5**3 (g.6986A>G; rs776746) [26] alleles and only noncarriers of the *ABCB1* c.1199A and *CYP3A5* g.6986A (*CYP3A5* expressor) alleles were recruited.

Four women and seven men homozygous for the *ABCB1* CGC haplotype, mean $\pm$ SD age 26 ± 4 years (range, 22-37 years), height 178 ± 9 cm (158-189 cm), and weight 74 ± 11 kg (63-95 kg), and five women and six men homozygous for the TTT haplotype, age 24 ± 2 years (21-27 years), height 175 ± 10 cm (158-186 cm), and weight 69 ± 12 kg (51-94 kg), participated in the study. Each participant's health was ascertained by medical history, clinical examination, and laboratory tests. None was on any continuous medication and none was a tobacco smoker.

Study design

The study protocol was approved by the Coordinating Ethics Committee of the Helsinki and Uusimaa Hospital District, and the National Agency for Medicines. Following an overnight fast, the subjects ingested a single 150-mg dose of aliskiren (Rasilez;

Novartis, Horsham, UK) with 150 ml water. A standardized warm meal was served 4 h after aliskiren, and a standardized light meal after 7 h and 10 h. As a safety assessment, systolic and diastolic blood pressures were measured from the forearm with an automatic oscillometric blood pressure monitor (Omron M5-I, Omron Healthcare Europe BV, Hoofddorp, The Netherlands), with the participant in a sitting position, prior to and 2, 4, 7, 9, 12, and 24 h after aliskiren ingestion. The participants were under direct medical supervision for 12 h after the administration of aliskiren. No significant differences in blood pressures were observed between the haplotype groups (data not shown). Use of other drugs was prohibited for one week and use of grapefruit products for three days before aliskiren administration and 72 h thereafter.

Timed blood samples (5 or 10 ml each) for aliskiren concentration measurements were drawn into tubes containing ethylenediaminetetraacetic acid (EDTA) prior to and 0.5, 1, 2, 3, 4, 5, 7, 9, 12, 24, 34, 48, and 72 h after aliskiren administration. Blood samples for the determination of plasma renin activity (5 ml each) were drawn before the administration of aliskiren and 4 h and 24 h thereafter into chilled EDTA tubes, which were placed on ice immediately after sampling. Plasma was separated within 30 min. Urine was collected up to 12 h after aliskiren ingestion. Urine aliquots and plasma were stored at -70°C until analysis.

Determination of aliskiren concentrations and renin activity

Plasma concentrations of aliskiren were quantified by use of solid phase extraction and liquid chromatography-tandem mass spectrometry (LC/MS/MS) [10]. An aliquot of 700 µl of each plasma sample was mixed with 100 µl of 10% phosphoric acid and 50 µl

internal standard solution (1000 ng/ml acebutolol in 1:1 methanol:water). Then the mixture was loaded to the extraction cartridge (Oasis MCX 1cc/30 mg, Waters, Milford, MA, USA), which had been pre-conditioned with 1 ml of methanol and 1 ml of 1% acetic acid. The cartridge was subsequently washed with 1 ml of methanol and 1 ml of 1% acetic acid. The analytes were eluted out with two times 0.6 ml of 2% ammonium hydroxide in methanol. The eluent was dried at 50°C under a nitrogen stream. The residue was dissolved in 100 µl of the mobile phase, and 15 µl of the sample was injected to the LC/MS/MS system. The chromatographic separation was carried out on an AtlantisT3 column (2.1×100 mm, 3 µm) protected by an AtlantisT3 (2.1×10 mm, 3 µm) guard cartridge (Waters). The mobile phase consisted of (A) 2 mmol/l ammonium acetate (pH 4.90) and (B) methanol. The total running time was 15 minutes with the mobile phase gradient of 0.5 minutes at 70% A, 0.5 minutes to 5% A, 3 minutes at 5% A, 0.1 minutes to 70% A, and 10.9 minutes at 70% A. The flow rate of the mobile phase was 250 µl/min. The column temperature was 40°C. Mass spectra were obtained using an Applied Biosystems SCIEX API 2000 Q Trap mass spectrometer operated in positive ion mode (Sciex Division of MDS Inc, Toronto, ON, Canada). The ion transitions were mass-to-charge ratio (m/z) 552.4 to m/z 436.3 for aliskiren and m/z 337.3 to m/z 116.3 for acebutolol.

For the determination of aliskiren concentrations in urine, an aliquot of 300 µl of each urine sample was mixed with 300 µl of fresh frozen plasma, 100 µl of 10% phosphoric acid, and 50 µl internal standard solution (1000 ng/ml acebutolol in 1:1 methanol:water). The extraction, chromatographic separation, and mass spectrometric detection were performed as described for plasma aliskiren. The lower limit of

quantification was 0.25 ng/ml for plasma and 9 ng/ml for urine aliskiren. The calibration curve for plasma aliskiren was linear over the range 0.25-500 ng/ml ($r>0.999$, weighting $1/x$) and for urine aliskiren over the range 9-1800 ng/ml ($r>0.999$, weighting $1/x$). The extraction recoveries were $> 80\%$. The intra-day coefficients of variation were $\leq 1.2\%$ at relevant concentrations in plasma and $\leq 4.8\%$ in urine ($n=6$). Plasma renin activity was determined by radioimmunoassay of generated angiotensin I with a commercially available method (RENCTK; DiaSorin, Saluggia, Italy) at HUSLAB laboratory (Helsinki, Finland).

Pharmacokinetics and pharmacodynamics

Aliskiren pharmacokinetics were characterized by peak plasma concentration ($C_{\max}$), time to $C_{\max}$ ($t_{\max}$), elimination half-life ($t_{1/2}$), AUC_{0-72} , $AUC_{0-\infty}$, the amount of aliskiren excreted into urine from 0 to 12 h (A_e), and the renal clearance (Cl_{renal}).

Pharmacokinetic parameters were calculated with non-compartmental methods using MK-Model, version 5.0 (Biosoft, Cambridge, UK). The terminal log-linear part of each concentration-time curve was identified visually. The elimination rate constant (k_e) was determined by linear regression analysis of the log-linear part of the plasma drug concentration-time curve. The $t_{1/2}$ was calculated by the equation $t_{1/2} = \ln 2 / k_e$. The AUC values were calculated by a combination of the linear and log-linear trapezoidal rules with extrapolation to infinity, when appropriate, by division of the last measured concentration by k_e . The Cl_{renal} of aliskiren was calculated by the equation $Cl_{\text{renal}} = A_e / AUC_{0-12}$. Aliskiren pharmacodynamics were characterized by plasma renin activity at 4 h and 24 h after aliskiren intake.

Statistical analysis

The data were analyzed using SPSS 17.0 (SPSS, Chicago, IL, USA). The C_{max} , AUC, Ae, and Cl_{renal} data were logarithmically transformed before statistical analysis. Statistical comparisons were made between the *ABCB1* haplotypes using analysis of variance (ANOVA). Body weight was set as a covariate for C_{max} , AUC, and Ae [27]. The t_{max} data were compared with the Mann-Whitney test. Differences were considered statistically significant when $P<0.05$. The geometric mean ratio between the TTT and CGC homozygotes with 95% confidence interval (CI) was calculated for the C_{max} , AUC, Ae, and Cl_{renal} values, and the mean difference with 95% CI for the $t_{1/2}$ and renin activity variables. The number of subjects was estimated to be sufficient to detect a 50% difference in the $AUC_{0-\infty}$ of aliskiren between the *ABCB1* CGC and TTT homozygotes, with a power of 80% (α -level 5%).

RESULTS

ABCB1 haplotypes had no significant effect on the pharmacokinetics or renin-inhibiting effect of aliskiren (Table 1, Fig. 1). The geometric mean ratio (95% CI) of aliskiren $C_{\max}$ and $AUC_{0-\infty}$ in participants homozygous for the TTT haplotype to those in participants homozygous for the CGC haplotype were 1.14 (0.66, 1.96; $P=0.631$) and 1.01 (0.58, 1.76; $P=0.960$), respectively. As a *post hoc* analysis, fractional AUC values (AUC_{0-1} , AUC_{0-2} , AUC_{0-3} , AUC_{0-4} , and AUC_{0-5}) were also calculated for aliskiren, but no tendency for a difference between the haplotype groups was observed in these values ($P>0.3$, data not shown). Aliskiren $C_{\max}$, $AUC_{0-\infty}$, A_e , and Cl_{renal} varied 14.1-, 16.2-, 10.4-, and 3.1-fold, respectively, between individual participants.

DISCUSSION

In the present study, the *ABCB1* haplotypes had no significant effect on the pharmacokinetics or renin-inhibiting effect of the P-glycoprotein substrate aliskiren. However, these genetic results do not indicate that the P-glycoprotein transporter could not be otherwise important for the pharmacokinetics of aliskiren. *In vitro*, aliskiren is a high affinity substrate of P-glycoprotein (K_m 2.1 $\mu\text{mol/l}$), as evidenced by studies in P-glycoprotein-expressing Sf9 cell membrane vesicles [2]. *In vivo*, inhibitors and inducers of P-glycoprotein markedly affect aliskiren pharmacokinetics [10, 11, 14].

It is noteworthy that *ABCB1* genotypes have explained only a fraction of the variability in P-glycoprotein expression, and the effects of *ABCB1* genotypes and haplotypes on drug pharmacokinetics have been modest [15, 17, 28, 29]. For example, the AUCs of simvastatin acid and atorvastatin have been 60% and 55% larger, respectively, in subjects homozygous for the *ABCB1* TTT haplotype compared with subjects homozygous for the CGC haplotype [17]. Studies investigating the effects of the *ABCB1* c.3435C>T SNP on the pharmacokinetics of the P-glycoprotein prototype substrate digoxin have revealed conflicting results, with a meta-analysis suggesting that the *ABCB1* c.3435C>T SNP does not affect digoxin pharmacokinetics [12, 15, 28-31]. It has been suggested that the *ABCB1* c.3435C>T SNP, or its combination with the c.1236C>T or c.2677G>T/A SNP, may alter the substrate specificity of P-glycoprotein [16]. Therefore, the effect of the *ABCB1* genotype on the pharmacokinetics of different P-glycoprotein substrates may vary.

The present study had sufficient power to detect a 50% difference in the $AUC_{0-\infty}$ of aliskiren between *ABCB1* haplotypes, and thus a small difference cannot be ruled out. However, it is unlikely that a smaller than 50% difference in aliskiren $AUC_{0-\infty}$ would be of clinical importance. For example, a 76% increase in aliskiren AUC by ketoconazole was not considered clinically relevant [2]. Nevertheless, the mean $AUC_{0-\infty}$ values of aliskiren were almost identical in the two haplotype groups.

Similarly to previous reports [25], a high interindividual variability was observed in the C_{max} and AUC values of aliskiren. The present results suggest that common *ABCB1* haplotypes do not explain this variability. The large differences in aliskiren pharmacokinetics might be explained by variability in gastrointestinal physiology, and other genetic and non-genetic factors affecting the activity of P-glycoprotein or CYP3A, or other transporters or enzymes involved in aliskiren pharmacokinetics [2].

In conclusion, the common *ABCB1* haplotypes c.1236C-c.2677G-c.3435C and c.1236T-c.2677T-c.3435T have no effects on the pharmacokinetics or pharmacodynamics of aliskiren.

ACKNOWLEDGEMENTS

We thank Ms Kaisa Kurkinen, MSc, for aliskiren concentration measurements, and Ms Eija Mäkinen-Pulli, Ms Lisbet Partanen, and Mr Jouko Laitila for skilled assistance. This study was supported by grants from the Sigrid Jusélius Foundation (Helsinki, Finland).

COMPETING INTEREST

None to declare.

For Peer Review

References

- 1 Waldmeier F, Glaenzel U, Wirz B, Oberer L, Schmid D, Seiberling M, Valencia J, Riviere GJ, End P, Vaidyanathan S (2007) Absorption, distribution, metabolism, and elimination of the direct renin inhibitor aliskiren in healthy volunteers. *Drug Metab Dispos* 35 (8): 1418-1428
- 2 Vaidyanathan S, Camenisch G, Schuetz H, Reynolds C, Yeh CM, Bizot MN, Dieterich HA, Howard D, Dole WP (2008) Pharmacokinetics of the oral direct renin inhibitor aliskiren in combination with digoxin, atorvastatin, and ketoconazole in healthy subjects: the role of P-glycoprotein in the disposition of aliskiren. *J Clin Pharmacol* 48 (11): 1323-1338
- 3 Fromm MF (2004) Importance of P-glycoprotein at blood-tissue barriers. *Trends Pharmacol Sci* 25 (8): 423-429
- 4 Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, Dahlin A, Evers R, Fischer V, Hillgren KM, Hoffmaster KA, Ishikawa T, Keppler D, Kim RB, Lee CA, Niemi M, Polli JW, Sugiyama Y, Swaan PW, Ware JA, Wright SH, Wah Yee S, Zamek-Gliszczyński MJ, Zhang L (2010) Membrane transporters in drug development. *Nat Rev Drug Discov* 9 (3): 215-236
- 5 Olkkola KT, Backman JT, Neuvonen PJ (1994) Midazolam should be avoided in patients receiving the systemic antimycotics ketoconazole or itraconazole. *Clin Pharmacol Ther* 55 (5): 481-485

6 Jalava KM, Partanen J, Neuvonen PJ (1997) Itraconazole decreases renal
clearance of digoxin. *Ther Drug Monit* 19 (6): 609-613

7 Wang EJ, Lew K, Casciano CN, Clement RP, Johnson WW (2002)
Interaction of common azole antifungals with P glycoprotein. *Antimicrob Agents
Chemother* 46 (1): 160-165

8 Rao US, Scarborough GA (1994) Direct demonstration of high affinity
interactions of immunosuppressant drugs with the drug binding site of the human P-
glycoprotein. *Mol Pharmacol* 45 (4): 773-776

9 Kajosaari LI, Niemi M, Neuvonen M, Laitila J, Neuvonen PJ, Backman JT
(2005) Cyclosporine markedly raises the plasma concentrations of repaglinide. *Clin
Pharmacol Ther* 78 (4): 388-399

10 Tapaninen T, Backman JT, Kurkinen KJ, Neuvonen PJ, Niemi M (2010)
Itraconazole, a P-glycoprotein and CYP3A4 inhibitor, markedly raises the plasma
concentrations and enhances the renin-inhibiting effect of aliskiren. *J Clin Pharmacol*
Doi:10.1177/0091270010365885

11 European Public Assessment Report for Rasilez. Available from
URL:<http://www.emea.europa.eu/humandocs/Humans/EPAR/rasilez/rasilez.htm>.
Accessed Mar 11, 2010

- 1
2
3
4
5 12 Greiner B, Eichelbaum M, Fritz P, Kreichgauer HP, von Richter O,
6
7 Zundler J, Kroemer HK (1999) The role of intestinal P-glycoprotein in the interaction of
8
9 digoxin and rifampin. *J Clin Invest* 104 (2): 147-153
10
11
12
13
14 13 Niemi M, Backman JT, Fromm MF, Neuvonen PJ, Kivistö KT (2003)
15
16 Pharmacokinetic interactions with rifampicin: clinical relevance. *Clin Pharmacokinet* 42
17
18 (9): 819-850
19
20
21
22
23 14 Tapaninen T, Neuvonen PJ, Niemi M (2010) Rifampicin reduces the
24
25 plasma concentrations and the renin-inhibiting effect of aliskiren. *Eur J Clin Pharmacol*
26
27 66 (5): 497-502
28
29
30
31
32 15 Hoffmeyer S, Burk O, von Richter O, Arnold HP, Brockmöller J, John A,
33
34 Cascorbi I, Gerloff T, Roots I, Eichelbaum M, Brinkmann U (2000) Functional
35
36 polymorphisms of the human multidrug-resistance gene: multiple sequence variations
37
38 and correlation of one allele with P-glycoprotein expression and activity in vivo. *Proc*
39
40 *Natl Acad Sci U S A* 97 (7): 3473-3478
41
42
43
44
45 16 Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar
46
47 SV, Gottesman MM (2007) A "silent" polymorphism in the MDR1 gene changes
48
49 substrate specificity. *Science* 315 (5811): 525-528
50
51
52
53
54 17 Keskitalo JE, Kurkinen KJ, Neuvonen PJ, Niemi M (2008) ABCB1
55
56 haplotypes differentially affect the pharmacokinetics of the acid and lactone forms of
57
58 simvastatin and atorvastatin. *Clin Pharmacol Ther* 84 (4): 457-461
59
60

18 Wang D, Johnson AD, Papp AC, Kroetz DL, Sadée W (2005) Multidrug
resistance polypeptide 1 (MDR1, ABCB1) variant 3435C>T affects mRNA stability.
Pharmacogenet Genomics 15 (10): 693-704

19 Kim RB, Leake BF, Choo EF, Dresser GK, Kubba SV, Schwarz UI,
Taylor A, Xie HG, McKinsey J, Zhou S, Lan LB, Schuetz JD, Schuetz EG, Wilkinson
GR (2001) Identification of functionally variant MDR1 alleles among European
Americans and African Americans. Clin Pharmacol Ther 70 (2): 189-199

20 Kroetz DL, Pauli-Magnus C, Hodges LM, Huang CC, Kawamoto M,
Johns SJ, Stryke D, Ferrin TE, DeYoung J, Taylor T, Carlson EJ, Herskowitz I,
Giacomini KM, Clark AG (2003) Sequence diversity and haplotype structure in the
human ABCB1 (MDR1, multidrug resistance transporter) gene. Pharmacogenetics 13
(8): 481-494

21 Cascorbi I, Gerloff T, Johne A, Meisel C, Hoffmeyer S, Schwab M,
Schaeffeler E, Eichelbaum M, Brinkmann U, Roots I (2001) Frequency of single
nucleotide polymorphisms in the P-glycoprotein drug transporter MDR1 gene in white
subjects. Clin Pharmacol Ther 69 (3): 169-174

22 Leschziner GD, Andrew T, Pirmohamed M, Johnson MR (2007) ABCB1
genotype and PGP expression, function and therapeutic drug response: a critical review
and recommendations for future research. Pharmacogenomics J 7 (3): 154-179

- 23 Kajinami K, Brousseau ME, Ordovas JM, Schaefer EJ (2004)
Polymorphisms in the multidrug resistance-1 (MDR1) gene influence the response to
atorvastatin treatment in a gender-specific manner. *Am J Cardiol* 93 (8): 1046-1050
- 24 Fiegenbaum M, da Silveira FR, Van der Sand CR, Van der Sand LC,
Ferreira ME, Pires RC, Hutz MH (2005) The role of common variants of ABCB1,
CYP3A4, and CYP3A5 genes in lipid-lowering efficacy and safety of simvastatin
treatment. *Clin Pharmacol Ther* 78 (5): 551-558
- 25 Vaidyanathan S, Jarugula V, Dieterich HA, Howard D, Dole WP (2008)
Clinical pharmacokinetics and pharmacodynamics of aliskiren. *Clin Pharmacokinet* 47
(8): 515-531
- 26 Kivistö KT, Niemi M, Schaeffeler E, Pitkälä K, Tilvis R, Fromm MF,
Schwab M, Eichelbaum M, Strandberg T (2004) Lipid-lowering response to statins is
affected by CYP3A5 polymorphism. *Pharmacogenetics* 14 (8): 523-525
- 27 Jarugula V, Yeh CM, Howard D, Bush C, Keefe DL, Dole WP (2010)
Influence of body weight and gender on the pharmacokinetics, pharmacodynamics, and
antihypertensive efficacy of aliskiren. *J Clin Pharmacol*
Doi:10.1177/0091270009359525
- 28 Verstuyft C, Schwab M, Schaeffeler E, Kerb R, Brinkmann U, Jaillon P,
Funck-Brentano C, Becquemont L (2003) Digoxin pharmacokinetics and MDR1
genetic polymorphisms. *Eur J Clin Pharmacol* 58 (12): 809-812

29 John A, Köpke K, Gerloff T, Mai I, Rietbrock S, Meisel C, Hoffmeyer S,
Kerb R, Fromm MF, Brinkmann U, Eichelbaum M, Brockmüller J, Cascorbi I, Roots I
(2002) Modulation of steady-state kinetics of digoxin by haplotypes of the P-
glycoprotein MDR1 gene. *Clin Pharmacol Ther* 72 (5): 584-594

30 Sakaeda T, Nakamura T, Horinouchi M, Kakumoto M, Ohmoto N, Sakai
T, Morita Y, Tamura T, Aoyama N, Hirai M, Kasuga M, Okumura K (2001) MDR1
genotype-related pharmacokinetics of digoxin after single oral administration in healthy
Japanese subjects. *Pharm Res* 18 (10): 1400-1404

31 Chowbay B, Li H, David M, Cheung YB, Lee EJ (2005) Meta-analysis of
the influence of MDR1 C3435T polymorphism on digoxin pharmacokinetics and
MDR1 gene expression. *Br J Clin Pharmacol* 60 (2): 159-171

Table 1 Pharmacokinetic variables of and plasma renin activity response to a single 150-mg oral dose of aliskiren in healthy subjects homozygous for the *ABCB1* c.1236C-c.2677G-c.3435C (CGC/CGC genotype; $n=11$) or c.1236T-c.2677T-c.3435T (TTT/TTT genotype; $n=11$) haplotype

Variable	ABCB1 genotype		Geometric mean ratio (95% CI)	P value
	CGC/CGC	TTT/TTT		
Aliskiren				
C _{max} (ng/ml)	98 (67, 143)	111 (76, 163)	1.14 (0.66, 1.96)	0.631
t _{max} (h)	1 (0.5-5)	0.5 (0.5-5)		0.495
t _{1/2} (h)	23.9 (19.0, 28.8)	22.5 (17.6, 27.4)	-1.4 (-8.3, 5.5)*	0.684
AUC ₀₋₇₂ (ng·h/ml)	419 (289, 608)	429 (296, 622)	1.02 (0.60, 1.74)	0.927
AUC _{0-∞} (ng·h/ml)	443 (301, 652)	449 (306, 661)	1.01 (0.58, 1.76)	0.960
Ae (mg)	0.499 (0.335, 0.742)	0.516 (0.347, 0.767)	1.03 (0.58, 1.83)	0.905
Cl _{renal} (l/h)	1.75 (1.48, 2.06)	1.71 (1.45, 2.01)	0.98 (0.77, 1.23)	0.842
Plasma renin activity (ng/ml/h)				
At baseline	1.73 (0.96, 2.50)	2.38 (1.61, 3.15)	0.66 (-0.43, 1.75)*	0.222
At 4 h	0.01 (-0.09, 0.11)	0.09 (-0.01, 0.19)	0.08 (-0.06, 0.22)*	0.251
At 24 h	1.08 (0.51, 1.65)	1.36 (0.79, 1.92)	0.28 (-0.52, 1.08)*	0.479

$C_{\max}$, AUC, and Ae data are estimated marginal mean (95% CI) adjusting for weight; $t_{1/2}$, Cl_{renal} , and plasma renin activity are estimated marginal mean (95% CI); $t_{\max}$ are median (range). CI, Confidence interval; $C_{\max}$, peak plasma concentration; $t_{\max}$, time to $C_{\max}$; $t_{1/2}$, elimination half-life; AUC_{0-72} , area under the plasma concentration-time curve from 0 to 72 h; $AUC_{0-\infty}$, area under the plasma concentration-time curve from time 0 to infinity; Ae, amount excreted into urine within 12 h; Cl_{renal} renal clearance; Baseline, before the administration of aliskiren.

* Mean difference (95% CI).

FIGURE LEGEND

Fig. 1 Weight-adjusted mean $\pm$ SEM plasma concentrations of aliskiren (A) and mean $\pm$ SEM plasma renin activity (B) after a single 150-mg oral dose of aliskiren in subjects homozygous for the *ABCB1* c.1236C-c.2677G-c.3435C ($n=11$; *open circles*) or *ABCB1* c.1236T-c.2677T-c.3435T ($n=11$; *solid circles*) haplotype. The inset in part A shows aliskiren plasma concentration data on a semi-logarithmic scale

Effect of *ABCB1* haplotypes on the pharmacokinetics and
renin-inhibiting effect of aliskiren

Tuija Tapaninen, Pertti J. Neuvonen and Mikko Niemi

Department of Clinical Pharmacology, University of Helsinki and Helsinki University
Central Hospital, Helsinki, Finland

Correspondence and requests for reprints:

Mikko Niemi, MD
Department of Clinical Pharmacology,
Helsinki University Central Hospital
PO Box 705, FI-00029 HUS, Finland
Fax: +358-9-471 74039
E-mail: mikko.niemi@helsinki.fi

Word count: 2153 words (excluding the title page, summary, references, tables, and
figures)

Deleted: 1540

A revised manuscript ([EJCP 2010-0093](#)) for *European Journal of Clinical
Pharmacology*, including 21 pages, 1 table, and 1 figure ([April 2010](#)).

Deleted: 19

Deleted: March

ABSTRACT

Purpose

This study aimed to investigate possible effects of *ABCB1* haplotypes on the pharmacokinetics and renin-inhibiting effect of aliskiren.

Deleted: genotype

Methods

Eleven healthy volunteers homozygous for the *ABCB1* c.1236C-c.2677G-c.3435C (CGC) haplotype and eleven homozygous for the c.1236T-c.2677T-c.3435T (TTT) haplotype ingested a single 150-mg dose of aliskiren. Plasma aliskiren concentrations were measured up to 72 h, its excretion into urine up to 12 h, and plasma renin activity up to 24 h.

Deleted: with

Deleted: /C

Deleted: /G

Deleted: /C

Deleted: /CGC

Deleted: genotype

Deleted: with

Deleted: /T

Deleted: /T

Deleted: /T

Deleted: /TTT

Deleted: genotype

Deleted: genotype

Results

The *ABCB1* haplotypes had no significant effect on the pharmacokinetics or renin-inhibiting effect of aliskiren. The geometric mean ratio (95% confidence interval) of aliskiren peak plasma concentration and area under the plasma concentration-time curve from 0 h to infinity in participants homozygous for the TTT haplotype to those in participants homozygous for the CGC haplotype were 1.14 (0.66, 1.96; $P=0.631$) and 1.01 (0.58, 1.76; $P=0.960$), respectively.

Deleted: with

Deleted: /TTT

Deleted: genotype

Deleted: with

Deleted: /CGC

Deleted: genotype

Conclusions

These data suggest that the *ABCB1* c.1236C-c.2677G-c.3435C and c.1236T-c.2677T-c.3435T haplotypes have no clinically meaningful effect on the pharmacokinetics of aliskiren.

KEYWORDS

Pharmacokinetics, Pharmacogenetics, *ABCB1* haplotypes, Aliskiren

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

The direct renin inhibitor aliskiren is poorly absorbed and has a low oral bioavailability of 2-3% [1]. Aliskiren is mainly eliminated by biliary excretion [1]. In addition, about 0.4% of the oral dose is excreted unchanged into urine and about 1.4% undergoes oxidative biotransformation via CYP3A4 [1]. Aliskiren is a substrate of P-glycoprotein (also known as ABCB1) [2], an efflux transporter expressed in tissues with an excretory or barrier function [3, 4]. The P-glycoprotein and CYP3A4 inhibitors, itraconazole and cyclosporine [5-9], have raised the area under the plasma concentration-time curve (AUC) of aliskiren by about 6.5-fold and 5-fold, respectively [10, 11], supporting a major role of P-glycoprotein in limiting the intestinal absorption of aliskiren. Moreover, rifampicin, an inducer of P-glycoprotein and CYP3A4 [12, 13], has reduced the AUC of aliskiren by 56% [14].

A common synonymous single nucleotide polymorphism (SNP) in the *ABCB1* gene encoding P-glycoprotein, c.3435C>T, has been variably associated with reduced expression and function of P-glycoprotein *in vitro* and increased plasma concentrations of P-glycoprotein substrate drugs *in vivo* [15-17]. The mechanism underlying these associations is not completely understood, but it has been suggested that c.3435C>T affects *ABCB1* mRNA stability [18]. Alternatively, the associations may be explained by linkage disequilibrium between c.3435C>T and a functional sequence variation, or that c.3435C>T and sequence variations in linkage disequilibrium with it form a haplotype conferring altered P-glycoprotein activity [16, 19]. The *ABCB1* c.3435C>T

Deleted: ABCB1 (also known as P-glycoprotein) is an efflux transporter expressed in tissues with an excretory or barrier function [1, 2].

Deleted: ABCB1

Deleted: ABCB1

Deleted: 3

Deleted: 5

Deleted: 6

Deleted: ABCB1

Deleted: 4

Deleted: 7

SNP is in strong linkage disequilibrium with the c.1236C>T (synonymous) and c.2677G>T/A (p.Ala893Ser/Thr) SNPs within the same gene [17, 19, 20].

Deleted: 5

Deleted: 7

Deleted: 8

The variant alleles c.1236T, c.2677T, and c.3435T are common in Europeans and European Americans, with reported frequencies of occurrence ranging from 39 to 46%, 39 to 47%, and 54 to 60%, respectively [17, 19-22]. In the Finnish population, the two most common haplotypes based on these SNPs are c.1236T-c.2677T-c.3435T (TTT) and c.1236C-c.2677G-c.3435C (CGC), with frequencies of 42.7% and 34.4%, respectively [17]. The *ABCB1* TTT haplotype has been associated, for example, with increased plasma concentrations of and enhanced lipid-lowering response to simvastatin (acid) and atorvastatin, compared with the *ABCB1* CGC haplotype [17, 23, 24].

Deleted: 5

Deleted: 7

Deleted: 10

Deleted: 5

Deleted: 5

Deleted: 11

Deleted: 12

Considerable interindividual variability exists in the pharmacokinetics of aliskiren [25], but the background of this variability is unknown. Our hypothesis was that genetic variability in *ABCB1* affects the pharmacokinetics of aliskiren. Therefore, we investigated the effects of the *ABCB1* CGC and TTT haplotypes on the pharmacokinetics and pharmacodynamics of aliskiren.

Deleted: The antihypertensive drug aliskiren is a substrate of ABCB1 [13]. It is poorly absorbed and has a low oral bioavailability of 2-3% [14]. Aliskiren is mainly eliminated by biliary excretion [14]. In addition, about 0.4% of the oral dose is excreted unchanged into urine and about 1.4% undergoes oxidative biotransformation via CYP3A4 [14]. The ABCB1 and CYP3A4 inhibitors, itraconazole and cyclosporine [15-19], have raised the AUC of aliskiren by about 6.5-fold and 5-fold, respectively [20, 21], supporting a major role of ABCB1 in limiting the intestinal absorption of aliskiren. Moreover, rifampicin, an inducer of ABCB1 and CYP3A4 [22, 23], has reduced the AUC of aliskiren by 56% [24].¶

Deleted: /CGC

Deleted: /TTT

Deleted: genotypes

METHODS

Subjects

Twenty-two healthy White Finnish volunteers participated in this study after giving written informed consent. They were recruited from a group of more than 700 subjects genotyped for the *ABCB1* c.1236C>T (rs1128503), c.2677G>T/A (rs2032582), and c.3435C>T (rs1045642) SNPs [17]. To minimize variability in aliskiren pharmacokinetics due to other genetic variants, all subjects were genotyped for the functionally significant *ABCB1* c.1199G>A (p.Ser400Asn; rs2229109) [17] and *CYP3A5**3 (g.6986A>G; rs776746) [26] alleles and only noncarriers of the *ABCB1* c.1199A and *CYP3A5* g.6986A (*CYP3A5* expressor) alleles were recruited.

Deleted: 5

Deleted: [5]

Deleted: [26]

Four women and seven men homozygous for the *ABCB1* CGC haplotype, mean $\pm$ SD age 26 ± 4 years (range, 22-37 years), height 178 ± 9 cm (158-189 cm), and weight 74 ± 11 kg (63-95 kg), and five women and six men homozygous for the TTT haplotype, age 24 ± 2 years (21-27 years), height 175 ± 10 cm (158-186 cm), and weight 69 ± 12 kg (51-94 kg), participated in the study. Each participant's health was ascertained by medical history, clinical examination, and laboratory tests. None was on any continuous medication and none was a tobacco smoker.

Deleted: with

Deleted: /CGC

Deleted: genotype

Deleted: (

Deleted: with

Deleted: /TTT

Deleted: genotype

Deleted: (

Study design

The study protocol was approved by the Coordinating Ethics Committee of the Helsinki and Uusimaa Hospital District, and the National Agency for Medicines. Following an overnight fast, the subjects ingested a single 150-mg dose of aliskiren (Rasilez;

Novartis, Horsham, UK) with 150 ml water. A standardized warm meal was served 4 h after aliskiren, and a standardized light meal after 7 h and 10 h. As a safety assessment, systolic and diastolic blood pressures were measured from the forearm with an automatic oscillometric blood pressure monitor (Omron M5-I, Omron Healthcare Europe BV, Hoofddorp, The Netherlands), with the participant in a sitting position, prior to and 2, 4, 7, 9, 12, and 24 h after aliskiren ingestion. The participants were under direct medical supervision for 12 h after the administration of aliskiren. No significant differences in blood pressures were observed between the haplotype groups (data not shown). Use of other drugs was prohibited for one week and use of grapefruit products for three days before aliskiren administration and 72 h thereafter.

Timed blood samples (5 or 10 ml each) for aliskiren concentration measurements were drawn into tubes containing ethylenediaminetetraacetic acid (EDTA) prior to and 0.5, 1, 2, 3, 4, 5, 7, 9, 12, 24, 34, 48, and 72 h after aliskiren administration. Blood samples for the determination of plasma renin activity (5 ml each) were drawn before the administration of aliskiren and 4 h and 24 h thereafter into chilled EDTA tubes, which were placed on ice immediately after sampling. Plasma was separated within 30 min. Urine was collected up to 12 h after aliskiren ingestion. Urine aliquots and plasma were stored at -70°C until analysis.

Deleted: Timed ethylenediaminetetraacetic acid blood samples (5-15 ml each) were drawn prior to and up to 72 h after aliskiren ingestion. Plasma was separated within 30 min. Urine was collected up to 12 h after aliskiren ingestion. The samples were stored at -70°C until analysis. ¶

Determination of aliskiren concentrations and renin activity

Plasma concentrations of aliskiren were quantified by use of solid phase extraction and liquid chromatography-tandem mass spectrometry (LC/MS/MS) [10]. An aliquot of 700 µl of each plasma sample was mixed with 100 µl of 10% phosphoric acid and 50 µl

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

internal standard solution (1000 ng/ml acebutolol in 1:1 methanol:water). Then the mixture was loaded to the extraction cartridge (Oasis MCX 1cc/30 mg, Waters, Milford, MA, USA), which had been pre-conditioned with 1 ml of methanol and 1 ml of 1% acetic acid. The cartridge was subsequently washed with 1 ml of methanol and 1 ml of 1% acetic acid. The analytes were eluted out with two times 0.6 ml of 2% ammonium hydroxide in methanol. The eluent was dried at 50°C under a nitrogen stream. The residue was dissolved in 100 µl of the mobile phase, and 15 µl of the sample was injected to the LC/MS/MS system. The chromatographic separation was carried out on an AtlantisT3 column (2.1×100 mm, 3 µm) protected by an AtlantisT3 (2.1×10 mm, 3 µm) guard cartridge (Waters). The mobile phase consisted of (A) 2 mmol/l ammonium acetate (pH 4.90) and (B) methanol. The total running time was 15 minutes with the mobile phase gradient of 0.5 minutes at 70% A, 0.5 minutes to 5% A, 3 minutes at 5% A, 0.1 minutes to 70% A, and 10.9 minutes at 70% A. The flow rate of the mobile phase was 250 µl/min. The column temperature was 40°C. Mass spectra were obtained using an Applied Biosystems SCIEX API 2000 Q Trap mass spectrometer operated in positive ion mode (Sciex Division of MDS Inc, Toronto, ON, Canada). The ion transitions were mass-to-charge ratio (m/z) 552.4 to m/z 436.3 for aliskiren and m/z 337.3 to m/z 116.3 for acebutolol.

For the determination of aliskiren concentrations in urine, an aliquot of 300 µl of each urine sample was mixed with 300 µl of fresh frozen plasma, 100 µl of 10% phosphoric acid, and 50 µl internal standard solution (1000 ng/ml acebutolol in 1:1 methanol:water). The extraction, chromatographic separation, and mass spectrometric detection were performed as described for plasma aliskiren. The lower limit of

Deleted: Aliskiren plasma and urine concentrations were quantified using an Applied Biosystems SCIEX Q Trap LC/MS/MS system (Sciex Division of MDS, Toronto, ON, Canada) [20]. Acebutolol served as an internal standard.

quantification was 0.25 ng/ml for plasma and 9 ng/ml for urine aliskiren. The calibration curve for plasma aliskiren was linear over the range 0.25-500 ng/ml ($r>0.999$, weighting $1/x$) and for urine aliskiren over the range 9-1800 ng/ml ($r>0.999$, weighting $1/x$). The extraction recoveries were $> 80\%$. The intra-day coefficients of variation were $\leq 1.2\%$ at relevant concentrations in plasma and $\leq 4.8\%$ in urine ($n=6$).

Plasma renin activity was determined by radioimmunoassay of generated angiotensin I with a commercially available method (RENCTK; DiaSorin, Saluggia, Italy) at HUSLAB laboratory (Helsinki, Finland).

Deleted: measured

Deleted: using

Deleted: radioimmunoassay

Pharmacokinetics and pharmacodynamics

Aliskiren pharmacokinetics were characterized by peak plasma concentration ($C_{\max}$), time to $C_{\max}$ ($t_{\max}$), elimination half-life ($t_{1/2}$), AUC_{0-72} , $AUC_{0-\infty}$, the amount of aliskiren excreted into urine from 0 to 12 h (A_e), and the renal clearance (Cl_{renal}).

Deleted: areas under the plasma concentration-time curve from 0 to 72 h (

Deleted:) and 0 h to infinity (

Deleted:)

Pharmacokinetic parameters were calculated with non-compartmental methods using MK-Model, version 5.0 (Biosoft, Cambridge, UK). The terminal log-linear part of each concentration-time curve was identified visually. The elimination rate constant (k_e) was determined by linear regression analysis of the log-linear part of the plasma drug concentration-time curve. The $t_{1/2}$ was calculated by the equation $t_{1/2} = \ln 2 / k_e$. The AUC values were calculated by a combination of the linear and log-linear trapezoidal rules with extrapolation to infinity, when appropriate, by division of the last measured concentration by k_e . The Cl_{renal} of aliskiren was calculated by the equation $Cl_{\text{renal}} = A_e / AUC_{0-12}$. Aliskiren pharmacodynamics were characterized by plasma renin activity at 4 h and 24 h after aliskiren intake.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Statistical analysis

The data were analyzed using SPSS 17.0 (SPSS, Chicago, IL, USA). The C_{max}, AUC, Ae, and Cl_{renal} data were logarithmically transformed before statistical analysis.

Deleted: genotypes

Statistical comparisons were made between the *ABCB1* haplotypes using analysis of variance (ANOVA). Body weight was set as a covariate for C_{max}, AUC, and Ae [27].

The t_{max} data were compared with the Mann-Whitney test. Differences were considered

Deleted: /TTT

statistically significant when $P < 0.05$. The geometric mean ratio between the TTT and

Deleted: /CGC

CGC homozygotes with 95% confidence interval (CI) was calculated for the C_{max},

Deleted: groups

AUC, Ae, and Cl_{renal} values, and the mean difference with 95% CI for the t_{1/2} and renin

activity variables. The number of subjects was estimated to be sufficient to detect a 50%

Deleted: /CGC

difference in the AUC_{0-∞} of aliskiren between the *ABCB1* CGC and TTT homozygotes,

Deleted: /TTT

with a power of 80% (α -level 5%).

Deleted: groups

RESULTS

ABCB1 [haplotypes](#) had no significant effect on the pharmacokinetics or renin-inhibiting effect of aliskiren (Table 1, Fig. 1). The geometric mean ratio (95% CI) of aliskiren $C_{\max}$ and $AUC_{0-\infty}$ in participants [homozygous for](#) the TTT [haplotype](#) to those in participants [homozygous for](#) the CGC [haplotype](#) were 1.14 (0.66, 1.96; $P=0.631$) and 1.01 (0.58, 1.76; $P=0.960$), respectively. As a *post hoc* analysis, fractional AUC values (AUC_{0-1} , AUC_{0-2} , AUC_{0-3} , AUC_{0-4} , and AUC_{0-5}) were also calculated for aliskiren, but no tendency for a difference between the [haplotype](#) groups was observed in these values ($P>0.3$, data not shown). Aliskiren $C_{\max}$, $AUC_{0-\infty}$, A_e , and Cl_{renal} varied 14.1-, 16.2-, 10.4-, and 3.1-fold, respectively, between individual participants.

Deleted: genotype

Deleted: with

Deleted: /TTT

Deleted: genotype

Deleted: with

Deleted: /CGC

Deleted: genotype

Deleted: genotype

DISCUSSION

In the present study, the *ABCB1* [haplotypes](#) had no significant effect on the pharmacokinetics or renin-inhibiting effect of the [P-glycoprotein](#) substrate aliskiren. However, these genetic results do not indicate that the [P-glycoprotein](#) transporter could not be otherwise important for the pharmacokinetics of aliskiren. *In vitro*, aliskiren is a high affinity substrate of [P-glycoprotein](#) (K_m 2.1 $\mu\text{mol/l}$), as evidenced by studies in [P-glycoprotein](#)-expressing Sf9 cell membrane vesicles [2]. *In vivo*, inhibitors and inducers of [P-glycoprotein](#) markedly affect aliskiren pharmacokinetics [10, 11, 14].

Deleted: genotype

Deleted: ABCB1

Deleted: ABCB1

Deleted: ABCB1

Deleted: ABCB1

Deleted: 13

Deleted: ABCB1

Deleted: 20

Deleted: 21

Deleted: 24

It is noteworthy that *ABCB1* genotypes have explained only a fraction of the variability in [P-glycoprotein](#) expression, and the effects of *ABCB1* genotypes and haplotypes on drug pharmacokinetics have been modest [15, 17, 28, 29]. For example, the AUCs of simvastatin acid and atorvastatin have been 60% and 55% larger, respectively, in subjects [homozygous for](#) the *ABCB1* TTT [haplotype](#) compared with subjects [homozygous for](#) the CGC [haplotype](#) [17]. Studies investigating the effects of the *ABCB1* c.3435C>T SNP on the pharmacokinetics of the [P-glycoprotein](#) prototype substrate digoxin have revealed conflicting results, with a meta-analysis suggesting that the *ABCB1* c.3435C>T SNP does not affect digoxin pharmacokinetics [12, 15, 28-31].

Deleted: ABCB1

Deleted: 3

Deleted: 5

Deleted: with

Deleted: /TTT

Deleted: genotype

Deleted: with

Deleted: /CGC

Deleted: genotype

Deleted: 5

Deleted: ABCB1

Deleted: 3

Deleted: 22

It has been suggested that the *ABCB1* c.3435C>T SNP, or its combination with the c.1236C>T or c.2677G>T/A SNP, may alter the substrate specificity of [P-glycoprotein](#) [16]. Therefore, the effect of the *ABCB1* genotype on the pharmacokinetics of different [P-glycoprotein](#) substrates may vary.

Deleted: ABCB1

Deleted: 4

Deleted: ABCB1

The present study had sufficient power to detect a 50% difference in the $AUC_{0-\infty}$ of aliskiren between *ABCB1* [haplotypes](#), and thus a small difference cannot be ruled out.

Deleted: genotypes

However, it is unlikely that a smaller than 50% difference in aliskiren $AUC_{0-\infty}$ would be of clinical importance. For example, a 76% increase in aliskiren AUC by ketoconazole was not considered clinically relevant [2]. Nevertheless, the mean $AUC_{0-\infty}$ values of aliskiren were almost identical in the two [haplotype](#) groups.

Deleted: 13

Deleted: genotype

Similarly to previous reports [25], a high interindividual variability was observed in the C_{max} and AUC values of aliskiren. The present results suggest that common *ABCB1* haplotypes do not explain this variability. The large differences in aliskiren pharmacokinetics might be explained by variability in gastrointestinal physiology, and other genetic and non-genetic factors affecting the activity of [P-glycoprotein](#) or CYP3A, or other transporters or enzymes involved in aliskiren pharmacokinetics [2].

Deleted: ABCB1

Deleted: 13

In conclusion, the common *ABCB1* haplotypes c.1236C-c.2677G-c.3435C and c.1236T-c.2677T-c.3435T have no effects on the pharmacokinetics or pharmacodynamics of aliskiren.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGEMENTS

We thank Ms Kaisa Kurkinen, MSc, for aliskiren concentration measurements, and Ms Eija Mäkinen-Pulli, Ms Lisbet Partanen, and Mr Jouko Laitila for skilled assistance. This study was supported by grants from the Sigrid Jusélius Foundation (Helsinki, Finland).

COMPETING INTEREST

None to declare.

For Peer Review

References

- 1 [Waldmeier F, Glaenzel U, Wirz B, Oberer L, Schmid D, Seiberling M, Valencia J, Riviere GJ, End P, Vaidyanathan S \(2007\) Absorption, distribution, metabolism, and elimination of the direct renin inhibitor aliskiren in healthy volunteers. Drug Metab Dispos 35 \(8\): 1418-1428](#)
- 2 [Vaidyanathan S, Camenisch G, Schuetz H, Reynolds C, Yeh CM, Bizot MN, Dieterich HA, Howard D, Dole WP \(2008\) Pharmacokinetics of the oral direct renin inhibitor aliskiren in combination with digoxin, atorvastatin, and ketoconazole in healthy subjects: the role of P-glycoprotein in the disposition of aliskiren. J Clin Pharmacol 48 \(11\): 1323-1338](#)
- 3 [Fromm MF \(2004\) Importance of P-glycoprotein at blood-tissue barriers. Trends Pharmacol Sci 25 \(8\): 423-429](#)
- 4 [Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, Dahlin A, Evers R, Fischer V, Hillgren KM, Hoffmaster KA, Ishikawa T, Keppler D, Kim RB, Lee CA, Niemi M, Polli JW, Sugiyama Y, Swaan PW, Ware JA, Wright SH, Wah Yee S, Zamek-Gliszczynski MJ, Zhang L \(2010\) Membrane transporters in drug development. Nat Rev Drug Discov 9 \(3\): 215-236](#)
- 5 [Olkola KT, Backman JT, Neuvonen PJ \(1994\) Midazolam should be avoided in patients receiving the systemic antimycotics ketoconazole or itraconazole. Clin Pharmacol Ther 55 \(5\): 481-485](#)

Deleted: 1

Deleted: 2

6 [Jalava KM, Partanen J, Neuvonen PJ \(1997\) Itraconazole decreases renal
clearance of digoxin. *Ther Drug Monit* 19 \(6\): 609-613](#)

7 [Wang EJ, Lew K, Casciano CN, Clement RP, Johnson WW \(2002\)
Interaction of common azole antifungals with P glycoprotein. *Antimicrob Agents
Chemother* 46 \(1\): 160-165](#)

8 [Rao US, Scarborough GA \(1994\) Direct demonstration of high affinity
interactions of immunosuppressant drugs with the drug binding site of the human P-
glycoprotein. *Mol Pharmacol* 45 \(4\): 773-776](#)

9 [Kajosaari LI, Niemi M, Neuvonen M, Laitila J, Neuvonen PJ, Backman JT
\(2005\) Cyclosporine markedly raises the plasma concentrations of repaglinide. *Clin
Pharmacol Ther* 78 \(4\): 388-399](#)

10 [Tapaninen T, Backman JT, Kurkinen KJ, Neuvonen PJ, Niemi M \(2010\)
Itraconazole, a P-glycoprotein and CYP3A4 inhibitor, markedly raises the plasma
concentrations and enhances the renin-inhibiting effect of aliskiren. *J Clin Pharmacol*
Doi:10.1177/0091270010365885](#)

11 [European Public Assessment Report for Rasilez. Available from
URL:<http://www.emea.europa.eu/humandocs/Humans/EPAR/rasilez/rasilez.htm>.
Accessed Mar 11, 2010](#)

12 [Greiner B, Eichelbaum M, Fritz P, Kreichgauer HP, von Richter O, Zundler J, Kroemer HK \(1999\) The role of intestinal P-glycoprotein in the interaction of digoxin and rifampin. J Clin Invest 104 \(2\): 147-153](#)

13 [Niemi M, Backman JT, Fromm MF, Neuvonen PJ, Kivistö KT \(2003\) Pharmacokinetic interactions with rifampicin: clinical relevance. Clin Pharmacokinet 42 \(9\): 819-850](#)

14 [Tapaninen T, Neuvonen PJ, Niemi M \(2010\) Rifampicin reduces the plasma concentrations and the renin-inhibiting effect of aliskiren. Eur J Clin Pharmacol 66 \(5\): 497-502](#)

15 [Hoffmeyer S, Burk O, von Richter O, Arnold HP, Brockmöller J, John A, Cascorbi I, Gerloff T, Roots I, Eichelbaum M, Brinkmann U \(2000\) Functional polymorphisms of the human multidrug-resistance gene: multiple sequence variations and correlation of one allele with P-glycoprotein expression and activity in vivo. Proc Natl Acad Sci U S A 97 \(7\): 3473-3478](#)

Deleted: 3

16 [Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar SV, Gottesman MM \(2007\) A "silent" polymorphism in the MDR1 gene changes substrate specificity. Science 315 \(5811\): 525-528](#)

Deleted: 4

17 [Keskitalo JE, Kurkinen KJ, Neuvonen PJ, Niemi M \(2008\) ABCB1 haplotypes differentially affect the pharmacokinetics of the acid and lactone forms of simvastatin and atorvastatin. Clin Pharmacol Ther 84 \(4\): 457-461](#)

Deleted: 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[18](#) Wang D, Johnson AD, Papp AC, Kroetz DL, Sadée W (2005) Multidrug resistance polypeptide 1 (MDR1, ABCB1) variant 3435C>T affects mRNA stability. *Pharmacogenet Genomics* 15 (10): 693-704

Deleted: 6

[19](#) Kim RB, Leake BF, Choo EF, Dresser GK, Kubba SV, Schwarz UI, Taylor A, Xie HG, McKinsey J, Zhou S, Lan LB, Schuetz JD, Schuetz EG, Wilkinson GR (2001) Identification of functionally variant MDR1 alleles among European Americans and African Americans. *Clin Pharmacol Ther* 70 (2): 189-199

Deleted: 7

[20](#) Kroetz DL, Pauli-Magnus C, Hodges LM, Huang CC, Kawamoto M, Johns SJ, Stryke D, Ferrin TE, DeYoung J, Taylor T, Carlson EJ, Herskowitz I, Giacomini KM, Clark AG (2003) Sequence diversity and haplotype structure in the human ABCB1 (MDR1, multidrug resistance transporter) gene. *Pharmacogenetics* 13 (8): 481-494

Deleted: 8

[21](#) Cascorbi I, Gerloff T, Johne A, Meisel C, Hoffmeyer S, Schwab M, Schaeffeler E, Eichelbaum M, Brinkmann U, Roots I (2001) Frequency of single nucleotide polymorphisms in the P-glycoprotein drug transporter MDR1 gene in white subjects. *Clin Pharmacol Ther* 69 (3): 169-174

Deleted: 9

[22](#) Leschziner GD, Andrew T, Pirmohamed M, Johnson MR (2007) ABCB1 genotype and PGP expression, function and therapeutic drug response: a critical review and recommendations for future research. *Pharmacogenomics J* 7 (3): 154-179

Deleted: 10

Deleted: 11

[23](#) Kajinami K, Brousseau ME, Ordovas JM, Schaefer EJ (2004)

Polymorphisms in the multidrug resistance-1 (MDR1) gene influence the response to atorvastatin treatment in a gender-specific manner. *Am J Cardiol* 93 (8): 1046-1050

Deleted: 12

[24](#) Fiegenbaum M, da Silveira FR, Van der Sand CR, Van der Sand LC,

Ferreira ME, Pires RC, Hutz MH (2005) The role of common variants of ABCB1, CYP3A4, and CYP3A5 genes in lipid-lowering efficacy and safety of simvastatin treatment. *Clin Pharmacol Ther* 78 (5): 551-558

[25](#) Vaidyanathan S, Jarugula V, Dieterich HA, Howard D, Dole WP (2008)

Clinical pharmacokinetics and pharmacodynamics of aliskiren. *Clin Pharmacokinet* 47 (8): 515-531

26 Kivistö KT, Niemi M, Schaeffeler E, Pitkälä K, Tilvis R, Fromm MF,

Schwab M, Eichelbaum M, Strandberg T (2004) Lipid-lowering response to statins is affected by CYP3A5 polymorphism. *Pharmacogenetics* 14 (8): 523-525

27 Jarugula V, Yeh CM, Howard D, Bush C, Keefe DL, Dole WP (2010)

Influence of body weight and gender on the pharmacokinetics, pharmacodynamics, and antihypertensive efficacy of aliskiren. *J Clin Pharmacol*

Doi:10.1177/0091270009359525

28 Verstuyft C, Schwab M, Schaeffeler E, Kerb R, Brinkmann U, Jaillon P,

Funck-Brentano C, Becquemont L (2003) Digoxin pharmacokinetics and MDR1 genetic polymorphisms. *Eur J Clin Pharmacol* 58 (12): 809-812

Deleted: 13 . Vaidyanathan S, Camenisch G, Schuetz H, Reynolds C, Yeh CM, Bizot MN, Dieterich HA, Howard D, Dole WP (2008) Pharmacokinetics of the oral direct renin inhibitor aliskiren in combination with digoxin, atorvastatin, and ketoconazole in healthy subjects: the role of P-glycoprotein in the disposition of aliskiren. *J Clin Pharmacol* 48 (11): 1323-1338 ¶
 14 . Waldmeier F, Glaenzel U, Wirz B, Oberer L, Schmid D, Seiberling M, Valencia J, Riviere GJ, End P, Vaidyanathan S (2007) Absorption, distribution, metabolism, and elimination of the direct renin inhibitor aliskiren in healthy volunteers. *Drug Metab Dispos* 35 (8): 1418-1428 ¶
 15 . Olkkola KT, Backman JT, Neuvonen PJ (1994) Midazolam should be avoided in patients receiving the systemic antimycotics ketoconazole or itraconazole. *Clin Pharmacol Ther* 55 (5): 481-485 ¶
 16 . Jalava KM, Partanen J, Neuvonen PJ (1997) Itraconazole decreases renal clearance of digoxin. *Ther Drug Monit* 19 (6): 609-613 ¶
 17 . Wang EJ, Lew K, Casciano CN, Clement RP, Johnson WW (2002) Interaction of common azole antifungals with P glycoprotein. *Antimicrob Agents Chemother* 46 (1): 160-165 ¶
 18 . Rao US, Scarborough GA (1994) Direct demonstration of high affinity interactions of immunosuppressant drugs with the drug binding site of the human P-glycoprotein. *Mol Pharmacol* 45 (4): 773-776 ¶
 19 . Kajosaari LI, Niemi M, Neuvonen M, Laitila J, Neuvonen PJ, Backman JT (2005) Cyclosporine markedly raises the plasma concentrations of repaglinide. *Clin Pharmacol Ther* 78 (4): 388-399 ¶
 20 . Tapaninen T, Backman JT, Kurkinen KJ, Neuvonen PJ, Niemi M (2010) Itraconazole, a P-glycoprotein and CYP3A4 inhibitor, markedly raises the plasma concentrations and enhances the renin-inhibiting effect of aliskiren. *J Clin Pharmacol* in press ¶
 21 . European Public Assessment Report for Rasilez. Available from URL: <http://www.emea.europa.eu/humandocs/Humans/EPAR/rasilez/rasilez.htm>. Accessed Mar 11, 2010 ¶
 22 . Greiner B, Eichelbaum M, Fritz P, Kreichgauer HP, von Richter O, Zundler J, Kroemer HK (1999) The role of intestinal P-glycoprotein in the interaction of digoxin and rifampin. *J Clin Invest* 104 (2): 147-153 ¶
 23 . Niemi M, Backman JT, Fromm MF, Neuvonen PJ, Kivistö KT (2003) Pharmacokinetic interactions with rifampicin : clinical relevance. *Clin Pharmacokinet* 42 (9): 819-850 ¶ ... [1]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

29 John A, Köpke K, Gerloff T, Mai I, Rietbrock S, Meisel C, Hoffmeyer S,
Kerb R, Fromm MF, Brinkmann U, Eichelbaum M, Brockmöller J, Cascorbi I, Roots I
(2002) Modulation of steady-state kinetics of digoxin by haplotypes of the P-
glycoprotein MDR1 gene. *Clin Pharmacol Ther* 72 (5): 584-594

30 Sakaeda T, Nakamura T, Horinouchi M, Kakumoto M, Ohmoto N, Sakai
T, Morita Y, Tamura T, Aoyama N, Hirai M, Kasuga M, Okumura K (2001) MDR1
genotype-related pharmacokinetics of digoxin after single oral administration in healthy
Japanese subjects. *Pharm Res* 18 (10): 1400-1404

31 Chowbay B, Li H, David M, Cheung YB, Lee EJ (2005) Meta-analysis of
the influence of MDR1 C3435T polymorphism on digoxin pharmacokinetics and
MDR1 gene expression. *Br J Clin Pharmacol* 60 (2): 159-171

Table 1 Pharmacokinetic variables of and plasma renin activity response to a single 150-mg oral dose of aliskiren in healthy subjects homozygous for the *ABCB1* c.1236C-c.2677G-c.3435C (CGC/CGC genotype; $n=11$) or c.1236T-c.2677T-c.3435T (TTT/TTT genotype; $n=11$) haplotype

Variable	ABCB1 genotype		Geometric mean ratio (95% CI)	P value
	CGC/CGC	TTT/TTT		
Aliskiren				
C _{max} (ng/ml)	98 (67, 143)	111 (76, 163)	1.14 (0.66, 1.96)	0.631
t _{max} (h)	1 (0.5-5)	0.5 (0.5-5)		0.495
t _{1/2} (h)	23.9 (19.0, 28.8)	22.5 (17.6, 27.4)	-1.4 (-8.3, 5.5)*	0.684
AUC ₀₋₇₂ (ng·h/ml)	419 (289, 608)	429 (296, 622)	1.02 (0.60, 1.74)	0.927
AUC _{0-∞} (ng·h/ml)	443 (301, 652)	449 (306, 661)	1.01 (0.58, 1.76)	0.960
Ae (mg)	0.499 (0.335, 0.742)	0.516 (0.347, 0.767)	1.03 (0.58, 1.83)	0.905
Cl _{renal} (l/h)	1.75 (1.48, 2.06)	1.71 (1.45, 2.01)	0.98 (0.77, 1.23)	0.842
Plasma renin activity (ng/ml/h)				
At baseline	1.73 (0.96, 2.50)	2.38 (1.61, 3.15)	0.66 (-0.43, 1.75)*	0.222
At 4 h	0.01 (-0.09, 0.11)	0.09 (-0.01, 0.19)	0.08 (-0.06, 0.22)*	0.251
At 24 h	1.08 (0.51, 1.65)	1.36 (0.79, 1.92)	0.28 (-0.52, 1.08)*	0.479

$C_{\max}$, AUC , and A_e data are estimated marginal mean (95% CI) adjusting for weight; $t_{1/2}$, Cl_{renal} , and plasma renin activity are estimated marginal mean (95% CI); $t_{\max}$ are median (range). CI, Confidence interval; $C_{\max}$, peak plasma concentration; $t_{\max}$, time to $C_{\max}$; $t_{1/2}$, elimination half-life; AUC_{0-72} , area under the plasma concentration-time curve from 0 to 72 h; $AUC_{0-\infty}$, area under the plasma concentration-time curve from time 0 to infinity; A_e , amount excreted into urine within 12 h; Cl_{renal} renal clearance; Baseline, before the administration of aliskiren.

* Mean difference (95% CI).

Deleted: with

Deleted: /C

Deleted: /G

Deleted: /C

Deleted: /T

Deleted: /T

Deleted: /T

Deleted: genotype

Deleted: //
mean difference

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGEND

Fig. 1 Weight-adjusted mean $\pm$ SEM plasma concentrations of aliskiren (A) and mean $\pm$ SEM plasma renin activity (B) after a single 150-mg oral dose of aliskiren in subjects homozygous for the *ABCB1* c.1236C-c.2677G-c.3435C (*n*=11; open circles) or *ABCB1* c.1236T-c.2677T-c.3435T (*n*=11; solid circles) haplotype. The inset in part A shows aliskiren plasma concentration data on a semi-logarithmic scale

- Deleted: with
- Deleted: /C
- Deleted: /G
- Deleted: /C
- Deleted: /T
- Deleted: /T
- Deleted: /T
- Deleted: genotype

For Peer Review

Page 18: [1] Deleted

tapatui20

4/20/2010 3:35:00 PM

- 13 Vaidyanathan S, Camenisch G, Schuetz H, Reynolds C, Yeh CM, Bizot MN, Dieterich HA, Howard D, Dole WP (2008) Pharmacokinetics of the oral direct renin inhibitor aliskiren in combination with digoxin, atorvastatin, and ketoconazole in healthy subjects: the role of P-glycoprotein in the disposition of aliskiren. *J Clin Pharmacol* 48 (11): 1323-1338
- 14 Waldmeier F, Glaenzel U, Wirz B, Oberer L, Schmid D, Seiberling M, Valencia J, Riviere GJ, End P, Vaidyanathan S (2007) Absorption, distribution, metabolism, and elimination of the direct renin inhibitor aliskiren in healthy volunteers. *Drug Metab Dispos* 35 (8): 1418-1428
- 15 Olkkola KT, Backman JT, Neuvonen PJ (1994) Midazolam should be avoided in patients receiving the systemic antimycotics ketoconazole or itraconazole. *Clin Pharmacol Ther* 55 (5): 481-485
- 16 Jalava KM, Partanen J, Neuvonen PJ (1997) Itraconazole decreases renal clearance of digoxin. *Ther Drug Monit* 19 (6): 609-613
- 17 Wang EJ, Lew K, Casciano CN, Clement RP, Johnson WW (2002) Interaction of common azole antifungals with P glycoprotein. *Antimicrob Agents Chemother* 46 (1): 160-165

- 1
2
3 18 Rao US, Scarborough GA (1994) Direct demonstration of high affinity
4 interactions of immunosuppressant drugs with the drug binding site of the human P-
5 glycoprotein. Mol Pharmacol 45 (4): 773-776
6
7
8
9
10
11
12 19 Kajosaari LI, Niemi M, Neuvonen M, Laitila J, Neuvonen PJ, Backman JT
13 (2005) Cyclosporine markedly raises the plasma concentrations of repaglinide. Clin
14 Pharmacol Ther 78 (4): 388-399
15
16
17
18
19
20
21 20 Tapaninen T, Backman JT, Kurkinen KJ, Neuvonen PJ, Niemi M (2010)
22 Itraconazole, a P-glycoprotein and CYP3A4 inhibitor, markedly raises the plasma
23 concentrations and enhances the renin-inhibiting effect of aliskiren. J Clin Pharmacol in
24 press
25
26
27
28
29
30
31
32
33 21 European Public Assessment Report for Rasilez. Available from
34 URL:<http://www.emea.europa.eu/humandocs/Humans/EPAR/rasilez/rasilez.htm>.
35
36 Accessed Mar 11, 2010
37
38
39
40
41
42 22 Greiner B, Eichelbaum M, Fritz P, Kreichgauer HP, von Richter O, Zundler
43 J, Kroemer HK (1999) The role of intestinal P-glycoprotein in the interaction of digoxin
44 and rifampin. J Clin Invest 104 (2): 147-153
45
46
47
48
49
50
51 23 Niemi M, Backman JT, Fromm MF, Neuvonen PJ, Kivistö KT (2003)
52 Pharmacokinetic interactions with rifampicin : clinical relevance. Clin Pharmacokinet 42
53 (9): 819-850
54
55
56
57
58
59
60

24 Tapaninen T, Neuvonen PJ, Niemi M (2010) Rifampicin reduces the plasma
concentrations and the renin-inhibiting effect of aliskiren. Eur J Clin Pharmacol

Doi:10.1007/s00228-010-0796-3

For Peer Review