Predictions of Taylor's power law, density dependence and pink noise from a neutrally modelled time series

Petr Keil, T. Tomáš Herben, James Rosindell, David Storch

► To cite this version:

Petr Keil, T. Tomáš Herben, James Rosindell, David Storch. Predictions of Taylor's power law, density dependence and pink noise from a neutrally modelled time series. *Journal of Theoretical Biology*, 2010, 265 (1), pp.78. 10.1016/j.jtbi.2010.04.014 . hal-00598093

HAL Id: hal-00598093

<https://hal.science/hal-00598093>

Submitted on 4 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript

Predictions of Taylor's power law, density dependence and pink noise from a neutrally modelled time series

Petr Keil, Tomáš Herben, James Rosindell, David Storch

PII: S0022-5193(10)00197-9
DOI: doi:10.1016/j.jtbi.2010.04.014
Reference: YJTBI5963

www.elsevier.com/locate/jtbi

To appear in: *Journal of Theoretical Biology*

Received date: 16 December 2009
Revised date: 12 April 2010
Accepted date: 12 April 2010

Cite this article as: Petr Keil, Tomáš Herben, James Rosindell and David Storch, Predictions of Taylor's power law, density dependence and pink noise from a neutrally modelled time series, *Journal of Theoretical Biology*, doi:[10.1016/j.jtbi.2010.04.014](https://doi.org/10.1016/j.jtbi.2010.04.014)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Predictions of Taylor's power law, density dependence and pink
noise from a neutrally modelled time series**

Authors: Petr Keil^{1,*}, Tomáš Herben^{2,3}, James Rosindell⁴ & David Storch^{1,5}

¹*Department of Ecology, Faculty of Science, Charles University, Vinicna 7, 12800*

Praha 2, Czech Republic, e-mail: pkeil@seznam.cz

tel: 00420 – 776 371 840

²*Institute of Botany, Academy of Sciences of the Czech Republic, Průhonice, 252 43,*

Czech Republic, e-mail: herben@site.cas.cz

³*Department of Botany, Faculty of Science, Charles University, Benátská 2, Praha 2,*

128 01, Czech Republic

⁴*Institute of Integrative and Comparative Biology, Faculty of Biological Sciences,*

University of Leeds, Leeds LS2 9JT, United Kingdom, e-mail:

j.l.rosindell@leeds.ac.uk

⁵*Center for Theoretical Study, Charles University in Prague and the Academy of*

Sciences of the Czech Republic, Jilská 1, 110 00 Praha 1, Czech Republic, e-mail:

storch@cts.cuni.cz

** Corresponding author*

21 **Abstract**

22

23 There has recently been increasing interest in neutral models of biodiversity and their
24 ability to reproduce the patterns observed in nature, such as species abundance
25 distributions. Here we investigate the ability of a neutral model to predict phenomena
26 observed in single-population time series, a study complementary to most existing
27 work that concentrates on snapshots in time of the whole community. We consider
28 tests for density dependence, the dominant frequencies of population fluctuation
29 (spectral density) and a relationship between the mean and variance of a fluctuating
30 population (Taylor's power law). We simulated an archipelago model of a set of
31 interconnected local communities with variable mortality rate, migration rate,
32 speciation rate, size of local community and number of local communities. Our
33 spectral analysis showed 'pink noise': a departure from a standard random walk
34 dynamics in favor of the higher frequency fluctuations which is partly consistent with
35 empirical data. We detected density dependence in local community time series but
36 not in metacommunity time series. The slope of the Taylor's power law in the model
37 was similar to the slopes observed in natural populations, but the fit to the power law
38 was worse. Our observations of pink noise and density dependence can be attributed
39 to the presence of an upper limit to community sizes and to the effect of migration
40 which distorts temporal autocorrelation in local time series. We conclude that some of
41 the phenomena observed in natural time series can emerge from neutral processes, as
42 a result of random zero-sum birth, death and migration. This suggests the neutral
43 model would be a parsimonious null model for future studies of time series data.

44

45 **Key words:** random walk, spectral analysis, mean-variance scaling, stochasticity,
 46 neutral theory of biodiversity, autocorrelation, 1/f noise

47

48

49 **Introduction**

50

51 The neutral theory of biodiversity (NTB; Hubbell, 2001; Chave, 2004; Alonso et al.,
 52 2006; Leigh, 2007; Rosindell et al., 2010) has revolutionized the way we think about
 53 species coexistence over the last decade. The assumptions of the theory are
 54 controversial (Tilman, 2004; Leibold and McPeck, 2006) and some argue that there is
 55 limited empirical support (McGill et al., 2006), but the theory still has huge potential
 56 as a null model (Gotelli and McGill, 2006), serving as an essential first approximation
 57 to reality and a base for more complex theories. Even with its apparently
 58 oversimplified set of assumptions, the NTB is capable of predicting realistic species-
 59 abundance distributions (Hubbell, 2001; Etienne, 2005) and species-area curves
 60 (Rosindell and Cornell, 2007, 2009), as well as many other ecological patterns
 61 (Herben, 2004, 2009). Most of the current research is focused on the community-level
 62 predictions of a single snapshot in time at equilibrium (Chave et al., 2002; McGill,
 63 2003; Turnbull et al., 2005; Volkov et al., 2005; Dornelas, et al. 2006). Dynamical
 64 behaviour of populations in NTB has rarely been studied. Azale et al. (2006) used
 65 ratios of population sizes between two snapshots of time to estimate species turnover
 66 rates and extinction times of trees in Barro Colorado Island, Panama. Mutshinda et al.
 67 (2008) produced the only study (as far as we are aware) that examines sufficiently
 68 long real-world time series from the perspective of NTB. They used the Bayesian
 69 statistical framework and a neutral model of community dynamics (zero-sum

assumption relaxed) to show that sizes of natural communities fluctuate more than expected under neutrality. However, Mutshinda et al. (2008) focused only on the dynamics of local community size and they did not examine time series of individual species.

There are some characteristics of natural time-series that are often used as summary statistics and can be employed here in making comparisons with the time series produced by a neutral model. For example, natural time series have been shown to obey some degree of stabilizing density dependence (Woiwod and Hanski, 1992; Brook and Bradshaw, 2006; Sibly et al., 2007). Natural populations also follow a specific relationship between mean population size (m) and variance of population size (s^2) called Taylor's Power Law (TPL) (Taylor et al., 1983; Tokeshi, 1995; Ballantyne and Kerkhoff, 2007) which has the form $s^2 \propto m^b$. In natural populations, the exponent b of TPL is usually observed to satisfy $1 < b < 2$ (Kendal, 2004). Natural time series can be considered as being the net result of a number of independent fluctuations each with a distinct frequency and amplitude. The amplitude (or 'spectral density') $S(f)$ is a function of the frequency f and often satisfies $S(f) \propto 1/f^\gamma$ (Halley, 1996). The exponent γ usually falls between 0 and 2 for natural time series (Arino and Pimm, 1995; Halley, 1996; Inchausti and Halley, 2001, 2002), a property known as 'pink noise'.

The null model for discrete population dynamics is an unbounded *random walk* of population size (N) and is described as

$$\ln N_{t+1} = \ln N_t + e_t \quad (1)$$

where e_t varies at random according to a given distribution with zero mean and is independent on population size (Murdoch, 1994; Hanski et al., 1996). The model is a discrete first-order Markov chain in which all the information that can influence the

95 future state ($t+1$) is fully captured in the present state (t). Random walk time series are
 96 often termed “brownian” or autocorrelated noise. Random walk is synonymous for a
 97 completely density-independent process in ecology, it produces a slope of $b \approx 2$ in
 98 Taylor’s power law and spectral scaling exponent of $\gamma \approx 2$.

99 Although the term random walk is sometimes used to describe the population
 100 time series in neutral models (Hubbell, 2001; Alonso et al., 2006), there are reasons
 101 why neutral models can behave differently from the random walk as defined above.
 102 Firstly, there is usually a constraint on the total number of individuals (the ‘zero-sum’
 103 rule) which imposes a clear upper limit to any population fluctuations (Hubbell, 2001;
 104 Hubbell and Lake, 2003). Secondly, immigration of individuals into the local
 105 community can cause a locally extinct species to reappear (zero is not necessarily
 106 absorbing; Hubbell, 2001) and finally immigration can distort the temporal
 107 autocorrelation of N_t (the number of individuals belonging to the species of interest in
 108 a local community). In the presence of immigration, N_{t+1} for a given local community
 109 does not depend solely on N_t for that community (as in the autocorrelated random
 110 walk), it also depends on the N_t for all the other local communities (the
 111 metacommunity).

112 In this paper we ask if a simple neutral model can produce time series that are
 113 similar to natural time series. In particular, we show that populations in the neutral
 114 model can produce realistic slopes of the Taylor’s power law, have spectral properties
 115 similar to natural populations and seem to reveal density-dependency more frequently
 116 than expected in a random walk. Additionally, we explore the influence of the NTB
 117 parameters (size of local communities, number of local communities, mortality rate,
 118 migration and mutation rate) on these properties of population time series.

119

Methods

The simulation model

Instead of the classic neutral model used by Hubbell (2001) and Volkov et al. (2003), we simulate a *metacommunity* consisting of an archipelago of L isolated *local communities* interconnected by migration at rate m (see Fig. 1 below and Volkov et al., 2007). Each local community contains J individuals all of which behave in an identical manner irrespective of the species to which they belong. The archipelago model enabled us to obtain data on abundances and their variances for one species at multiple local communities, which is necessary for the analysis of Taylor's Power Law. We used the simulation approach because it enabled us to record the species identity of every individual at each time step and across all local communities. Furthermore, we wished to collect time series of limited lengths that are comparable to those of empirically collected time series in order to accurately mimic any sampling effects present in empirical data. This would be particularly difficult to approach analytically. Our simulations were conducted using a 'forwards in time' approach. A 'backwards in time' or coalescence approach can be much faster (Rosindell et al., 2008), but the main strengths of this approach are only present when collecting a snapshot in time of a sample from a very large or infinite metacommunity. We are instead collecting single species time series data and sampling all individuals from a relatively small metacommunity: a scenario where coalescence is far less beneficial.

Four processes operate in the local community at each step: mortality (d ; $0 < d < 1$), local replacement (l ; $0 < l < 1$), immigration from outside of the local community (m ; $m = 1 - l$) and speciation (s ; $0 < s < 1$) in that order (Fig. 1, Table 1). To model mortality, $J \times d$ individuals were randomly removed from each local

community in every time step with all individuals having an equal probability of removal. These removed individuals were then replaced by offspring from others in the local community and other local communities (immigration). Local replacement was modeled by drawing individuals to reproduce at random from the $J(1-d)$ individuals that survived mortality until $J \times d \times l$ offspring have been created. The remaining $J \times d \times m$ dead individuals were replaced by immigration from all of the other local communities. The probability of immigration of a species is given by its relative abundance in the whole metacommunity (excluding the local community in focus).

A local population can occasionally give rise to a new species (a speciation event), this happens with a fixed probability at each time step for each local population of a species. When speciation occurs, individuals of the given species in the local community were randomly split into two parts, one of which forms the initial abundance of a new species (a version of *random-fission* model; Hubbell, 2001; Etienne and Haegeman, 2010; Haegeman and Etienne, 2010). Pilot testing of a *point mutation* model (Hubbell, 2001) showed that the choice of a speciation model does not affect the outcome of the simulations, this is not entirely surprising because for a single species population time series, speciation can be regarded as another form of mortality: individuals belonging to our focal species that switch to a new species reduce the population of our focal species in the same way as dead individuals. Each simulation was launched with only one species that occurred in all local communities and was left running for 5000 steps which we found was always sufficient for the system to reach a steady state. The last 100 steps of the simulation were then taken and used for analyses.

Spectral analysis

We performed 200 simulations for each combination of model parameters (Table 1). From each simulation we randomly selected one species and from this species we used two time series: (1) time series of population sizes within a randomly selected *local community* and (2) time series of population sizes within the whole *metacommunity*. In cases where the species went extinct in one or both time series we randomly selected another one thereby conditioning the time series as being drawn from a locally extant species. For these time series we conducted spectral analysis (Halley, 1996; pages 680-681 in Legendre and Legendre, 2003) and estimated the value of the exponent γ in $S(f) \propto 1/f^\gamma$ where $S(f)$ is spectral density at given frequency f ($0 \leq f \leq f_{n/2}$; n is length of the time series). Values of $S(f)$ were obtained through Fast Fourier Transform of the log-transformed time series data. The spectral exponent γ was estimated as minus the regression slope of the $\log(S(f))$ versus $\log(f)$. This procedure is exactly the same as used by Inchausti and Halley (2001) and hence our estimates of γ are directly comparable to theirs (Fig. 2). The value of γ can vary from $\gamma < 0$ ('blue noise') through $\gamma = 0$ ('white noise'), $\gamma \approx 1$ ('pink noise') and $\gamma = 2$ ('brown noise' or random walk) to $\gamma > 2$ ('black noise'). We use the term 'whitening' of the spectra for a deviation of γ from 2 towards 0.

To assess effects of the neutral model parameters on γ we fitted a multiple ANOVA model with γ as the response variable and five explanatory variables: mortality rate, migration rate, speciation rate, number of local communities and number of individuals. Since we were dealing with simulated data we only report R^2 s and do not report any P -values because these could be modified arbitrarily by changing number of simulations. Comparisons were made with the values of γ measured in natural populations taken from the Global Population Dynamics

195 Database (kindly provided by Pablo Inchausti; see also Inchausti and Halley, 2001,
196 2002).

197

198 ***Testing for density dependence***

199 Using the same data, we conducted three distinct tests for density dependence in the
200 time series: Pollard's randomization test (Pollard et al., 1987), Dennis and Taper's
201 parametric bootstrap maximum likelihood test (PLBR; Dennis and Taper, 1994) and
202 Bulmer's test (Bulmer, 1974). The Pollard's test is a distribution-free likelihood ratio
203 randomization test in which a random walk (Eq. 1) or a random walk with trend are
204 density independent hypotheses, whereas the density dependent hypothesis is defined
205 as a first-order *linear* autoregression model (stochastic Gompertz model):

$$206 \quad \ln N_{t+1} = \ln N_t + r + \beta \ln N_t + e_t \quad (\beta \neq 1) \quad (2)$$

207 where r and β are model parameters. The Bulmer's test is a parametric
208 predecessor of the Pollard's test and has been criticized for weak performance in cases
209 where there was a trend in the data ($r \neq 0$). The PLBR test is a likelihood ratio test
210 which uses the first-order *non-linear* autoregression model (stochastic logistic model)
211 as the density dependent hypothesis:

$$212 \quad \ln N_{t+1} = \ln N_t + r + \beta N_t + e_t \quad (3)$$

213 All of these tests have been criticized for disputable performance when there is
214 a census error affecting the estimation of population abundance in the field (Shenk et
215 al., 1998; Freckleton et al., 2006), but this is not the case for our simulated data. All
216 three tests gave nearly identical results and we therefore only present the output from
217 Pollard's randomization test (Pollard et al., 1987) which is 'Pollard's P ': the
218 probability of observing the empirical data in the case of the density independent
219 hypothesis (Eq. 1). Following the approach of Woiwod and Hanski (1992) we

normalized the distribution of Pollard's P by logit transformation. To assess effects of the neutral model parameters on Pollard's P we again fitted a multiple ANOVA model in which we used the values of logit-transformed Pollard's P as the response variable.

Taylor's power law

To test if populations in our model obey the Taylor's Power Law (TPL) we performed 50 simulations for each combination of parameters (Table 1). For each species in the simulations we created a metapopulation matrix (MPM) of abundances in which rows represent the last 100 time steps of the simulation and columns represent the populations in different local communities. We excluded all MPMs in which the species had gone globally extinct. We studied the TPL in its temporal context (Taylor and Woivod, 1982) which means that variances and means are calculated over each column (one column is one local population) in the MPM – when plotted, each point in the log-log plot therefore represents one local population. Using rows instead of columns in the MPM would yield the TPL in a spatial context, but the temporal context is more germane to this time series study. A linear regression was fitted in each of the log-log plots and the TPL slope (parameter b ; Taylor, 1961; Tokeshi, 1995) and the coefficients of determination (R^2) were calculated. As with our other results, the simulated values of TPL slopes and R^2 s were compared with empirically observed values (taken from Taylor and Woivod, 1980, 1982).

Results

Spectral analysis

245 Values of the scaling exponent γ estimated from our simulations are summarized in
 246 Figure 2. For local communities, the median value of γ was 1.59 (2.5% and 97.5%
 247 quantiles were 1.26 and 2.19 respectively). The value of γ was only influenced by
 248 mortality rate and migration rate (Table 2, Fig. 3) with higher mortality and higher
 249 migration each producing time series that were closer to pink noise ($\gamma < 2$) (Fig. 2).
 250 On the metacommunity scale, the median of γ was 1.75 (2.5% and 97.5% quantiles
 251 were 1.05 and 2.07 respectively). We did not detect any influence of the model
 252 parameters on the value of γ at the metacommunity scale (Table 2, Fig. 3). Our
 253 simulations produced γ values falling within a narrower interval than is generally
 254 observed in nature (Fig. 2), however, they departed from 2 (the value for an
 255 unbounded random walk) in the same direction as for natural populations.

256 In general, the population time series emerging from our neutral model had
 257 lower γ than would be expected for an autocorrelated random walk ($\gamma = 2$). This
 258 could partly be an artifact caused by the insufficient length of time series. It was
 259 recently shown that random walks of length ~ 100 actually give estimates of between
 260 1.5 and 1.8 instead of 2 because of the missing low-frequency fluctuations
 261 (Miramontes and Rohani, 2002). Nevertheless, since empirically collected time series
 262 are rarely longer than 100 sequential observations, our simulations of time series of
 263 similar lengths are amply justified to purposefully include the effects of sampling over
 264 a limited time scale. Since we also showed that the values of γ tended to be lower at
 265 local scales when comparing time series of equal lengths, we can still conclude that
 266 some whitening of spectra at local community scales is caused by more than just the
 267 length of the sampled time series.

268

269 *Testing for density dependence*

On the scale of local communities the median of Pollard's P was 0.14 (2.5% and 97.5% quantiles were 0 and 0.82 respectively) indicating a distribution skewed towards the lower values of Pollard's P . We also found that the values of Pollard's P were dependent on mortality rate and migration rate (Table 2, Fig. 4). Similarly to spectral analysis, simulations with higher mortality rate and higher migration rate produced time series with lower values of Pollard's P (Fig. 4). On the metacommunity scale the median value of Pollard's P was 0.5 (2.5% and 97.5% quantiles being 0.01 and 0.97 respectively). This indicates no clear signal of density dependence on the metacommunity scale. We did not detect any influence of the other model parameters on the values of logit-transformed Pollard's P (Table 2, Fig. 4). Pollard's P varied with relative abundance of species in local communities but not in metacommunities (Fig. 6).

Taylor's power law (TPL)

The median slope of TPL was 1.73 (2.5% and 97.5% quantiles were 0.059 and 2.49 respectively). The coefficients of determination (R^2) of power law regressions varied, following skewed, non-normal distributions (Figs. 2, 5), having generally lower values to what has been observed in nature (Fig. 2). The effects of model parameters on the slope of TPL was not as distinct as in spectral analysis and density dependence analysis; we detected a weak influence of number of local communities and mortality rate on the slopes of TPL (Table 2, Fig. 5). Both the slopes and R^2 s showed negative dependence on the mean relative abundance of species (Fig. 7).

Discussion

295 *Spectral analysis and density dependence*

296 The observed whitening of spectra in our model is not what we would expect under
 297 random walk dynamics. One possible reason for this is the effect of *migration* which
 298 can cause the spectral whitening. A population in an isolated local community (low or
 299 no immigration from outside) follows dynamics similar to an autocorrelated random
 300 walk ($\gamma \approx 2$) in which there is little or no dependence on the abundances in other local
 301 communities. However, after introducing immigration from outside of the local
 302 community, the autocorrelative nature of the process is broken and the species
 303 abundance in the whole metacommunity becomes important. The influence of single
 304 local community dynamics on the metacommunity are likely to be small (inversely
 305 proportional to the number of local communities), but in contrast the influence of the
 306 metacommunity on local community dynamics can be substantial (proportional to the
 307 migration rate). Hence, the migration causes the local time series to be whitened (non-
 308 autocorrelated in time) while the metacommunity time series remains “brownian”
 309 (autocorrelated in time). The *mortality* rate can also cause spectral whitening by
 310 adding random population fluctuations at high frequencies, but not at low ones
 311 subsequently lowering γ . We expect that this effect of mortality rate would not be
 312 apparent in very long time series (> 400 time steps; Miramontes and Rohani, 2002).
 313 However, such long time series are nearly never available in ecology and we wanted
 314 to keep our time series comparably long to the real-world ones.

315 The spectral analysis results are mirrored by the results of the tests for density
 316 dependence. Density dependence is modeled either as stochastic Gompertz (Eq. 2) or
 317 stochastic logistic (Eq. 3) models in the tests we used (Shenk et al., 1998, see also
 318 Methods).

It has been recognized for a long time (Eberhardt, 1970; Freckleton et al., 2006) that a sequence of uncorrelated random numbers (“white noise”) tends to follow a pattern consistent with density dependence. If you take white noise time series and plot changes of $\ln$ population size ($\ln N_{t+1} - \ln N_t$) against $\ln N_t$ (where N_t is the number of individuals at time t) you will get a linear negative relationship, corresponding (spuriously) to the density dependent Gompertz model (Eq. 2) with $\beta \approx -1$. Moreover, white noise can be very similar to the output of the stochastic logistic model (Eq. 3) with $\beta < 0$. We have learned that the whitening of a spectra means bringing it from $\gamma \approx 2$ down towards 0, which is white noise. We also know that migration (by breaking the autocorrelation structure) and mortality rate (by increasing spectral density at high frequencies in time series of limited length) cause spectral whitening. The detection of density dependence can also be attributed to the same mechanisms, although most ecologists would consider such density dependence to be spurious (Freckleton et al., 2006).

Uncorrelated noise in the form of census error also causes spurious detection of density dependence has long been known to population ecologists (Shenk et al., 1998; Freckleton et al., 2006). Moreover, Freckleton et al. (2006) showed an example of a population that follows a random walk but consists of two subpopulations each of which apparently shows density dependence because there is random migration between them. That immigration can cause time series to resemble a sequence of uncorrelated random numbers was also noted by Anderson et al. (1982) and this is indeed the case for our neutral model.

There should also be real density dependence in our model. In an isolated community, the closer N_t gets to zero (extinction) or to J (monodominance) the lower the probability of N_{t+1} differing from N_t will be. When a species almost fills the local

community, its probability of increasing its abundance in the next time step is small, because it is unlikely that one of the few individuals belonging to a different species will be selected for mortality. Similarly, when the abundance of a species is low, its probability of reproducing is low because after mortality it is not likely to be selected to reproduce. The highest rate of fluctuations and lowest incidence of density dependence is then expected when a population has a relative abundance of 0.5 (Hubbell, 2001). We detected this form of density dependence only in local communities at low relative abundances (Fig. 6). Species with high relative abundances were rare in our simulations and hence we were unable to clearly show the real density dependence for them (Fig. 6) even though in theory it must exist.

Taylor's power law (TPL)

The archipelago neutral model produced slopes of Taylor's Power Law with median close to the range observed in natural populations ($1 < b < 2$), but with higher incidence of low coefficients of determination. In order to explain this we need to summarize the mean-variance scaling properties of simple stochastic models. If a species undergoes an unrestricted random walk (Eq. 1), the TPL has a slope $b = 1$ at very low population densities and gradually changes to $b = 2$ at a certain density (Anderson et al., 1982; Tokeshi, 1995; Keeling, 2000). The population density at which b starts to change from 1 to 2 depends on the relative magnitude of stochastic changes because of the high incidence of extinctions at low population densities which lower the variance. This effect can explain $b < 2$ at very low densities.

Hubbell (2001) shows what happens with the mean-variance scaling if the stochastic population fluctuations are restricted by an upper boundary (the zero-sum rule) and are fed by immigration, although Hubbell does not use the term Taylor's

369 power law. Hubbell (2001) predicts hump-shaped parabolic relationship between
 370 abundance of species in a metacommunity and the variance of a local population size
 371 undergoing the zero-sum ecological drift. He also hypothesizes that most of the
 372 species in nature have low relative abundances and hence should occur in the left part
 373 (relative abundance < 0.2) of the parabola which is nearly linear in the log-log space,
 374 resembling the power-law with slope of 2. Although Hubbell's (2001) reasoning
 375 focuses on the relationship between mean and variance in a set of species within one
 376 community (one point in the mean-variance plot represents one species) it can apply
 377 to the set of single-species local populations in our model: the mean relative
 378 abundances of species in local communities were rarely higher than 0.2 in our
 379 simulations and indeed, there is a much better fit of the TPL for species with low
 380 abundances (Fig. 7). The poor fit at higher relative abundances is then a result of
 381 fitting the linear TPL on the actually parabolic mean-variance relationship.
 382 Consequently, as the mean relative abundance of species increases, the slope of the
 383 TPL decreases (Fig. 7).

384

385 *Comparison with empirical data and conclusions*

386 Our model produces qualitatively similar patterns to those observed in nature. This is
 387 perhaps surprising given the assumptions made by the neutral model, but it does again
 388 highlight its use as a powerful null model.

389 There is a general agreement that natural time series are not "brownian" (Fig.
 390 1; Arino and Pimm, 1995; Inchausti and Halley, 2001, 2002) – they clearly fall into
 391 the whitened range of scaling exponents ($0 < \gamma < 2$; Fig. 2). Our simulations indicate
 392 that neutral models have the potential to explain part of this observed whitening,
 393 although this potential may be limited. Although higher migration rates can result in

time series with more realistic values of γ , our migration rates are similar (10%) or much higher (30%) than what is usually observed in natural metacommunities (see Volkov et al., 2003 and Mutshinda et al., 2008). Mutshinda et al. (2008) already showed that sizes of local communities fluctuate more than is expected from neutral theory. This resonates with our simulations of individual species time series which show rather high values of γ (i.e. lower importance of high-frequency fluctuations) when compared to empirical data. Recent progress in population ecology suggests that a promising explanation for these high fluctuations of natural populations might be environmental stochasticity (Lande et al., 2006; Mutshinda et al., 2009) which is absent in neutral models, but not impossible to include in future and more general versions of the original theory (Allen and Savage, 2007).

Realistic slopes of TPL can be produced by a number of different mechanisms (Tokeshi, 1995; Keeling, 2000; Kendal, 2004; Ballantyne and Kerkhoff, 2007) and it is therefore difficult to infer any definitive underlying mechanisms from its shape (Mitzenmacher, 2003). Here we add neutral models to the list of possible mechanisms that can produce realistic TPL slopes, although not necessarily realistic fits.

In conclusion, our results show that realistic slopes of Taylor's power law and density dependence can emerge from a simple neutral model, whilst spectral colours are less likely, but still possible to be reproduced by the neutral model. Still some other natural phenomena remain unexplained by our model, such as the possibility for very small γ closer to white noise. Our findings suggest the interesting possibility that the properties of population time series observed in nature are in part due to simple factors such as sampling effects and dispersal limitation, which would then represent the most parsimonious explanation of observed patterns. The patterns of population fluctuations observed in nature may not reflect resource-driven species-specific

dynamics but instead processes at the metacommunity scale, with an important effect of stochastic mortality and random immigration of individuals into local communities. The neutral model encompasses these factors without being overly complex and we therefore suggest that neutral models make more powerful null models for time series data compared to alternatives such as Brownian random walks.

Acknowledgements

We are grateful to Rampal Etienne, Stephen Cornell, William E. Kunin and Samuel Zajicek for discussions on the subject and to one anonymous referee for helpful comments. Pablo Inchausti generously provided us the GPPD data on spectral exponents. The study was supported by the EU FP7 project SCALES (project 226852), by the Grant Agency of Charles University (GAUK 106108), by the Grant Agency of the Academy of Sciences of the CR (IAA601970801) and by the Czech Ministry of Education (LC06073, MSM0021620845). James Rosindell was supported with funding from the EPSRC (grant EP/F043112/1). Petr Keil was supported with Marie Curie Early Stage Training fellowship at University of Leeds (UK).

References

- Allen, A.P., Savage, V.M. 2007. Setting the absolute tempo of biodiversity dynamics. *Ecol. Lett.* 10, 637–646.
- Alonso, D., Etienne, R.S., McKane, A.J. 2006. The merits of neutral theory. *Trends Ecol. Evol.* 21, 451–457., doi:10.1016/j.tree.2006.03.019.

- 443 Anderson, R.M., Gordon, D.M., Crawley, M.J., Hassell, M.P. 1982. Variability in the
444 abundance of animal and plant species. *Nature* 296, 245-296.
- 445 Arino, A., Pimm, S.L. 1995. On the nature of population extremes. *Evol. Ecol.* 9, 429-
446 443.
- 447 Azaele, S., Pigolotti, S., Banavar, J.R., Maritan, A. 2006. Dynamical evolution of
448 ecosystems. *Nature* 444, 926-928.
- 449 Ballantyne, F., Kerkhoff, A.J. 2007. The observed range for temporal mean-variance
450 scaling exponents can be explained by reproductive correlation. *Oikos* 116,
451 174-180.
- 452 Brook, B.W., Bradshaw, C.J.A. 2006. Strength of evidence for density dependence in
453 abundance time series of 1198 species. *Ecology* 87, 1445-1451.
- 454 Bulmer, M.G. 1974. A statistical analysis of the 10-year cycle in Canada. *J. Anim.*
455 *Ecol.* 43, 701-718.
- 456 Chave, J. 2004. Neutral theory and community ecology. *Ecol. Lett.* 7, 241-253.
- 457 Dennis, B., Taper, M.L. 1994. Density dependence in time series observations of
458 natural populations: Estimation and testing. *Ecol. Monogr.* 64, 205-224.
- 459 Dornelas, M., Connolly, S.R., Hughes, T.P. 2006. Coral reef diversity refutes the
460 neutral theory of biodiversity. *Nature* 440, 80-82.
- 461 Eberhardt, L.L. 1970. Correlation, regression and density dependence. *Ecology* 51,
462 306-310.
- 463 Etienne, R.S. 2005. A new sampling formula for neutral biodiversity. *Ecol. Lett.* 8,
464 253-260.
- 465 Etienne, R.S., Haegeman, B. 2010. The neutral theory of biodiversity with random
466 fission speciation. *Theor. Ecol.* 'in press'.

- 467 Freckleton, R.P., Watkinson, A.R., Green, R.E., Sutherland, W.J. 2006. Census error
468 and the detection of density dependence. *J. Anim. Ecol.* 75, 837-851.
- 469 Gotelli, N.J., McGill, B.J. 2006. Null versus neutral models: what's the difference?
470 *Ecography* 29, 793-800.
- 471 Haegeman, B., Etienne, R.S. 2010. Self-consistent approach for neutral community
472 models with speciation. *Phys. Rev. E* 81, 031911.
- 473 Halley, J.M. 1996. Ecology, evolution and 1/f-noise. *Trends Ecol. Evol.* 11, 33-37.,
474 doi:10.1016/0169-5347(96)81067-6.
- 475 Hanski, I., Foley, P., Hassell, M. 1996. Random walks in a metapopulation: how
476 much density dependence is necessary for long-term persistence? *J. Anim.*
477 *Ecol.* 65, 274-282.
- 478 Herben, T., Mandák, B., Bímová, K., Münzbergová, Z. 2004 Invasibility and species
479 richness of a community: a neutral model and a survey of published data.
480 *Ecology* 85, 3223–3233.
- 481 Herben, T. 2009. Invasibility of neutral communities. *Basic Appl. Ecol.* 10, 197–207.,
482 doi:10.1016/j.baae.2008.08.006.
- 483 Hubbell, S.P. 2001. The unified neutral theory of biodiversity and biogeography.
484 Princeton Univ. Press, Princeton.
- 485 Hubbell, S.P., Lake, J.K. 2003. The neutral theory of biodiversity an biogeography,
486 and beyond. In: Blackburn, T.M., Gaston, K.J. (Eds.), *Macroecology:*
487 *Concepts and Consequences.* Blackwell, Oxford, pp. 45-63.
- 488 Inchausti, P., Halley, J. 2001. Investigating long-term ecological variability using the
489 global population dynamics catabase. *Science* 293, 655-657.
- 490 Inchausti, P., Halley, J. 2002. The long-term temporal variability and spectral colour
491 of animal populations. *Evol. Ecol. Res.* 4, 1033-1048.

- 492 Keeling, M.J. 2000. Simple stochastic models and their power-law type behaviour.
 493 Theor. Popul. Biol. 58, 21-31., doi:10.1006/tpbi.2000.1475.
- 494 Kendal, W.S. 2004. Taylor's ecological power law as a consequence of scale invariant
 495 exponential dispersion models. Ecol. Complex. 1, 193-209.,
 496 doi:10.1016/j.ecocom.2004.05.001.
- 497 Lande, R., Engen, S., Saether, B.-E. 2006. Stochastic population dynamics in ecology
 498 and conservation. Oxford. Univ. Press, Oxford.
- 499 Legendre, P., Legendre L. 2003. Numerical Ecology. Elsevier, Amsterdam.
- 500 Leibold, M.A., McPeck, M.A. 2006. Coexistence of the niche and neutral perspectives
 501 in community ecology. Ecology 87, 1399-1410.
- 502 Leigh, E.G. 2007. Neutral theory: a historical perspective. J. Evolution. Biol. 20,
 503 2075-2091.
- 504 McGill, B.J. 2003. A test of the unified neutral theory of biodiversity. Nature 422,
 505 881-885.
- 506 McGill, B.J., Maurer, B.A., Weiser, M.D. 2006. Empirical evaluation of neutral
 507 theory. Ecology 87, 1411-1423.
- 508 Miramontes, O., Rohani, P. 2002. Estimating $1/f^\alpha$ scaling exponents from short time-
 509 series. Physica D 166, 147-154., doi:10.1016/S0167-2789(02)00429-3.
- 510 Mitzenmacher, M. 2003. A brief history of generative models for power laws and
 511 lognormal distributions. Internet Mathematics 1, 226-251.
- 512 Murdoch, W.W. 1994. Population regulation in theory and practice. Ecology 75, 271-
 513 287.
- 514 Mutshinda, C.M., O'Hara, R.B.O., Woivod, P. 2008. Species abundance dynamics
 515 under neutral assumptions: a Bayesian approach to the controversy. Funct.
 516 Ecol. 22, 340-347.

- 517 Mutshinda, C.M., O'Hara, R.B.O., Woiwod, P. 2009. What drives community
518 dynamics? *P. Roy. Soc. B-Biol. Sci.* 276, 2923-2929.
- 519 Pollard, E., Lakhani, K.H., Rothery, P. 1987. The detection of density-dependence
520 from a series of annual censuses. *Ecology* 68, 2046-2055.
- 521 Rosindell, J., Cornell, S.J. 2007. Species-area relationships from a spatially explicit
522 neutral model in an infinite landscape. *Ecol. Lett.* 10, 586-595.
- 523 Rosindell, J., Cornell, S.J. 2009. Species-area curves, neutral models, and long-
524 distance dispersal. *Ecology* 90, 1743-1750.
- 525 Rosindell, J., Cornell, S.J., Hubbell, S.P., Etienne, R.S. 2010. Protracted speciation
526 revitalizes the neutral theory of biodiversity. *Ecol. Lett.* 'in press'.
- 527 Rosindell, J., Wong, Y., Etienne, R.S. 2008. Coalescence models for spatial neutral
528 ecology. *Ecol. Inform.* 90, 1743-1750., doi:10.1016/j.ecoinf.2008.05.001.
- 529 Shenk, T.M., White, G.C., Burnham, K.P. 1998. Sampling-variance effects on
530 detecting density dependence from temporal trends in natural populations.
531 *Ecol. Monogr.* 68, 445-463.
- 532 Sibly, R.M., Barker, D., Hone, J., Pagel, M. 2007. On the stability of populations of
533 mammals, birds, fish and insects. *Ecol. Lett.* 10, 970-976.
- 534 Taylor, L.R., Woiwod, I.P. 1980. Temporal stability as a density-dependent species
535 characteristic. *J. Anim. Ecol.* 49, 209-224.
- 536 Taylor, L.R., Woiwod, I.P. 1982. Comparative synoptic dynamics. I. Relationships
537 between inter- and intra-specific spatial and temporal variance/mean
538 population parameters. *J. Anim. Ecol.* 51, 879-906.
- 539 Taylor, L.R., Taylor, R.A.J., Woiwod, I.P., Perry, J.N. 1983. Behavioural dynamics.
540 *Nature* 303, 801-804.

- 541 Tilman, D. 2004. Niche tradeoffs, neutrality, and community structure: A stochastic
542 theory of resource competition, invasion, and community assembly. *P. Natl.*
543 *Acad. Sci. USA* 101, 10854-10861.
- 544 Tokeshi, M. 1995. On the mathematical basis of the variance-mean power
545 relationship. *Researches on Population Ecology* 37, 43-48.
- 546 Turnbull, L.A., Manley, L., Rees, M. 2005. Niches, rather than neutrality, structure a
547 grassland pioneer guild. *P. Roy. Soc. B-Biol. Sci.* 272, 1357-1364.
- 548 Volkov, I., Banavar, J.R., Hubbell, S.P., Martian, A. 2003. Neutral theory and relative
549 species abundance in ecology. *Nature* 424, 1035-1037.
- 550 Volkov, I., Banavar, J.R., He, F., Hubbell, S.P., Maritan, A. 2005. Density
551 dependence explains tree species abundance and diversity in tropical forests.
552 *Nature* 438, 658-661.
- 553 Volkov, I., Banavar, J.R., Hubbell, S.P., Maritan, A. 2007. Patterns of relative species
554 abundance in rainforests and coral reefs. *Nature* 450, 45-49.
- 555 Woiwod, I.P., Hanski, I. 1992. Patterns of density dependence in moths and aphids. *J.*
556 *Anim. Ecol.* 61, 619-629.
- 557

Figure legends

Figure 1 Schematic illustration of the “archipelago” model. Note that we performed our simulations using much larger numbers of local communities than illustrated here. See the methods for detailed description of the model.

Figure 2 Some aspects of real-world populations and those produced by our neutral model. Empirical slopes of Taylor’s Power Law and coefficients of determination of TPL of 429 species of aphids, moths and birds were taken from Taylor & Woiwod (1980, 1982). Empirical values of scaling exponent γ of power spectra of 544 time series were taken from the Global Population Dynamics Database (GPDD, kindly provided by Pablo Inchausti; see also Inchausti and Halley, 2001, 2002). Box and whisker plots show medians, quartiles and outliers.

Figure 3 Results of the spectral analysis of population time series produced by a neutral model. γ is the scaling exponent in $S(f) \propto 1/f^\gamma$ where $S(f)$ is spectral density and f is frequency. Displayed are the effects of the neutral model’s parameters (Tab. 1) on γ as a box and whisker plot with median and quartiles. A random walk of sufficient length would have $\gamma = 2$.

Figure 4 Results of the testing for density dependence in population time series produced by neutral model as a box and whisker plot with median and quartiles. Pollard’s P is the probability of obtaining the data in case of density independence (Eq. 1; Pollard et al., 1987).

582 **Figure 5** The slopes of temporal Taylor's power law (TPL) as produced by neutral
 583 models with various levels of model parameters. Solid horizontal lines delimit the
 584 range observed in natural populations. Shown as a box and whisker plot with median
 585 and quartiles shown, outliers are not displayed.

586
 587 **Figure 6** Dependence of Pollard's P on relative abundance of species in local
 588 communities and in metacommunities. Shades of grey represent density of points in
 589 hexagonal bins.

590
 591 **Figure 7** Dependence of slope and coefficient of determination (R^2) of Taylor's
 592 Power Law on mean relative abundance of species in local communities. Shades of
 593 grey represent density of points in hexagonal bins.

4. Figure

4. Figure

Table 1 An overview of the parameters and their levels used in our neutral model simulations. See material and methods for detailed description of the exact function of each parameter.

Parameter	Description	Levels used in simulations
Number of local communities	number of local communities that make up the metacommunity	100, 20
Size of each local community (<i>J</i>)	number of individuals in each local community	2000, 500
Mortality rate (<i>d</i>)	mortality rate of individuals per time step in the local community	0.6, 0.3
Migration rate (<i>m</i>)	proportion of vacant positions in the community replaced by migration from other local communities	0.1, 0.3
Speciation rate (<i>s</i>)	speciation rate per species per step	2×10^{-6} , 3×10^{-6}

Table 2 The strength of the effect of each parameter on the value of γ and the logit-transformed Pollard's P . Parameter γ describes the 'colour' of the spectra of a time series. Pollard's P is the probability of obtaining the data in case of density independence (random walk). The values of R^2 were estimated using a multiple ANOVA. Figures in bold show values higher than 0.001.

	Local communities		Metacommunities		
	R^2 (γ)	R^2 (logit of Pollard's P)	R^2 (γ)	R^2 (logit of Pollard's P)	R^2 (slope of TPL)
Mortality rate	0.045	0.078	<0.001	<0.001	0.019
Migration rate	0.11	0.19	<0.001	<0.001	0.0014
Speciation rate	<0.001	<0.001	<0.001	<0.001	<0.001
# of local communities	<0.001	0.014	<0.001	<0.001	0.048
Size of local community	<0.001	<0.001	<0.001	<0.001	0.0015