Risk profiles and penetrance estimations in multiple endocrine neoplasia type 2A caused by germline RET mutations located in exon 10

Karin Frank-Raue, Lisa A. Rybicki, Zoran Erlic, Heiko Schweizer, Aurelia Winter, Ioana Milos, Sergio Pa Toledo, Rodrigo A Toledo, Marcos R Tavares, Maria Alevizaki, et al.

► To cite this version:

Karin Frank-Raue, Lisa A. Rybicki, Zoran Erlic, Heiko Schweizer, Aurelia Winter, et al.. Risk profiles and penetrance estimations in multiple endocrine neoplasia type 2A caused by germline RET mutations located in exon 10. Human Mutation, 2010, 32 (1), pp.51. 10.1002/humu.21385 . hal-00599477

HAL Id: hal-00599477

<https://hal.science/hal-00599477>

Submitted on 10 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Risk profiles and penetrance estimations in multiple endocrine neoplasia type 2A caused by germline RET mutations located in exon 10

Journal:	<i>Human Mutation</i>
Manuscript ID:	humu-2010-0302.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	27-Aug-2010
Complete List of Authors:	<p>Frank-Raue, Karin; Endocrine Practice and Molecular Laboratory Rybicki, Lisa; Cleveland Clinic, Department of Quantitative Health Sciences and Taussig Cancer Institute Erlic, Zoran; University of Freiburg Medical Centre, Section of Preventive Medicine, Department of Nephrology Schweizer, Heiko; University of Freiburg Medical Centre, Section of Preventive Medicine, Department of Nephrology Winter, Aurelia; University of Freiburg Medical Centre, Section of Preventive Medicine, Department of Nephrology Milos, Ioana; University of Freiburg Medical Centre, Section of Preventive Medicine, Department of Nephrology Toledo, Sergio; Hospital das Clinicas, University of Sao Paulo School of Medicine, Endocrine Genetics Unit (LIM-25), Endocrinology, Department of Internal Medicine Toledo, Rodrigo; Hospital das Clinicas, University of Sao Paulo School of Medicine, Endocrine Genetics Unit (LIM-25), Endocrinology, Department of Internal Medicine Tavares, Marcos; Hospital das Clinicas, University of Sao Paulo School of Medicine, Head and Neck Division, Department of Surgery Alevizaki, Maria; Alexandra Hospital, University School of Medicine, Endocrine Unit, Department of Medical Therapeutics Mian, Caterina; University of Padua, Operative Unit of Endocrinology Department of Surgical and Medical Sciences Siggelkow, Heide; University of Göttingen, Department of Endocrinology Hüfner, Michael; University of Göttingen, Department of Endocrinology Wohlk, Nelson; Universidad de Chile, Endocrine Section, Hospital del Salvador Opocher, Giuseppe; University of Padova, Padova, Familial Cancer Clinic, Veneto Institute of Oncology, IRCCS & Department of Medical and Surgical Sciences Dvořáková, Šárka; Institute of Endocrinology Prague, Department</p>

	<p>of Molecular Endocrinology Bendlova, Bela; Maria Sklodowska Curie Memorial Center and Institute of Oncology, Department of Nuclear Medicine and Oncological Endocrinology Czetwertynska, Małgorzata; Maria Sklodowska Curie Memorial Center and Institute of Oncology, Department of Endocrinology Skasko, Elżbieta; Maria Sklodowska Curie Memorial Center and Institute of Oncology, Department of Endocrinology Barontini, Marta; Hospital de Niños R. Gutiérrez, Centro de Investigaciones Endocrinológicas Sanzo, Gabriela; Hospital de Niños R. Gutiérrez, Centro de Investigaciones Endocrinológicas Vorländer, Christian; Bürgerhospita, Department of Endocrine Surgery Maia, Ana; Hospital de Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Thyroid Section, Endocrine Division Patocs, Attila; Semmelweis University, Molecular Medicine Research Group, Hungarian Academy of Sciences and 2nd Department of Medicine Links, Thera; University Medical Center Groningen, Department of Endocrinology de Groot, Jan; University Medical Center Groningen, Department of Endocrinology Kerstens, Michiel; University Medical Center Groningen, Department of Endocrinology Valk, Gerlof D.; University Medical Center Utrecht, Department of Internal Medicine Miehle, Konstanze; Universityhospital of Leipzig, Department of Endocrinology Musholt, Thomas; University Medical Center of the Johannes Gutenberg University Mainz, Endocrine Surgery, Department of General and Abdominal Surgery Biarnes, Josefina; Hospital Dr Josep Trueta de Girona, Endocrinology Unit Damjanovic, Svetozar; University of Belgrade, Institute of Endocrinology, Diabetes & Metabolic Diseases, Medical School Muresan, Mihaela; Hôpital N.-D. de Bon Secours, Department of Endocrinology Wüster, Christian; Private Praxis for Endocrinology Fassnacht, Martin; University Hospital, Department of Medicine I Peczkowska, Mariola; Institute of Cardiology, Department of Hypertension Fauth, Christine; Innsbruck Medical University, Division of Clinical Genetics, Department of Medical Genetics, Molecular and and Clinical Pharmacology Golcher, Henriette; Friedrich-Alexander Universityhospital of Erlangen, Department of Surgery Walter, Martin; Institute of Nuclear Medicine, University Hospital Pichl, Josef; St.Theresien-Hospital Raue, Friedhelm; Endocrine Practice and Molecular Laboratory Eng, Charis; Lerner Research Institute and Taussig Cancer Institute, Cleveland Clinic, Genomic Medicine Institute Neumann, Hartmut; Medical University Clinic, Nephrology</p>
Key Words:	Multiple endocrine neoplasia type 2, medullary thyroid carcinoma, pheochromocytoma, genotype-phenotype correlation

SCHOLARONE™
Manuscripts

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

HuMu 2010-0302 Revised

**Risk profiles and penetrance estimations in multiple endocrine neoplasia type
2A caused by germline *RET* mutations located in exon 10**

Karin Frank-Raue^{1*}, Lisa A. Rybicki^{2*}, Zoran Erlic^{3*}, Heiko Schweizer^{3*}, Aurelia Winter³, Ioana Milos^{3,4}, Sergio PA Toledo⁵, Rodrigo A Toledo⁵, Marcos R Tavares⁶, Maria Alevizaki⁷, Caterina Mian⁸, Heide Siggelkow⁹, Michael Hüfner¹⁰, Nelson Wohllk¹¹, Giuseppe Opocher¹², Šárka Dvořáková¹³, Bela Bendlova¹⁴, Małgorzata Czetwertynska¹⁵, Elżbieta Skasko¹⁵, Marta Barontini¹⁶, Gabriela Sanso¹⁶, Christian Vorländer¹⁷, Ana Luiza Maia¹⁸, Attila Patocs¹⁹, Thera P. Links²⁰, Jan Willem de Groot²⁰, Michiel N. Kerstens²⁰, Gerlof D Valk²¹, Konstanze Miehle²², Thomas J. Musholt²³, Josefina Biarnes²⁴, Svetozar Damjanovic²⁵, Mihaela Muresan²⁶, Christian Wüster²⁷, Martin Fassnacht²⁸, Mariola Peczkowska²⁹, Christine Fauth³⁰, Henriette Golcher³¹, Martin A Walter³², Josef Pichl³³, Friedhelm Raue^{1**}, Charis Eng^{34**}, Hartmut PH Neumann^{3**} for the International *RET* Exon 10 Consortium

*These authors contributed equally to this study.

** These authors are senior authors.

¹Endocrine Practice and Molecular Laboratory, Heidelberg, Germany

²Department of Quantitative Health Sciences and Taussig Cancer Institute, Cleveland Clinic; Cleveland, OH, 44195; USA

³Section of Preventive Medicine, Department of Nephrology, University of Freiburg

Medical Centre, Freiburg im Breisgau, Germany

⁴Department of Endocrinology of the University of Medicine and Pharmacy “Victor Babes” Timisoara, Romania

⁵ Endocrine Genetics Unit (LIM-25), Endocrinology, Department of Internal Medicine, Hospital das Clinicas, University of Sao Paulo School of Medicine, Sao Paulo, Brazil

⁶Head and Neck Division, Department of Surgery, Hospital das Clinicas, University of Sao Paulo School of Medicine, Sao Paulo, Brazil

⁷Endocrine Unit, Department of Medical Therapeutics, Alexandra Hospital, University School of Medicine, Athens, Greece

⁸Operative Unit of Endocrinology Department of Surgical and Medical Sciences University of Padua, Italy

⁹Department of Gastroenterology and Endocrinology, University of Göttingen and Endokrinologikum Göttingen, Germany

¹⁰Endokrinologikum Göttingen, Germany

¹¹Endocrine Section, Hospital del Salvador, Universidad de Chile, Santiago de Chile, Chile

¹²Familial Cancer Clinic, Veneto Institute of Oncology, IRCCS & Department of Medical and Surgical Sciences, University of Padova, Padova, Italy

¹³Department of Molecular Endocrinology of the Institute of Endocrinology Prague, Prague, Czech Republic

¹⁴Department of Nuclear Medicine and Oncological Endocrinology, Maria Sklodowska Curie Memorial Center and Institute of Oncology, Warsaw, Poland

- Seite 3 von 24

²⁹Department of Hypertension, Institute of Cardiology, Warsaw, Poland

³⁰Division of Clinical Genetics, Department of Medical Genetics, Molecular and and
Clinical Pharmacology, Innsbruck Medical University, Innsbruck, Austria

³¹Department of Surgery, Friedrich-Alexander Universityhospital of Erlangen,
Erlangen, Germany

³²Institute of Nuclear Medicine, University Hospital, Basel, Switzerland

³³St. Theresien-Hospital, Nuernberg, Germany

³⁴Genomic Medicine Institute, Lerner Research Institute, Taussig Cancer Institute,
and Stanley Shalom Zielony Nursing Institute, Cleveland Clinic, Cleveland, OH, USA

Acknowledgement:

This study is funded in part by grants from the Deutsche Krebshilfe (70-3313-Ne 1 to HPHN), the Deutsche Forschungsgemeinschaft (NE 571/5-3 to HPHN), the European Union (LSHC-CT-2005-518200 to HPHN).

RAT was supported by a Sao Paulo State Research Foundation-FAPESP grant (2009/15386-6), and SPAT by a National Research Foundation-CNPq grant (310614/09)

CE was a recipient of the Doris Duke Distinguished Clinical Scientist Award, holds the Sondra J. and Stephen R. Hardis Endowed Chair of Cancer Genomic Medicine at the Cleveland Clinic, and is the American Cancer Society Clinical Research Professor, generously funded in part by the F. M. Kirby Foundation.

ALM is a recipient of a National Research Foundation-CNPq grant (305629/07).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Short Title: MEN 2 in Patients with RET Exon 10 Mutations

Key words: Multiple endocrine neoplasia type 2, medullary thyroid carcinoma, pheochromocytoma, genotype-phenotype correlation

Corresponding address:

Hartmut P.H. Neumann, MD

Medizinische Universitätsklinik,

Abteilung Innere Medizin 4, Sektion für Präventive Medizin

Hugstetter Str. 55

D 79106 Freiburg, Germany

Phone: 0761 270 3578, Fax: 0761 270 3778

Email: hartmut.neumann@uniklinik-freiburg.de

Formatted: English U.K.

Formatted: English U.S.

Formatted: English U.S.

Formatted: English U.S.

Field Code Changed

Abstract

Multiple endocrine neoplasia type 2 is characterized by germline mutations in *RET*. For exon 10, comprehensive molecular and corresponding phenotypic data are scarce. The International *RET* Exon 10 Consortium, comprising 27 centers from 15 countries, analyzed patients with *RET* exon 10 mutations for clinical risk profiles. Presentation, age-dependent penetrance and stage-at-presentation of medullary thyroid carcinoma (MTC), pheochromocytoma and hyperparathyroidism were studied. A total of 340 subjects from 103 families, age 2-86, were registered. There were 21 distinct single nucleotide germline mutations located in codons 609 (45 subjects), 611 (50), 618 (94), and 620 (151). MTC was present in 263 registrants, pheochromocytoma in 54 and hyperparathyroidism in 8 subjects. Of the patients with MTC, 53% were detected when asymptomatic, and amongst those with pheochromocytoma, 54%. Penetrance for MTC was 4% by age 10, 25% by 25 and 80% by 50. Codon-associated penetrance by age 50 ranged from 60% (codon 611) to 86% (620). More advanced stage and increasing risk of metastases correlated with mutation in codon position (609->620) near the juxtamembrane domain. Our data provide rigorous bases for timing of pre-morbid diagnosis and personalized treatment/prophylactic procedure decisions depending on specific *RET* exon 10 codons affected.

Word count: 194

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Multiple endocrine neoplasia type 2 (MEN 2) is an autosomal dominant multi-glandular tumor syndrome, affecting derivatives from neural crest. MEN 2 is caused by germline mutations of the *RET* proto-oncogene (OMIM164761), encoding a transmembrane receptor tyrosine kinase expressed by the C-cells of the thyroid, adrenal medullary cells, and enteric autonomic ganglia (Elisei, et al., 2007; Eng C, 1996 ; Zbuk and Eng, 2007). When *RET* mutations were found in the large majority of MEN 2 patients and a *RET* genotype-clinical phenotype found (Eng C, 1996), this was the paradigm for the practice of clinical cancer genetics (Kloos, et al., 2009; Zbuk and Eng, 2007). Accurate molecular diagnosis, pre-morbid predictive testing and gene-informed medical management became possible.

Deleted: (Kloos, et al., 2009; Zbuk and Eng, 2007; Eng Ann NY Acad Sci 2010 in press)

Based on combinations of clinical manifestations, MEN 2 is historically classified into three clinical subtypes, MEN 2A (MIM# 171400), MEN 2B (MIM# 162300), and familial medullary thyroid carcinoma (FMTC, MIM# 155240). The components of MEN 2A are medullary thyroid carcinoma (MTC), pheochromocytoma and parathyroid adenoma and/or hyperplasia (HPT). In MEN 2B, constitutional abnormalities such as marfanoid habitus and neuromas of the tongue and the intestine are present whereas clinical parathyroid disease is absent. FMTC refers to the familial occurrence of MTC without other lesions (Kloos, et al., 2009). MEN 2B is considered more aggressive than MEN 2A or FMTC, because the age-of-onset of the neoplasias is a mean 10 years earlier than that of MEN 2A, and metastases from MTC occur as early as 3 years of age (Kloos, et al., 2009).

The spectrum of germline *RET* mutations in MEN 2 are encompassed by 64 distinct mutations that lie in 32 codons belonging to 8 exons (Kloos, et al., 2009) . It is clear that the MEN 2B-defining mutations, M918T and A883F, portend an aggressive

course, with management appropriately aggressive and pre-emptive (Kloos, et al., 2009). It is also obvious and relatively consistent that the exon 11 C634R, the most common mutation seen in MEN 2A, predicts for the development of MTC, pheochromocytoma and HPT, at relatively young ages, and so management is tailored accordingly (Eng C, 1996 ; Frank-Raue, et al., 2006; Kloos, et al., 2009; Neumann, et al., 2002). However, apart from vague generalizations such as non-634 mutations are not as aggressive, perhaps have later ages of onset and decreased penetrance, the breadth of clinical presentations and course, and the less-than-firm data preclude clinicians from altering medical management based on precise single-codon/specific missense mutation genotype. For instance, studies have shown that mutations at codon 634, the most frequently mutated in MEN 2, result in a receptor tyrosine kinase that is constitutively active by the tendency of mutant *RET* monomers to undergo dimerization, a situation that simulates ligand binding. Whereas cysteine residues in wild-type *RET* form intramolecular disulfide bonds, mutation of a cysteine residue leaves an unpaired residue. Unpaired cysteines from two mutant *RET* monomers can dimerize by formation of a disulfide bond. Taking into account the clinical presentation, we know that the C634R mutation is highly penetrant and associated with the more severe forms of MEN 2A (Eng C, 1996 ; Kloos, et al., 2009). Similarly, while investigating pheochromocytoma presentations in our Registry, we noted that MTC does not always occur before other component neoplasias in those with C634W mutations (Neumann, et al., 2007). Therefore, we believe that single nucleotide genotype-specific clinical risk profiles comprising risk of developing each component neoplasia, the ages-at-onset, penetrance and clinical course might be useful for tailoring more precise clinical care.

In contrast to well-defined risk profiles for carriers of the C634R, C634Y, C634W,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A883F, and M918T mutations, mutations located in the codons of *RET* exon 10 are relatively rare (Mian, et al., 2009; Moers, et al., 1996; Siggelkow, et al., 2001), and their anecdotal penetrance and ages-of-onset have resulted in appropriately cautious and tentative non-consensual clinical guidelines (Brandi, et al., 2001; Kloos, et al., 2009). Here, concerted effort from an international consortium have resulted in *RET* exon 10 genotyping, documented phenotypes-at-presentation and longitudinal clinical data with the purpose of generating codon- and nucleotide-specific genotype-adjusted neoplasia risk profiles for exon 10 mutations.

Patients and Methods

Patients

The inclusion criterion for registrants was a proven carrier status of a germline mutation in exon 10 of the *RET* gene. In addition, relatives of index registrants were also registered if an MTC or a pheochromocytoma was diagnosed histologically. The ascertainment of carriers of such mutations was performed in 3 steps. First, *RET* exon 10 mutation carriers were identified from existing registrants in the European-American Pheochromocytoma Registry (EAPR), which has been previously described (Erlic, et al., 2009; Neumann, et al., 2002; Neumann, et al., 2004; Neumann, et al., 2007; Schiavi, et al., 2005). Second, all clinicians and researchers who had contributed to this registry or to our previous risk estimation study of *RET* C634W carriers were contacted (Milos, et al., 2008). Finally, all known centers with clinical activities in MEN 2 or one of the component neoplasias (eg, MTC or pheochromocytoma) were asked to contribute any relevant data to this study.

Clinical Data

Recording of demographic and clinical information was performed using a uniform data-format for all registrants. Demographic data included year of birth, gender, ethnic background and country of residence. Clinical data were recorded for thyroid, adrenal and parathyroid glands, and additional features of MEN 2, as follows.

For all three components, i.e., MTC, pheochromocytoma and HPT, the age(s)-at-diagnosis were registered using the age at histological diagnosis. Criteria for absence of MTC included normal basal serum calcitonin or histological absence in surgical thyroid specimens. Absence of pheochromocytoma was documented either by normal CT or MRI of the adrenal glands or by normal 24-hour catecholamine excretion. Normal serum calcium and parathyroid hormone levels denoted an absence of hyperparathyroidism.

We differentiated symptomatic patients and subjects who were identified by screening procedures (screening assessment). Screening procedures included genetic screening of relatives after a *RET* mutation was identified and verified in an index case. This was followed by clinical screening if the mutation was also found in that relative. In addition, relatives may have already had clinical screening following the identification of MEN 2 or one of its principal components in an index case.

For histopathology of the thyroid, we recorded presence or absence of C-cell hyperplasia and of MTC. For lymph node status, we recorded presence or absence of metastases at thyroid surgery. Distant metastases were recorded at thyroid surgery or any time during follow up.

For MTC tumor stages, we used the TNM classification of 2002 and the AJCC

Seite 10 von 24

1
2 classification (DeLellis RA, 2004; Greene LF, 2002; Sobin LH, 2002). T1N0M0 for
3
4 MTC<2cm in diameter, T2N0M0 for MTC of 2-4cm in diameter, T3 >4cm either
5
6 limited to the thyroid or with minimal invasion to the sternocleidomastoid muscle or
7
8 perithyroidal soft tissue, T4a any size with invasion to subcutaneous tissue, larynx,
9
10 trachea, esophagus or recurrent laryngeal nerve, T4b invasion of prevertebral fascia,
11
12 carotid artery or mediastinal vessels, N1 for lymph node metastases and M1 for
13
14 distant metastases. In cases of persistent or clearly elevated calcitonin levels after
15
16 thyroidectomy including regional lymph node dissection, we classified 6 patients with
17
18 distant metastases, although radiological evidence was not present.

19
20
21 This study has been approved by the respective institutional review boards for
22
23 human subjects' protection in accordance with the ethical standards of each country
24
25 and center.

26
27
28
29
30 **Statistical Analysis**

31
32
33 Categorical variables were summarized as frequency counts and percentages.
34
35 Continuous variables were summarized as the median and range. MTC stage,
36
37 cure/no-cure, and presence/absence of metastasis were compared among codons.
38
39 Comparisons were made with the Chi-square test which determines, if variables differ
40
41 between at least two of the four codons; and with the Cochran-Mantel-Haenszel
42
43 mean score test (TNM stage) or Cochran-Armitage trend test (cure, metastasis).
44
45 These trend tests account for the ordering of codons by distance from the
46
47 transmembrane domain and assess trends in stage, cure or metastasis. Risk factors
48
49 for MTC were assessed via Chi-square tests and logistic regression analysis. Three
50
51 candidate MTC risk factors were assessed: gender, age, and codon. All three

variables were included in a multivariate logistic regression model whether or not they were statistically significant. Age-dependent penetrance estimates of MTC, pheochromocytoma and HPT were calculated using the Kaplan-Meier method. Penetrance estimates were compared among codons using the log-rank test and with Cox proportional hazards analysis. The log-rank analysis assesses penetrance differences among codons, while the Cox analysis assesses trends across codons. Some patients had incomplete data regarding the different manifestations and were excluded from penetrance estimates. Thus, the penetrance data for MTC are based on 340 subjects, pheochromocytoma 319 subjects, and HPT 299 subjects. All analyses were done using SAS® software (SAS® Institute, Inc., Cary, NC, USA). All statistical tests were two-sided, and $P \leq 0.05$ was used to indicate statistical significance.

Results

Overall, 340 individuals, 199 females and 141 males, had germline mutations in exon 10 of the *RET* gene and their clinical phenotypes and penetrance were analyzed for purposes of this study. The mutations affected only cysteine codons 609, 611, 618 and 620 (Table 1). The 340 subjects, 255 European and 85 South American (comprising 280 of Caucasian, 20 of Hispanic, 3 of Italian-German-Hispanic, 37 of Indian-Caucasian and 1 of Gypsy origin), belonged to 103 different families and hailed from 14 different countries (Table 1). Each of the centers contributed from 1 to 43 cases. The center specific distributions of cases amongst the codons is shown in Table 2.

Deleted: 4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Codon 609 mutations were found in 45 subjects from 13 families. Codon 611 mutations were noted in 50 subjects from 12 families. Codon 618 mutations were found in 94 subjects from 37 families. Codon 620 mutations were identified in 151 subjects from 41 families (Table 1).

Phenotypic Spectrum of *RET* Exon 10 Mutations

MTC was the most common neoplasia, with 263 affected (77%). In the remaining 23%, diagnosis of MTC was excluded by objective means. The median age at diagnosis of MTC was 35 (range 4-86). Of note, symptomatic ascertainment of MTC occurred in 47% of the 263 patients whereas in 53%, their MTC was detected by screening.

Pheochromocytoma was documented in 54 of 319 subjects (17%). The median age at diagnosis of pheochromocytoma was 42 years (17-80). Pheochromocytoma was observed in individuals with mutations of all four codons. However, pheochromocytoma was not observed in carriers of 7 of the 22 distinct mutations. The frequencies of pheochromocytoma varied, occurring in 26% of those patients with codon 609 mutations, 10% codon 611, 23% codon 618 and 13% codon 620 (Table 1). All pheochromocytomas were adrenal in location. Bilateral pheochromocytoma was present in 15 of 54 (28%) affected subjects. None had malignant pheochromocytoma. Symptomatic pheochromocytoma occurred in 29 (54%) of the 54 patients; 24 of these 29 also had MTC. Among these 24, pheochromocytoma was diagnosed at least one year prior to MTC in 6 patients (25%), within 1 year of MTC in 8 (33%), and at least one year after MTC in the remaining 10 (42%).

HPT was present in 8 of 299 subjects (2.7%). Median age at diagnosis of HPT was 46 years (range 28-82). Of the 8 with HPT, 7 were ascertained by screening at

an asymptomatic stage. Although sample size is small, HPT was found in 4 out of 22 distinct mutations residing in all four exon 10 codons.

Regarding codon-specific neoplasias seen in our MEN 2 series, MTC and pheochromocytoma were present in carriers of germline mutations of each of the four codons. Notably, codon 609 mutation carriers as a group have similar frequencies of MTC and pheochromocytoma as the groups carrying mutations in codons 611, 618 and 620 ($p=0.2$, Fisher 2-tailed exact test). However, if we analyzed the frequencies of MTC and pheochromocytoma for each distinct mutation, FMTC was found in patients carrying 6 specific missense mutations distributed across all 4 codons (Table 1), although sample sizes per nucleotide change are small. In contrast, none of the 340 subjects had signs of MEN 2B.

Age-related Penetrance for *RET* Exon 10 Mutations

Age-related penetrance for each MEN 2 component neoplasia was estimated for all the subjects as well as for each of the 4 cysteine codons. Overall for all exon 10 mutations, considering both symptomatic and screening ascertained cases, 50% penetrance was achieved by the age of 36 years for MTC, by 68 years for pheochromocytoma, and by 82 years for HPT (Fig. 1).

Age-related penetrance of MTC for mutations in codons 609, 611, 618, and 620 is shown in detail in table 3. Fifty % penetrance was achieved by the ages of 40, 44, 35, and 34 years for mutations in codons 609, 611, 618 and 620, respectively. Penetrance curves were significantly different among the four codons ($p=0.018$, log-rank test; Fig. 2), and there was also a significant trend across the four codons ($p=0.010$, Cox analysis) whereas within each of the affected codons, penetrance for MTC was not different for the given mutations (data not shown). The latter may be due to small sample sizes due to subset analysis.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We also compared age-related penetrance between symptomatic and screening ascertainment for MTC and found they were not different ($p=0.90$). The ages at 50% penetrance for MTC was 36 years among those symptomatically ascertained and 35 years among those ascertained by screening. Similarly, the age-related penetrance at age 50 years was 83% and 76%, respectively.

For 319 subjects in whom pheochromocytoma was assessed, age-related penetrance at 50 years was 23% (Fig. 1 and Table 3). Penetrance curves were nearly different among codons ($p=0.08$, log-rank test; Fig. 3), but there was no trend in pheochromocytoma penetrance across the four codons ($p=0.99$, Cox analysis). Pheochromocytoma ascertainment was only known in 156 patients; however, penetrance was significantly different between pheochromocytoma ascertained from symptomatic assessment and by screening ($p<0.001$; Fig. 4). Fifty % penetrance was seen by 40 years for 29 symptomatic patients, and by 65 years for 127 screened patients, while penetrance by age 50 was 76% and 23%, respectively.

MTC Tumor Stage

TNM classification was recorded for 300 subjects. Age at diagnosis of any MTC is summarized by codon in table 1. The youngest age at manifestation of MTC in these patient groups is 4, 14, 5, and 6 years for codons 609, 611, 618, and 620, respectively. The youngest age having lymph node metastasis was 21, 29, 9, and 16 years for the respective codons (Table 3). Penetrance for localized MTC was relatively high (67.3%) by the age of 60 for all exon 10 mutations with the highest penetrance for regional nodal involvement and distant metastases for codon 620 mutations (Table 3). Penetrance differed significantly among codons ($p=0.005$, log-rank test), and there was also a trend across the four codons ($p=0.003$, Cox analysis). In addition penetrance for the very early stage of C cell hyperplasia (CCH),

as well for distant metastases (M1) are shown in table 3. CCH penetrance was not different among codons ($p=0.16$, log-rank test), nor was there a trend across codons ($p=0.10$, Cox analysis). M1 penetrance was not quite different among codons ($p=0.12$, log-rank test), but there was a trend across codons ($p=0.020$, Cox analysis). [Yes, this is fine to be added here as well]

Comparing MTC stages, i.e. metastasized or not metastasized and cured or not cured among codons using the Chi-square test showed a significant difference among codons (Table 4). When ordered by codons, 609, 611, 618 and 620, higher stages ($p<0.001$), increased the likelihood of metastasis ($p=0.002$), and decreased the chance of cure ($p<0.001$). Death caused by MTC was documented in 0, 1, 6, and 11 patients in the codons 609, 611, 618, and 620, respectively.

Gender, age, and mutated codon were examined as possible risk factors for MTC. In univariable Chi-square analysis, female gender ($p=0.008$) and older age ($p<0.001$) were associated with higher risk of MTC. Specific codon mutated was not quite significant ($p=0.06$); there appeared to be no trend across codons, but rather codon 618 mutation had the highest risk of MTC (76% MTC for codon 609, 72% for 611, 87% for 618, and 74% for 620). In multivariable analysis, gender becomes non-significant, age remains significant ($p<0.001$), and codon is not quite significant, but again, codon 618 has the highest risk of MTC ($p=0.07$).

Discussion

Because the prevalence of *RET* exon 10 mutations, even all together, is much lower than that of the exon 11 codon 634 mutations, information on the natural history and age-related penetrance for component neoplasias remained scarce and was provided mostly by reports from single centers or on single large families (Calva, et al., 2009; Mian, et al., 2009; Moers, et al., 1996; Siggelkow, et al., 2001). The

Deleted: Lisas suggestion for the sentence highlighted in blue: Penetrance differed significantly among codons ($p=0.005$, log-rank test), and there was also a trend across the four codons ($p=0.003$, Cox analysis). In addition penetrance for the very early stage of C cell hyperplasia (CCH), as well for distant metastases (M1) are shown in table 3. CCH penetrance was not different among codons ($p=0.16$, log-rank test), nor was there a trend across codons ($p=0.10$, Cox analysis). M1 penetrance was not quite different among codons ($p=0.12$, log-rank test), but there was a trend across codons ($p=0.020$, Cox analysis).¶

Deleted: 2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

American Thyroid Association (ATA) has, in particular, addressed this issue as a point of concern. Guidelines for diagnosis and management of MTC and other components of MEN 2 have reached only a weak consensus level for these management guidelines for carriers of mutations in exon 10 (Kloos, et al., 2009). In this present study, we summarize and analyze comprehensive data for subjects carrying such *RET* exon 10 mutations based on a multicenter, multiethnic and multinational series of 340 individuals who all have been registered using identical criteria. Thus, we are able to obtain molecular-based neoplasia risk profiles and codon-specific age-related penetrance on which to base genetic counseling and medical decision-making.

Our current series demonstrates that 15 distinct mutations involving the 4 codons of *RET* exon 10 can all be associated with MEN 2A. In contrast, the FMTC phenotype was seen to be associated with 6 distinct mutations, also involving all the 4 codons (Table 1). The studied number of carriers with these 7 mutations ranges from 2 to 26 (mean 11) per mutation. Therefore, the lack of pheochromocytoma in these subjects may be due to sample size and/or penetrance.

Missense mutations nearer the N-terminus are associated more with HSCR based on a clear mechanism (Santoro, et al., 1999). The first comprehensive genotype-phenotype study on MEN 2 suggested that as high as 1/3 of all carriers of mutations in codons 618 and 620 had HSCR (Eng C, 1996). However, the sample sizes for exon 10 mutations in that series was relatively small. In even smaller series and single family reports, the prevalence of HSCR in codon 620 mutation carriers approached 50% (Butter, et al., 2007). In the present series, HSCR was only present in 20/267 informative carriers (7.5%), 2/37 (5.4%) in those with mutations in codon 609, 0/44 in codon 611, 1/56 (1.8%) in codon 618, and 17/130 (13.1%) in codon 620 (data not shown). This is a lower prevalence than initially predicted, however, it is

tempting to speculate if other intra-*RET* modifying variants, whether in cis or trans, would modulate the HSCR phenotype in such individuals (Borrego, et al., 1998; Borrego, et al., 2000; Borrego, et al., 1999; Borrego, et al., 2003; Fernandez, et al., 2003).

Even before *RET* was identified, the classic penetrance of MTC in MEN 2A is cited as 70% by age 70 by symptomatic presentation and 100% by age 70 by stimulated calcitonin screening (Ponder, et al., 1988a; Ponder, et al., 1988b). Not surprisingly, age-related penetrance for codon 634 mutations approach >85% by age 80 for MTC and >50% by age 80 for pheochromocytoma (Iihara, et al., 1997; Kloos, et al., 2009; Milos, et al., 2008; Modigliani, et al., 1998). Mutations in exon 10 were believed to have much lower age-related penetrances. While our series showed that penetrance for MTC was 50% at age 36 for exon 10 mutations which is lower than the 50% by age 28 for the C634W mutation (Milos, et al., 2008), both exon 10 and 11 mutations reach >80% penetrance by age 50 and almost 100% by age 70. Interestingly, the penetrance for distant metastases from MTC was 9% for p.C634W compared to 36% for codon 620 mutations (Table 3). In contrast, the age-related penetrance of exon 10 mutations for pheochromocytomas was systematically lower than that for codon 634 mutations, such as for p.C634W (33% vs 70% by age 60, Table 3). Notably, there is a significant difference in pheochromocytoma penetrance between those that were symptomatically ascertained compared to those ascertained by screening. This underscores the great importance of clinical screening for pheochromocytoma even in exon 10 mutation carriers, beginning in the late teens or early 20s. Our data support the ATA guidelines recommending initiation of screening for pheochromocytoma at the age of 20 in all mutations except those in codons 918 and 634 respectively; for these mutations, screening for pheochromocytoma is recommended beginning by the age of 8 years. Although sample size was small, the

Field Code Changed

Formatted: German Germany

Formatted: German Germany

1
2 penetrance for HPT was 34% by age 60 for p.C634W compared to 3.9% by age 60
3
4 for exon 10 mutations (Table 3).
5

6 The results of this multicenter study of 340 patients carrying exon 10
7
8 mutations show differences in aggressiveness of MTC depending on the mutated
9
10 codon. This results in earlier age at manifestation, more advanced tumor stage at
11
12 manifestation and decreased chance of cure when ordered by codons (620, 618,
13
14 611, and 609). In this patient group, age is an independent risk factor for the
15
16 manifestation of MTC. Position of mutation was associated with metastatic disease,
17
18 with the lowest likelihood amongst those with codon 609 mutations, and the highest,
19
20 codon 620 mutations. This is important for the decisions regarding timing of
21
22 premorbid diagnoses and treatment/prophylactic procedure in *RET* gene carriers, as
23
24 age is an independent predictor of aggressive disease at thyroidectomy: MTC is
25
26 more aggressive and might develop earlier in codons 618 and 620. This supports the
27
28 clinical impression that patients with *RET* mutations in codons 609 and 611 are older
29
30 at presentation (Calva, et al., 2009; Hansen, et al., 2000; Mian, et al., 2009;
31
32 Siggelkow, et al., 2001), and have more indolent tumors than patients with *RET*
33
34 mutations at 618 and 620 (Machens, et al., 2009), but there are notable exceptions
35
36 as shown in our systematic study. It is important to point out that also a patient with
37
38 codons 609 mutation can present with MTC by age 4.

39 Multivariate analysis of our data reveals that age at presentation is associated
40
41 with higher likelihood of metastases from MTC, a finding that is not surprising. What
42
43 is also not surprising is that those who were ascertained symptomatically present at
44
45 more advanced stages compared to those ascertained by screening, where a full 1/3
46
47 presents with normal histology or CCH only. Taken together, our observations
48
49 strongly support clinical surveillance and/or timely prophylactic thyroidectomy even in
50
51 exon 10 mutation carriers, which is certainly notated in the ATA MTC Practice
52

Guidelines. However, the age to begin these remain unclear as the impression up until recently is exon 10 mutations do not normally confer early ages of onset of MTC. In our current series, however, the youngest age at diagnosis for MTC was 4 years (in a codon 609 mutation carrier) and for MTC with regional nodal metastases 9 years (in a codon 618 mutation carrier), respectively. Not surprisingly, therefore, those with codons 618 and 620 mutations had the highest prevalence of metastatic MTC, even higher than those with p.C634W. The youngest age for distant metastases amongst exon 10 mutation positive individuals was aged 13 years. The more typical age of MTC development for the exon 10 genotype reveals significant differences in aggressiveness of MTC manifestation e.g. higher penetrance of lymph node metastases and distant metastases in codons 618/620 in comparison to exon 609/611. The suggestion of the ATA guidelines to consider prophylactic surgery in patients carrying exon 10 codon 618 and 620 mutations at age of 5 years seems to be supported by our data. In patients with codon 609 and 611 prophylactic thyroidectomy possibly might be postponed under careful yearly clinical and laboratory examinations.

In summary, we have presented a large series of uniformly collected and registered cases with germline exon 10 mutations in *RET*, and have analyzed neoplastic risk profiles and age-related penetrance for each component neoplasia, which will help evidence-based genetic counseling and medical management.

Word count: 3844

References

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[Borrego S, Eng C, Sanchez B, Saez ME, Navarro E, Antinolo G. 1998. Molecular analysis of the ret and GDNF genes in a family with multiple endocrine neoplasia type 2A and Hirschsprung disease. J Clin Endocrinol Metab 83\(9\):3361-4.](#)

[Borrego S, Ruiz A, Saez ME, Gimm O, Gao X, Lopez-Alonso M, Hernandez A, Wright FA, Antinolo G, Eng C. 2000. RET genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease. J Med Genet 37\(8\):572-8.](#)

[Borrego S, Saez ME, Ruiz A, Gimm O, Lopez-Alonso M, Antinolo G, Eng C. 1999. Specific polymorphisms in the RET proto-oncogene are over-represented in patients with Hirschsprung disease and may represent loci modifying phenotypic expression. J Med Genet 36\(10\):771-4.](#)

[Borrego S, Wright FA, Fernandez RM, Williams N, Lopez-Alonso M, Davuluri R, Antinolo G, Eng C. 2003. A founding locus within the RET proto-oncogene may account for a large proportion of apparently sporadic Hirschsprung disease and a subset of cases of sporadic medullary thyroid carcinoma. Am J Hum Genet 72\(1\):88-100.](#)

[Brandi ML, Gagel RF, Angeli A, Bilezikian JP, Beck-Peccoz P, Bordi C, Conte-Devolx B, Falchetti A, Gheri RG, Libroia A and others. 2001. Guidelines for diagnosis and therapy of MEN type 1 and type 2. J Clin Endocrinol Metab 86\(12\):5658-71.](#)

[Butter A, Gagne J, Al-Jazaeri A, Emran MA, Deal C, St-Vil D. 2007. Prophylactic thyroidectomy in pediatric carriers of multiple endocrine neoplasia type 2A or familial medullary thyroid carcinoma: mutation in C620 is associated with Hirschsprung's disease. J Pediatr Surg 42\(1\):203-6.](#)

[Calva D, O'Dorisio TM, Sue O'Dorisio M, Lal G, Sugg S, Weigel RJ, Howe JR. 2009. When is prophylactic thyroidectomy indicated for patients with the RET codon 609 mutation? Ann Surg Oncol 16\(8\):2237-44.](#)

[DeLellis RA LR, Heitz PU, Eng C \(eds\). . 2004. World Healthe Organization Classifican of Tumours. Pathology and genetics of tumours of endocrine organs.. IARC Press: Lyon](#)

[Elisei R, Romei C, Cosci B, Agate L, Bottici V, Molinaro E, Sculli M, Miccoli P, Basolo F, Grasso L and others. 2007. RET genetic screening in patients with medullary thyroid cancer and their relatives: experience with 807 individuals at one center. J Clin Endocrinol Metab 92\(12\):4725-9.](#)

[Eng C CD, Schuffenecker I, Lenoir G, Cote G, Gagel RF, van Amstel HK, Lips CJ, Nishisho I, Takai SI, Marsh DJ, Robinson BG, Frank-Raue K, Raue F, Xue F, Noll WW, Romei C, Pacini F, Fink M, Niederle B, Zedenius J, Nordenskjold M, Komminoth P, Hendy GN, Mulligan LM & et al. . 1996 The relationship between specific RET proto-oncogene mutations and disease phenotype in multiple endocrine neoplasia type 2. International RET mutation consortium analysis. JAMA 276:1575-1579.](#)

[Erlic Z, Rybicki L, Peczkowska M, Golcher H, Kann PH, Brauckhoff M, Mussig K, Muresan M, Schaffler A, Reisch N and others. 2009. Clinical predictors and algorithm for the genetic diagnosis of pheochromocytoma patients. Clin Cancer Res 15\(20\):6378-85.](#)

[Fernandez RM, Antinolo G, Eng C, Borrego S. 2003. The RET C620S mutation causes multiple endocrine neoplasia type 2A \(MEN2A\) but not Hirschsprung disease \(HSCR\) in a family cosegregating both phenotypes. Hum Mutat 22\(5\):412-5.](#)

[Frank-Raue K, Buhr H, Dralle H, Klar E, Senninger N, Weber T, Rondot S, Hoppner W, Raue F. 2006. Long-term outcome in 46 gene carriers of hereditary medullary thyroid carcinoma after prophylactic thyroidectomy: impact of individual RET genotype. Eur J Endocrinol 155\(2\):229-36.](#)

[Greene LF PD, Fritz DA, Batch M, Haller DG, Morrow M.. 2002. American Joint Committee on Cancer \(AJCC\). Cancer Staging Manual 6th edition](#)

[Hansen HS, Topping H, Godballe C, Jager AC, Nielsen FC. 2000. Is thyroidectomy necessary in RET mutations carriers of the familial medullary thyroid carcinoma syndrome? Cancer 89\(4\):863-7.](#)

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt, German Germany

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt

- liihara M, Yamashita T, Okamoto T, Kanbe M, Yamazaki K, Egawa S, Yamaguchi K, Obara T. 1997. A nationwide clinical survey of patients with multiple endocrine neoplasia type 2 and familial medullary thyroid carcinoma in Japan. *Jpn J Clin Oncol* 27(3):128-34.
- Kloos RT, Eng C, Evans DB, Francis GL, Gagel RF, Gharib H, Moley JF, Pacini F, Ringel MD, Schlumberger M and others. 2009. Medullary thyroid cancer: management guidelines of the American Thyroid Association. *Thyroid* 19(6):565-612.
- Machens A, Hauptmann S, Dralle H. 2009. Modification of multiple endocrine neoplasia 2A phenotype by cell membrane proximity of RET mutations in exon 10. *Endocr Relat Cancer* 16(1):171-7.
- Mian C, Barollo S, Zamboni L, Pennelli G, Bernante P, Pelizzo MR, Nacamulli D, Mantero F, Girelli ME, Opocher G. 2009. Characterization of the largest kindred with MEN2A due to a Cys609Ser RET mutation. *Fam Cancer* 8(4):379-82.
- Milos IN, Frank-Raue K, Wohllk N, Maia AL, Pusiol E, Patocs A, Robledo M, Biarnes J, Barontini M, Links TP and others. 2008. Age-related neoplastic risk profiles and penetrance estimations in multiple endocrine neoplasia type 2A caused by germ line RET Cys634Trp (TGC>TGG) mutation. *Endocr Relat Cancer* 15(4):1035-41.
- Modigliani E, Cohen R, Campos JM, Conte-Devolx B, Maes B, Boneu A, Schlumberger M, Bigorgne JC, Dumontier P, Leclerc L and others. 1998. Prognostic factors for survival and for biochemical cure in medullary thyroid carcinoma: results in 899 patients. The GETC Study Group. *Groupe d'etude des tumeurs a calcitonine. Clin Endocrinol (Oxf)* 48(3):265-73.
- Moers AM, Landsvater RM, Schaap C, Jansen-Schillhorn van Veen JM, de Valk IA, Blijham GH, Hoppener JW, Vroom TM, van Amstel HK, Lips CJ. 1996. Familial medullary thyroid carcinoma: not a distinct entity? Genotype-phenotype correlation in a large family. *Am J Med* 101(6):635-41.
- Neumann HP, Bausch B, McWhinney SR, Bender BU, Gimm O, Franke G, Schipper J, Klisch J, Althoefer C, Zerres K and others. 2002. Germ-line mutations in nonsyndromic pheochromocytoma. *N Engl J Med* 346(19):1459-66.
- Neumann HP, Pawlu C, Peczkowska M, Bausch B, McWhinney SR, Muresan M, Buchta M, Franke G, Klisch J, Bley TA and others. 2004. Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations. *Jama* 292(8):943-51.
- Neumann HP, Vortmeyer A, Schmidt D, Werner M, Erlic Z, Cascon A, Bausch B, Januszewicz A, Eng C. 2007. Evidence of MEN-2 in the original description of classic pheochromocytoma. *N Engl J Med* 357(13):1311-5.
- Ponder BA, Finer N, Coffey R, Harmer CL, Maisey M, Ormerod MG, Pembrey ME, Ponder MA, Rosswick P, Shalet S. 1988a. Family screening in medullary thyroid carcinoma presenting without a family history. *Q J Med* 67(252):299-308.
- Ponder BA, Ponder MA, Coffey R, Pembrey ME, Gagel RF, Telenius-Berg M, Semple P, Easton DF. 1988b. Risk estimation and screening in families of patients with medullary thyroid carcinoma. *Lancet* 1(8582):397-401.
- Santoro M, Carlomagno F, Melillo RM, Billaud M, Vecchio G, Fusco A. 1999. Molecular mechanisms of RET activation in human neoplasia. *J Endocrinol Invest* 22(10):811-9.
- Schiavi F, Boedeker CC, Bausch B, Peczkowska M, Gomez CF, Strassburg T, Pawlu C, Buchta M, Salzmann M, Hoffmann MM and others. 2005. Predictors and prevalence of paraganglioma syndrome associated with mutations of the SDHC gene. *Jama* 294(16):2057-63.
- Siggelkow H, Melzer A, Nolte W, Karsten K, Hoppner W, Hufner M. 2001. Presentation of a kindred with familial medullary thyroid carcinoma and Cys611Phe mutation of the RET proto-oncogene demonstrating low grade malignancy. *Eur J Endocrinol* 144(5):467-73.
- Sobin LH WC. 2002. TNM classification of malignant tumours. 6th edition Wiley, John and sons. Hoboken
- Zbuk KM, Eng C. 2007. Cancer phenomics: RET and PTEN as illustrative models. *Nat Rev Cancer* 7(1):35-45.

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt, German Germany

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt, German Germany

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt, German Germany

Formatted: Font: (Default) Calibri, 11 pt, Complex Script Font: Calibri, 11 pt

Formatted: Indent: Before: 0 pt, First line: 0 pt

Figure Legends

Figure 1. Penetrance of medullary thyroid carcinoma (MTC), pheochromocytoma (Pheo) and hyperparathyroidism (HPT) in 340 carriers of *RET* exon 10 germline mutations.

Figure 2. Penetrance of medullary thyroid carcinoma (MTC) of 340 carriers adjusted for germline mutations of the codons 609 (n=45 carriers), 611 (n=50 carriers), 618 (n=94 carriers), and 620 (n=151 carriers).

Figure 3. Penetrance of pheochromocytoma (Pheo) of 319 carriers adjusted for germline mutations of the codons 609 (n=43 carriers), 611 (n=50 carriers) 618 (n=84 carriers), and 620 (n=142 carriers).

Figure 4. Penetrance of pheochromocytoma (Pheo) adjusted for detection by symptomatic pheochromocytoma (Symptomatic) and by screening-detected pheochromocytoma (Screening).

Deleted: Borrego S, Eng C, Sanchez B, Saez ME, Navarro E, Antinolo G. 1998. Molecular analysis of the ret and GDNF genes in a family with multiple endocrine neoplasia type 2A and Hirschsprung disease. *J Clin Endocrinol Metab* 83(9):3361-4.¶

Borrego S, Ruiz A, Saez ME, Gimm O, Gao X, Lopez-Alonso M, Hernandez A, Wright FA, Antinolo G, Eng C. 2000. RET genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease. *J Med Genet* 37(8):572-8.¶

Borrego S, Saez ME, Ruiz A, Gimm O, Lopez-Alonso M, Antinolo G, Eng C. 1999. Specific polymorphisms in the RET proto-oncogene are over-represented in patients with Hirschsprung disease and may represent loci modifying phenotypic expression. *J Med Genet* 36(10):771-4.¶

Borrego S, Wright FA, Fernandez RM, Williams N, Lopez-Alonso M, Davuluri R, Antinolo G, Eng C. 2003. A founding locus within the RET proto-oncogene may account for a large proportion of apparently sporadic Hirschsprung disease and a subset of cases of sporadic medullary thyroid carcinoma. *Am J Hum Genet* 72(1):88-100.¶

Brandi ML, Gagel RF, Angeli A, Bilezikian JP, Beck-Peccoz P, Bordi C, Conte-Devolx B, Falchetti A, Gheri RG, Libroia A and others. 2001. Guidelines for diagnosis and therapy of MEN type 1 and type 2. *J Clin Endocrinol Metab* 86(12):5658-71.¶

Butter A, Gagne J, Al-Jazaeri A, Emran MA, Deal C, St-Vil D. 2007. Prophylactic thyroidectomy in pediatric carriers of multiple endocrine neoplasia type 2A or familial medullary thyroid carcinoma: mutation in C620 is associated with Hirschsprung's disease. *J Pediatr Surg* 42(1):203-6.¶

Calva D, O'Dorisio TM, Sue O'Dorisio M, Lal G, Sugg S, Weigel RJ, Howe JR. 2009. When is prophylactic thyroidectomy indicated for patients with the RET codon 609 mutation? *Ann Surg Oncol* 16(8):2237-44.¶

DeLellis LA, Heitz PU, Eng C (eds). . 2004. World Health Organization Classification of Tumours. Pathology and genetics of tumours of endocrine organs. . IARC Press: Lyon ¶

Elisei R, Romei C, Cosci B, Agate L, Bottici V, Molinaro E, Sculli M, Miccoli P, Basolo F, Grasso L and others. 2007. RET genetic screening in patients with medullary thyroid cancer and their relatives: experience with 807 individuals at one center. *J Clin Endocrinol Metab* 92(12):4725-9.¶

Eng C CD, Schuffenecker I, Lenoir G, Cote G, Gagel RF, van Amstel HK, Lips CJ, Nishisho I, Takai SI, Marsh DJ, Robinson BG, Frank-Raue K, Raue F, Xue F, Noll WW, Romei C, Pacini F, Fink M, N[... [1]

Formatted: German Germany

Formatted: German Germany

Formatted: German Germany

Formatted: Right: 56.7 pt, Bottom: 70.85 pt, Width: 841.9 pt, Height: 595.3 pt

For Peer Review

Page 23: [1] Deleted	User	8/27/2010 9:25:00 AM
Borrego S, Eng C, Sanchez B, Saez ME, Navarro E, Antinolo G. 1998. Molecular analysis of the ret and GDNF genes in a family with multiple endocrine neoplasia type 2A and Hirschsprung disease. J Clin Endocrinol Metab 83(9):3361-4.		
Borrego S, Ruiz A, Saez ME, Gimm O, Gao X, Lopez-Alonso M, Hernandez A, Wright FA, Antinolo G, Eng C. 2000. RET genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease. J Med Genet 37(8):572-8.		
Borrego S, Saez ME, Ruiz A, Gimm O, Lopez-Alonso M, Antinolo G, Eng C. 1999. Specific polymorphisms in the RET proto-oncogene are over-represented in patients with Hirschsprung disease and may represent loci modifying phenotypic expression. J Med Genet 36(10):771-4.		
Borrego S, Wright FA, Fernandez RM, Williams N, Lopez-Alonso M, Davuluri R, Antinolo G, Eng C. 2003. A founding locus within the RET proto-oncogene may account for a large proportion of apparently sporadic Hirschsprung disease and a subset of cases of sporadic medullary thyroid carcinoma. Am J Hum Genet 72(1):88-100.		
Brandi ML, Gagel RF, Angeli A, Bilezikian JP, Beck-Peccoz P, Bordi C, Conte-Devolx B, Falchetti A, Gheri RG, Libroia A and others. 2001. Guidelines for diagnosis and therapy of MEN type 1 and type 2. J Clin Endocrinol Metab 86(12):5658-71.		
Butter A, Gagne J, Al-Jazaeri A, Emran MA, Deal C, St-Vil D. 2007. Prophylactic thyroidectomy in pediatric carriers of multiple endocrine neoplasia type 2A or familial medullary thyroid carcinoma: mutation in C620 is associated with Hirschsprung's disease. J Pediatr Surg 42(1):203-6.		
Calva D, O'Dorisio TM, Sue O'Dorisio M, Lal G, Sugg S, Weigel RJ, Howe JR. 2009. When is prophylactic thyroidectomy indicated for patients with the RET codon 609 mutation? Ann Surg Oncol 16(8):2237-44.		
DeLellis RA LR, Heitz PU, Eng C (eds). . 2004. World Healthe Organization Classifican of Tumours. Pathology and genetics of tumours of endocrine organs. . IARC Press: Lyon		
Elisei R, Romei C, Cosci B, Agate L, Bottici V, Molinaro E, Sculli M, Miccoli P, Basolo F, Grasso L and others. 2007. RET genetic screening in patients with medullary thyroid cancer and their relatives: experience with 807 individuals at one center. J Clin Endocrinol Metab 92(12):4725-9.		
Eng C CD, Schuffenecker I, Lenoir G, Cote G, Gagel RF, van Amstel HK, Lips CJ, Nishisho I, Takai SI, Marsh DJ, Robinson BG, Frank-Raue K, Raue F, Xue F, Noll WW, Romei C, Pacini F, Fink M, Niederle B, Zedenius J, Nordenskjoeld M, Komminoth P, Hendy GN, Mulligan LM & et al. . 1996 The relationship between specific RET proto-oncogene mutations and disease phenotype in multiple endocrine neoplasia type 2. International RET mutation consortium analysis. JAMA 276:1575-1579.		
Erlic Z, Rybicki L, Peczkowska M, Golcher H, Kann PH, Brauckhoff M, Mussig K, Muresan M, Schaffler A, Reisch N and others. 2009. Clinical predictors and algorithm for the genetic diagnosis of pheochromocytoma patients. Clin Cancer Res 15(20):6378-85.		
Fernandez RM, Antinolo G, Eng C, Borrego S. 2003. The RET C620S mutation causes multiple endocrine neoplasia type 2A (MEN2A) but not Hirschsprung disease (HSCR) in a family cosegregating both phenotypes. Hum Mutat 22(5):412-5.		
Frank-Raue K, Buhr H, Dralle H, Klar E, Senninger N, Weber T, Rondot S, Hoppner W, Raue F. 2006. Long-term outcome in 46 gene carriers of hereditary medullary thyroid carcinoma after prophylactic thyroidectomy: impact of individual RET genotype. Eur J Endocrinol 155(2):229-36.		

- Greene LF PD, Fritz DA, Batch M, Haller DG, Morrow M. . 2002. American Joint Committee on Cancer (AJCC). Cancer Staging Manual 6th edition
- Hansen HS, Topping H, Godballe C, Jager AC, Nielsen FC. 2000. Is thyroidectomy necessary in RET mutations carriers of the familial medullary thyroid carcinoma syndrome? *Cancer* 89(4):863-7.
- Iihara M, Yamashita T, Okamoto T, Kanbe M, Yamazaki K, Egawa S, Yamaguchi K, Obara T. 1997. A nationwide clinical survey of patients with multiple endocrine neoplasia type 2 and familial medullary thyroid carcinoma in Japan. *Jpn J Clin Oncol* 27(3):128-34.
- Kloos RT, Eng C, Evans DB, Francis GL, Gagel RF, Gharib H, Moley JF, Pacini F, Ringel MD, Schlumberger M and others. 2009. Medullary thyroid cancer: management guidelines of the American Thyroid Association. *Thyroid* 19(6):565-612.
- Machens A, Hauptmann S, Dralle H. 2009. Modification of multiple endocrine neoplasia 2A phenotype by cell membrane proximity of RET mutations in exon 10. *Endocr Relat Cancer* 16(1):171-7.
- Mian C, Barollo S, Zambonin L, Pennelli G, Bernante P, Pelizzo MR, Nacamulli D, Mantero F, Girelli ME, Opocher G. 2009. Characterization of the largest kindred with MEN2A due to a Cys609Ser RET mutation. *Fam Cancer* 8(4):379-82.
- Milos IN, Frank-Raue K, Wohllk N, Maia AL, Pusiol E, Patocs A, Robledo M, Biarnes J, Barontini M, Links TP and others. 2008. Age-related neoplastic risk profiles and penetrance estimations in multiple endocrine neoplasia type 2A caused by germ line RET Cys634Trp (TGC>TGG) mutation. *Endocr Relat Cancer* 15(4):1035-41.
- Modigliani E, Cohen R, Campos JM, Conte-Devolx B, Maes B, Boneu A, Schlumberger M, Bigorgne JC, Dumontier P, Leclerc L and others. 1998. Prognostic factors for survival and for biochemical cure in medullary thyroid carcinoma: results in 899 patients. The GETC Study Group. Groupe d'etude des tumeurs a calcitonine. *Clin Endocrinol (Oxf)* 48(3):265-73.
- Moers AM, Landsvater RM, Schaap C, Jansen-Schillhorn van Veen JM, de Valk IA, Blijham GH, Hoppener JW, Vroom TM, van Amstel HK, Lips CJ. 1996. Familial medullary thyroid carcinoma: not a distinct entity? Genotype-phenotype correlation in a large family. *Am J Med* 101(6):635-41.
- Neumann HP, Bausch B, McWhinney SR, Bender BU, Gimm O, Franke G, Schipper J, Klisch J, Althoefer C, Zerres K and others. 2002. Germ-line mutations in nonsyndromic pheochromocytoma. *N Engl J Med* 346(19):1459-66.
- Neumann HP, Pawlu C, Peczkowska M, Bausch B, McWhinney SR, Muresan M, Buchta M, Franke G, Klisch J, Bley TA and others. 2004. Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations. *Jama* 292(8):943-51.
- Neumann HP, Vortmeyer A, Schmidt D, Werner M, Erlic Z, Cascon A, Bausch B, Januszewicz A, Eng C. 2007. Evidence of MEN-2 in the original description of classic pheochromocytoma. *N Engl J Med* 357(13):1311-5.
- Ponder BA, Finer N, Coffey R, Harmer CL, Maisiey M, Ormerod MG, Pembrey ME, Ponder MA, Rosswick P, Shalet S. 1988a. Family screening in medullary thyroid carcinoma presenting without a family history. *Q J Med* 67(252):299-308.
- Ponder BA, Ponder MA, Coffey R, Pembrey ME, Gagel RF, Telenius-Berg M, Semple P, Easton DF. 1988b. Risk estimation and screening in families of patients with medullary thyroid carcinoma. *Lancet* 1(8582):397-401.
- Santoro M, Carlomagno F, Melillo RM, Billaud M, Vecchio G, Fusco A. 1999. Molecular mechanisms of RET activation in human neoplasia. *J Endocrinol Invest* 22(10):811-9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Schiavi F, Boedeker CC, Bausch B, Peczkowska M, Gomez CF, Strassburg T, Pawlu C, Buchta M, Salzmann M, Hoffmann MM and others. 2005. Predictors and prevalence of paraganglioma syndrome associated with mutations of the SDHC gene. *Jama* 294(16):2057-63.

Siggelkow H, Melzer A, Nolte W, Karsten K, Hoppner W, Hufner M. 2001. Presentation of a kindred with familial medullary thyroid carcinoma and Cys611Phe mutation of the RET proto-oncogene demonstrating low grade malignancy. *Eur J Endocrinol* 144(5):467-73.

Sobin LH WC. 2002. TNM classification of malignant tumours. 6th edition Wiley, John and sons. Hoboken

Zbuk KM, Eng C. 2007. Cancer phenomics: RET and PTEN as illustrative models. *Nat Rev Cancer* 7(1):35-45.

For Peer Review

177x127mm (300 x 300 DPI)

177x127mm (300 x 300 DPI)

177x127mm (300 x 300 DPI)

177x127mm (300 x 300 DPI)

Table 1 Demographic and molecular data of the study population

Codon	Nucleotide	Protein	Families n	Carriers n	Gender f/m	Age at MTC diagnosis: median (range)	MTC n (%)	Pheo n (%)	HPT n (%)	HSCR n (%)	Phenotype	Age at N1M0 n (range)	Age at M1 n (range)
609	c.1825T>C	C609R	2	7	7/0	42 (29-43)	4/7 (57)	2/7 (29)	0	0	MEN2A	1 (40)	0
	c.1825T>G	C609G	3	6	1/5	48 (4-54)	5/6 (83)	4/5 (80)	0	0	MEN2A	2 (33-48)	0
	c.1826G>A	C609Y	4	8	7/1	33 (21-49)	7/8 (88)	0	0	2/8 (25)	FMTC,HSRC	2 (21-48)	0
	c.1826G>C	C609S	2	18	6/12	37 (15-86)	14/18 (78)	4/17 (24)	2/17 (12)	0	MEN2A	1 (38)	1 (62)
	c.1826G>T	C609F	2	6	5/1	45 (36-63)	4/6 (67)	1/6 (17)	0	0	MEN2A	1 (54)	0
	All		13	45	26/19	37 (4-86)	34/45 (76)	11/43 (26)	2/44 (5)	2/37 (5)	MEN2A,HSRC	7 (21-54)	1 (62)
611	c.1832G>A	C611Y	7	15	11/4	35 (15-65)	9/15 (60)	4/15 (27)	1/15 (7)	0	MEN2A	4 (29-64)	1 (65)
	c.1833C>G	C611W	2	5	4/1	37 (14-66)	5/5 (100)	1/5 (20)	0	0	MEN2A	1 (49)	1 (37)
	c.1832G>T	C611F	2	26	12/14	44 (25-69)	19/26 (73)	0	0	0	FMTC	1 (44)	5 (27-69)
	c.1832G>T c.1833C>T	C611F	1	4	3/1	53 (36-64)	3/4 (75)	0	0	0	FMTC	0	0
	All		12	50	30/20	42 (14-69)	36/50 (72)	5/50 (10)	1/44 (2)	0	MEN2A	6 (29-64)	7 (27-69)
618	c.1852T>A	C618S	4	13	8/5	35 (18-59)	10/13 (77)	3/12 (25)	0	0	MEN2A	2 (42-59)	2 (31-39)
	c.1852T>C	C618R	17	28	22/6	35 (10-65)	28/28 (100)	11/28 (39)	0	1/12 (8)	MEN2A	11 (21-45)	7 (13-65)
	c.1852T>G	C618G	3	6	4/2	30 (21-57)	5/6 (83)	0	0	0	FMTC	0	1 (57)
	c.1853G>C	C618S	7	14	4/10	27 (9-63)	12/14 (86)	2/13 (15)	0	0	MEN2A	4 (9-50)	0

	c.1853G>A	C618Y	2	6	4/2	33 (25-55)	6/6 (100)	1/3 (33)	0	0	MEN2A	0	0
	c.1853G>T	C618F	4	27	9/18	45 (5-72)	21/27 (78)	2/21 (10)	2/21 (10)	0	MEN2A	3 (43-51)	3 (40-72)
	all		37	94	51/43	35 (5-72)	82/94 (87)	19/84 (23)	2/78 (3)	1/56 (2)	MEN2A	20 (9-59)	13 (13-72)
620	c.1858T>C	C620R	24	101	57/44	29 (6-73)	79/101 (78)	15/98 (15)	3/92 (3)	12/86 (14)	MEN2A,HSCR	27 (16-73)	20 (18-64)
	c.1858T>G	C620G	1	1	0/1	22	7/13 (54)	1/1 (100)	0	4/12 (33)	MEN2A	0	0
	c.1859G>A	C620Y	7	19	14/5	36 (18-76)	14/19 (74)	0	0	1/15 (7)	FMTC,HSCR	6 (31-60)	0
	c.1859G>C	C620S	4	20	9/11	40 (14-73)	14/20 (70)	3/20 (15)	0	0	MEN2A,HSCR	5 (28-73)	0
	c.1860C>G	C620W	1	2	2/0	40 (40)	1/2 (50)	0	0	0	FMTC	0	0
	c.1859G>T	C620F	4	8	7/1	35 (27-58)	3/8 (38)	2/7 (29)	0	0	MEN2A	1 (58)	0
	all		41	151	92/59	31 (6-76)	111/151 (74)	19/142 (13)	3/133 (2)	17/130 (13)	MEN2A,HSCR	39 (16-73)	20 (18-64)
All	all		104	340	199/141	35 (4-86)	263/340 (77)	54/319 (17)	8/299 (3)	20/267 (7)	MEN2A,HSCR	72 (9-73)	41 (13-72)

Table 2
Cases (families) with mutation per exon

Center	Total of cases	Exon 609	Exon 611	Exon 618	Exon 620
Athens	27			4 (3)	23 (6)
Belgrade	5			2 (2)	3 (2)
Budapest	8	7 (2)	1 (1)		
Buenos Aires	16		6 (2)	6 (3)	4 (2)
Erlangen	2	2 (1)			
Frankfurt	10		10 (3)		
Freiburg	10			9 (2)	1 (1)
Girona	5	5 (1)			
Göttingen	24		24 (1)		
Groningen	20	1 (1)		16 (1)	3 (1)
Heidelberg	32	2 (1)		17 (11)	13 (8)
Innsbruck	1			1 (1)	
Leipzig	7	6 (1)			1 (1)
Mainz	8			8 (3)	
Nancy	3		2 (1)	1 (1)	
Nürnberg	1	1 (1)			
Padova	43	14 (1)		10 (4)	19 (2)
Porto Allegre	9			9 (3)	
Prague	13	2 (1)	2 (1)		9 (3)
Santiago de Chile	14			2 (1)	12 (2)
Sao Paolo	43		1 (1)		42 (3)
Utrecht	6		2 (1)	4 (2)	
Warsaw	30	5 (4)		4 (3)	21 (7)
Würzburg	3		2 (1)		1 (1)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3 Penetrance data for RET codon 10 mutations compared to the C634W mutation of codon 11 adjusted for C cell hyperplasia (CCH), MTC stage T1-4N0M0, stage T1-4N1M0, and stage T1-4N0or1M1, and for pheochromocytoma (pheo) and hyperparathyroidism (HPT)

Table 3 (part 1)

CCH

All Exon 10 Codons	5,4	9,6	10,4	13,9	13,9	16,7
Codon 609	0	5,6	5,6	11,9	11,9	33,9
Codon 611	4,3	6,4	9,0	9,0	9,0	9,0
Codon 618	3,4	8,2	8,2	8,2	8,2	8,2
Codon 620	8,6	12,6	13,7	20,9	20,9	20,9
C634W	5,3	9,9	9,9	9,9	9,9	9,9
AGE →	10	20	30	40	50	60

MTC T1-4N0M0

All Exon 10 Codons	2,9	10,2	26,6	42,0	57,0	67,3
Codon 609	2,4	11,2	27,0	46,6	69,2	74,3
Codon 611	0	7,1	15,6	34,0	44,3	59,4
Codon 618	1,2	8,7	31,1	43,7	60,2	72,8
Codon 620	5,3	12,0	27,0	42,2	54,6	61,6
C634W	6,8	26,0	36,1	49,6	65,9	76,1
AGE →	10	20	30	40	50	60

Table 3 continued (1)

MTC T1-4N1M0

All Exon 10 Codons	0,3	2,3	11,0	25,4	43,2	51,3
Codon 609	0	0	3,4	19,8	34,4	45,3
Codon 611	0	0	3,2	6,6	24,6	24,6
Codon 618	1,2	2,5	8,8	20,4	41,5	52,5
Codon 620	0	3,8	17,9	38,9	55,0	64,0
C634W	0	0	16,3	26,2	41,6	41,6
AGE →	10	20	30	40	50	60

MTC T1-4N0or1M1

All Exon 10 Codons	0	0,8	5,1	11,4	18,4	27,4
Codon 609	0	0	0	0	0	0
Codon 611	0	0	2,9	7,0	7,0	23,9
Codon 618	0	1,2	4,3	11,8	20,2	28,1
Codon 620	0	0,9	8,2	16,5	29,3	35,7
C634W	0	0	5,6	8,8	8,8	8,8
AGE →	10	20	30	40	50	60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3 continued (2)

PHEO

All Exon 10 Codons	0	0,8	5,1	12,0	23,1	33,2
Codon 609	0	2,8	5,7	9,4	26,9	47,0
Codon 611	0	0	0	3,1	15,4	15,4
Codon 618	0	0	5,1	13,5	24,8	42,5
Codon 620	0	0,9	6,9	16,0	22,9	26,3
C634W	0	1,4	19,6	42,6	67,1	70,1
AGE →	10	20	30	40	50	60

HPT

All Exon 10 Codons	0	0	0,5	1,8	3,9	3,9
Codon 609	0	0	0	0	5,6	5,6
Codon 611	0	0	0	0	5,6	5,6
Codon 618	0	0	1,9	1,9	1,9	1,9
Codon 620	0	0	1,3	3,2	3,2	3,2
C634W	0	0	2,6	10,7	20,6	33,8
AGE →	10	20	30	40	50	60

Table 2 Comparison of MTC-tumor-stage and outcome in the 4 mutated codons in exon 10

	609 n (%)	611 n (%)	618 n (%)	620 n (%)	p-value
Normal histology	5/40 (12)	4/42 (10)	2/87 (2)	2/131 (2)	0.004 ^a <0.001 ^b
CCH only	4/40 (10)	4/42 (10)	7/87 (8)	22/131 (17)	
MTC - T1-4,N0,M0	23/40 (58)	21/42 (50)	45/87 (52)	48/131 (37)	
MTC - N1,M0	7/40 (18)	6/42 (14)	20/87 (23)	39/131 (30)	
MTC - N1,M1	1/40 (2)	7/42 (17)	13/87 (15)	20/131 (15)	
MTC, last info - cured ^c	26/32 (81)	25/34 (74)	39/72 (54)	51/104 (49)	0.002 ^a <0.001 ^b
MTC - metastasized ^c	4/33 (12)	8/33 (24)	24/76 (32)	43/104 (41)	0.011 ^a 0.002 ^b

^aTest of general association among codon groups ^bTest of trend across codons,

^c number of observations are different due to missing values