Impact of tropical land use change on soil organic carbon stocks – a meta-analysis

Journal:	<i>Global Change Biology</i>
Manuscript ID:	GCB-10-0535
Wiley - Manuscript type:	Review
Date Submitted by the Author:	28-Jun-2010
Complete List of Authors:	Don, Axel; Johann Heinrich von Thünen Institute, Institute of Agricultural Climate Research Schumacher, Jens; Friedrich Schiller University, Institute of Stochastic Freibauer, Annette; Johann Heinrich von Thünen Institute, Institute of Agricultural Climate Research
Keywords:	Land use change, Soil organic carbon, Deforestation, Afforestation, Subsoil, Bulk density, Tropics
Abstract:	Land use changes are the second largest source of human induced greenhouse gas emission, mainly due to deforestation in the tropics and sub-tropics. CO ₂ emissions result from biomass and soil organic carbon (SOC) losses and may be offset with afforestation programs. However, the effect of land use changes on SOC is poorly quantified due to insufficient data quality (only SOC concentrations and no SOC stocks, shallow sampling depth) and representativeness. In a global meta-analysis, 385 studies on land use change in the tropics were explored to estimate the SOC stock changes for all major land use change types. The highest SOC losses were caused by conversion of primary forest into cropland (-25%) and perennial crops (-30%) but forest conversion into grassland also reduced SOC stocks by 12%. Secondary forests stored less SOC than primary forests (-9%) underlining the importance of primary forests for C stores. SOC losses are partly reversible if agricultural land is afforested (+29%) or under cropland fallow (+32%) and with cropland conversion into grassland (+26%). Data on soil bulk density are critical in order to estimate SOC stock changes because i) the bulk density changes with land use and needs to be accounted for when calculating SOC stocks and ii) soil sample mass has to be corrected for bulk density changes in order to compare land use types on the same basis of soil mass. Without soil mass correction, land use change effects would have been underestimated by 28%. Land use change impact on SOC was not restricted to the surface soil, but relative changes

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	were equally high in the subsoil, stressing the importance of sufficiently deep sampling.

SCHOLARONE™
Manuscripts

For Review Only

1 REVIEW

2 Impact of tropical land use change on soil organic carbon stocks – a meta-analysis

3 Running title: Soil organic carbon and land use change

4
5 Axel Don^{1)*}, Jens Schumacher²⁾, Annette Freibauer¹⁾

6
7 1) Johann Heinrich von Thünen-Institute, Institute of Agricultural Climate Research,
8 Bundesallee 50, 38116 Braunschweig, Germany

9 2) Friedrich Schiller University, Institute of Stochastic, Jena, Germany

10
11 * Corresponding author:

12 Phone: ++49-531-596-2641

13 Fax: ++49-531-596-2645

14 Email: axel.don@vti.bund.de

15
16 **Keywords:** Land use change, Soil organic carbon, Tropics, Deforestation, Afforestation,
17 Subsoil, Bulk density

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Land use changes are the second largest source of human induced greenhouse gas emission, mainly due to deforestation in the tropics and sub-tropics. CO₂ emissions result from biomass and soil organic carbon (SOC) losses and may be offset with afforestation programs. However, the effect of land use changes on SOC is poorly quantified due to insufficient data quality (only SOC concentrations and no SOC stocks, shallow sampling depth) and representativeness. In a global meta-analysis, 385 studies on land use change in the tropics were explored to estimate the SOC stock changes for all major land use change types. The highest SOC losses were caused by conversion of primary forest into cropland (-25%) and perennial crops (-30%) but forest conversion into grassland also reduced SOC stocks by 12%. Secondary forests stored less SOC than primary forests (-9%) underlining the importance of primary forests for C stores. SOC losses are partly reversible if agricultural land is afforested (+29%) or under cropland fallow (+32%) and with cropland conversion into grassland (+26%). Data on soil bulk density are critical in order to estimate SOC stock changes because i) the bulk density changes with land use and needs to be accounted for when calculating SOC stocks and ii) soil sample mass has to be corrected for bulk density changes in order to compare land use types on the same basis of soil mass. Without soil mass correction, land use change effects would have been underestimated by 28%. Land use change impact on SOC was not restricted to the surface soil, but relative changes were equally high in the subsoil, stressing the importance of sufficiently deep sampling.

1 Introduction

Land use changes in the tropics are responsible for 12-20% of the human induced greenhouse gas emissions and are expected to remain the second largest source of greenhouse gas emission also for the future (IPCC, 2007, van der Werf *et al.*, 2009). The dominant type of land use change is the conversion of forest to agricultural systems with continuously high rates of 13 million ha being deforested per year (FAO, 2006). Governmental measures to reduce deforestation have been effective only in some countries such as Costa Rica and India during the last years. The destruction of primary forest causes a rapid biomass carbon (C) loss that is accompanied by a C loss from soils. A shift from higher to lower average total ecosystem C stocks increases in atmospheric CO₂. Soils are major carbon stores in tropical areas, with 36-60% of ecosystem C in forests being stored in soils (Dixon *et al.*, 1994, FAO, 2006, Malhi *et al.*, 1999). Tropical soils are estimated to emit 0.2 Gt C yr⁻¹ due to land use changes, accounting for 10-30% of the total C emission from deforestation (Houghton, 1999, Achard *et al.*, 2004). In contrast, other land use changes may lead to increased soil organic carbon (SOC) stocks, e.g., if cropland is converted into grassland or afforested (Paul *et al.*, 2002, Guo *et al.*, 2002). However, the estimates of SOC losses and gains are subject to large errors and methodological biases (Goidts *et al.*, 2009) and the susceptibility of SOC to land use change in tropical soils is insufficiently quantified. The estimated errors of the IPCC default values (Good Practice Guidelines LULUCF) for SOC stock changes after land use change are three to four times higher for tropical than for temperate regions (Penman *et al.*, 2003). The reduction of land use changes that lead to C losses from soils and biomass could be a substantial and economically sound contribution to reduce greenhouse gas emissions (Kindermann *et al.*, 2008). Avoided deforestation activities could reduce anthropogenic C emission by 0.8 to 1.3 Gt C yr⁻¹. The most cost effective way to reduce C emissions can be achieved, if under a full carbon accounting scheme all major effects of human activities are included and reported. The land use induced changes in biomass and soil organic carbon

1
2
3 65 (SOC) stocks are the major uncertainty in such accounting schemes and in life cycle
4
5 66 assessments of tropical agricultural products.
6
7
8 67
9
10 68 SOC changes are controlled by i) the decomposition rate of SOC, e.g., due to changes in
11
12 69 microclimate, and ii) alterations in the quantity and quality of C cycled through the system
13
14 70 (Juo *et al.*, 1996). Land use directly affects both microclimate and quantity, quality and the
15
16 71 pathways of C input. Moreover, erosion is controlled by land use and land management and
17
18 72 may decrease SOC stocks in agricultural systems compared with forests. Erosion may be a
19
20 73 major pathway of SOC loss on the plot scale on insufficiently aggregated soils typical for
21
22 74 tropical regions (van Noordwijk *et al.*, 1997, Berhe *et al.*, 2007). On the other hand,
23
24 75 erodibility generally decreases with increasing topsoil SOC content, which stresses the
25
26 76 importance of SOC for soil fertility and productivity (Feller *et al.*, 1997). This is especially
27
28 77 true for tropical regions where nutrient poor, highly weathered soils are often managed with
29
30 78 few external inputs of nutrients and C. Tropical SOC stocks may be more susceptible to
31
32 79 perturbations such as land use changes with twice as high SOC turnover than in temperate
33
34 80 regions (Trumbore, 1993, Six *et al.*, 2002, Penman *et al.*, 2003). Higher temperature and soil
35
36 81 moisture regimes enhance decomposition rates and thus may speed up SOC losses. Highly
37
38 82 weathered soils, e.g., Oxisols and Ultisols, cover 60 to 70% of tropical land area. In these
39
40 83 soils low activity clays are predominant and provide less mineral surfaces for physical
41
42 84 protection and stabilisation of SOC (Feller *et al.*, 1997). However, there is an ongoing
43
44 85 discussion about whether climatic factors or the differences in soil mineralogy and land use
45
46 86 history contribute most to distinct tropical SOC dynamics (Paul *et al.*, 2008, Zinn *et al.*, 2005,
47
48 87 Zinn *et al.*, 2007, Feller *et al.*, 1997).
49
50
51 88
52
53 89 During the last years, many new research programs and projects have aimed to improve the
54
55 90 understanding on carbon fluxes and balances in tropical soils. Hundreds of new studies were
56
57
58
59
60

published but have never been analysed together. Moreover, insufficient sampling depth and missing correction for differences in bulk density after land use change may have lead to significant bias in previous studies (Ellert *et al.*, 1995, Baker *et al.*, 2007). Differences in rooting depth and tillage on croplands directly influence the C distribution in the soil profile. Shallow sampling misses C which is incorporated below the topsoil and may lead to overestimations of land use change effects on soil C. Thus, the objective of this study was to gather the existing high quality data sets on land use change effects and SOC for the tropics to derive new estimates beyond site specific values and including also subsoil horizons. More than 380 old and new data sets were compiled and quantitatively analysed. This study provides the first estimate of tropical soil C stock changes after land use change for the depth 0-30 cm, the soil depth that has to be reported under UNFCCC, and below.

2 Material and Methods

2.1 Data sources and compilation

Data from 385 studies from 153 published and mostly peer reviewed publications on the influence of land use changes on soil organic carbon were compiled. Data were derived from 39 different tropical countries covering all continents ranging from semi-arid regions such as southern Africa and northern Australia to the humid tropics along the equator. Twelve different land use change types were classified and investigated covering all land use transitions occurring in the tropical zone (Tab. 1). Most studies were conducted in paired plot design using the “space for time” approach. Since SOC may reach a new equilibrium only after several years or decades, there was almost no study with a time series going back to prior land use change conditions. Thus, for each paired site, it has to be assumed that soil conditions were similar before the land use change. Studies were rejected for the data compilation if the different land use types were i) confounded by different soil types as, e.g., indicated by significant differences in texture, ii) sampled for different soil depth or iii)

reporting only short term effects (< 5 years). For chronosequences only data from the longest treatments of land use change were used in this study. With the exclusion of short term studies, the influence of the time period since land use change on the estimated SOC changes was minimized and not detectable anymore. Data on SOC stocks, bulk density and the associated meta-data were compiled. Organic layers (forest floor) are rare in tropical forests and the few existing data sets did not allow them to be included in this analysis.

In the current study, primary forest is defined as natural vegetation without apparent and reported human impacts. The primary forest vegetation class also comprises natural vegetations with shrubland and non-managed grassland with savannah-like characters, e.g., the South American Cerrado. It has to be noted that there are only few remaining totally undisturbed tropical forests leading to a rather broad definition of “primary forest” in many studies and consequently also in our study (Lugo *et al.*, 1993). Secondary forests are managed forests and forests regrown after destruction or partial exploitation of the primary forest. Natural successions and fallow older than 7 years were classified as secondary forest. “Grassland” comprises pastures but no natural grasslands, since natural grasslands are mostly savannah type grasslands with tree and shrub vegetation. Additionally, there is no harvested fraction of net primary production on natural grasslands, which directly affects the C dynamics. Croplands are classified as “perennial crops,” such as sugar cane and coffee plantations, and “croplands,” with annual crops such as maize and beans. Both cropland types and grasslands were described as agricultural systems.

2.2 Data treatment and missing bulk densities

For 81% of the reported data, SOC stocks were directly available or calculated as

$$\text{SOC stock [Mg ha}^{-1}\text{]} = \sum_{i=1}^n \text{SOC concentration [Mg Mg}^{-1}\text{]} * \text{bulk density [Mg m}^{-3}\text{]} * \text{soil volume [m}^3 \text{ ha}^{-1}\text{]} \text{ (Eq 1),}$$

1
2
3 142 where n was the number of soil layers, varying for each study. Bulk density data were
4
5 143 available for 52% of all reported soil horizons. Most studies reported SOC stocks (81%) but
6
7
8 144 19% of the studies reported only SOC concentrations. Two strategies can be used to handle
9
10 145 the problem of missing bulk density data: either all studies without bulk density
11
12 146 measurements are excluded from the meta-analysis, or estimated mean bulk densities replace
13
14
15 147 the unknown values to convert SOC concentrations into stocks. To quantify the difference in
16
17 148 accuracy between the two approaches we first conducted the meta-analysis (see below) based
18
19 149 only on those studies with bulk density measurements. For the second approach the data set
20
21 150 was divided in two sub datasets comprising the studies with and without bulk density
22
23 151 measurements. For those studies lacking own bulk density data, the bulk densities before and
24
25 152 after land use change were simulated as two-dimensional truncated normal random vectors
26
27 153 separately for each land use change type. For these Monte Carlo simulations the means and
28
29 154 covariances were derived from the studies with bulk density measurements. The truncation
30
31 155 was necessary to avoid unrealistic (e.g., negative) values for bulk density caused by the
32
33 156 unbounded normal distribution. The standard deviation of the estimated mean effect size from
34
35 157 10000 repeated simulations provided a direct estimate of the uncertainty introduced by using
36
37 158 an estimated mean bulk density instead of true measurements. Finally, estimates of mean
38
39 159 effect sizes and their standard errors were obtained as weighted averages of the estimates
40
41 160 from the two sub datasets. These Monte Carlo simulations revealed that for all land use
42
43 161 change types the estimated uncertainty can be reduced by including also the studies that
44
45 162 reported only SOC concentrations after converting them with weighted mean bulk densities
46
47 163 into SOC stocks. Thus, we decided to also include studies that report only SOC
48
49 164 concentrations.
50
51
52
53
54
55
56
57 165
58
59 166 SOC stocks were corrected to an equivalent soil mass on both land use types (Ellert *et al.*,
60
167 1995). Weighted mean bulk densities for each land use change type were used if bulk density

data were not available to perform this correction. Data reported as SOM concentrations were converted to SOC by multiplying with a conversion factor of 0.58 (Mann, 1986). Studies were restricted to mineral soils. Wetlands soils such as peatlands and paddy soils were not included in this analysis, mainly due to an insufficient number of studies on these soil types to obtain an adequate representation compared to non-wetland soils. Soil horizons down to max 100 cm were included in the analysis. The following relevant meta-data were also included in the compilation: time since conversion (age), clay content (texture), soil type, mean annual precipitation, mean annual temperature, soil sampling depth and other management factors (tillage, species, fertilisation etc.). If some of these data were not available, data were not estimated by interpolation or transfer functions and the study was excluded from the corresponding analysis. Only for climatic data were other sources used such as long term climate records of the region.

2.2 Meta-analysis

The simplest measure of effect size δ commonly employed in a meta-analysis is the difference between control group mean μ_c (before land use change) and treatment group mean μ_e (after land use change). We used both the absolute effect size $\delta_{abs,i} = \mu_{e,i} - \mu_{c,i}$ and the relative effect size $\delta_{rel,i} = (\mu_{e,i} - \mu_{c,i}) / \mu_{c,i} * 100\%$. To account for the different accuracy of the heterogeneous set of studies, the mean effect size for the different land use change types was estimated as weighted mean with the optimal weights being inversely proportional to the variance of the single-study effect sizes. To estimate the optimal weights we had to estimate these variances. Two sources of variability contribute to the uncertainty of the effect sizes, the within-study variability derived from sampling and analytical errors and the between-study variability derived from differences in climate, soil, plant species and land management between studies. Since 73% of all studies did not report any measure of variability/accuracy for their estimated means $\mu_{c,i}$ and $\mu_{e,i}$, we used the available information to estimate the underlying within-study

variances $\sigma_{within_{c,i}}^2$ and $\sigma_{within_{e,i}}^2$ as a power function of SOC stocks ($R^2=0.67$ for a linear regression on a log-log scale). The standard errors of reported SOC stocks estimates for a single study are then determined by the differing sample sizes n_i and can be estimated as:

$$\hat{\sigma}_{\mu_{within,i}} = \frac{\hat{\sigma}_{within,i}}{\sqrt{n_i}} \quad \text{Eq (2)}$$

for both means, μ_c and μ_e , respectively. The main uncertainty is derived from the field heterogeneity and therefore, the sample size n is the number of collected soil samples, even though they have been pooled to compound samples before analysis. Between-study variability was estimated by the moment estimator

$$\hat{\sigma}_{between}^2 = \frac{1}{k-1} \sum_{i=1}^k (\mu_i - \bar{\mu})^2 - \frac{1}{k} \sum_{i=1}^k \frac{\hat{\sigma}_{within,i}^2}{n_i} \quad \text{Eq (3)}$$

resulting in the following estimated variance for the absolute effect size $\delta_{abs,i}$:

$$\hat{\sigma}_{abs,i}^2 = \hat{\sigma}_{between,c}^2 + \hat{\sigma}_{\mu_{c,within,i}}^2 + \hat{\sigma}_{between,e}^2 + \hat{\sigma}_{\mu_{e,within,i}}^2 - 2\hat{cov}(\mu_{c,i}, \mu_{e,i}) \quad \text{Eq (4)}$$

The mean absolute effect size δ_{abs} was then estimated as the weighted average

$$\hat{\delta}_{abs} = \frac{\sum_{i=1}^k \left(\frac{1}{\hat{\sigma}_{abs,i}^2} \cdot \hat{\delta}_{abs,i} \right)}{\sum_{i=1}^k \frac{1}{\hat{\sigma}_{abs,i}^2}} \quad \text{Eq (5)}$$

For the variance estimation of the mean relative effect sizes $\delta_{rel,i}$, we used the same weights

$$\hat{\delta}_{rel} = \frac{\sum_{i=1}^k \left(\frac{1}{\hat{\sigma}_{abs,i}^2} \cdot \hat{\delta}_{rel,i} \right)}{\sum_{i=1}^k \frac{1}{\hat{\sigma}_{abs,i}^2}} \quad \text{Eq (6)}$$

Estimates of standard errors for the weighted means were obtained by nonparametric bootstrap based on 1000 bootstrap samples.

Four different sampling depths were selected in order to investigate the effects of land use change for different soil depth: Topsoil (0-10 cm), 0-30 cm depth (ploughing horizon),

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

subsoil (>20 cm depth) and full depth (total sampling depth of each study). Studies reporting only data for some soil depth classes were however included for the analysis of the reported soil depth class which led to different numbers of studies for the four soil depth classes. There was no significant influence of the maximum sampling depth on the relative SOC changes (as fixed and variable effect in the general linear models; $F_{133,4,1} = 0.097$, $p=0.76$), indicating that the variability between different studies is as large as between different maximum sampling depths.

2.3 Statistical analysis

The influence of the following variables on SOC change and bulk density change was investigated on a core data set which contained all of the following variables comprising 377 land use pairs: Mean annual temperature, annual precipitation, soil mineralogy (using the three clay type classes: high activity clay, low activity clay, allophanic clay), region/continent, sampling depth and SOC stocks before land use change. Additionally, the methodological parameters maximum sampling depth and number of sample replicates were used as independent variables. We checked the effects with mixed linear models including the author of the studies as a random effect. Since no author-specific effects could be found, we used classical general linear models for the further analysis. Data are presented in the text as $F_{\text{Sum of squares, degrees of freedom}}$ and the p-value. Statistical analysis was performed using R software.

3 Results

3.1 Data quality and mass correction

The analysis of land use effects on soil carbon is hampered by the high heterogeneity of the data set including different sampling methods, sampling intervals and missing meta-data such as the land use history. Most important for improved estimates of SOC changes is the availability of bulk density data in order to account for SOC changes on an area basis and to

be able to correct data for different soil mass sampled in land use types. The correction for different soil mass (bulk density) of land use types increased the land use change effect by an average of 28% (Tab. 2). Pedo-transfer functions (Post *et al.*, 2000, De Vos *et al.*, 2005, Mann, 1986) are rarely able to take these effects into account with sufficient accuracy and we were able to predict bulk density in our study only with a correlation coefficient of 0.67 (data not shown). However, the uncertainty of the estimated SOC change could be decreased by combining studies with and without density measurements. The fraction of studies reporting only SOC concentrations per land use change type was between 6 and 36%, indicating that the majority of studies reported bulk densities and stocks. The uncertainty has been reduced by on average 52% as compared to a meta-analysis only comprising studies with bulk density managements. With our approach, we take the uncertainty derived from an incomplete data set (e.g. missing bulk density data) and the uncertainty of SOC and bulk density measurements into account. The within-study uncertainty depends on the sampling and analytical errors and the soil heterogeneity in the field and could be reduced by increasing numbers of soil samples (Eq. 2).

Land use change caused changes in the bulk density of slightly lower magnitude like SOC changes and in the reverse direction (Fig. 1, Tab. 2). Since organic carbon has an inherent low bulk density, SOC concentrations directly affect soil bulk density (Lal *et al.*, 2001). However, land use changes affect bulk density beyond this effect due to compaction by animal trampling, machinery and loosening by tillage. The cultivation of forests caused bulk density to increase by 5 to 23% with the strongest increase occurring in the surface, lessening with depth with no significant effects below 20 cm depth (Fig. 1). Surprisingly, even tillage on croplands did not decrease bulk density, but cropland bulk density was always higher than grassland and forest bulk density.

Rates of land use change vary widely among tropical regions but the number of existing studies did not reflect land use change rates. Malaysia and Indonesia are among the countries with the highest emissions due to land use changes (Houghton, 2003, FAO, 2006). However, the region of South Eastern Asia was undersampled with only 11 studies reporting quantitative data on land use effects on SOC. Most regions in Africa – except for Nigeria – were also undersampled, whereas good data coverage has been reached in Central and South America, especially in Ecuador, Costa Rica and Brazil.

3.2 Primary forest to agricultural land

The conversion of native vegetation to agricultural systems caused the highest SOC losses among all land use change types (Tab. 1, Fig. 2 and 3). Native vegetation such as primary forest and native grassland stored among the highest amounts of SOC ($80 \pm 9 \text{ Mg SOC ha}^{-1}$, mean sampling depth 36 cm). Conversion of primary forest to cropland (-25%) caused twice as high SOC losses than its conversion to grassland (-12%). The relative SOC loss in the subsoil was similar on grasslands but not significant for croplands due to a great variability caused by different management practices and crop types. With the cultivation of primary forests, soils were compacted and bulk density increased by 14 and 18% for grasslands and croplands, respectively. Especially if forests are converted to grasslands the correction for different soil mass exerted a strong influence on the estimated SOC changes, switching this land use change from almost C neutral to a significant C source (Tab. 2).

42% of the variability between data sets could be explained with different land use change types and the climatic factors precipitation and temperature. SOC losses increased with increasing temperature and for conversion of forest into grassland also with increasing precipitation (Tab. 3 and 4). For forest cultivation we found no uniform influence of precipitation on SOC changes and a higher uncertainty of the models as compared to forest conversion to grassland.

291

292 **3.3 Secondary forest to agricultural land**

293 Similar to primary forests also secondary forests' conversion to agriculture systems led to

294 SOC losses. However, relative SOC loss (for all depths) was less compared to primary

295 forests, indicating a higher vulnerability of SOC in primary forests than in secondary forests

296 to land use changes. Surface soil SOC stocks remain unchanged when secondary forest was

297 converted to grassland (Fig. 3). Grasslands are characterised by a steep C gradient with soil

298 depth leading to high surface soil SOC stocks. In contrast, a smaller fraction of total SOC is

299 stored in the surface soil of secondary forests. SOC losses after deforestation were

300 significantly affected by climatic factors, in particular moisture conditions (mean annual

301 precipitation) and temperature (Tab. 3, Fig. 4). Surprisingly, we found no effect of the clay

302 content ($F_{7750,5}$; $p=0.58$; Fig. 5) and soil type ($F_{53213,30}$ $p=0.45$) on SOC losses. However,

303 beside temperature and precipitation the differences between clay types (low activity, high

304 activity allophonic) exerted a significant effect on SOC changes (Tab. 3). Land use change

305 types, climate factors and clay type could explain 55% of the SOC change variance leaving

306 almost half of the data set variability unexplained. The effect of different management

307 practices for croplands (e.g., tillage vs. no tillage) could not be investigated due to a very

308 small number of studies covering both land use change and different management practices.

310 **3.4 Primary to secondary forest**

311 Management of primary forest with wood extraction and planting of productive tree species

312 caused a mean SOC loss of 7% or 9 Mg SOC ha⁻¹ (Tab. 1). One major difference between

313 primary and secondary forest is the SOC distribution in the soil profile (Fig. 2 and 3), with a

314 higher surface SOC fraction in primary forest compared to secondary forests. 7 Mg SOC ha⁻¹

315 (-15%) were lost in the upper 10 cm only after conversion of primary forest to secondary

316 forest. However, there was no significant SOC change below 20 cm depth. SOC losses were

1
2
3 317 accompanied by a 6% increase in the soil bulk density leading to 28% higher SOC changes
4
5 318 after mass correction as compared with no mass corrected data (Tab. 2).
6
7
8 319

10 320 **3.5 Afforestation and fallow**

12 321 SOC losses due to deforestations were partly reversible by afforestations of croplands or
13
14 322 grasslands. Estimated mean SOC stock gains for afforestations were even higher than SOC
15
16 323 losses from deforestations, with highest SOC gains in afforestations on former croplands (+
17
18 324 50%) compared to afforestations on former grassland (18%, Tab. 1). Afforested grassland
19
20 325 stored particular low amounts of SOC ($60 \pm 9 \text{ Mg ha}^{-1}$) since afforestations were mainly
21
22 326 conducted on degraded grasslands or on marginal land with intrinsic low SOC storage
23
24 327 capacity (Tab. 1). Similarly, the termination of cropping leading to natural succession (fallow)
25
26 328 on croplands took place mainly on degraded land as indicated by low SOC stocks (Tab. 1).
27
28 329 Fallow increased SOC stocks by 32% indicating a rapid recovery of SOC stocks.
29
30
31
32
33

34 330
35
36 331 **3.6 Conversion of grassland to cropland and vice versa**

38 332 Typical land use change cascades in the tropics are first the conversion of forest into
39
40 333 grassland for cattle grazing and at a later stage the conversion of grassland into cropland.
41
42 334 Cropland establishment on grasslands reduced SOC stocks by 6 Mg C ha^{-1} (-10%) but this
43
44 335 effect was restricted to the uppermost soil horizon. Subsoil below 20 cm depth was not
45
46 336 significantly affected by these land use changes due to high C input with tillage. Several
47
48 337 studies reported lower subsoil SOC stock in grassland compared to croplands (Fujisaka 1998,
49
50 338 Huges, 2000, Freitas 2000). Cropland conversion or re-conversion to grassland increased
51
52 339 SOC stocks by 8 Mg C ha^{-1} (+26%), which is more than the SOC loss after cropland
53
54 340 establishment on grassland. Similar to the afforestation of croplands, this indicates that
55
56 341 croplands management causes SOC losses leading to lower initial SOC stocks of croplands
57
58
59 342 before conversion into grasslands (Tab. 1).
60

343

344 3.7 Perennial crops and plantations

345 The conversion of primary forests to perennial crops caused an even higher C loss of than the
346 conversion to cropland (-30%, Tab. 1). In contrast, the conversion of secondary forests to
347 perennial crops seems to hardly affect SOC stocks. This may be partly explained by higher
348 mean SOC stock before land use change in primary forests than in secondary forests of
349 studies reporting conversion into perennial crops. For all land use change types SOC losses
350 and gains were weakly positively correlated with SOC stock before land use change. Most
351 data on soil carbon on perennial crops were reported from sugar cane plantations (28% of
352 studies including perennial crops), fruit tree plantations (including banana) (12%) and cacao
353 plantations (9%). These findings are based on 35 studies and indicate that a permanent
354 vegetation cover does not always prevent SOC losses under intensive management when SOC
355 rich forests are converted to perennial crop plantations. SOC changes may be different in low
356 input agro-forestry type perennial croplands which are not well covered in this meta-analysis.

357

358 4 Discussion

359 4.1 Deforestation and afforestation

360 A large number of studies on land use change effects were conducted during the last years,
361 25% of the studies in this meta-analysis were published during the last five years and 67%
362 during the last 10 years. Moreover, it was only recently that more studies also included deeper
363 soil horizons down to 100 cm depth. Former reviews calculated higher global and tropical
364 SOC stock changes after cultivation of forests compared to our study (Davidson *et al.*, 1993,
365 Guo *et al.*, 2002, Detwiler, 1986, Paustian *et al.*, 1997, Amundson, 2001). This can partly be
366 explained with an improved data quality and quantity, e.g., with deeper sampling and more
367 data on bulk density changes. Detwiler (1986) found twice as high SOC losses after
368 deforestation than reported in our study (-20% for forest to grassland and -40% for forest to

1
2
3 369 cropland). A range of SOC losses from -24 to -43% was reported for cultivated tropical soils
4
5 370 (Davidson *et al.*, 1993). The IPCC guidelines set a default value of -31 and -42% SOC for dry
6
7
8 371 and wet tropical regions after forest cultivation, respectively, which is higher than calculated
9
10 372 in our study (Tab. 5). Our study confirmed the impact of soil moisture and precipitation on
11
12 373 SOC dynamics with higher SOC changes in regions with higher precipitation for most land
13
14 374 use change types (Tab. 3, Fig. 4). Soils in humid regions maybe more vulnerable to land use
15
16 375 changes than in dryer regions (Brown *et al.*, 1990, Amundson, 2001). The impact of
17
18 376 precipitation seems to be stronger when forests are converted into grasslands than for forest
19
20 377 conversion into cropland (Tab. 5). In a global analysis Guo and Gifford (2002) found
21
22 378 conversion of forest to grassland to increase SOC stocks by 9% (2002), no SOC change has to
23
24 379 be assumed as default value under IPCC and is reported in other reviews (Lugo *et al.*, 1993,
25
26 380 Cerri *et al.*, 2004, Penman *et al.*, 2003). In contrast, we found tropical forests lost 12% SOC
27
28 381 after grassland establishment (Tab. 1). These differences can be partly attributed to improved
29
30 382 data quality and the application of a soil mass correction which accounts for changes in
31
32 383 different bulk densities in different land use types (Ellert *et al.*, 1995, Gifford *et al.*, 2003, de
33
34 384 Moraes *et al.*, 1996). Detwiler (1986) tried already to overcome the problem of different soil
35
36 385 mass but had to rely on calculated and not measured bulk densities. Soils were compacted by
37
38 386 10 and 16% due to forest conversion into grassland and cropland, respectively. Most tropical
39
40 387 grasslands are under higher grazing pressure, a higher biomass fraction is exported (harvest)
41
42 388 and fertilizer input is low compared to temperate grasslands. Improved grassland management
43
44 389 with the application of fertilizers would help to increase productivity and SOC stocks
45
46 390 compared to extensive pastures (Soussana *et al.*, 2007, Ammann *et al.*, 2007). Roots are a
47
48 391 more effective pathway to build up SOC stocks than foliar litter, which explains high
49
50 392 grassland SOC (Lugo *et al.*, 1993, Rasse *et al.*, 2005) and relatively small SOC losses if
51
52 393 forests are converted to grasslands as compared to croplands.
53
54
55
56
57
58
59
60
394

A major proportion of total SOC change occurred during the first few years after cultivation of forest, indicating that these soils contain large amounts of labile SOC, potentially vulnerable to degradation upon human-induced land-use changes (Solomon *et al.*, 2007). A new equilibrium of SOC has been reached most often within 3 to 10 years (Houghton, 1999, Feller *et al.*, 1997, Detwiler, 1986, Davidson *et al.*, 1993). Other studies found 20 to 40 years (Solomon *et al.*, 2007, Sa *et al.*, 2001, Riezebos *et al.*, 1998, Cerri *et al.*, 2007). In our study, the average time period since deforestation was 22 years, and 33 years since afforestation, indicating that major parts of SOC changes are captured within this time period.

Reforestation and afforestation were found to successfully recover SOC stocks (Silver *et al.*, 2000, Post *et al.*, 2000, Bashkin *et al.*, 1998). Cropland afforestation increased SOC stocks by 33 Mg ha⁻¹ which is slightly lower than the mean SOC accumulation of 41 Mg ha⁻¹ after 80 years reported from Silver *et al.* (2000). SOC gains with afforestation were higher than SOC losses after deforestation (Tab. 1). Forest establishment has mainly been performed with highly productive tree species like eucalyptus with a low litter quality and high recalcitrance. High SOC accumulation in secondary forest may be also fostered by a low initial SOC content in the afforested degraded agricultural land (Lugo *et al.*, 1993). Agricultural management on highly weathered soils often lead to a rapid decline in soil fertility, leaving degraded land for forest regrowth, as it is part of the traditional shifting cultivation system.

4.2 Agricultural systems

Low SOC stocks in croplands have important implications for crop production since organic matter supplies most of the nitrogen and parts of the phosphorous taken up by unfertilized crops (Sanchez, 1976). SOC is essential for the retention of nutrients and water in highly weathered soils with low cation exchange capacity. Tropical regions cover very different stages of agricultural mechanisation and development with various management options on

1
2
3 421 croplands, including organic amendments and different tillage practice, and a high number of
4
5 422 different crops. Thus, the estimated effect of cultivation can only set a mean value for regions
6
7
8 423 but cannot be applied to specific field sites. Climatic and soil parameters could only explain
9
10 424 55% of the data variability. For agricultural systems, the biomass fraction left for SOC build-
11
12 425 up (crop residuals) is strongly controlled by management practices including the selection of
13
14 426 crop species. Improved cropland management may partly offset SOC losses due to
15
16
17 427 deforestation (Lugo *et al.*, 1993); 13% of croplands included in this meta-analysis reported
18
19 428 similar or higher SOC stocks than in forests. Additional effort with field data collection is
20
21 429 necessary to quantify the effect of different management options on a global scale. Moreover,
22
23 430 insufficient sampling depths were found to obscure conclusions on management and land use
24
25
26 431 effects on the SOC balance (Baker *et al.*, 2007).

27
28
29 432
30
31 433 Regular soil disturbance during tillage or harvest is one of the main reasons for low cropland
32
33 434 SOC stocks (Lal, 1998). Grasslands, pastures and perennial crops, unlike croplands, maintain
34
35 435 a permanent vegetation cover and a high root turnover leading to high SOC input. We found
36
37 436 surprisingly high SOC losses after primary forest was converted to perennial cropland or
38
39 437 grassland. The amount of crop residuals returned to the soil directly affect SOC, and most
40
41 438 perennial crops such as sugar cane plantations, are managed with high intensity and high
42
43 439 biomass export (Graham *et al.*, 2002). Similar to other cropland types, different management
44
45 440 options and land use history determined the amount of SOC loss after cultivation of primary
46
47 441 forests and, on the other hand, the sequestration potential if perennial cropland and grassland
48
49 442 is afforested.

50
51 443
52
53
54
55
56
57 444 **4.3 Soil characteristics and erosion**
58
59 445 Differences in SOC change are expected to be attributable to soil parameters such as soil type,
60
446 texture and clay mineral type (de Moraes *et al.*, 1996, Feller *et al.*, 1997, Hartemink, 1997,

Davidson *et al.*, 1993). Clay type was found to explain only additional 13% of the SOC variability, beside land use change type, temperature and precipitation (Tab. 3). No influence on SOC change could be attributed to soil type and clay content (Fig. 5), which is in line with findings from Davidson *et al.* (1993). Soil parameters' influence maybe obscured by dominant other factors such as climate and management. Additionally, the data availability was low with only 22% of all studies reporting clay content. Highly weathered soils, such as Ferralsols were found to loose more SOC after cultivation than other soil types (Hartemink, 1997), but this could not be confirmed in our study.

Erosion is a major factor affecting SOC stocks that is directly related to land use and forest clearing (Nye *et al.*, 1964, van Noordwijk *et al.*, 1997, Wairiu *et al.*, 2003). Soils under low vegetation cover (agricultural systems, conventional tillage), on steep slopes and under high precipitation intensity are most prone to erosion. However, adequate data was not available in this meta- analysis to assess the proportion of erosion-triggered SOC loss. Some areas may even gain SOC with deposition of eroded material, leaving the question open of whether erosion decreases or increases the terrestrial C sink (Berhe *et al.*, 2007, Lal, 2003, Van Oost *et al.*, 2007).

4.4 SOC changes in the surface soil and the subsoil

The SOC in topsoil is supposed to be more prone to land use change and other perturbations than subsoil (Siband, 1974, Veldkamp *et al.*, 2003, Veldkamp, 1994). We found equally high relative subsoil SOC stock changes compared with surface soil horizons after conversion of native forests to agriculture systems (Fig. 2 and 4). Native forests stored higher amounts of subsoil C which are lost upon cultivation compared with secondary forests. The mean soil sampling depth did not contribute to an explanatory model indicating that the relative SOC change is only weakly related to the soil depth. In contrast, absolute SOC changes

1
2
3 473 significantly decrease with soil depth owing to decreasing absolute SOC stocks in deeper
4
5 474 horizons. In fact, tillage may even increase subsoil SOC stocks in croplands due to C rich
6
7
8 475 topsoil being mixed with deeper horizons (Hughes *et al.*, 2000, Fujisaka *et al.*, 1998). A
9
10 476 sampling depth as deep as the tillage depth is the minimum to quantify land use change
11
12 477 effects. Our results indicate that at least the conversion of native forests also affects subsoil
13
14 478 SOC below 20 cm depth and a comprehensive assessment should also include subsoil
15
16
17 479 horizons, if possible down to 100 cm depth. In order to estimate subsoil SOC changes with
18
19 480 land use change it is even more important to ensure comparable soil intrinsic conditions on
20
21 481 paired or chronosequence sites since SOC stabilisation in the subsoil is highly dependent on
22
23 482 soil mineralogy, texture and other soil parameters. The high variability of subsoil SOC change
24
25
26 483 may be a result of the variability of these soil intrinsic parameters.
27
28
29 484

30
31
32 485 **4.5 Bulk density change and its impact of SOC stock estimates**

33
34 486 The relative changes in bulk density were almost as high as the relative SOC changes, e.g.,
35
36 487 cultivation of forest increased bulk density by 16 % (Fig. 1, Tab. 2). Bulk density changes are
37
38 488 important to account for SOC stocks changes, since SOC stocks linearly depend on both SOC
39
40 489 concentration and bulk density. Moreover, bulk density change causes a sampling bias if
41
42 490 sampling of each land use type is performed at the same sampling depth (Ellert *et al.*, 1995,
43
44 491 Gifford *et al.*, 2003, Davidson *et al.*, 1993). If bulk density increased with land use change,
45
46 492 the soil is compacted and sampling down to the same sample sampling depth would lead to
47
48 493 higher sampled soil mass than in the corresponding land use type. Since soil mass and soil
49
50 494 carbon are ultimately linked, sampling of more soil mass results in higher SOC stocks
51
52 495 (Davidson *et al.*, 1993). Thus, the effect of land use change is underestimated, in our study by
53
54 496 an average 28% (Tab. 2). This can only be completely corrected if bulk density data prior to
55
56 497 land use change are available and it can partly be corrected if bulk density data were recorded
57
58 498 after land use change for both land use types (Lee *et al.*, 2009). We found mass correction
59
60

strongly influencing the effect size (SOC change) with up to three times higher mean SOC changes than estimated without mass correction. However, Monte Carlo simulations revealed that the exclusion of studies that report only SOC concentration would increase the uncertainty of the estimated SOC change by 52%. The high diversity of soil types, climate conditions, vegetation and management types call for as many studies as possible to be included in such meta-analysis, even though not all studies provide the full parameter set. This confirms earlier findings that fewer bulk density than SOC concentration measurements are necessary to estimate SOC stocks (Don *et al.*, 2007). Coefficient of variation (CV) of all studies was 2.7 times lower for bulk density than for SOC concentration (29 and 81% for bulk density and SOC concentration, respectively) indicating that even with few bulk density measurements, uncertainties on land use change effect can be reduced considerably.

5 Conclusions

The conversion of forest, especially primary forests into agricultural systems always lead to SOC losses, but losses are reversible to a high degree if, e.g., agricultural land is afforested or properly managed. For the SOC balance of a land use system, the harvested fraction of net primary production seems to be more important than its disturbance frequency, e.g., with tillage or climate or soil characteristics.

Mass correction of SOC stock estimates is crucial in order to estimate land use change effects since land use change is always accompanied by bulk density changes. The comparison of SOC stocks based on different soil mass deeply confound estimates of SOC changes. This meta-analysis provides soil mass-corrected estimates to improve the current UNFCCC default values. Mean SOC changes were smaller than reported in previous reviews even though soil mass correction increased land use change effects on SOC by 28% on average.

The global data coverage does not mirror the current hot spots of land use changes. New effort are needed to quantify the effect of land use changes in South East Asia and Africa, also

1
2
3 525 taking to account carbon rich wetland forests and degradation cascades within land use
4
5 526 classes.
6
7
8 527
9
10 528 **Acknowledgements**
11
12 529 We thank the library of the Max Planck Institute for Biogeochemistry (Jena) and of the
13
14 530 Johann Heinrich von Thünen Institute (Braunschweig) for their support. We thank Jana
15
16 531 Niebuhr for her help with extracting the uncertainties from the studies. This study was partly
17
18 532 financed by the Federal Ministry of Education and Research (BMBF) (FKZ 01LG0801A).
19
20 533
21
22
23
24 534 **References**
25
26 535 Achard F, Eva HD, Mayaux P, Stibig HJ, Belward A (2004) Improved estimates of net carbon
27
28 536 emissions from land cover change in the tropics for the 1990s. *Global Biogeochemical*
29
30 537 *Cycles*, **18**.
31
32
33 538 Ammann C, Flechard CR, Leifeld J, Neftel A, Fuhrer J (2007) The carbon budget of newly
34
35 539 established temperate grassland depends on management intensity. *Agriculture*
36
37 540 *Ecosystems & Environment*, **121**, 5-20.
38
39
40 541 Amundson R (2001) The carbon budget in soils. *Annual Review of Earth and Planetary*
41
42 542 *Sciences*, **29**, 535-562.
43
44
45 543 Baker JM, Ochsner TE, Venterea RT, Griffis TJ (2007) Tillage and soil carbon sequestration -
46
47 544 What do we really know? *Agriculture Ecosystems & Environment*, **118**, 1-5.
48
49
50 545 Bashkin MA, Binkley D (1998) Changes in soil carbon following afforestation in Hawaii.
51
52 546 *Ecology*, **79**, 828-833.
53
54
55 547 Berhe AA, Harte J, Harden JW, Torn MS (2007) The significance of the erosion-induced
56
57 548 terrestrial carbon sink. *Bioscience*, **57**, 337-346.
58
59
60

- 1
2
3 549 Brown S, Lugo AE (1990) Effects of Forest Clearing and Succession on the Carbon and
4
5 550 Nitrogen-Content of Soils in Puerto-Rico and Us Virgin Islands. *Plant and Soil*, **124**,
6
7 551 53-64.
8
9
10 552 Cerri CEP, Easter M, Paustian K, *et al.* (2007) Predicted soil organic carbon stocks and
11
12 553 changes in the Brazilian Amazon between 2000 and 2030. *Agriculture Ecosystems &*
13
14 554 *Environment*, **122**, 58-72.
15
16
17 555 Cerri CEP, Paustian K, Bernoux M, Victoria RL, Melillo JM, Cerri CC (2004) Modeling
18
19 556 changes in soil organic matter in Amazon forest to pasture conversion with the
20
21 557 Century model. *Global Change Biology*, **10**, 815-832.
22
23
24 558 Davidson EA, Ackerman IL (1993) Changes in Soil Carbon Inventories Following
25
26 559 Cultivation of Previously Untilled Soils. *Biogeochemistry*, **20**, 161-193.
27
28
29 560 de Moraes JFL, Volkoff B, Cerri CC, Bernoux M (1996) Soil properties under Amazon forest
30
31 561 and changes due to pasture installation in Rondonia, Brazil. *Geoderma*, **70**, 63-81.
32
33
34 562 De Vos B, Van Meirvenne M, Quataert P, Deckers J, Muys B (2005) Predictive quality of
35
36 563 pedotransfer functions for estimating bulk density of forest soils. *Soil Science Society*
37
38 564 *of America Journal*, **69**, 500-510.
39
40
41 565 Detwiler RP (1986) Land-Use Change and the Global Carbon-Cycle - the Role of Tropical
42
43 566 Soils. *Biogeochemistry*, **2**, 67-93.
44
45
46 567 Dixon RK, Brown S, Houghton RA, Solomon AM, Trexler MC, Wisniewski J (1994) Carbon
47
48 568 Pools and Flux of Global Forest Ecosystems. *Science*, **263**, 185-190.
49
50
51 569 Don A, Schumacher J, Scherer-Lorenzen M, Scholten T, Schulze E-D (2007) Spatial and
52
53 570 vertical variation of soil carbon at two grassland sites - Implications for measuring soil
54
55 571 carbon stocks. *Geoderma*, **141**, 272-282.
56
57
58 572 Ellert BH, Bettany JR (1995) Calculation of organic matter and nutrients stored in soils under
59
60 573 contrasting management regimes. *Canadian Journal of Soil Science*, **75**, 529-538.
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

- 575 Feller C, Beare MH (1997) Physical control of soil organic matter dynamics in the tropics.
576 *Geoderma*, **79**, 69-116.
- 577 Fujisaka S, Castilla C, Escobar G, Rodrigues V, Veneklaas EJ, Thomas R, Fisher M (1998)
578 The effects of forest conversion on annual crops and pastures: Estimates of carbon
579 emissions and plant species loss in a Brazilian Amazon colony. *Agriculture*
580 *Ecosystems & Environment*, **69**, 17-26.
- 581 Gifford RM, Roderick ML (2003) Soil carbon stocks and bulk density: spatial or cumulative
582 mass coordinates as a basis of expression? *Global Change Biology*, **9**, 1507-1514.
- 583 Goidts E, van Wesemael B, Crucifix M (2009) Magnitude and sources of uncertainties in soil
584 organic carbon (SOC) stock assessments at various scales. *European Journal of Soil*
585 *Science*, **60**, 723-739.
- 586 Graham MH, Haynes RJ, Meyer JH (2002) Soil organic matter content and quality: effects of
587 fertilizer applications, burning and trash retention on a long-term sugarcane
588 experiment in South Africa. *Soil Biology & Biochemistry*, **34**, 93-102.
- 589 Guo LB, Gifford RM (2002) Soil carbon stocks and land use change: a meta analysis. *Global*
590 *Change Biology*, **8**, 345-360.
- 591 Hartemink AE (1997) Soil fertility decline in some major soil groupings under permanent
592 cropping in Tanga region, Tanzania. *Geoderma*, **75**, 215-229.
- 593 Houghton RA (1999) The annual net flux of carbon to the atmosphere from changes in land
594 use 1850-1990. *Tellus Series B-Chemical and Physical Meteorology*, **51**, 298-313.
- 595 Houghton RA (2003) Revised estimates of the annual net flux of carbon to the atmosphere
596 from changes in land use and land management 1850-2000. *Tellus Series B-Chemical*
597 *and Physical Meteorology*, **55**, 378-390.
- 598 Hughes RF, Kauffman JB, Cummings DL (2000) Fire in the Brazilian Amazon 3. Dynamics
599 of biomass, C, and nutrient pools in regenerating forests. *Oecologia*, **124**, 574-588.

- 1
2
3 600 IPCC (2007) *IPCC Special Report on Land Use, Land-Use Change And Forestry*. University
4
5 601 Press, Cambridge.
6
7
8 602 Juo ASR, Manu A (1996) Chemical dynamics in slash-and-burn agriculture. *Agriculture*
9
10 603 *Ecosystems & Environment*, **58**, 49-60.
11
12 604 Kindermann G, Obersteiner M, Sohngen B, *et al.* (2008) Global cost estimates of reducing
13
14 605 carbon emissions through avoided deforestation. *Proceedings of the National*
15
16 606 *Academy of Sciences of the United States of America*, **105**, 10302-10307.
17
18 607 Lal R (1998) Land use and soil management effects on soil organic matter dynamics on
19
20 608 alfisols in western Nigeria. *Soil Processes and the Carbon Cycle*, 109-126.
21
22 609 Lal R (2003) Soil erosion and the global carbon budget. *Environment International*, **29**, 437-
23
24 610 450.
25
26 611 Lal R, Kimble JM (2001) Importance of Soil Bulk Density and Methods of Its Importance. In
27
28 612 *Assessment methods for soil carbon* (eds Lal R, Kimble JM, Follett RF, Stewart BA),
29
30 613 pp. 31-44. Lewis Publishers, London.
31
32 614 Lee JH, Hopmans JW, Rolston DE, Baer SG, Six J (2009) Determining soil carbon stock
33
34 615 changes: simple bulk density corrections fail. *Agriculture, Ecosystems & Environment*,
35
36 616 **134**, 251-256.
37
38 617 Lugo AE, Brown S (1993) Management of Tropical Soils as Sinks or Sources of Atmospheric
39
40 618 Carbon. *Plant and Soil*, **149**, 27-41.
41
42 619 Malhi Y, Baldocchi DD, Jarvis PG (1999) The carbon balance of tropical, temperate and
43
44 620 boreal forests. *Plant Cell and Environment*, **22**, 715-740.
45
46 621 Mann LK (1986) Changes in Soil Carbon Storage after Cultivation. *Soil Science*, **142**, 279-
47
48 622 288.
49
50 623 Nye PH, Greenland DJ (1964) Changes in the soil after clearing tropical forest. *Plant and*
51
52 624 *Soil*, **21**, 101-112.
53
54
55
56
57
58
59
60

- 625 Paul KI, Polglase PJ, Nyakuengama JG, Khanna PK (2002) Change in soil carbon following
626 afforestation. *Forest Ecology and Management*, **168**, 241-257.
- 627 Paul S, Flessa H, Veldkamp E, Lopez-Ulloa M (2008) Stabilization of recent soil carbon in
628 the humid tropics following land use changes: evidence from aggregate fractionation
629 and stable isotope analyses. *Biogeochemistry*, **87**, 247-263.
- 630 Paustian K, Andren O, Janzen HH, *et al.* (1997) Agricultural soils as a sink to mitigate CO₂
631 emissions. *Soil Use and Management*, **13**, 230-244.
- 632 Penman J, Gytarsky M, Hiraishi T, *et al.* (2003) In *Intergovernmental Panel on Climate*
633 *Change, National Greenhouse Gas Inventories Programme*. Institute for Global
634 Environmental Strategies, Kanagawa, Japan.
- 635 Post WM, Kwon KC (2000) Soil carbon sequestration and land-use change: processes and
636 potential. *Global Change Biology*, **6**, 317-327.
- 637 Rasse DP, Rumpel C, Dignac MF (2005) Is soil carbon mostly root carbon? Mechanisms for a
638 specific stabilisation. *Plant and Soil*, **269**, 341-356.
- 639 Riezebos HT, Loerts AC (1998) Influence of land use change and tillage practice on soil
640 organic matter in southern Brazil and eastern Paraguay. *Soil & Tillage Research*, **49**,
641 271-275.
- 642 Sa JCdM, Cerri CC, Dick WA, Lal R, Venske Filho SP, Piccolo MC, Feigl BE (2001)
643 Organic matter dynamics and carbon sequestration rates for a tillage chronosequence
644 in a Brazilian Oxisol. *Soil Science Society of America Journal*, **65**, 1486-1499.
- 645 Siband P (1974) Evolution of soil characters and fertility in a red soil in Casamance.
646 *Agronomie Tropicale*, **29**, 1228-1248.
- 647 Silver WL, Ostertag R, Lugo AE (2000) The potential for carbon sequestration through
648 reforestation of abandoned tropical agricultural and pasture lands. *Restoration*
649 *Ecology*, **8**, 394-407.

- 650 Six J, Feller C, Denef K, Ogle SM, Sa JCD, Albrecht A (2002) Soil organic matter, biota and
651 aggregation in temperate and tropical soils - Effects of no-tillage. *Agronomie*, **22**, 755-
652 775.
- 653 Solomon D, Lehmann J, Kinyangi J, *et al.* (2007) Long-term impacts of anthropogenic
654 perturbations on dynamics and speciation of organic carbon in tropical forest and
655 subtropical grassland ecosystems. *Global Change Biology*, **13**, 511-530.
- 656 Soussana JF, Allard V, Pilegaard K, *et al.* (2007) Full accounting of the greenhouse gas (CO₂,
657 N₂O, CH₄) budget of nine European grassland sites. *Agriculture Ecosystems &*
658 *Environment*, **121**, 121-134.
- 659 Trumbore SE (1993) COMPARISON OF CARBON DYNAMICS IN TROPICAL AND
660 TEMPERATE SOILS USING RADIOCARBON MEASUREMENTS. *Global*
661 *Biogeochemical Cycles*, **7**, 275-290.
- 662 van der Werf GR, Morton DC, DeFries RS, *et al.* (2009) CO₂ emissions from forest loss.
663 *Nature Geoscience*, **2**, 737-738.
- 664 van Noordwijk M, Cerri C, Woomer PL, Nugroho K, Bernoux M (1997) Soil carbon
665 dynamics in the humid tropical forest zone. *Geoderma*, **79**, 187-225.
- 666 Van Oost K, Quine TA, Govers G, *et al.* (2007) The impact of agricultural soil erosion on the
667 global carbon cycle. *Science*, **318**, 626-629.
- 668 Veldkamp E (1994) Organic-Carbon Turnover in 3 Tropical Soils under Pasture after
669 Deforestation. *Soil Science Society of America Journal*, **58**, 175-180.
- 670 Veldkamp E, Becker A, Schwendenmann L, Clark DA, Schulte-Bisping H (2003) Substantial
671 labile carbon stocks and microbial activity in deeply weathered soils below a tropical
672 wet forest. *Global Change Biology*, **9**, 1171-1184.
- 673 Wairiu M, Lal R (2003) Soil organic carbon in relation to cultivation and topsoil removal on
674 sloping lands of Kolombangara, Solomon Islands. *Soil & Tillage Research*, **70**, 19-27.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

675 Zinn YL, Lal R, Bigham JM, Resck DVS (2007) Edaphic controls on soil organic carbon
676 retention in the Brazilian Cerrado: Texture and mineralogy. *Soil Science Society of*
677 *America Journal*, **71**, 1204-1214.

678 Zinn YL, Lal R, Resck DVS (2005) Changes in soil organic carbon stocks under agriculture
679 in Brazil. *Soil & Tillage Research*, **84**, 28-40.

680

681

For Review Only

List of Tables

Table 1: Mean absolute and relative SOC stocks changes and reported minimum and maximum relative SOC stock changes for different land use change types. Additionally, SOC stocks before land use change, mean full available sampling depth (on average 32 cm), mean time interval between the two land use systems and the number of studies included in this meta analysis is displayed with standard error of the mean in brackets.

Land use change (LUC) type	Absolute SOC change	Relative SOC change			SOC prior LUC	Sampling depth	Time after LUC	Number of studies
	Mg ha ⁻¹	%	Min %	Max %	Mg ha ⁻¹	cm	Years	
Primary forest to grassland	-12.6 (±3.0)	-12.1 (±2.3)	-73	51	73 (±7)	36 (±3)	25 (±3)	93
Primary forest to cropland	-20.1 (±5.2)	-25.2 (±3.3)	-80	58	83 (±9)	36 (±4)	28 (±4)	56
Primary forest to perennial crops	-32.0 (±3.5)	-30.3 (±2.7)	-62	6	105 (±20)	48 (±8)	49 (±12)	20
Primary forest to secondary forest	-12.6 (±2.4)	-8.6 (±2.0)	-64	72	91 (±9)	39 (±4)	28 (±3)	71
Secondary forest to grassland	-11.0 (±3.4)	-6.4 (±2.5)	-71	72	85 (±6)	43 (±3)	27 (±2)	66
Secondary forest to cropland	-25.8 (±6.9)	-21.3 (±4.1)	-74	53	88 (±12)	39 (±5)	36 (±7)	26
Second. forest to perennial crops	-5.6 (±3.0)	-2.4 (±4.2)	-46	243	90 (±17)	51 (±9)	23 (±4)	15
Grassland to secondary forest	12.4 (±6.1)	17.5 (±8.0)	-35	282	60 (±9)	35 (±6)	28 (±4)	32
Cropland to secondary forest	33.2 (±10.5)	50.3 (±11.9)	-63	67	70 (±9)	44 (±6)	32 (±7)	25
Grassland to cropland	-6.0 (±5.7)	-10.4 (±6.1)	-41	167	64 (±15)	38 (±11)	22 (±5)	15
Cropland to grassland	7.6 (±5.8)	25.7 (±11.1)	-32	362	61 (±17)	40 (±10)	21 (±6)	16
Cropland to fallow	8.9 (±2.9)	32.2 (±16.1)	-73	51	43 (±7)	20 (±2)	≤ 7	21

1
2
3 689 Table 2: Effect of mass correction on SOC stock change estimates and relative changes in
4
5 690 bulk density after land use change per land use change type for studies reporting bulk density
6
7
8 691 data ($\pm$ SE in brackets).
9

Land use change type	SOC stock changes [Mg ha ⁻¹]		Relative bulk
	with mass correction	no mass correction	density changes [%]
Primary forest to grassland	-12.1 ($\pm$ 2.3)	-4.9 ($\pm$ 2.5)	14.0 ($\pm$ 2.2)
Primary forest to cropland	-25.2 ($\pm$ 3.3)	-22.3 ($\pm$ 3.1)	17.8 ($\pm$ 3.5)
Primary forest to perennial crops	-30.3 ($\pm$ 2.7)	-23.2 ($\pm$ 2.7)	22.8 ($\pm$ 6.2)
Primary forest to secondary forest	-8.6 ($\pm$ 2)	-6.7 ($\pm$ 2.1)	5.7 ($\pm$ 2.7)
Secondary forest to grassland	-6.4 ($\pm$ 2.5)	-4.1 ($\pm$ 2.6)	5.4 ($\pm$ 2.3)
Secondary forest to cropland	-21.3 ($\pm$ 4.1)	-19.2 ($\pm$ 4.1)	11.6 ($\pm$ 4.4)
Grassland to secondary forest	17.5 ($\pm$ 8.0)	13.1 ($\pm$ 8.3)	-6.4 ($\pm$ 3.8)
Cropland to secondary forest	50.3 ($\pm$ 11.9)	42.9 ($\pm$ 11.5)	5.0 ($\pm$ 5.4)
Grassland to cropland	-10.4 ($\pm$ 6.1)	-8.8 ($\pm$ 6.2)	-2.7 ($\pm$ 5.8)
Cropland to grassland	25.7 ($\pm$ 11.1)	25.5 ($\pm$ 10.4)	1.9 ($\pm$ 5.3)
Cropland to fallow	32.2 ($\pm$ 16.1)	27.3 ($\pm$ 15.8)	14.0 ($\pm$ 2.2)

Table 3: General linear model with degrees of freedom (Df), sum of squares (sum of Sq), F-value, P-value. LUC= land use change type, MAT= Mean annual temperature, MAP= Mean annual precipitation, max depth= maximum sampling depth [cm], significance codes: '***' <0.001, '**' 0.001, '*' 0.05, 'n.s.' not significant.

Models	Df	Sum of Sq	F	P-value		Explained variance [%]
rel. SOC change ~ LUC	5	113095	38.39	<0.000	***	23
rel. SOC change ~ LUC+MAT	1	992	1.68	0.196	n.s.	
rel. SOC change ~ LUC * MAT	6	15093	4.27	<0.000	***	33
rel. SOC change ~ LUC * MAT * MAP	12	16108	2.27	0.009	**	42
rel. SOC change ~ LUC * MAT * MAP *						55
Clay mineral type	34	44853	2.24	<0.000	***	
rel. SOC change ~ LUC	5	117882	29.23	<0.000	***	
rel. SOC change ~ LUC + max depth	1	213	0.26	0.608	n.s.	
rel. SOC change ~ LUC * max depth	5	14199	1.95	1.955	n.s.	
rel. SOC stock change ~ LUC	4	141916	20.35	<0.000	***	
rel. SOC stock change ~ LUC + Soil type	9	10463	0.67	0.739	n.s.	
rel. SOC stock change ~ LUC * Soil type	45	42750	0.28	0.280	n.s.	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 4: Relative SOC stocks changes ($\pm$ standard error) for different climate and soil conditions derived from a general linear model (Tab. 3).

Land use change	MAT [°C]	MAP [mm]	rel. SOC change
Forest to grassland	20	1000	-1.2 (± 1.3)
		2000	-2.7 (± 0.2)
		4000	-5.6 (± 0.1)
	23	1000	-4.2 (± 1.5)
		2000	-6.4 (± 3.6)
		4000	-10.8 (± 5.8)
	26	1000	-7.2 (± 3.1)
		2000	-10.2 (± 6.2)
		4000	-16.0 (± 9.3)
Forest to cropland	20	1000	-29.5 (± 19.3)
		2000	-25.8 (± 17.0)
		4000	-18.2 (± 6.5)
	23	1000	-29.9 (± 20.4)
		2000	-28.4 (± 19.6)
		4000	-25.2 (± 14.7)
	26	1000	-30.3 (± 19.9)
		2000	-30.9 (± 21.3)
		4000	-32.2 (± 20.3)

MAT: Mean annual temperature

MAP: Mean annual precipitation

Table 5: Fraction of original soil carbon stock for 0-30 cm depth remaining after land use change. Revised default values for tropical regions from the IPCC Good Practice Guidelines (GPG) LULUCF (Penman *et al.*, 2003) and from this meta-analysis.

SOC stock change	Climate regime	This meta analysis	Uncertainty [%]	Revised GPG default	Error [%]
LUC native vegetation to cropland	Dry	0.76	2	0.69	38
LUC native vegetation to cropland	Wet	0.68	7	0.58	42
LUC native vegetation to grassland		0.91	3	1	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

List of Figures

Figure 1: Weighted average relative bulk density change [%] for different soil depth for different land use change types derived from all studies reporting bulk density measurements.

Figure 2: Absolute SOC stock changes [Mg C ha⁻¹] for different soil depth for different land use change types. The different depth increments are covered by different numbers of studies.

Figure 3: Relative SOC stock changes [%] for different soil depth for different land use change types. The different depth increments are covered by different numbers of studies.

Figure 4: SOC stock change [Mg ha⁻¹] after conversion to grassland (open symbols) and cropland (filled symbols) vs. Mean Annual Precipitation [mm].

Figure 5: SOC stock change [Mg ha⁻¹] after conversion to grassland (filled symbols) and cropland (open symbols) vs. mean content clay of the soil [%].

Weighted average relative bulk density change [%] for different soil depth for different land use change types derived from all studies reporting bulk density measurements.

169x96mm (600 x 600 DPI)

Absolute SOC stock changes [Mg C ha⁻¹] for different soil depth for different land use types.
The different depth increments are covered by different numbers of studies.
186x105mm (600 x 600 DPI)

Relative SOC stock changes [%] for different soil depth for different land use change types. The different depth increments are covered by different numbers of studies.
173x104mm (600 x 600 DPI)

SOC stock change [Mg ha⁻¹] after conversion to grassland (open symbols) and cropland (filled symbols) vs. Mean Annual Precipitation [mm].
171x150mm (600 x 600 DPI)

SOC stock change [Mg ha^{-1}] after conversion to grassland (filled symbols) and cropland (open symbols) vs. mean content clay of the soil [%].
154x139mm (600 x 600 DPI)