Direct and indirect effects of nitrogen deposition on species composition change in calcareous grasslands

Leon JI van den Berg, Philippine Vergeer, Tim Cg Rich, Simon M Smart, Dan Guest, Mike R Ashmore

► To cite this version:

Leon JI van den Berg, Philippine Vergeer, Tim Cg Rich, Simon M Smart, Dan Guest, et al.. Direct and indirect effects of nitrogen deposition on species composition change in calcareous grasslands. Global Change Biology, 2010, 17 (5), pp.1871. 10.1111/j.1365-2486.2010.02345.x . hal-00599530

HAL Id: hal-00599530

<https://hal.science/hal-00599530>

Submitted on 10 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Direct and indirect effects of nitrogen deposition on species composition change in calcareous grasslands

Journal:	<i>Global Change Biology</i>
Manuscript ID:	GCB-10-0584
Wiley - Manuscript type:	Primary Research Articles
Date Submitted by the Author:	16-Jul-2010
Complete List of Authors:	van den Berg, Leon; Radboud University Nijmegen, Aquatic Ecology and Environmental Biology; University of York, Environment Vergeer, Philippine; Radboud University Nijmegen, Department of Ecogenomics Rich, Tim; National Museum Wales, Dept. Biodiversity & Systematic Biology Smart, Simon; CEH, Lancaster Guest, Dan; Countryside Council for Wales Ashmore, Mike R; University of York, Environment
Keywords:	Nitrogen deposition, species diversity, soil acidity, Festuco Brometalia grasslands, management
Abstract:	<p>Atmospheric nitrogen (N) deposition has been identified as a major threat to biodiversity, but field surveys of its effects have rarely focussed on sites which are actively managed to maintain characteristic species. We analysed permanent quadrat data from 106 plots in nature reserves on calcareous grassland sites in the UK collected during a survey between 1990 and 1993 and compared the data with the results from re-survey of 48 of these plots between 2006 and 2009.</p> <p>N deposition showed no significant spatial association with species richness, species diversity, or the frequency of species adapted to low nutrient conditions in the 1990-1993 dataset. However, temporal analysis showed that N deposition was significantly associated with changes in Shannon diversity and evenness. In plots with high rates of N deposition, there was a decrease in species diversity and evenness, a decline in the frequency of characteristic calcareous grassland species, and a lower number of rare and scarce species. Since all sites had active management to maintain a high diversity and characteristic species, our results imply that even focussed management on nature conservation objectives cannot prevent adverse effects of high rates of N deposition.</p> <p>Structural equation modelling was used to compare different causal</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	mechanisms to explain the observed effects. For change in Shannon diversity, direct effects of N deposition were the dominant mechanism and there was an independent effect of change in grazing intensity. In contrast, for change in herb species number, indirect effects on soil acidity, linked to both N and S deposition, were more important than direct effects of N deposition.

SCHOLARONE™
Manuscripts

For Review Only

Title:

Direct and indirect effects of nitrogen deposition on species composition change in calcareous grasslands

Running Title:

Ndeposition effects on calcareous grasslands

Leon JL van den Berg^{1,2*}, Philippine Vergeer³, Tim CG Rich⁴, Simon M Smart⁵, Dan Guest⁶, Mike R Ashmore¹

*Corresponding author

¹Environment Department, University of York, York YO10 5DD, United Kingdom

^{2*}*current address*: Aquatic Ecology and Environmental Biology, Radboud University Nijmegen, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands Tel: +0031- 24 3652337, email: l.vandenberg@science.ru.nl

³Department of Ecogenomics, Radboud University Nijmegen, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands

⁴Welsh National Herbarium (NMW), Dept. Biodiversity & Systematic Biology, National Museum Wales, Cardiff CF10 3NP, United Kingdom

⁵Centre for Ecology and Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster LA1 4AP, United Kingdom

⁶Cyngor CefnGwladCymru/ Countryside Council for Wales, FforddPenrhos, Bangor Gwynedd LL57 2BX, United Kingdom

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Keywords**

2 Nitrogen deposition, species diversity, soil acidity, *FestucoBrometalia* grasslands,

3 management

4

For Review Only

Abstract

Atmospheric nitrogen (N) deposition has been identified as a major threat to biodiversity, but field surveys of its effects have rarely focussed on sites which are actively managed to maintain characteristic species. We analysed permanent quadrat data from 106 plots in nature reserves on calcareous grassland sites in the UK collected during a survey between 1990 and 1993 and compared the data with the results from re-survey of 48 of these plots between 2006 and 2009.

N deposition showed no significant spatial association with species richness, species diversity, or the frequency of species adapted to low nutrient conditions in the 1990-1993 dataset.

However, temporal analysis showed that N deposition was significantly associated with changes in Shannon diversity and evenness. In plots with high rates of N deposition, there was a decrease in species diversity and evenness, a decline in the frequency of characteristic calcareous grassland species, and a lower number of rare and scarce species. Since all sites had active management to maintain a high diversity and characteristic species, our results imply that even focussed management on nature conservation objectives cannot prevent adverse effects of high rates of N deposition.

Structural equation modelling was used to compare different causal mechanisms to explain the observed effects. For change in Shannon diversity, direct effects of N deposition were the dominant mechanism and there was an independent effect of change in grazing intensity. In contrast, for change in herb species number, indirect effects on soil acidity, linked to both N and S deposition, were more important than direct effects of N deposition.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Introduction

2
3
4
5
6
7
8 Atmospheric deposition of nitrogen (N) and sulphur (S) compounds increased dramatically in
9
10 Europe in the second half of the 20th century, mainly as a result of increased fossil fuel
11
12 combustion and intensive agriculture (Galloway, 1995). Despite a strong decline in S
13
14 emissions and deposition in most parts of Europe since the 1990's (Vestreng *et al.* 2007), N
15
16 deposition has not declined to the same extent. Both N and S deposition has continued to
17
18 increase in other parts of the world, especially in parts of India and China (Galloway &
19
20 Cowling, 2002; Dentener *et al.* 2006).
21
22
23
24
25
26
27 N deposition can result in severe eutrophication via the direct input of nutrients but also
28
29 acidification via indirect processes such as nitrification (Galloway, 1995; Bobbink *et al.*
30
31 1998). N deposition has been identified as one of the most important global drivers of
32
33 biodiversity change in terrestrial ecosystems (Vitousek *et al.* 1997; Sala *et al.* 2000, Phoenix *et*
34
35 *al.* 2006), and has been linked to changes in plant species composition in a wide range of
36
37 terrestrial biomes from the Arctic to the tropics (e.g. Bobbink *et al.* 2010). Some of the most
38
39 affected terrestrial habitats are nutrient-poor semi-natural habitats such as heathlands and
40
41 grasslands. Calcareous grasslands are floristically amongst the most species rich communities
42
43 of North-Western Europe, containing many rare and endangered plant species (e.g. Willems,
44
45 1990). Their biodiversity is threatened by changes in management (e.g. Tansley, 1922;
46
47 Willems, 1978), habitat fragmentation (e.g. Fisher & Stöcklin, 1997) and disturbance (e.g.
48
49 Smart *et al.* 2003a), among other factors.
50
51
52
53
54
55
56
57
58 Experimental studies in Dutch calcareous grasslands have shown that increased N deposition
59
60 results in the loss of characteristic plant species and an increase of highly competitive grass

species such as *Brachypodium pinnatum* (e.g. Willems 1990; Bobbink *et al.* 1998). In the UK, changes in species composition, a reduction in flowering, and a reduction in the forb/grass ratio have been shown in a long-term N-addition experiment (Carroll *et al.* 2003), but encroachment by *B. pinnatum* or other highly competitive grasses such as *Bromus erectus* was not observed, possibly due to co-limitation by phosphorus (Morecroft *et al.* 1994; Wilson *et al.* 1995; Carroll *et al.* 2003; Phoenix *et al.* 2003).

Stevens *et al.* (2004) found a significant negative spatial relationship between N deposition and species richness in a national survey of one acidic grassland community in the UK. More recently, analysis of data from a national survey covering all plant communities in the UK corroborated these findings for both heathlands and acid grasslands, but failed to find a significant relationship between N deposition and species richness in calcareous grasslands, even when interacting environmental factors such as rainfall and altitude were included in the models (Maskell *et al.* 2010). However, significantly more species characteristic of fertile soil conditions, and a higher grass/forb ratio, was found in calcareous grassland plots with higher N deposition. These findings suggest that the responses of calcareous grasslands to increased N deposition differ from those of other semi-natural ecosystems, such as acid grasslands and heathlands. This may be due to several factors, including (a) the importance of co-limitation by phosphorus, (b) active management and conservation to protect characteristic plant species and (c) the occurrence on shallow well buffered soils that are less sensitive to acidification caused by N and S deposition.

In addition, the limitations of the studies undertaken to date need to be considered.

Experimental studies measure change over a relatively short period of time with N deposition as the only variable factor. Field surveys which explore the spatial associations between N

1
2
3 1 deposition and species richness, or other response variables, do not allow an assessment of
4
5 2 whether these are due to long-term historical deposition or recent changes in deposition, such
6
7
8 3 as those simulated in experimental studies. Although long-term vegetation change in
9
10 4 calcareous grasslands in the UK since 1950 has been described by Bennie *et al.* (2006), their
11
12 5 analysis was not designed to assess the effect of N deposition, and there are no field surveys
13
14 6 relating recent changes in species composition or species numbers to atmospheric N
15
16 7 deposition, using plots that can be relocated with confidence. A further limitation to the
17
18 8 interpretation of field survey data is the multi-collinearity between predictor variables such as
19
20 9 N deposition and climate variables, and earlier studies have highlighted the limitations that
21
22 10 multiple regression or covariance analysis have in predicting the relative effects of the
23
24 11 explanatory variables on dependent variables such as species richness (e.g. Stevens *et al.*
25
26 12 2004, Maskell *et al.* 2010).
27
28
29
30
31
32
33
34 14 We investigated the direct, indirect and interacting effects of atmospheric N deposition, other
35
36 15 confounding environmental variables, and site management on species richness and species
37
38 16 diversity in calcareous grasslands in the UK. We based our study on a network of 128
39
40 17 permanently marked plots in areas of calcareous grassland spread across the UK that were
41
42 18 specifically established in the early 1990s, with the aim of monitoring long-term impacts of
43
44 19 climate change and air pollution, by John Rodwell and colleagues (Rich *et al.* 1993). Between
45
46 20 2006 and 2009, 48 plots were revisited and additional soil measurements were made. This
47
48 21 allowed us to assess the effect of N deposition and other factors on both spatial variation and
49
50 22 recent temporal changes in species richness and other relevant variables.
51
52
53
54
55 23
56
57 24 In order to test if changes in species richness and composition are significantly affected by
58
59 25 atmospheric N deposition in a matrix of effects of factors such as management, climate, S

deposition and soil conditions, we analysed the data using both mixed effect models and structural equation modelling (SEM). The SEM models allow explanatory variables to be placed in 'series' rather than 'parallel', using sequential arrangements that reflect hypothetical causal chains linking the effects. Such models provide insights that are not possible with standard covariance analyses into the relative importance of different direct and indirect effects of N deposition.

We tested three major hypotheses:

1) Ndeposition is a significant explanatory driver of spatial variation in species richness, species diversity and the frequency of individual species in calcareous grasslands.

2) Ndeposition is a significant explanatory driver of change over the last 15 years in species richness, species diversity and the frequency of individual species in calcareous grasslands.

3) Recent changes in species richness, species diversity and the frequency of individual species in calcareous grasslands can be explained by a combination of the effects of management, direct effects of N deposition, and indirect effects of S and N deposition via soil acidification.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Materials and Methods

Calcareous grassland survey: 1990-1993 survey

A large national survey of calcareous grasslands was carried out between 1990 and 1993 (Rich *et al.* 1993). The survey method was specifically designed to be repeatable with a minimum of observer bias. All the selected sites were nature reserves with active management to maintain species diversity and characteristic species; 56 sites were selected, with 1-6 plots on each site, resulting in a total of 128 plots. Sites were chosen to cover a wide geographical range (Fig. 1), a large climatic gradient, a wide range in deposition of air pollutants and all types of calcareous grasslands known in the UK, using the National Vegetation Classification (NVC) (Rodwell, 1998). On most sites, plots were chosen on both south and north-facing slopes to account for effects of aspect. Each plot covered an area of 144 m² (typically measuring 12×12 m or more rarely 6×24 m) and was established in a floristically homogeneous stand of calcareous grassland. Individual plots were permanently marked at unique places with buried copper coils. Detailed field notes, describing the exact location of the plots, photographs, sketch maps and measurements from fixed features were also recorded, to ensure successful relocation of each plot in future surveys. The presence/absence of all vascular plants was recorded in 36 50×50 cm quadrats positioned within each permanent plot. Soil pH of the top 10 cm was determined in the field by mixing soil into a slurry with distilled water (1:1 vol:vol), equilibrating for 5 minutes, and measuring using a calibrated pH meter. Soil type, soil depth, slope, aspect and sward height were recorded for each plot. For each site, the size, its protection status and detailed information on local management (grazing intensity) were noted. Data from the original reports could not be recovered for 7 plots (4 sites), resulting in available data for 121 plots.

1

2
3
4
5
6
7 2006-2009 survey

8
9
10 3 In 2006, 35 out of the 56 sites were selected for re-survey (48 plots), covering a wide range of
11
12 4 Ndeposition. Sites were selected to reduce the effects of spatial autocorrelation in atmospheric
13
14 5 N deposition (e.g. by selecting high and low Ndeposition sites at similar latitudes). Sites and
15
16 6 plots with NVC communities CG1, CG2, CG10, CG11 and CG12 were targeted where
17
18 7 possible to reduce variation due to initial vegetation composition. Only six of the plots selected
19
20 8 for re-survey were identified as supporting different NVC communities (CG3, CG8, CG9),
21
22 9 mainly due to the presence of grasses such as *Bromus erectus* (CG3) or *Sesleria albicans*
23
24 10 (CG8&9). In broad terms, plots in the lowlands of Wales and England focussed on stands of
25
26 11 CG1 and CG2, while those in the uplands of Wales and Scotland focussed on stands of
27
28 12 CG10, CG11 or CG12.

29
30
31
32
33
34 13 Metal detectors were used for relocation of the plots. In each plot, 36 quadrats were
35
36 14 positioned in exactly the same pattern as in 1990-1993 and in each quadrat the
37
38 15 presence/absence of all vascular plants was recorded. One soil sample, composed of 5 sub-
39
40 16 samples, was taken from the top 10 cm of each plot using an auger (3 cm diameter). Soil
41
42 17 samples were stored at 4 °C until further analysis. Sward height and soil depth were recorded
43
44 18 at each plot and detailed information on the local management (grazing intensity) for the past
45
46 19 15 years was obtained from local managers at the plot level.

50
51
52 2053
54
55 21 Soil analysis56
57
58 22 Soil samples from the 2006-2009 survey were thoroughly mixed prior to extraction.59
60 23 Extractions were performed on fresh soil and corrected for moisture content after drying for

1
2
3 1 24h at 105°C. A portion of 35g of fresh soil was mixed with 100ml bi-distilled water for
4
5 2 water soluble extraction or 100ml 0.2M NaCl for the exchangeable fraction. Extracts were
6
7 3 shaken for 1h at 100rpm after which the pH was measured in the water fraction with a
8
9 4 Radiometer type PHM 82 pH meter. Extracts were filtered (0.2µm) and stored at 4°C until
10
11 5 analysis. Plant available P was measured after Olsen-P extraction (Olsen *et al.* 1954); 5 g of
12
13 6 fresh soil was mixed with 100 ml NaHCO₃ and shaken for 30 minutes at 100 rpm. Extracts
14
15 7 were filtered (0.2µm) and stored at 4°C until analysis. NO₃⁻ and NH₄⁺ were analysed using a
16
17 8 Bran and LuebbeAutoanalyser 3 (Bran and Luebbe, Norderstedt, Germany). Concentrations
18
19 9 of base cations (Ca²⁺, Na⁺, Mg²⁺ and K⁺) and Al³⁺ and plant available P were determined in
20
21 10 the NaCl and Olsen extracts respectively using an ICP Spectrometer (IRIS Intrepid II,
22
23 11 Thermo Electron Corporation, Franklin, MA). Organic matter contents of the soils were
24
25 12 determined by loss on ignition (LOI, 4h, 550°C).
26
27
28
29
30
31
32
33

34 14 *Plant attributes*

35
36 15 The names of all plant species in the 1990-1993 survey were checked and if necessary
37
38 16 updated to enable a comparison with the recent 2006-2009 survey. All data werethen linked
39
40 17 to the attribute list PLANTATT (Hill *et al.* 2004).Stace (1997) was followed for
41
42 18 nomenclature. We used Ellenberg indicator values modified for the UK (Hill *et al.* 2004) to
43
44 19 characterise plant species. For each plot the average, frequency weighted, Ellenberg value
45
46 20 was calculated in order to characterise plots according to the species they contained. Species
47
48 21 were grouped into eutrophic (Ellenberg N>5), oligotrophic (Ellenberg N<3), acidophobic
49
50 22 (Ellenberg R>5), and acidophilic (Ellenberg R<3) species. The total number of scarce and
51
52 23 rare species, as denoted in the PLANTATT database, was recorded for each plot.
53
54 24 Speciesrichness (S) was recorded as the total number of vascular plant species in each plot.
55
56 25 For each plot the species diversity was calculated as the Shannon index (H'): $H' = -\sum(p_i \ln p_i)$,
57
58
59
60

where p_i is the number of occurrences of the i^{th} species in the 36 quadrats expressed as a proportion of the total number of occurrences of all species in those quadrats (Magurran 2004). In addition, Shannon's equitability or the Evenness (E_H) was calculated as $E_H = H'/\ln(S)$. For each species the frequency of occurrence was calculated as $n_i/36$, where n_i is the number of quadrats in which the species occurred in plot i .

Survey data selection

In the soils of the plots revisited in 2006-9, pH varied widely, ranging from shallow rendzinas of pH 8.5 to peaty soils overlaying calcareous rock of pH 4.2. The six plots with the lowest pH had in general very different vegetation (e.g. more acidophilic species such as *Calluna vulgaris* or *Juncus effusus*), a very high soil organic matter and low base cation content. In order to relate atmospheric N deposition and other environmental parameters to species composition and richness in calcareous grasslands which share similar characteristics and to reduce confounding effects, only those plots with a soil $\text{pH}_{(1990-1993)} > 6.0$ were included in the analysis. This resulted in 106 plots from the 1990-1993 data and 42 plots which could be used for the comparison between 1990-1993 and 2006-2009. For analysis at the individual species level, changes in species frequency were analysed on a subset of the data; only those species which occurred in at least 11 (25%) or more of the 42 plots in both surveys were included. This resulted in a dataset for only 52 species. Grass to herb ratio was calculated as the ratio between the numbers of grass and herb species at the plot level.

Plot and site specific parameters

Modelled N and S deposition data for each site were obtained from the Centre of Ecology and Hydrology (CEH) for two periods: 1993-1995 and 2006-2008 for the surveys of 1990-1993

1
2
3 1 and 2006-2009 respectively. The Ndeposition ranged from 7.3 to 40.7 kg N ha⁻¹ y⁻¹ while
4
5 2 Sdeposition ranged from 8.0 to 27.6 kg S ha⁻¹ y⁻¹. Climate data for each site were obtained
6
7 3 from the 5km gridded climate database held by the UK Meteorological Office
8
9 4 (www.metoffice.co.uk). 10 year averaged data for the periods 1981-1991 and 1997-2006
10
11 5 respectively were used for the 1990-1993 and 2006-2009 surveys, including mean minimum
12
13 6 winter temperature, mean maximum summer temperature, mean annual temperature and
14
15 7 mean annual rainfall. The aspect of each plot was converted to a 0-1 scale using
16
17 8 $\sin(a/(360/\pi))$, where a = the aspect in degrees. This resulted in a gradient from North =0 to
18
19 9 South =1, with East and West both at 0.5. Where information on stocking was available from
20
21 10 local site managers, grazing intensity for each plot was calculated in terms of livestock units
22
23 11 (LU) per hectare per year using the conversion factors provided in Attwood & Heavey (1964).
24
25 12 For the 9 plots where detailed stocking was not provided we estimated the grazing intensity
26
27 13 from the sward height, based on the highly significant correlation ($r^2=0.58$) between sward
28
29 14 height and livestock units obtained from the other plots.
30
31
32
33
34
35
36
37
38
39
40
41

42 16 *Statistical analysis*

43 17 Principal component analysis (PCA) in CANOCO (TerBraak, 1988) was used to determine
44
45 18 the variables explaining most of the variation between the plots. For each survey, we selected
46
47 19 the strongest, non-correlated, variables for further analysis.
48
49
50

51 21 Possible combinations of predictor variables were explored to assess the effects of
52
53 22 Ndeposition, soil conditions, climate data and grazing pressure on species richness and
54
55 23 species composition. Data from the 1990-1993 survey were used in the spatial analysis, while
56
57 24 data from both surveys were used in the temporal analysis. All models were explicitly tested
58
59
60

for spatial autocorrelation in the response variable and residuals by inspection of semi-variograms. In the presence of spatial autocorrelation, generalized linear mixed-effect models (GLMM) were used to control for this spatial autocorrelation. In these models, a correlation structure was added to correct for spatial autocorrelation. Correlation structures such as corExp, and corSpher were used with the “form=~Easting+Northing” argument in the correlation option to calculate the Euclidean distances (using Pythagoras theorem) between sites with coordinates given by Easting and Northing. The models were ranked according to the Akaike Information Criterion (AIC) and only the models and corresponding correlation structure with the lowest AIC, representing the ‘best fitting models’, were selected. When no spatial autocorrelation was observed, multiple linear regression models were used. Again, models were ranked according to AIC and only the models with the lowest AIC were selected. To test for multicollinearity, variance inflation factors (VIF) were computed for each predictor variable in the model. Predictor variables that were highly correlated ($VIF > 10$; Gujarati, 1995) were not analysed in the same model; in those situations only the predictor variable that explained most of the variance was selected.

For the change in Shannon diversity H' , the ecological significance of the effect was quantified by dividing the data into two groups, one with N deposition below the lowest, and one with N deposition above the highest, critical load of N deposition for this vegetation type (< 15 and > 25 kg N ha⁻¹ y⁻¹ respectively, Bobbink *et al.*, 2003). The Cohen d-value was then

$$d = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{(s_1^2 + s_2^2)}{2}}}$$

calculated according to: , where $\bar{x}_1$ and $\bar{x}_2$ are the means of the data for sites with high and low N deposition respectively and s_1 and s_2 the standard deviations in these groups. The calculated value of d was checked against Cohen’s classification (Cohen, 1988). We also compared the predicted change in H' from our models at sites with N

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 deposition above 25 kg ha⁻¹ y⁻¹ with the value of H' in a reference dataset representing an
2 highly undesirable endpoint that could be realised given continued high N deposition when
3 grazing and/or cutting is maintained. We selected botanical data (n=327) from within the
4 randomly located plots of the Countryside Survey(CS) of Great Britain which independent
5 land cover mapping showed could be referenced to the Neutral Grassland Broad Habitat
6 (Jackson 2000). The change in H' from 1990-93 to 2006-09 in the high N deposition plots
7 was then expressed as a percentage of the difference between the 1990-93 value and that of
8 the reference dataset, an approach similar to that used by Smart *et al.* (2003b).The analyses
9 were done in the 'R' (version 2.9.0) statistical and programming environment (R
10 Development Core Team, 2008).

11

12 Structural equation models (SEM) were used to further examine the relationships between N
13 deposition, soil conditions, climate data and grazing pressure on species richness and species
14 composition. The software package AMOS 17 was used to design the SEM models and
15 calculate path coefficients, squared multiple correlations, direct and indirect effects and
16 model fit. A significant χ^2 -statistic indicates that the SEM model provides a poor fit. χ^2 -
17 statistics and four non- χ^2 -distributed fitting functions (Comparative Fit Index, CFI; Normed
18 Fit Index, NFI; Tucker-Lewis Index, TLI; Incremental Fit Index, IFI) were compared in order
19 to test the goodness-of-fit of the models. In the SEM models, soil nutrient concentrations
20 were represented by "mineral N" (NH₄⁺ + NO₃⁻).

Results

Analysis of 1990-1993 survey

Linear regression between the total number of vascular plant species and Ndeposition did not reveal a significant relationship in the 1990-1993 dataset (Fig. 2a), but Ndeposition did show a slight positive relationship with the grass to herb ratio (Fig. 2b). Almost no scarce and rare species were observed at plots with Ndeposition above $25 \text{ kg N ha}^{-1} \text{ y}^{-1}$ (Fig. 2c); the average number of scarce and rare species was 1.6 in the 77 plots with deposition below $25 \text{ kg N ha}^{-1} \text{ y}^{-1}$ compared to 0.3 in the 29 plots with deposition above $25 \text{ kg N ha}^{-1} \text{ y}^{-1}$ (ANOVA, $df = 1$, $F = 13.62$, $p < 0.001$).

Minimum winter temperature and maximum summer temperature were highly correlated; only maximum summer temperature was included in the model, as this was the stronger predictor according to the PCA. For similar reasons, organic content of the soil, Olsen P and slope (which were all strongly correlated with pH) and mean annual rainfall (which was strongly correlated with maximum summer temperature) were excluded from the analysis. Grazing pressure entered the analysis, but was not a significant explanatory variable for any of the dependant parameters (Table 1a).

Maximum summer temperature and soil depth were the most important variables explaining variation in species composition and species richness in the 1990-1993 dataset (Table 1a). Spatial variation in species richness, the number of eutrophic species, the number of herb species, and H' could be explained solely by soil depth and maximum summer temperature, with positive relationships with both variables in all cases.

1
2
3 1 Ndeposition did not have a significant effect on speciesrichness or on the number of
4
5 2 eutrophic or oligotrophic species. The frequency weighted average Ellenberg N value was
6
7 3 also not significantly related toN deposition, but was best explained by the maximum summer
8
9 4 temperature.The only variable for which Ndeposition entered the final model was the grass to
10
11 5 herb ratio, for which a positive effect, albeit at the $p<0.1$ level,was recorded.A significant
12
13 6 negative effect of soil pH was found on the number of acidophilic species, in a model that
14
15 7 also included a negative effect of maximum summer temperature. The frequency weighted
16
17 8 Ellenberg R values increased with soil pH, in a model that also included a positive effect of
18
19 9 maximum summer temperature.
20
21
22
23
24
25

26
27 11 *Analysis of change between surveys*
28

29 12 The average soil pH in the 1990-1993 dataset was 7.41. The average soil pH in the more
30
31 13 recent survey wasmeasured more accurately and was 7.14. The significant difference (paired
32
33 14 t-test: $t=2.52$, $df=35$, $p=0.016$)was most likely due to different measurement methods and
34
35 15 cannot be interpreted in terms of environmental change.
36
37
38
39
40

41 17 A significant, but slight, decrease in speciesrichness, independent of Ndeposition, was
42
43 18 observed between 1990-1993 (average = 41) and 2006-2009 (average=39) (paired t-test $t=-$
44
45 19 2.274 , $p=0.028$, $df=41$).The change in both H' and E_H over this period was significantly
46
47 20 negatively related to Ndeposition,with a general increase in H' and E_H at low Ndeposition and
48
49 21 a decrease at high Ndeposition (Fig. 3).Ndeposition was not significantly related to the
50
51 22 change in any of the other dependent variables.
52
53
54
55
56

57 23
58 24 Instrong contrast to the spatial analysis for the 1990-93 data, maximum summer temperature
59
60 25 and aspect did not enter the best fit model for the change in any of the dependent variables

(Table 1b). In contrast to the spatial models, soil depth entered the models for change in number of oligotrophic, acidophilic and acidophobic species, and was never significant.

Soil pH showed a significant positive relationship with the change in the species richness, i.e. there tended to be an increase in plots with a higher soil pH. In addition, soil pH was associated positively with change in the number of herb species and negatively with change in grass to herb ratios. The mineral N content was also weakly positively associated with change in the number of herb species. However, the changes in the number of acidophilic species, the number of acidophobic species, and the cover weighted Ellenberg R index were not related significantly to soil pH or any other variable.

The models showed a significant negative effect of N deposition on changes in H' and evenness (Table 1b). However, N deposition was not significantly related to change in the number of eutrophic or oligotrophic species, or change in Ellenberg N index. The change in grazing was significantly negatively related to the change in frequency weighted Ellenberg N index, while the change in the number of oligotrophic species showed a positive relationship to the mineral N content of the soil.

Structural equation modelling

Structural equation models were used to further investigate the direct and indirect effects of N deposition for two variables for which change between 1990-1993 and 2006-2009 was not spatially auto-correlated: H' and number of herb species. Using maximum likelihood estimation, we observed a χ^2 of 4.488 for the model for H' ($df=7$, $p=0.722$) for a goodness-of-fit test of the null hypothesis that the covariance matrix implied by the (expected) model reproduces the observed covariance matrix. Failure to reject the null hypothesis indicates that

1
2
3 1 the model was a good fit. In addition, the CFI of 1.00, the NFI of 0.94, the TLI of 1.14 and the
4
5 2 IFI of 1.04 that we calculated all indicated a good model fit.
6
7 3
8
9
10 4 As expected from the linear models (Table 1b), the SEM model showed that N deposition is a
11
12 5 significant explanatory parameter for the changes in H' of the plots (Fig. 4a, Table 2a). A
13
14 6 direct negative standardised path coefficient of -0.302 (Fig. 4a) on the change in H' was
15
16 7 observed. Standardised path coefficients represent the standardised change in a response
17
18 8 variable per standardised unit change in a predictor variable, i.e. when N deposition
19
20 9 increased by one standard deviation, change in H' decreased by 0.302 standard deviation. In
21
22 10 terms of ecological significance, this can be regarded as a moderate effect (Chin 1998). In
23
24 11 addition, there was a positive relationship between soil pH and increase in H'. Mineral N
25
26 12 content exerted an indirect effect on the changes via soil pH, as well as a direct effect. N
27
28 13 deposition was positively related to mineral N content. S deposition had no direct effect but a
29
30 14 negative indirect effect via soil pH (Fig. 4a, Table 2a). Grazing pressure had an independent
31
32 15 direct negative effect on change in H'.
33
34 16
35
36
37
38
39
40
41 17 The SEM model for herb species showed that N deposition only weakly explains the
42
43 18 change between 1990-1993 and 2006-2009 (Fig. 4b, Table 2b) and that this was an indirect
44
45 19 effect via soil pH. A much stronger negative effect of S deposition on the change in herb
46
47 20 species was observed (Table 2b), also indirectly via soil pH. Grazing pressure did not have an
48
49 21 effect and was eliminated from this model. Mineral N content exerted an indirect effect via
50
51 22 soil pH (Fig. 4b, Table 2b).
52
53 23
54
55
56
57 24 *Change in frequency of individual species*
58
59
60

1 Some of the 52 individual species that met our selection criteria showed very distinct changes
2 in frequency between the two surveys. We divided the plots into three groups based on the
3 2006-8 N deposition data, and calculated the mean change in frequency of individual species
4 for plots in each deposition class. Species which declined or increased in frequency over all
5 plots by more than 15% are listed in Table 3, with the change in frequency split between the
6 three deposition categories. For the three species with an increase in frequency at the highest
7 N deposition, the overall relationship with N deposition was not significant. However, the five
8 species which declined in frequency in the highest class of N deposition (*Galium sterneri*,
9 *Linum catharticum*, *Briza media*, *Gentianella amarella* and *Campanula rotundifolia*) all
10 showed a significant correlation between the decline in frequency and N deposition.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

A key finding from our study is that the effects of N deposition, and of other driving variables, are quite different when considering the spatial variation between sites in 1990-93, and when considering the temporal change between 1990-93 and 2006-9. Here we consider the results from the spatial analysis and the analysis of temporal change in turn, placing our findings into the wider context of the role of Ndeposition and other factors in influencing the species richness, composition and diversity of calcareous grasslands.

Spatial variation

We found that N deposition did not explain spatial variation in species richness, diversity, or the frequency of eutrophic species. This is consistent with a study in 1998 of 94 calcareous grassland plots in the UK, in which there was no significantrelationship between N deposition, or climatic variables, and species richness (Maskell *et al.* 2010), even though significant relationships between N deposition and species richness were found in the same survey in acid grasslands and heathlands. These findings contrast with results from experimental studies in which clear N fertilization effects have been shown in calcicolous habitats, with fast-growing eutrophic species outcompeting slow growing sensitive species (e.g. Bobbink, 1991; Carroll *et al.* 2003; van den Berg *et al.* 2005). Maskell *et al.* (2010) argued that their relatively small sample size (94 plots),and the correlation between N deposition and climatic parameters, may have masked the significance of Ndeposition as an explanatory variable. Our sample size is similar (106 plots) to that of Maskell *et al.* (2010), but a more likely explanation for the lack of significant relationships with Ndeposition is that all of our sites are nature reserves, at which the management is aimed specifically at the conservation of high species diversity and in which grazing regimes are implemented to

1 prevent the dominance of eutrophic species. Grazing at moderate stocking rates maintains
 2 species diversity by reducing the number of competitive species (Collins *et al.* 1998; Olff &
 3 Ritchie, 1998; Klimek *et al.* 2007). The absence of a management effect may result from the
 4 management regimes at all the sites being successfully focussed on preserving species
 5 richness.

6
 7 Despite this, the frequency of rare and scarce species was significantly reduced, by a factor of
 8 5, at sites with N deposition above 25 kg ha⁻¹ y⁻¹. These relationships are difficult to test in a
 9 rigorous way in multivariate models, as rare species, such as *Epipactis atrorubens*,
 10 *Triniaglauca* and *Pulsatilla vulgaris*, not only have a highly restricted geographic distribution
 11 but are also typically highly localised on the sites where they do occur. Fisher and Stöcklin
 12 (1997) and Bennie *et al.* (2006) both found a negative correlation between the initial
 13 frequency of a plant species and the extinction rate in calcareous grasslands over a period of
 14 several decades, over which Bennie *et al.* (2006) also found an increase in Ellenberg fertility
 15 index, although there was not a significant relationship between extinction rates and the
 16 Ellenberg fertility index. Our results suggest a link between N deposition and the occurrence
 17 of scarce and rare species, which may be due to local extinctions over a period of decades.

18
 19 Maximum summer temperature, along with soil depth, explained most of the variation in
 20 species richness and the frequency of different types of species; species richness was higher
 21 in plots with a higher maximum temperature. Plots with deeper soils tended to occupy
 22 moderate to low slopes and had a higher organic content ($r^2 = -0.14$, $p = 0.022$) and a higher
 23 moisture content ($r^2 = 0.35$, $p < 0.001$). The higher organic content and the fact that deeper
 24 soils are more resistant to drought stress may be important factors in explaining the higher
 25 species richness. Duckworth *et al.* (2000) found that temperature and soil organic matter are

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 amongst the strongest parameters explaining spatial variation in vegetation composition of
2 Atlantic European calcareous grasslands. In addition, Bennie *et al.* (2006) found that
3 calcareous grassland plots on deeper soils were more likely to be invaded by (often
4 mesotrophic) plant species, which may explain the higher species richness.

6 *Comparison between surveys*

7 The species richness was lower in 2006-2009 than in 1990-1993 with an average loss of 2
8 species over 15 years; this small overall change may represent differences between surveyors
9 in the two periods. Change in species richness was not associated with N deposition,
10 but changes in H' and evenness were significantly negatively related to N deposition. The
11 Cohen's d value for the change in H' was 0.99, representing a medium to strong ecological
12 effect (Cohen 1988). Progression towards an undesirable reference dataset in high N
13 deposition plots showed the reduction in H' between 1990-1993 and 2006-2009 represented a
14 6.4% movement towards the H' value typical of British Neutral Grasslands. This implies that
15 a much longer period would be needed to reach the low values of H' typical of neutral
16 grasslands. However, many typical calcareous grassland species may be lost well before this
17 value of H' is reached, and we conclude that the change in H' at high N deposition sites does
18 represent a significant ecological change.

20 H' is a measure of diversity that accounts for both species richness and evenness of the
21 species present in the sample (Magurran, 2004). N deposition did not significantly affect
22 changes in species richness but did significantly affect changes in evenness, indicating that
23 changes in H' were driven primarily by changes in the evenness of the plots.

In addition, there was a significant decline in the frequency of five individual species when N deposition exceeded $25 \text{ kg ha}^{-1} \text{ y}^{-1}$. All five individual species (*Linum catharticum*, *Briza media*, *Gentianella amarella*, *Campanula rotundifolia* and *Galium sterneri*) have a low Ellenberg nutrient score. The decline in frequency in these high N deposition plots was substantial (on average 22%) over a relatively short period of 15 years despite the focussed management on these nature reserves, and indicates that local extinctions of these species could occur in the future unless N deposition declines. Both *C. rotundifolia* and *B. media* are regarded as poor competitors (Leishman, 1999; Dixon, 2002; Hutchings *et al.* 2003) and for *C. rotundifolia*, lower frequencies have been found at sites with high N deposition (Stevens *et al.* 2006; Maskell *et al.* 2010). *Linum catharticum* and *G. amarella* are respectively annual and bi-annual species which may strongly depend on gap formation in the swards for germination (e.g. Turnbull *et al.* 2005). These results are corroborated by an earlier study on calcareous grasslands by Bennie *et al.* (2006) who found strong decreases in frequency over the period 1950-2000 of *C. rotundifolia*, *L. catharticum* and *B. media*.

The change in species richness was positively related to soil pH, as was herb species number, indicating that plots of higher buffering capacity tended to gain species and plots with slightly lower pH tended to lose species. Despite the fact that calcareous soils are highly buffered, experimental studies have shown that calcareous grasslands also leach base cations when N deposition is increased, as a result of soil exchange processes and plant uptake of nitrogen (Horswillet *et al.* 2008). An increase of soil pH due to the reduction in acid deposition over the last 15 years (Emmett *et al.* 2010) may have contributed to increased species richness, especially in those plots which were least affected by base cation depletion.

Importance of direct and indirect effects of N deposition

1
2
3 1 The use of structural equation models allowed us to identify and compare the comparative
4
5 2 strengths of direct effects of N deposition and indirect effects of both N and S deposition,
6
7
8 3 through soil pH, using hypothesised causal relationships that are consistent with experimental
9
10 4 studies. The responses of vegetation to changing N deposition are complex and involve
11
12 5 several mechanisms (Bobbink *et al.* 2010), and therefore statistical methods, such as
13
14 6 structural equation modelling, that incorporate this complexity and aim to disentangle
15
16 7 different direct and indirect effects through potential causal structures provide powerful new
17
18 8 tools to complement more conventional regression analysis.
19
20 9
21
22
23
24 10 The results show very different models for change in H' and for change in herb species
25
26 11 numbers. Changes in H' were dominated by a direct negative effect of N deposition; although
27
28 12 indirect negative effects of S and N deposition via soil pH were also significant, they had a
29
30 13 much smaller contribution. The negative direct effect of N deposition on change in H' was
31
32 14 also found for changes in evenness in the mixed effect models, suggesting that eutrophication
33
34 15 rather than acidification is the dominant mechanism. In contrast, the direct effect of N
35
36 16 deposition on change in herb species numbers was not significant, and was small compared to
37
38 17 indirect effects via soil pH, indicating that acidification was the dominant mechanism.
39
40 18
41
42
43
44
45 19 Soil pH, which had a significant positive relationship with both changes in H' and changes in
46
47 20 herb species numbers, was itself negatively influenced by S deposition and mineral N
48
49 21 content. The strong acidifying nature of S in deposition compared to N was revealed by the
50
51 22 stronger negative effects that were exerted by S deposition on soil pH. Calcareous grasslands
52
53 23 have been shown to retain most of the deposited N in the system with little leaching to deeper
54
55 24 soil layers or groundwater, thereby increasing mineral N content of the top soils (Phoenix *et*
56
57 25 *al.* 2008). The increased mineral N in the soils in turn can have a significant negative effect

on soil pH (e.g. Horswillet *et al.* 2008). This is consistent with the positive relationships between N deposition and mineral N content and the significant negative relationships between the mineral N content and soil pH in our study. These findings, using a different method of analysis, are consistent with earlier studies of acid grasslands in which negative effects of soil acidity on herb species numbers were recorded (e.g. Stevens *et al.* 2006; Duprè *et al.* 2010, Stevens *et al.* 2010).

A recent survey of changes in a local UK flora over the last 50 years showed the positive effects of the designation of conservation sites on maintaining biodiversity, especially in calcareous grasslands (Walker *et al.* 2009). In our structural equation model, change in grazing also significantly improved the model fit, with a decrease in grazing pressure being directly associated with a decrease in H'. A decrease in grazing pressure was also associated with an increase in Ellenberg N score. These results may reflect an effect of poor management on some sites over recent years, since the spatial analysis suggested that grazing management in 1990-93 was optimal overall and hence did not affect spatial variation in species richness or diversity.

Conclusions

Critical loads for adverse effects of N deposition for calcareous grasslands, based on field experiments, have been set as a range of 15-25 kg N ha⁻¹y⁻¹ (Bobbink *et al.* 2003; Bobbink *et al.* 2010), with the range indicating the variation in sensitivity. Our results provide the first support for this critical load range from field data, as, in the plots where the critical load range is exceeded, there is evidence of a decrease in species diversity and evenness, a decline in the frequency of characteristic species, and a lower number of rare and scarce species. These effects are likely to be due to both direct effects of N deposition and indirect effects

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 through changes in soil pH. Since all sites had active management to maintain a high diversity
2 and characteristic species, our results imply that even focussed management on nature
3 conservation objectives cannot prevent adverse effects of high rates of N deposition. Given
4 the high and increasing rates of N deposition in many parts of the world, N deposition is a
5 wide-scale threat to the conservation of species-rich grassland communities, and their
6 characteristic species.

For Review Only

Acknowledgements

This study was supported by the UK Natural Environment Research Council under award number NE/E00413. The original establishment of the plots in 1990-93 was supported by the then UK Department of the Environment under contract PECD 7/12/22. We thank Countryside Council for Wales, Natural England, and the Scottish Natural Heritage for granting permission to access and sample the sites.

For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

1 **Tables**

2 **Table 1:**

3 Effect sizes of the predictor variables (expressed as the coefficient $\pm$ s.d.), for the subset of predictors that were included in the model with the
4 lowest AIC for each response variable, for (a) the survey in 1990-93 and (b) the change between the surveys in 2006-9 and 1990-93. Predictors
5 significant at $p<0.1$, $p<0.05$, $p<0.01$ and $p<0.001$ are indicated by +, *, ** and ***, respectively. Analyses were based on a maximum of 106
6 calcareous grassland surveys in 1993 and a maximum of 42 comparisons between 2009 and 1993. All analyses were based on vascular plants
7 only. The number of acidophilic species was $\log(x+1)$ transformed.

1

a) 1990-1993		Predictor variable					df ^{1,2}	AIC
	pH	Soil Depth	Aspect	Ndeposition	Max Temp summer	Grazing		
Species richness ¹		0.338*			1.442*		1, 103	770
		±0.121			± 0.616			
Eutrophic species ¹		0.274***			1.530		1, 103	715
		±0.093			±0.467***			
Oligotrophic species ¹			1.715				1, 104	706
			±1.218					
Acidophylic species	-0.507***				-0.166***		1, 103	176
	± 0.123				±0.025			
Acidophobic species ¹			2.815		2.296***		1, 103	714
			±1.095		± 0.513			
Herb species ¹		0.266**			1.501**		1, 103	726
		±0.097			±0.503			
Grass:herb ratio ¹				0.004+	-0.028***		1, 103	-119
				±0.002	±0.007			
WA Ellenberg N ¹					0.028**		1, 104	-79
					±0.009			
WA Ellenberg R	0.395***				0.115***		1, 103	109
	±0.090				± 0.018			
Shannon index		0.0133***			0.049***		1, 103	32
		±0.004			±0.013			
Evenness ¹		0.001					1, 104	-460
		±3.390E-4						

1

b) Change between 2006-2009-and 1990-1993	Predictor variable							df _{1,2}	AIC
	pH	Soil Depth	Mineral N soil	Aspect	Ndepositio n	Max Temp summer	Grazing		
Species richness	2.634* ±1.242							1,35	227
Eutrophic species							-1.434 ± 1.326	1,35	170
Oligotrophic species		-0.106 ±0.056	3.773E-4+ ± 2.003E-4					1,34	149
Acidophylic species		-0.032 ±0.021						1,35	77
Acidophobic species		0.089 ± 0.077						1,35	174
Herb species	2.427* ±0.9348		8.085E-04+ ±4.253E-04					1,34	203
Grass:herb ratio ¹	-0.056+ ±0.021							1,35	-61
WA Ellenberg N ¹							-0.213** ±0.073	1,35	-32
WA Ellenberg R ¹			-2.039E-5 ±1.505E-5					1,35	-38
Shannon index					-0.006* ±0.003			1,35	-48
Evenness ¹			-3.110E-6 ± 1.939E-6		-0.001* ±0.0004			1,34	-141

2 df_{1,2} is degrees of freedom of the tested effect term and the error term, respectively

3 ¹ models tested with lme rather than lm function

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2:

Summary of results from structural equation modelling, showing the decomposition of correlations into standardised direct, indirect and total effects of seven explanatory variables on (a) the difference in Shannon diversity index and (b) the difference in the number of herb species between 2006-2009 and 1990-1993. Total effects are the sum of direct and indirect effects. Direct effect coefficients are equal to the path coefficients shown in Fig. 4a and b.

a)	Difference in Shannon diversity between 2009 and 1993 (H2009-H1993)		
	Direct	Indirect	Total
Ndeposition	-0.302	-0.008	-0.310
SOx deposition	-	-0.037	-0.037
Mineral N soil	-0.022	-0.018	-0.040
Soil pH	0.075	-	0.075
Grazing pressure	0.063	-	0.063
b)	Difference in herb species between 2009 and 1993 (n2009-n1993)		
	Direct	Indirect	Total
Ndeposition	-	-0.014	-0.014
SOx deposition	-	-0.147	-0.147
Mineral N soil	-	-0.072	-0.072
Soil pH	0.298	-	0.298

Table 3:

Effect of N deposition class on the mean change in frequency of individual species (expressed as the difference of frequency between 2006-2009 and 1990-1993), for those species with a change in frequency of more or less that 0.15 (15%) in the highest Ndeposition class. Shown are the correlation coefficients (Cor.) between the change in relative frequency of the species and modelled Ndeposition based on 2006-8 data and the level of significance of the correlation.

Species	N deposition class (kgN ha ⁻¹ y ⁻¹)			Cor.	p
	0-15	15-25	25-35		
<i>Cynosurus cristatus</i>	0.08	0.12	0.53	0.04	0.809
<i>Holcus lanatus</i>	0.14	0.20	0.25	-0.08	0.611
<i>Polygala vulgaris</i>	0.23	0.06	0.17	-0.03	0.812
<i>Linum catharticum</i>	0.27	-0.02	-0.18	-0.34	0.026
<i>Briza media</i>	0.17	0.05	-0.21	-0.37	0.015
<i>Gentianella amarella</i>	-0.03	0.12	-0.24	-0.30	0.050
<i>Campanula rotundifolia</i>	0.05	-0.15	-0.25	-0.61	<0.001
<i>Galium sternerii</i>	0.11	0.04	-0.26	-0.53	<0.001

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Figure legends**

2 **Fig. 1**

3 Map of the UK showing the location of the 56 calcareous grassland sites that were surveyed.

4 Closed circles are sites that were surveyed between 1990 and 1993 only, open circles are sites
5 which were re-surveyed between 2006 and 2009. Outline map from Ordnance Survey UK.

For Review Only

Fig. 2

The relationship between total Ndeposition ($\text{kg ha}^{-1}\text{y}^{-1}$) in 1993-1995 and (a) the total number of vascular plant species in 1990-1993; (b) grass to herb ratio in 1990-1993; and (c) the total number of scarce and rare vascular plant species in 1990-1993.). The equation of the fitted line in Figure 2b is $y = 0.004x + 0.40$ ($r^2=0.03$; $p=0.053$).

For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 3

The relationship between the total Ndeposition ($\text{kg ha}^{-1}\text{y}^{-1}$) in 2006-2008 and the decrease between 1990-1993 and 2006-2009 of (a) the plot species diversity (H'); and (b) the plot species evenness. The equation of the fitted line in Figure 3a is $y = -0.008x + 0.19$ ($r^2=0.15$; $p=0.011$). The equation of the fitted line in Figure 3b is $y = -0.002x + 0.03$ ($r^2=0.21$; $p=0.002$).

Fig. 4a

Structural equation model depicting direct and indirect effects of N and S deposition, soil pH, mineral N soil and grazing on the change in Shannon diversity between 2006-2009 and 1990-1993. The width of each arrow is proportional to the standardised path coefficient (coefficients shown in *italic*). Positive path coefficients are indicated by solid lines and negative path coefficients are represented by dashed lines. Asterisks indicate effects that are significant at $p < 0.1$. Estimates of the proportion of total variance explained (squared multiple correlations) are shown in bold for each dependent variable. Variables and paths representing unmeasured residual variation are not indicated for simplicity.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 4b

Structural equation model depicting direct and indirect effects of N and S deposition, soil pH, and soil mineral N soil on the change in herb species numbers between 2006-2009 and 1990-1993. The width of each arrow is proportional to the standardised path coefficient (coefficients shown in *italic*). Positive pathcoefficients are indicated by solid lines and negative pathcoefficients are represented by dashed lines. Asterisks indicate effects that are significant at $p < 0.1$. Estimates of the proportion of total variance explained (squared multiple correlations) are shown in bold for each dependent variable. Variables and paths representing unmeasured residual variation are not indicated for simplicity.

References

- Attwood EA ,Heavey JF (1964) Determination of Grazing Livestock Units. *Irish Journal of Agricultural Research*, **3**, 249-251.
- Bennie J, Hill MO, Baxter R, Huntley B (2006) Influence of slope and aspect on long-term vegetation change in British chalk grasslands. *Journal of Ecology*, **94**, 355-368.
- Bobbink R (1991) Effects of nutrient enrichment in Dutch chalk grassland. *Journal of Applied Ecology*, **28**, 28-41.
- Bobbink R, Ashmore M, Braun S *et al.* (2003) Empirical nitrogenloads for natural and semi-natural ecosystems: 2002 update. In:Empirical Critical Loads for Nitrogen, Environmental DocumentationNo. 164 Air, (edsAcher mann B, Bobbink R), Swiss Agencyfor Environment, Forest and Landscape SAEFL, Berne.
- Bobbink R, Hornung M, Roelofs JGM (1998) The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. *Journal of Ecology*, **86**, 717-738.
- Bobbink R, Hicks K, Galloway J *et al.* (2010) Global Assessment of Nitrogen Deposition Effects on Terrestrial Plant Diversity: a synthesis. *Ecological Applications*, **20**, 30-59.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Carroll JA, Caporn SJM, Johnson D, Morecroft MD, Lee JA (2003) The interactions between
2 plant growth, vegetation structure and soil processes in semi-natural acidic and calcareous
3 grasslands receiving long-term inputs of simulated pollutant nitrogen deposition.
4 *Environmental Pollution*, **121**, 363-376.

5

6 Chin WW (1998) Issues and opinion on structural equation modelling.*MIS Quaterly*, **22**, 7-
7 16.

8

9 Cohen J (1988) Statistical power analysis for the behavioural sciences (2nd edition). New
10 Jersey. Lawrence Erlbaum Associates Ltd.

11

12 Collins SL, Knapp AK, Briggs JM, Blair JM, Steinauer EM (1998) Modulation of diversity
13 by grazing and mowing in native tallgrass prairie. *Science*, **280**, 745-747.

14

15 Dentener F, Drevet J, Lamarque JF *et al.* (2006) Nitrogen and sulfur deposition on regional
16 and global scales: A multimodel evaluation. *Global Biogeochemical Cycles*, **20**,
17 GB4003.doi:10.1029/2005GB002672

18

19 Dixon JM (2002). Biological Flora:*Briza media* L. *Journal of Ecology*, **90**, 737-752.

20

21 Duckworth JC, Bunce RGH, Malloch AJC (2000) Vegetation-environment relationships in
22 Atlantic European calcareous grasslands.*Journal of vegetation Science*, **11**, 15-22.

23

- 1 Duprè C, Stevens CJ, Ranke T *et al.* (2010) Changes in species richness and composition in
2 European acidic grasslands over the past 70 years: the contribution of cumulative
3 atmospheric nitrogen deposition. *Global Change Biology*, **16**, 344-357.
- 4
5 Emmett BA, Reynolds B, Chamberlain PM *et al.* (2010) Countryside Survey: Soils Report
6 from 2007. Technical Report No. 9/07 NERC/Centre for Ecology & Hydrology 192pp. (CEH
7 Project Number: C03259).
- 8
9 Fischer M, Stöcklin J (1997) Local extinctions of plants in remnants of extensively used
10 calcareous grasslands 1950-1985. *Conservation Biology*, **11**, 727-737.
- 11
12 Galloway JN (1995) Acid deposition: Perspectives in time and space. *Water Air and Soil
13 Pollution*, **85**, 15-24.
- 14
15 Galloway JN, Cowling EB (2002) Reactive nitrogen and the world: 200 years of change.
16 *Ambio*, **31**, 64-71.
- 17
18 Gujarati DN (1995) Basic Econometrics, (Third edition) McGraw-Hill, New York
- 19
20 Hill MO, Preston CD Roy DB (2004) *PLANTATT- Attributes of British and Irish Plants:
21 Status, Size, Life History, Geography and Habitats*. CEH Monks Wood, United Kingdom.
- 22
23 Horswill P, O'Sullivan O, Phoenix GK, Lee JA, Leake JR (2008) Base cation depletion,
24 eutrophication and acidification of species-rich grasslands in response to long-term simulated
25 nitrogen deposition. *Environmental Pollution*, **155**, 336-349.

1
2
3
4 1
5
6 2 Hutchings MJ, John EA, Wijesinghe DK (2003) Toward understanding the consequences of
7
8 3 soil heterogeneity for plant populations and communities. *Ecology*, **84**, 2322–2334.
9
10 4
11
12 5 Jackson, DL (2000). *Guidance on the interpretation of the Biodiversity Broad Habitat*
13 6 *Classification (terrestrial and freshwater types): Definitions and the relationship with other*
14 7 *habitat classifications*. JNCC Report No. 307. Peterborough, UK: Joint Nature Conservation
15 8 Committee.
16 9
17 10
18 11 Klimek S, Kemmermann ARG, Hofmann M, Isselstein J (2007) Plant species-richness and
19 12 composition in managed grasslands: The relative importance of field management and
20 13 environmental factors. *Biological Conservation*, **134**, 559-570.
21 14
22 15 Leishman, MR (1999) How well do plant traits correlate with establishment ability? Evidence
23 16 from a study of 16 calcareous grassland species. *New Phytologist*, **141**, 487-496.
24 17
25 18 Magurran AE (2004) *Measuring biological diversity*. Blackwell Publishing, Oxford.
26 19
27 20 Maskell LC, Smart SM, Bullock JM, Thompson K, Stevens CJ (2010) Nitrogen deposition
28 21 causes widespread loss of species-richness in British habitats. *Global Change Biology*, **16**,
29 22 671-679.
30 23
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1 Morecroft MD, Sellers EK, Lee JA (1994). An experimental investigation into the effects of
2 atmospheric nitrogen deposition on 2 semi-natural grasslands. *Journal of Ecology*, **82**, 475-
3 483.
- 4
- 5 Olff H, Ritchie ME (1998) Effects of herbivores on grassland plant diversity. *Trends in*
6 *Ecology and Evolution*, **13**, 261-265.
- 7
- 8 Olsen SR, Cole CV, Watanabe FS, Dean LA (1954) *Estimation of available phosphorus in*
9 *soils by extraction with sodium bicarbonate*. pp. 1-19, U.S. Dept. of Agriculture, Washington
10 DC
- 11
- 12 Phoenix GK, Booth RE, Leake JR, Read DJ, Grime JP, Lee JA (2003) Effects of enhanced
13 nitrogen deposition and phosphorus limitation on nitrogen budgets of semi-natural
14 grasslands. *Global Change Biology*, **9**, 1309-1321.
- 15
- 16 Phoenix GK, Hicks WK, Cinderby S *et al.* (2006) Atmospheric Nitrogen Deposition in World
17 Biodiversity Hotspots: the need for a greater global perspective in assessing N deposition
18 impacts. *Global Change Biology*, **12**, 470–476.
- 19
- 20 Phoenix GK, Johnson D, Grime JP, Booth RE (2008) Sustaining ecosystem services in
21 ancient limestone grassland: importance of major component plants and community
22 composition. *Journal of Ecology*, **96**, 894-902
- 23
- 24 R Development Core Team (2008) R: a language and environment for statistical computing.
25 Vienna, Austria: R Foundation for Statistical Computing, <http://www.R-project.org>.

1
2
3
4 1
5
6 2 Rich TCG, Cooper EA, Rodwell JS, Malloch AJC (1993) Effects of climate change and air
7
8 3 pollution on British calcicolous ecosystems.Final report to UK Department of the
9
10 4 Environment.
11
12
13 5
14
15 6 Rodwell JS (ed) (1998). *British Plant Communities, Vol. 3, Grasslands and Montane*
16
17 7 *Communities*. Cambridge University Press, Cambridge.
18
19
20 8
21
22 9 Sala OE, Chapin FS, Armesto JJ *et al.* (2000) Biodiversity-Global biodiversity scenarios for
23
24 10 the year 2100.*Science*, **287**, 1770-1774.
25
26
27 11
28
29 12 Smart SM, Clarke RT, van de Poll HM, Robertson EJ, Shield ER, Bunce RGH, Maskell LC
30
31 13 (2003a) National-scale vegetation change across Britain; an analysis of sample-based
32
33 14 surveillance data from the Countryside Surveys of 1990 and 1998. *Journal of Environmental*
34
35 15 *Management*, **67**, 239-254.
36
37
38 16
39 17 Smart SM, Robinson JC, Shield EJ, van de Poll HM (2003b) Locating eutrophication effects
40
41 18 across British vegetation between 1990 and 1998. *Global Change Biology***9**, 1763-1774.
42
43
44 19
45
46
47 20 Stace C (1997) *New Flora of the British Isles, second edition*. Cambridge University Press,
48
49 21 Cambridge.
50
51
52 22
53
54 23 Stevens CJ, Dise NB, Gowing DJG, Mountford JO (2006) Loss of forb diversity in relation to
55
56 24 nitrogen deposition in the UK: regional trends and potential controls. *Global Change Biology*,
57
58 25 **12**, 1823-1833.
59
60

- 1
2
3
4 1
5
6 2 Stevens CJ, Dise NB, Mountford JO, Gowing DJ (2004) Impact of nitrogen deposition on the
7
8 3 species-richness of grasslands. *Science*, **303**, 1876-1879.
9
10 4
11
12 5 Stevens CJ, Thompson K, Grime JP, Long CJ, Gowing DJG (2010) Contribution of
13
14 6 acidification and eutrophication to declines in species richness of calcifuge grasslands along a
15
16 7 gradient of atmospheric nitrogen deposition. *Functional Ecology*, **24**, 478-484.
17
18 8
19
20 9 Tansley AG (1922) Studies of the vegetation of the English chalk. II Early stages of
21
22 10 redevelopment of woody vegetation on chalk grassland. *Journal of Ecology*, **10**, 168-177.
23
24 11
25
26 12 TerBraak CJF (1988) CANOCO, an extension of Decorana to analyze species-environment
27
28 13 relationships. *Vegetatio*, **75**, 159-160.
29
30 14
31
32 15 Turnbull LA, Manley L, Rees M (2005) Niches, rather than neutrality, structure a grassland
33
34 16 pioneer guild. *Proceedings of the Royal Society Biological Sciences*, **272**, 1357-1364.
35
36 17
37
38 18 Van den Berg LJJ, Tomassen HBM, Roelofs JGM, Bobbink R (2005) Effects of nitrogen
39
40 19 enrichment on coastal dune grassland: A mesocosm study. *Environmental Pollution*, **138**, 77-
41
42 20 85.
43
44 21
45
46 22 Vestreng V, Myhre G, Fagerli H, Reis S, Tarrason L (2007) Twenty-five years of continuous
47
48 23 sulphur dioxide emission reduction in Europe. *Atmospheric Chemistry and Physics*, **7**, 3663-
49
50 24 3681.
51
52
53
54
55
56
57
58
59
60 25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Vitousek PM, Aber JD, Howarth RW *et al.* (1997) Human alteration of the global nitrogen
2 cycle: Sources and consequences. *Ecological Applications*, **7**, 737-750.
3
4 Walker KJ, Preston CD, Boon CR (2009) Fifty years of change in an area of intensive
5 agriculture: plant trait responses to habitat modification and conservation, Bedfordshire,
6 England. *Biodiversity and Conservation*, **18**, 3579-3613.
7
8 Willems JH (1978) Observations on North-West European limestone grassland communities
9 – Phytosociological and ecological notes on chalk grasslands of southern England. *Vegetatio*,
10 **37**, 141-149.
11
12 Willems JH (1990) Calcareous grasslands in Continental Europe. In: *Calcareous grasslands:
13 ecology and management. Proceedings of a BES/NCC Symposium at University of Sheffield,
14 14–16 September 1987* (eds Hillier SH, Walton DWH, Wells DA), pp 3-10. Bluntisham
15 Books, Bluntisham.
16
17 Wilson EJ, Wells TCE, Sparks TH (1995) Are calcareous grasslands in the UK under threat
18 from nitrogen deposition-an experimental determination of a critical load. *Journal of
19 Ecology*, **83**, 823-832.

58x95mm (150 x 150 DPI)

Figure 2

Figure 3

Figure 4a

Figure 4b