GENETIC EVIDENCE FOR THE PRESENCE OF TWO DISTINCT HANTAVIRUSES ASSOCIATED WITH APODEMUS MICE IN CROATIA AND ANALYSIS OF LOCAL STRAINS

Angelina Plyusnina, Lidija Cvetko Krajinović, Josip Margaletic, Jukka Niemimaa, Kirill Nemirov, Åke Lundkvist, Alemka Marcotić, Marica Miletić Medved, Tatjana Avšič-Županc, Heikki Henttonen, et al.

► To cite this version:

Angelina Plyusnina, Lidija Cvetko Krajinović, Josip Margaletic, Jukka Niemimaa, Kirill Nemirov, et al.. GENETIC EVIDENCE FOR THE PRESENCE OF TWO DISTINCT HANTAVIRUSES ASSOCIATED WITH APODEMUS MICE IN CROATIA AND ANALYSIS OF LOCAL STRAINS. *Journal of Medical Virology*, 2010, 83 (1), pp.108. 10.1002/jmv.21929 . hal-00599791

HAL Id: hal-00599791

<https://hal.science/hal-00599791>

Submitted on 11 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**GENETIC EVIDENCE FOR THE PRESENCE OF TWO DISTINCT
HANTAVIRUSES ASSOCIATED WITH APODEMUS MICE IN
CROATIA AND ANALYSIS OF LOCAL STRAINS**

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-10-1909.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	21-Jul-2010
Complete List of Authors:	<p>Plyusnina, Angelina; Haartman Inst, University of Helsinki, Dept Virology Cvetko Krajinović, Lidija; University Hospital for Infectious Diseases Margaletic, Josip; Faculty of Forestry, University of Zagreb Niemimaa, Jukka; Finnish Forest Research Institute, Vantaa Research Unit Nemirov, Kirill; Swedish Institute for Infectious Disease Control Lundkvist, Åke; Swedish Institute for Infectious Disease Control Marcotić, Alemka; University Hospital for Infectious Diseases Miletić Medved, Marica; Public Health Institute Brodsko-posavska County Avšič-Županc, Tatjana; Medical School, University of Ljubljana Henttonen, Heikki; Finnish Forest Research Institute, Vantaa Research Unit Plyusnin, Alexander; Haartman Inst, University of Helsinki, Virology</p>
Keywords:	HFRS , hantavirus , Dobrava virus, Saaremaa virus

SCHOLARONE™
Manuscripts

For J.Med.Virol.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 1 GENETIC EVIDENCE FOR THE PRESENCE OF TWO DISTINCT
3
4 2 HANTAVIRUSES ASSOCIATED WITH *APODEMUS* MICE IN CROATIA AND
5
6 3 ANALYSIS OF LOCAL STRAINS
7

8 4 (Short title: Dobrava and Saaremaa hantaviruses in Croatia)
9

10 5
11 6 Angelina Plyusnina¹, Lidija Cvetko Krajinovic², Josip Margaletic³, Jukka Niemimaa⁴,
12 7 Kirill Nemirov⁵, Åke Lundkvist⁵, Alemka Markotic², Marica Miletic Medved⁶,
13 8 Tatjana Avšič Županc⁷, Heikki Henttonen⁴, and Alexander Plyusnin^{1*}
14
15 9

16
17
18 10 ¹Department of Virology, Infection Biology Research Program, Haartman Institute,
19 11 University of Helsinki, Finland

20
21
22 12 ²University Hospital for Infectious Diseases, Zagreb, Croatia

23
24 13 ³Faculty of Forestry, University of Zagreb, Zagreb, Croatia

25
26 14 ⁴Finnish Forest Research Institute, Vantaa Research Unit, Vantaa, Finland

27
28 15 ⁵Swedish Institute for Infectious Disease Control, Stockholm, Sweden

29
30 16 ⁶Public Health Institute Brodsko-posavska County, Slavonski Brod, Croatia

31
32 17 ⁷Medical School, University of Ljubljana, Ljubljana, Slovenia
33

34
35
36 19 *Correspondence to:

37
38 20 Alexander Plyusnin

39
40 21 Department of Virology, Haartman Institute

41
42 22 University of Helsinki P.O. Box 21

43
44 23 FIN-00014, Finland

45
46 24 Phone: +358 9 19126486; Fax: +358 9 19126491

47
48 25 E-mail: alexander.plyusnin@helsinki.fi
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

In Europe, Dobrava-Belgrade (DOBV), Saaremaa (SAAV) and Puumala (PUUV) viruses are known to cause hemorrhagic fever with renal syndrome (HFRS). All three hantaviruses are now found in Croatia. Lung tissue samples of 315 *Apodemus* mice trapped in 2003-2004 were screened for the presence of hantaviral N-Ag and 20 mice (6.3%) were found either strongly positive or weak/suspected-positive. Partial sequences of hantavirus M and S segments were recovered by RT-PCR from six mice and subjected to (phylo)genetic analysis that revealed the presence of four novel strains of DOBV and one of SAAV. Curiously, one of the newly described DOBV strains was found in *A. agrarius* mouse, i.e. not in the traditional host, *A. flavicollis* mice, suggesting a spillover event. S segment sequences recovered previously from HFRS cases (Markotic et al., 2002) were confirmed as DOBV sequences; one of which appeared particularly close to the prototype Slovenian DOBV isolate. Taken together with earlier data on PUUV in Croatia, these results show a co-circulation of three European hantavirus pathogens in this country. So far, not a single SAAV sequence has been recovered from HFRS patients either in Croatia or neighbouring Slovenia and Hungary nor in Slovakia suggesting a somewhat lower frequency of acute SAAV infection in humans in this part of Europe than for example in the Baltics.

Key words: HFRS; hantavirus; Dobrava virus, Saaremaa virus.

INTRODUCTION

Hantaviruses constitute the *Hantavirus* genus in the family *Bunyaviridae* [Nichol et al., 2005]. They are enveloped viruses with tri-segmented RNA genome of negative polarity. Small (S), Medium (M) and Large (L) RNA segments encode, respectively, nucleocapsid (N) protein, a precursor for two surface glycoproteins Gn and Gc, and the RNA-dependent RNA polymerase (L protein). In some hantaviruses, the S segment encodes also the nonstructural protein NSs. The function(s) of this protein in the virus replication cycle remain to be determined but there is an evidence that it can counteract an interferon response [Jääskeläinen et al., 2007]. In nature, hantaviruses are maintained in rodents or insectivores infected persistently. When transmitted to humans, usually via aerosolized excreta, some rodent-borne hantaviruses can cause disease. In Europe, Dobrava-Belgrade (DOBV), Saaremaa (SAAV) and Puumala (PUUV) viruses cause hemorrhagic fever with renal syndrome (HFRS) of varying severity. In Asia, the main causative agents of HFRS are Hantaan and Seoul (SEOV) viruses. In the Americas, Sin Nombre, Andes and related viruses are associated with hantavirus (cardio)pulmonary syndrome (HCPS) [for a reviews, see Vapalahti et al., 2003; Schmaljohn and Hjelle, 1997]. Very little is known about hantaviruses in Africa [Klempa et al., 2006, 2007] and nothing is known about these viruses in Australia.

DOBV, SAAV, PUUV, and SEOV are found in Europe, often in co-circulation, and the first three viruses also in association with human HFRS cases [Vapalahti et al., 2003]. DOBV, harbored by yellow-necked mice (*Apodemus flavicollis*), is associated with severe HFRS mostly in the Balkans and Alpe-Adrian region [Papa et

For J.Med.Virol.

al., 1998; Avšič Županc et al., 1999; Markotic et al., 2002; Jakab et al., 2007]. The natural host for SAAV is the striped field mouse (*Apodemus agrarius*); the virus causes mild HFRS in the Baltics, Central Europe and the European part of Russia [Lundkvist et al., 1998; Plyusnin et al., 1999; Golovljova et al., 2000; Sibold et al., 2001; Klempa et al., 2004, 2008]. It should be noted that SAAV was considered initially a genetic variant of DOBV carried by *A. agrarius* [Plyusnin et al., 1997; Nemirov et al., 1999] and for a while there was a controversy in the taxonomy and terminology of SAAV. The virus is now recognized as a distinct species [for the latest taxonomy, see www.ictvonline.org] but, in some publications, it is still named "DOBV-Aa".

PUUV is carried by bank voles (*Myodes glareolus*) and causes HFRS, usually mild, in Northern and Central Europe, the Baltics, European part of Russia and Alpe-Adria [Settergren et al., 1989; Mustonen et al., 1994; Tkachenko et al., 1998; Golovljova et al., 2000, 2007; Heyman et al., 2001; Markotic et al., 2002; Cvetko et al., 2005; Avšič-Županc et al., 2007; Hoffmann et al., 2008].

The whole of Croatia except costal region and islands is endemic for HFRS. Serological and genetic findings in HFRS patients provided evidence that PUUV and Dobrava virus (DOBV) have been two major pathogenic hantaviruses circulating in Croatia and causing hantavirus infections in humans [Markotic et al. 2002, Cvetko et al., 2005]. Apathogenic Tula virus was detected in wild rodents in Croatia as well [Scharninghausen et al., 2002]. Croatia borders Slovenia to the northwest and has an extended border with Bosnia and Herzegovina, one of the largest HFRS regions in Europe [Markotic et al., 1995; Hukic et al., 2009]. Nowhere else is the co-existence of PUUV and DOBV as obvious as in Croatia, Bosnia and Herzegovina and Slovenia

For J.Med.Virol.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

96 [Markotic et al., 2002; Cvetko et al., 2007; Lundkvist et al., 1997; Avšic et al., 1999].
97 During the HFRS outbreak in Croatia in 2002 when more than 400 patients were
98 serologically confirmed as HFRS positive, PUUV was shown to be dominant
99 pathogen [Miletic-Medved et al. 2002; Kuzman et al. 2003].

100 Phylogenetic analysis of wild-type (wt) PUUV strains from the 2002-outbreak
101 area showed that they clustered together with the strains from Slovenia and Austria
102 forming distinct Alpe-Adrian genetic lineage [Cvetko et al., 2005]. Most recently,
103 Hungarian PUUV strains were added to this lineage [Plyusnina et al., 2009].

104 The knowledge on DOBV and SAAV genetic variants in Croatia is still
105 restricted to partial S segment sequences recovered from three HFRS patients
106 [Markotic et al., 2002]. To fill this gap, we attempted to recover and analyze DOBV
107 and SAAV sequences from carrier rodents, *Apodemus* mice, trapped during field
108 expeditions organized by the Faculty of Forestry, University of Zagreb and the
109 Finnish Forest Research Institute.

110

MATERIALS AND METHODS

Trapping of rodents. Rodents were collected during two expeditions, in north-east Croatia in May 2003 and the south-west parts of Croatia in September, 2004 at the following locations: Migalovci, Merolino, Loze, Nova Gradiška, Gratac/Mala Kapela, and Delnice/Mala Kapela. Animals were trapped with snap traps, set in the evening and checked early in the morning. Animals were placed in cooler box in the field and kept subsequently in a refrigerator until dissected on the same day. Lung tissue samples were taken from all animals and preserved in RNAlater reagent (Ambion, Applied Biosystems, Carlsbad, CA, USA) for genetic analyses. Also blood and serum samples were taken for further serological/epidemiological analyses by pressing the heart on filter paper.

Screening of rodent samples. Lung tissue samples were screened first for hantavirus nucleocapsid (N) protein antigen (N-Ag) by immuno(Western)blotting (WB) as described earlier [Plyusnin et al., 1995]. In brief, approximately 100 mg of tissue was placed into 400-500 µl of Laemmli sample buffer and homogenized by sonication. Aliquots of 10 µl were separated by electrophoresis in 10% sodium dodecyl sulphate-polyacrylamide gels and then immunoblotted with rabbit polyclonal antibody raised against DOBV. Swine anti-rabbit antibodies conjugated with the horse radish peroxidase (Dako, Glostrup, Denmark) were used as secondary antibodies.

Reverse transcription - polymerase chain reaction (RT-PCR) and sequencing.

RNA was extracted from lung tissue samples using the Tripure reagent (Boehringer Mannheim) following recommendations of the manufacturer. RNA was then

For J.Med.Virol.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

135 subjected to RT-PCR to recover partial M segment and S segment sequences,
136 (sequences of primers and other experimental details are available upon request).
137 PCR amplicons were gel-purified with QIAquick Gel Extraction -kit (QIAGEN,
138 GmbH, Germany) and sequenced directly using ABI PRISM™ Dye Terminator
139 sequencing kit (Perkin Elmer/ABI, NJ). SAAV and DOBV genome sequences
140 described in this paper have been deposited to the GenBank sequences database
141 under accession numbers FN813291-7.

142 **Mitochondrial DNA (mtDNA) analysis** was performed as described earlier
143 [Nemirov et al., 2002]. The analysis was done to confirm the species status,
144 particularly that of *A. flavicollis* and *A. sylvaticus* that sometimes are problematic to
145 identify. Briefly, DNA was extracted from lung tissue samples using the Tripure
146 reagent (Behringer Mannheim). A 427 nt-long PCR-product from the D-loop-
147 encoding region was amplified with primers 5'-CCACCATCAGCACCCAAAGCTG-
148 3' and 5'-CTGAAGTAAGAACCAGATGTCTG-3'. The product was purified from the
149 gel and subjected to direct sequencing. For comparison, mtDNA sequences of *A.*
150 *agrarius*, *A. flavicollis* and *A. sylvaticus* were retrieved from the GenBank nucleotide
151 databases.

152 **Phylogenetic analysis.** Multiple nucleotide sequence alignments were prepared
153 manually using the SeqApp 1.9a169 sequence editing program. Phylogenetic analysis
154 was performed using the PHYLIP program package [Felsenstein, 1993]. 500
155 bootstrap replicates were generated using the "Seqboot" program and submitted to
156 the distance matrix algorithm ("Dnadist" program), with the maximum likelihood
157 model for nucleotide substitutions). The resulting distance matrices were analysed
158 with either Neighbor-Joining (NJ) or Fitch-Margoliash (FM) tree-fitting algorithms

For J.Med.Virol.

159 ("Neighbor" and "Fitch" programs, respectively). The bootstrap support values were
160 calculated with the "Consense" program. Hantavirus sequences used for comparison
161 were recovered from the GenBank.

162

For Peer Review

RESULTS

Screening of rodents. In May, 2003, 362 rodents were trapped. Of those, 252 were *Apodemus* mice. In the field, 18 animals were classified preliminarily as *A. agrarius*, 66 as *A. flavicollis* and 151 as *A. sylvaticus*. Of 245 small mammals trapped in September 2004, 80 were classified as *Apodemus* mice. These included 75 yellow-necked mice (*A. flavicollis*) and five wood mice (*A. sylvaticus*). All 332 *Apodemus* mice were subjected to the WB-screening for hantaviral N-Ag and 20 mice (6.3%) were found either strongly positive or weak/suspected-positive. These mice, two *A. agrarius*, six *A. flavicollis*, and 12 *A. sylvaticus*, were trapped at four localities: Migalovci, Nova Gradiška, Loze and Mala Kapela. Lung tissue samples of these mice were further tested for the presence of SAAV/DOBV genome by RT-PCR designed to amplify partial M segment sequence, nucleotides (nt) 1673 to 1989. Six samples, two from *A. agrarius* and four from *A. flavicollis* were positive.

Genetic analysis of SAAV and DOBV strains. Six PCR-amplicons, two recovered from *A. agrarius* mice and four from *A. flavicollis* mice, were purified and sequenced. Hantaviral M segment sequences of all amplicons were recovered successfully and subjected to genetic analyses. Direct sequence comparison revealed that they form two clearly distinct groups. The first group included five sequences originated from samples ##33, 56 (these two sequences were identical and only one, #56, was taken to further analyses), and #64 of the May-2003 set and also samples ##21 and 109 of the September-2004 set. These sequences showed the highest level of identity (90-98%) to the corresponding sequences of DOBV strains from Hungary, Slovenia, Greece and Slovakia suggesting that they belong to this hantavirus

For J.Med.Virol.

species. The sequence identity to SAAV strains was substantially lower: 81-84%:
Consequently, the corresponding wild-type (wt) strains were designated as
DOBV/Croatia_Nova Gradiška/Af56/2003,
DOBV/Croatia_Nova Gradiška/Aa64/2003,
DOBV/Croatia_Gratak-MalaKapela/Af21/2004, and
DOBV/Croatia_Delnice-MalaKapela/Af109/2004,
or DOBV/Croatia56, DOBV/Croatia64, DOBV/Croatia21, and DOBV/Croatia109
for short.

The second group was represented by a single sequence recovered from the
sample #12 of the May-2003 set. This sequence was more closely related to SAAV
sequences originated from Hungary, Slovakia, Central Russia and Estonia (identity
of 90-97%) than to DOBV sequences (81-84%). The corresponding wt strain was
designated as SAAV/Croatia_Migalovci/Aa12/2003, or SAAV/Croatia12 for short.

Phylogenetic analysis of the partial M segment sequences confirmed this
grouping (Fig. 1). All new Croatian DOBV strains clustered together with other
DOBV strains and shared with them a well-supported common ancestor. In
contrast, the new Croatian SAAV strain appeared monophyletic with other SAAV
strains. Within the SAAV cluster, the Croatian strain shared the most recent
common ancestor (TMRCA) with the Hungarian strain and, in turn, these two
shared a more ancient common ancestor with the Slovakian strain. Within the
DOBV cluster, two strains (Croatia56 and Croatia64) grouped together, strain
Croatia109 shared TMRCA with the prototype DOBV strain from Slovenia while the
strain Croatia21 did not group with any particular strain. Interestingly, one of four

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

210 new DOBV strains, DOBV/Croatia64, was found in *A. agrarius* field mouse, i.e. not
211 in the traditional host, suggesting a spillover event.

212 Recovery of partial S segment sequences was also attempted on six RT-PCR-
213 positive samples but only two sequences were obtained: the longer (nt 172 to 1149),
214 for SAAV strain Croatia12 and the shorter (nt 387 to 940), for DOBV strain
215 Croatia56. This made possible to compare the M segment- and S segment- based
216 phylogenies and also to compare the newly recovered sequences to the earlier
217 reported DOBV sequences recovered from Croatian HFRS patients: DOBcro100kt,
218 DOBcro126pb, and DOBcro231lm [Markotic et al., 2002]. On the phylogenetic trees
219 calculated for both longer and shorter regions, the new Croatian DOBV and SAAV
220 strains were located within their respective species clusters (on Fig. 2, the tree
221 calculated for the shorter region, nt 387 to 853, to accommodate the above
222 mentioned Croatian and also the Greek HFRS-originated sequences, is shown).
223 Thus the topologies of M- and S-trees were similar. The SAAV strain Croatia12 was
224 situated within SAAV cluster, next to the group of SAAV strains from Slovakia. The
225 newly recovered, rodent-originated DOBV strain Croatia56 was placed within
226 DOBV-cluster, together with three sequences described previously from Croatian
227 HFRS patients and also with the sequence of prototype DOBV strain from Slovenia.
228 Five DOBV strains from Greece, together with one strain from Slovakia, form
229 another group. It should be noted that the bootstrap support values for both groups
230 were not convincingly high. Interestingly, one of the Croatian sequences derived
231 from a patient, DOBcro100kt, was particularly close to Slovenian sequence and
232 shared with this prototype sequence a reasonably well supported (69%) common
233 ancestor. Of three DOBV sequences recovered from Greek HFRS cases [Papa et al.,

For J.Med.Virol.

1998], two (Greece-PA and Greece -CG) were placed together with high bootstrap support (93%), while the third sequence, Greece-PR, clustered with two DOBV sequences recovered from *A. flavicollis* that were trapped in Ano Poroja, northern Greece [Papa et al., 2001].

238

For Peer Review

DISCUSSION

The results of this study confirmed the circulation of DOBV in Croatia and also presented the first evidence for the presence of SAAV in local *A. agrarius* mice. It was clarified that the sequences described previously from Croatian human HFRS-cases belong to DOBV, not to SAAV. Taken together with the earlier reports on PUUV in Croatia, both in rodents and in association with human HFRS cases [Markotic et al., 2002, Cvetko et al., 2005], the data of (phylo)genetic analyses suggested co-circulation of three pathogenic hantaviruses in this country (the evidence of apathogenic Tula virus was earlier recorded as well [Scharninghausen et al., 2002]). Thus, in that respect, Croatia resembles neighbouring Slovenia and Hungary and also Slovakia where different hantaviruses have been found within a relatively small territory [Avšič-Županc et al., 2000; 2007; Plyusnina et al., 2009; Sibold et al., 1999; 2001]. So far, not a single SAAV genome sequence has been recovered from HFRS patients in these countries suggesting a lower frequency of acute SAAV infection in humans in this part of Europe, in contrast to e.g. Estonia where SAAV-infections are more common than DOBV-infections [Golovljova et al., 2000, 2002, 2007]. Croatian HFRS patients exhibit mostly mild to moderate clinical picture of the disease associated with PUUV, and only a few of them have severe clinical picture associated with DOBV [Markotic et al., 2002]. However, there are some patients with cross-reactive IgG ELISA antibodies to both PUUV and DOBV and usually with moderate to mild clinical picture, which may be possibly linked to SAAV or some other related, but not-yet discovered hantavirus(es) [A. Markotic, unpublished results].

For J.Med.Virol.

Recovery of DOBV genome sequences from the tissue sample of *A. agrarius* mice #64 trapped in Nova Gradiška is of special interest. So far, all *A. agrarius* hantaviral sequences belonged to SAAV species (or, according to another suggested classification, to "DOB-Aa", i.e. the subtype of DOBV associated with *A. agrarius*). All hantavirus-positive rodents analyzed in this study were assigned to *A. flavicollis* or *A. agrarius* species using genetic analysis thus leaving no uncertainties. The most likely explanation for finding of DOBV in another rodent species is a spillover of the virus, the phenomenon registered occasionally for other hantaviruses [see, e.g. Klingström et al., 2002] but, before this study, not for DOBV. Most recently, spillover of SAAV from *A. agrarius* to *A. flavicollis* mice was described in Germany [Schlegel et al., 2009] thus our data on a spillover of DOBV from *A. flavicollis* to *A. agrarius* bring a pleasant symmetry to the case.

Another interesting fact is the existence of hantaviral sequences originated from Caucasian wood mice *A. ponticus* and one human HFRS-case from Krasnodar region of Russian Federation [Klempa et al., 2008]. These sequences form a well-supported group (Fig. 2). The *A. ponticus*- associated isolate Sochi/Ap shows a respectful 10% difference from DOBV and SAAV in GnGc protein sequence and also at least 4-fold differences in focus-reduction neutralization titers of some human sera [Klempa et al., 2008]. Should this virus be considered a distinct species?

During the last two decades the data on hantaviruses associated with different species of *Apodemus* mice in Europe have been accumulating with an increasing speed. Further progress in this field is expected.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements. The work was supported by The Academy of Finland,
Sigrid Juselius Foundation (Finland), projects 143-1430115-0103, 068-1430115-2119
and J-29-2007 funded by The Croatian Ministry of Science, Education and Sports,
and by EU grant GOCE-2003-010284 EDEN (<http://www.eden-fp6project.net/>).

The paper was catalogued by the EDEN steering committee as EDEN0213.

REFERENCES

Avšič-Županc T, Petrovec M, Furlan P, Kaps R, Elgh F, Lundkvist Å. 1999.
Hemorrhagic fever with renal syndrome in the Dolenjska region of Slovenia-A 10-
year survey. Clin Infect Dis 28: 860-865.

Avšič-Županc T, Nemirov K, Petrovec M, Trilar T, Poljak M, Vaheiri A, Plyusnin A.
2000. Genetic analysis of wild-type Dobrava hantavirus in Slovenia: co-existence of
two distinct genetic lineages within the same natural focus. J Gen Virol 81:1747-
1755.

Avšič-Županc T, Petrovec M, Dun D, Plyusnina A, Lundkvist Å, Plyusnin A. 2007.
Puumala hantavirus in Slovenia: analyses of S and M segment sequences
recovered from patients and rodents. Virus Res 123: 204-210.

Cvetko L, Markotic A, Plyusnina A, Margaletic J, Miletic-Medved M, Turk N,
Milas Z, Avšič-Županc T, Plyusnin A. 2005. HFRS and Puumala virus in Croatia
during 2002 outbreak. J Med Virol 77:290-294.

Felsenstein 1993. PHYLIP (Phylogeny Inference Package) version 3.5c. Distributed
by the author. Department of Genetics, University of Washington, Seattle.

For J.Med.Virol.

314 Golovljova I, Vasilenko V, Prukk T, Brus Sjölander K, Plyusnin A, Lundkvist Å.
315 2000. Puumala and Dobrava hantaviruses causing hemorrhagic fever with renal
316 syndrome in Estonia. Eur J Clin Microbiol Infect Dis 19:968-969.

318 Golovljova I, Brus Sjölander K, Lindegren G, Vene S, Vasilenko V, Plyusnin A,
319 Lundkvist Å. 2002. Hantaviruses in Estonia. J Med Virol 68:589-598.

321 Golovljova I, Vasilenko V, Mittzenkov V, Prukk T, Seppet E, Vene S, Plyusnin A,
322 Lundkvist Å. 2007. Characterization of Hemorrhagic Fever with Renal Syndrome
323 caused by hantaviruses, Estonia. Emerg Infect Dis 13:1773-1776.

325 Heyman P, Vervoort T, Escutenaire S, Degraeve E, Konings J, Vandenvelde C,
326 Verhagen R. 2001. Incidence of hantavirus infections in Belgium. Virus Res 77:71-
327 80.

329 Hukic M, Nikolic J, Valjevac A, Šeremet M, Tešic G, Markotic A. 2009. A
330 Serosurvey reveals Bosnia and Herzegovina as a Europe's hotspot in Hantavirus
331 seroprevalence. Epidemiol Infect 2:1-9.

333 ICTV website: www.ictvonline.org

335 Jakab F, Sebok J, Ferenczi E, Horváth G, Szucs G. 2007. First detection of Dobrava
336 hantavirus from a patient with severe haemorrhagic fever with renal syndrome by
337 SYBR Green-based real time RT-PCR. Scand J Infect Dis 39:902-906.

339 Jääskeläinen K, Kaukinen P, Minskaya ES, Plyusnina A, Vapalahti O, Elliott R,
340 Weber F, Vaheri A, Plyusnin, A. 2007. Tula and Puumala hantavirus NSs ORFs are
341 functional and the products inhibit activation of the interferon-beta promoter. J
342 Med Virol.79:1527-1536.

Formatted: Bullets and Numbering

1
2 344 Klempa B, Schütt M, Auste B, Labuda M, Ulrich R, Meisel H, Krüger DH. 2004.
3
4 345 First molecular identification of human Dobrava virus infection in Central Europe.
5 346 J Clin Microbiol 42:1322-1325.
6
7 347
8 348 Klempa B, Fichet-Calvet E, Lecompte E, Auste B, Aniskin V, Meisel H, Denys C,
9
10 349 Koivogui L, ter Meulen J, Krüger DH. 2006. Hantavirus in African wood mouse,
11 350 Guinea. Emerg Infect Dis 12: 838-840.
12
13 351
14 352 Klempa B, Fichet-Calvet E, Lecompte E, Auste B, Aniskin V, Meisel H, Barriere,
15 353 P, Koivogui L, ter Meulen J, Krüger DH. Novel hantavirus sequences in schrew,
16 354 Guinea. 2007. Emerg Infect Dis 13:520-522.
17
18 355
19 356 Klempa B, Tkachenko EA, Dzagurova TK, Yunicheva YV, Morozov VG, Okulova
20 357 NM, Slyusareva GP, Smirnov A, Kruger DH. 2008. Hemorrhagic fever with renal
21 358 syndrome caused by 2 lineages of Dobrava hantavirus, Russia. Emerg Infect Dis
22 359 14: 617-625.
23
24 360
25 361 Klingström J, Heyman P, Escutenaire S, Sjölander KB, De Jaegere F, Henttonen H,
26 362 Lundkvist A. 2002. Rodent host specificity of European hantaviruses: evidence of
27 363 Puumala virus interspecific spillover. J Med Virol 68:581-588.
28
29 364
30 365 Kuzman I, Puljiz I, Turcinov D, Markotic A, Turkovic B, Aleraj B, Andric Z,
31 366 Petkovic D, Tutek V, Herendic B, Iskra M, Pandak N, Misetic Z, Peric L, Jelaska D,
32 367 Majetic-Sekovanic M, Ledina D, Misic-Majerus L, Radonic R. 2003. [The biggest
33 368 epidemic of hemorrhagic fever with renal syndrome in Croatia (in Croatian,
34 369 Abstract - English)]. Acta Med Croatica 57:337-346.
35
36 370
37 371
38 372 Lundkvist Å, Vasilenko V, Golovljova I, Plyusnin A, Vaheri A. 1998. Human
39 373 Dobrava hantavirus infections in Estonia. Lancet 352:369.
40
41 374
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For J.Med.Virol.

- Markotic A, LeDuc JW, Hlaca D, Rabatic S, Šarcevic A, Dašić G, Gagro A, Kuzman I, Barac V, Avšić-Županc T, Beus I, Dekaris D 1996. Hantaviruses are a likely threat to NATO forces in Bosnia and Herzegovina and Croatia. *Nature Med* 2:269-270.
- Markotic A, Nichol ST, Kuzman I, Sanchez AJ, Ksiazek TG, Gagro A, Rabatic S, Zgorelec R, Avšić-Županc T, Beus I, Dekaris D. 2002. Characteristics of Puumala and Dobrava Infections in Croatia. *J Med Virol*; 66:542-551.
- Mustonen J, Brummer-Korvenkontio M, Hedman K, Pqasternak A, Pietilä K, Vaheri A. 1994. Nephropathia epidemica in Finland. A retrospective study of 126 cases. *Scand J Infect Dis* 26:7-13.
- Nemirov K, Vapalahti O, Lundkvist Å, Vasilenko V, Golovljova I, Plyusnina A, Niemimaa J, Laakkonen J, Henttonen H, Vaheri A, Plyusnin A. 1999. Isolation and characterization of Dobrava hantavirus carried by the striped field mouse (*Apodemus agrarius*) in Estonia. *J Gen Virol* 80:371-379.
- Nemirov K, Henttonen H, Vaheri A, Plyusnin A. 2002. Phylogenetic evidence for host switching in the evolution of hantaviruses carried by *Apodemus* mice. *Virus Res* 90: 207-215, 2002. (Corrigendum: *Virus Res* 92:125-126, 2003).
- Nichol ST, Beaty BJ, Elliott RM, Goldbach R, Plyusnin A, Schmaljohn CS, Tesh, RB. 2005. Bunyaviridae. In *Virus taxonomy*. VIIIth report of the International Committee on Taxonomy of Viruses (Fauquet CM, Mayo MA, Maniloff J, Desselberger U, Ball LA - Eds), pp. 695-716. Elsevier Academic Press, Amsterdam.
- Papa A, Johnson AM, Stockton PC, Bowen MD, Spiropoulou CF, Alexiou-Daniel S, Ksiazek TG, Nichol ST, Antoniadis A. 1998. Retrospective serologic and genetic study of the distribution of hantaviruses in Greece. *J Med Virol* 55: 321-327.

1
2 406 Papa A, Nemirov K, Henttonen H, Antoniandis A, Vaheri A, Plyusnin A,
3
4 407 Vapalahti O. 2001. Isolation of Dobrava virus from *A. flavicollis* in Greece. J Clin
5 408 Microbiol 39:2291-2293.
6
7 409
8 410 Plyusnin A, Cheng Y, Vapalahti O, Pejcoch M, Unar J, Jelinkova Z, Lehvslaiho H,
9 411 Lundkvist , Vaheri, A. 1995. Genetic variation in Tula hantaviruses;; sequence
10 412 analysis of the S and M segments of strains from Central Europe. Virus Res 39:
11 413 237-250.
12
13 414
14 415 Plyusnin A, Vapalahti O, Vasilenko V, Henttonen H, Vaheri A. 1997. Dobrava
15 416 hantavirus in Estonia: does the virus exist throughout Europe? Lancet 349:1369-
16 417 1370.
17
18 418
19 419 Plyusnin A, Nemirov K, Apekina N, Plyusnina A, Lundkvist , Vaheri A. 1999.
20 420 Dobrava hantavirus in Russia. Lancet 353:207.
21
22 421
23 422 Plyusnina A, Ferenczi E, Racz GR, Nemirov K, Lundkvist , Vaheri A, Vapalahti
24 423 O, Plyusnin A. 2009. Co-circulation of three pathogenic hantaviruses: Puumala,
25 424 Dobrava and Saaremaa hantaviruses in Hungary. J Med Virol 81:2045-2052.
26
27 425
28 426 Schmaljohn CS, Hjelle B. 1997. Hantaviruses: a global disease problem. Emerg
29 427 Infect Dis 3:95-104.
30
31 428
32 429 Settergren B, Juto P, Trollfors B, wadell G, Norrby SR. 1989. Clinical characteristics
33 430 of nephropathia epidemica in Sweden: prospective study of 74 cases. Rev Inf Dis
34 431 11:921-927.
35
36 432
37 433 Scharninghausen JJ, Pfeffer M, Meyer H, Davis DS, Honeycutt RL, Faulde M. 2002.
38 434 Genetic evidence for tula virus in *Microtus arvalis* and *Microtus agrestis* populations
39 435 in Croatia. Vector Borne Zoonotic Dis 2:19-27.
40 436

- Schlegel M, Klempa B, Auste B, Bemann M, Schmidt-Chanasit J, Büchner T, Groschup MH, Meier M, Balkema-Buschmann A, Zoller H, Krüger DH, Ulrich RG. 2009. Dobrava-Belgrade virus spillover infections, Germany. *Emerg Infect Dis* 15:2017-2020.
- Sibold C, Meisel H, Lundkvist Å, Schulz A, Cifire F, Ulrich R, Kozuch O, Labuda M, Krüger DH. 1999. Short report: simultaneous occurrence of Dobrava, Puumala, and Tula hantaviruses in Slovakia. *Am J Trop Med Hyg* 61:409-11.
- Sibold C, Ulrich R, Labuda M, Lundkvist Å, Martens H, Schutt M, Gerke P, Leitmeyer K, Meisel H, Krüger DH. 2001. Dobrava hantavirus causes hemorrhagic fever with renal syndrome in central Europe and is carried by two different *Apodemus* mice species. *J Med Virol* 63:158-167.
- Tkachenko E, Dekonenko A, Ivanov A, Dzagurova T, Ivanov L, Slonova R, Nirgaleeva R, Stepanenko A, Ivanidze E, Zagidullin I. 1999. Hemorrhagic fever with renal syndrome and hantaviruses in Russia. In: Dodet B, Saluzzo J - Eds. *Emergence and control of rodent-borne viral diseases*. Amsterdam: Elsevier, p.63-72.
- Vapalahti O, Mustonen J, Lundkvist Å, Henttonen H, Plyusnin A, Vaheri A. 2003. Hantavirus infections in Europe (Review). *Lancet Infect Dis* 3:653-661.

Legends to figures

Fig. 1. Phylogenetic tree of hantaviruses based on partial M segment sequences (nt 1749-2019). Abbreviations: **SNV**, Sin Nombre virus, strain NM H10; **NYV**, New York virus, strain H-NY1; **ELMCV**, El Moro Canyon virus, strain RM-97; **LANV**, Laguna Negra virus, strain 510B; **BAYV**, Bayou virus, strain Louisiana; **BCCV**, Black Creek Canal virus; **SEOV**, seoul virus, strain 80-39; **THAIV**, Thailand virus, strain 741; **HTNV**, Hantaan virus, strain 76-118; **SAAV**, Saaremaa virus; **DOBV**, Dobrava-Belgrade virus; **PHV**, Prospect Hill virus, strain PH-1; **TULV**, Tula virus, strain Moravia02v; **PUUV**, Puumala virus, strain Sotkamo; **TOPV**, Topografov virus, strain Ls136v; **KHAV**, Khabarovsk virus, strain MF-43.

Fig. 2. Phylogenetic tree of Dobrava, Saaremaa and related hantaviruses based on partial S segment sequences (nt 387-853). **ASV**, Amur-Soochong virus, strain Soochong1. For other abbreviations, see Fig. 1.