



HAL
open science

Reduced Statherin Reactivity of Human Submandibular Gland in Diabetes

Michela Isola, Paola Solinas, Ernesto Proto, Margherita Cossu, Maria Serenella Lantini

► **To cite this version:**

Michela Isola, Paola Solinas, Ernesto Proto, Margherita Cossu, Maria Serenella Lantini. Reduced Statherin Reactivity of Human Submandibular Gland in Diabetes. *Oral Diseases*, 2010, 17 (2), pp.217. 10.1111/j.1601-0825.2010.01725.x . hal-00599896

HAL Id: hal-00599896

<https://hal.science/hal-00599896>

Submitted on 11 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORAL DISEASES

Reduced Statherin Reactivity of Human Submandibular Gland in Diabetes

Journal:	<i>Oral Diseases</i>
Manuscript ID:	ODI-03-10-OM-1616.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	23-Apr-2010
Complete List of Authors:	Isola, Michela; University, Citomorphology Solinas, Paola; university,, Cytomorphology Proto, Ernesto; Otorhinolaryngology Unit Cossu, Margherita; university, Cytomorphology Lantini, Maria Serenella; university, Cytomorphology
Keywords:	Cell, Pathology



1
2
3
4
5 **Reduced Statherin Reactivity of Human Submandibular Gland in Diabetes**
6
7

8
9 Running head: **Statherin in Diabetic Submandibular Gland**
10

11 M. Isola¹, P. Solinas¹, E. Proto², M. Cossu¹, M.S. Lantini¹
12

13 ¹Department of Cytomorphology, University of Cagliari, Italy; ²Otorhinolaryngology Unit,
14 University Hospital, Cagliari, Italy.
15
16
17

18
19
20
21 Corresponding author: Michela Isola PhD

22 Department of Cytomorphology, University of Cagliari

23 Cittadella Universitaria, 09042 Monserrato (CA) ITALY

24
25 Phone: +39 070 6754029, Fax: +39 070 6754003
26
27

28 Email: misola@unica.it
29
30
31
32

33 Date of submission: 29/03/2010
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

BACKGROUND AND OBJECTIVE: Statherin is a salivary protein involved in the formation of enamel pellicle and in regulation of calcium homeostasis. Diabetes and other pathologies affect both salivary flow and protein secretion by salivary glands, causing increased susceptibility to mucosal infections, tooth demineralization, and caries. The purpose of this study was to compare the statherin expression in submandibular glands of healthy and diabetic subjects.

MATERIALS AND METHODS. Fragments of submandibular glands obtained from diabetic and non diabetic patients were fixed, dehydrated, embedded in Epon Resin and processed for the immunogold histochemistry. The results were statistically evaluated.

RESULTS. Specific statherin labeling was demonstrated in secretory granules of acinar cells in both diabetic and normal samples. The staining was much more intense in the latter compared to those of diabetics. The labeling density was quantified by evaluating the number and spatial distribution of gold particles within the granules. The number of gold particles was significantly lower in glands from diabetics than in control glands.

CONCLUSIONS. The results obtained suggest that a reduced statherin secretion by salivary glands might be partly responsible for a less effective protection of the oral tissues, resulting in an higher incidence of caries and oral infections associated with diabetes.

Key words: Statherin, Diabetes, Immunogold, Human

Introduction

Statherin is a small salivary protein (43 aa) rich in **negatively** charged aminoacids, whose highly polar N-terminus contains two phosphorylated serine residues which confer to the molecule a high affinity for hydroxyapatite (Schlesinger and Hay, 1977). Due to this distinctive property, it binds tooth enamel and form, with proline-rich proteins, the acquired enamel pellicle, an acellular layer which covers the entire surface of the dental crown (Raj *et al.*, 1992; Yao *et al.*, 2003). Adsorbed on enamel, it plays multiple activities related to the maintenance of the dental health. **The inhibitory activity of statherin** on spontaneous precipitation of calcium phosphate has fundamental importance in promoting salivary supersaturation that allows optimal regulation of tooth demineralization and remineralization (Raj *et al.*, 1992; García-Godoy and Hicks, 2008). Moreover, binding on enamel causes changes in the spatial conformation of statherin, so that the C-terminus becomes available to bind microorganisms forming the dental plaque (Gibbons and Hay, 1988). Therefore, statherin is generally included among the numerous salivary proteins that participate in the protection of the mouth by means of antimicrobial, anti-inflammatory or other regulatory activities, such as lysozyme, lactoferrin, immunoglobulins, histatin and defensins (Humphrey and Williamson, 2001; Edgerton *et al.*, 2002; Van Nieuw Amerongen and Veerman, 2002; Dodds *et al.*, 2005).

Recently we investigated the ultrastructural localization of this peptide in human normal salivary glands by a post-embedding immunogold staining technique (Isola *et al.*, 2008). A strong statherin reactivity was demonstrated over the serous granules of parotid and submandibular glands, confirming that these glands are an important source of statherin in human saliva.

Considering the high incidence of caries, tooth loss, periodontal disease and candidiasis in patients affected by diabetes (Lin *et al.*, 1999; Javed *et al.*, 2007; Patiño Marín *et al.*, 2008; Siudikiene *et al.*, 2008) **and given that no literature data document statherin concentration in diabetics**, we undertook the present study in order to verify, using submandibular glands, if salivary glands of diabetic subjects have abnormal statherin expression, which could be in part responsible for oral complications.

Materials and Methods

Transmission Electron Microscopy Tissue Preparation.

Fragments of submandibular glands were obtained from 8 consenting **adult** patients, aged 40–60 years, 4 affected by type 2 diabetes and 4 non diabetic, undergoing surgery **for the removal of tumors from the oral region** at the Otorhinolaryngology Clinic, University of Cagliari. **All samples investigated were taken avoiding the tissues adjacent to the tumors, and appeared normal by the histological analysis. The samples from non diabetics were the same examined in our previous paper (Isola et. al, 2008).** All procedures were approved by the local Institutional Committee for human experimentation at the ASL 8 (Azienda Sanitaria Locale 8), Cagliari. **Each sample was** cut into small pieces and fixed for 2 hr in a mixture of 3% paraformaldehyde and 1% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.2). Then, they were rinsed in cacodylate buffer added with 3.5% sucrose, dehydrated, and embedded in Epon Resin (Glycide Ether 100, Merk, Darmstadt). Ultrathin sections were collected on nickel grids and processed for the immunohistochemical analysis.

Immunohistochemical analysis.

The grids were treated with 1% bovine serum albumin (BSA) and 5% normal rabbit serum (NRS) in phosphate-buffered saline (PBS) solution to block non-specific binding. Then, they were incubated overnight at 4°C with a goat polyclonal antibody specific for statherin (Santa Cruz Biotechnology), diluted 1:50 in 1% BSA and 5% NRS. After rinsing, the grids were incubated for 1 hr at room temperature with the secondary antiserum, a rabbit anti-goat IgG **conjugated to 10 nm gold particles (Sigma) diluted 1:50 in BSA-PBS.** Controls were **incubated** with a non immune goat serum or omitting the primary antibody. After washing with PBS and distilled water, they were stained with uranyl acetate and bismuth subnitrate, and finally observed and photographed in a JEOL 100S transmission electron microscope. **Two grids from each patient were examined, and secretory cells rich in granules were randomly selected and photographed at low magnification.**

Statistical analysis.

Quantitative analysis of randomly chosen micrographs was done with Image Pro Plus (Media Cybernetics, Silver Spring, MD). The gold particles were counted inside 150 serous granules randomly selected from 40 electron micrographs (10 from each diabetic patient) and the labelling density was expressed as number of gold particles / μm^2 . The data values were expressed as mean \pm standard deviation (SD). To determine if the differences were significant, statistical analysis was done using the Student's t-test. The values $p < 0.05$ were considered significant. The proportions of granules with no, weak and intense labelling were also calculated and expressed as percent of the 150 granules.

All the results were then compared with those obtained in micrographs of normal samples following the same procedure. In addition, the statherin labelling in diabetics was estimated as % of statherin labelling in controls.

Results

Statherin reactivity was found in all the submandibular gland samples from both normal and diabetic subjects. The labeling was confined to the serous cells, whose secretory granules appeared the specific site of gold deposition (Figg. 1, 2)

Within the positive granules, gold particles decorated uniformly the dense component in both diabetic and normal samples. However, a weaker staining was easily appreciated in diabetic samples with respect to the controls (Figg. 1, 2).

Statistical analysis showed significant reduction of statherin expression in diabetes ($p < 0.001$), since immunoreactivity was decreased approximately 75% in the diabetic (mean 6,67 / $\mu\text{m}^2 \pm$ SD) versus normal glands (mean 24,25 / $\mu\text{m}^2 \pm$ SD) (Fig. 3). The relative amounts of positive and negative secretory granules also changed: statherin labelling was absent in 3% in normal samples, versus 22% in diabetics. Moreover, granules with intense labelling (10-49 gold particles / μm^2), were not found in diabetic cells, while represented the majority in normal samples (Fig. 4).

Discussion

This study first demonstrates that the statherin reactivity of submandibular glands is significantly decreased in diabetic status. This reduction is clearly demonstrated by the statistical analysis, but it is easily appreciable by the simple observation of the electron micrographs. On the other hand,

our samples showed none of the morphological alterations, specific for parotid glands, such as lipid accumulation, sialosis, reduction of acinar size (Baumann *et al.*, 1985; Carda *et al.*, 2005), probably reflecting the absence of xerostomia and other oral complications in the patients.

The reduced statherin immunoreactivity provides new evidence that salivary glands are affected by diabetes and is thus a further example of altered parameters in diabetic subjects (Hirsch, 2004; Venza *et al.*, 2006; Rao *et al.*, 2009; Rudney *et al.*, 2009). The dry mouth symptoms are undoubtedly effects of a reduction of the salivary flow (Dodds *et al.*, 2005; Sabino-Silva *et al.*, 2009). Great attention has been devoted to the protein composition of diabetic saliva, since protein alterations are generally believed to allow bacteria, bacterial plaque and caries to progress rapidly, significantly hampering the quality of life. Although literature data are often conflicting, the total protein content seems to remain unvaried, while the relative amounts of the different proteins appear significantly changed (Dodds *et al.*, 2000; Rao *et al.*, 2009). For example, lactoferrin and sIgA were increased, while a decrease is noticed for ghrelin (Aydin, 2007), salivatin (Kimura *et al.*, 2001), EGF and prolin-rich proteins (Kasayama *et al.*, 1989; Szczepanski *et al.*, 1998). Salivary amylase may also be affected (Baumann *et al.*, 1985; Aydin, 2007; Mednieks *et al.*, 2009). The decreased statherin reactivity shown here suggests that diabetes causes reduced statherin synthesis or changes in its intracellular processing and storage in submandibular glands. **Unfortunately, at present the biochemical measurement of salivary statherin in diabetics has not been performed to support this assumption. When confirmed, statherin decrease** will represent a further evidence that vulnerability of oral tissues may be partly dependent on a defective enamel pellicle, which poorly protect the teeth, and by an unbalanced calcium metabolism. Moreover, although a direct bactericidal activity of statherin has been still not demonstrated, its decrease could be partly responsible for the increased susceptibility to oral infections, since its ability to bind some bacterial species also would be reduced. Besides, the importance of this peptide in the maintenance of tooth integrity was clearly signified by the strong correlation between its high amounts and absence of dental caries (Vitorino *et al.*, 2005). Therefore, the present results could serve as a starting point for researches that will explain which metabolic signals determine salivary changes in the diabetic status and the relationship between the systemic disorders and the reduced salivary statherin.

Acknowledgments

We wish to thank Mrs Silvana Bernardini and Mr Alessandro Cadau for technical assistance. The investigation was supported by Fondazione Banco di Sardegna (FBS) and PRIN 2008.

References

Aydin S (2007) A comparison of ghrelin, glucose, alpha-amylase and protein levels in saliva from diabetics. *Biochem Mol Biol* 40, 29-35.

Baumann H, Meyer P, Winter K, Lorenz G (1985) Sialochemical and morphology studies of the submandibular gland of the diabetic patient-a contribution to diabetic sialadenosis. *Laryngol Rhinol Otol* 64, 542-546.

Carda C, Carranza M, Arriaga A, Díaz A, Peydró A, Gomez de Ferraris ME (2005) Structural differences between alcoholic and diabetic parotid sialosis. *Med Oral Patol Oral Cir Bucal* 10, 309-314.

Dodds MW, Johnson DA, Yeh CK (2005) Health benefits of saliva: a review. *J Dent* 33, 223-233.

Dodds MW, Yeh CK, Johnson DA (2000). Salivary alterations in type 2 (non-insulin-dependent) diabetes mellitus and hypertension. *Community Dent Oral Epidemiol* 28, 373-381.

Edgerton M, Koshlukova S, Araujo MWB, Patel RC, Dong J, Bruenn JA (2000). Salivary histatin 5 and human neutrophil defensin 1 kill *Candida albicans* via shared pathways. *Antimicrob Agents Chemother* 44, 3310-3316.

García-Godoy F, Hicks MJ (2008). Maintaining the integrity of the enamel surface: the role of dental biofilm, saliva and preventive agents in enamel demineralization and remineralization. *J Am Dent Assoc* 139, 25-34.

1
2
3 Gibbons RJ, Hay DI (1988). Human salivary acidic proline-rich proteins and statherin
4 promote the attachment of *Actinomyces viscosus* LY7 to apatitic surfaces. *Infect Immun* 56,
5 439-445.
6
7

8
9 Hirsch R (2004). Diabetes and periodontitis. *Aust Prescr* 27, 36-38.
10

11 Humphrey SP, Williamson RT (2001) A review of saliva: normal composition, flow and
12 function. *J Prost Dent* 85, 162-168.
13
14

15 Isola M, Cabras T, Inzitari R, Lantini MS, Proto E, Cossu M, et al. (2008) Electron
16 microscopic detection of statherin in secretory granules of human major salivary glands. *J*
17 *Anat* 212, 664-668.
18
19
20

21 Javed F, Näsström K, Benchimol D, Altamash M, Klinge B, Engström PE (2007)
22 Comparison of periodontal and socioeconomic status between subjects with type 2 diabetes
23 mellitus and non-diabetic controls. *J Periodontol* 78, 2112-2119.
24
25
26

27 Kasayama S, Ohba Y, Oka T (1989) Epidermal growth factor deficiency associated with
28 diabetes mellitus. *Proc Natl Acad Sci USA* 86, 7644-7648.
29
30
31

32 Kimura I, Sasamoto H, Sasamura T, Sugihara Y, Ohgaku S, Kobayashi M (2001) Reduction
33 of incretin-like salivatin in saliva from patients with type 2 diabetes and in parotid glands of
34 streptozotocin diabetic BALB/c mice. *Diabetes Obes Metab* 3, 254-258.
35
36
37

38 Lin BP, Taylor GW, Allen DJ, Ship JA (1999) Dental caries in older adults with diabetes
39 mellitus. *Spec Care Dentist* 19, 8-14.
40
41

42 Mednieks MI, Szczepanski A, Clark B, Hand AR (2009) Protein expression in salivary glands
43 of rats with streptozotocin diabetes. *Int J Exp Path* 90, 412-422.
44
45

46 Nieuw Amerongen AV, Veerman ECI (2002) Saliva – the defender of the oral cavity. *Oral*
47 *Dis* 8, 12-22.
48
49

50 Patiño Marín N, Loyola Rodríguez JP, Medina Solís CE, Pontigo Loyola AP, Reyes Macías
51 JF, Ortega Rosado JC, et al. (2008) Caries, periodontal disease and tooth loss in patients with
52 diabetes mellitus types 1 and 2. *Acta Odontol Latinoam* 21, 127-133.
53
54
55
56
57
58
59
60

1
2
3 Raj PA, Johnsson M, Levine MJ, Nancollas GH (1992) Salivary statherin. Dependence on
4 sequence, charge, hydrogen bonding potency, and helical conformation for adsorption to
5 hydroxyapatite and inhibition of mineralization. *J Biol Chem* 267, 5968-5976.
6
7

8
9 Rao PV, Reddy AP, Lu X, Dasari S, Krishnaprasad A, Biggs E, et al (2009) Proteomic
10 identification of salivary biomarkers of type-2 diabetes. *J Proteome Res* 8, 239-245.
11

12
13 Rudney JD, Staikov RK, Johnson JD (2009) Potential biomarkers of human salivary function:
14 a modified proteomic approach. *Arch Oral Biol* 54, 91-100.
15

16
17 Sabino-Silva R, Freitas HS, Lamers ML, Okamoto MM, Santos MF, Machado UF (2009)
18 Na⁺-Glucose cotransporter SGLT1 protein in salivary glands: potential involvement in the
19 diabetes-induced decrease in salivary flow. *Membr Biol* 228, 63-69.
20
21

22
23 Schlesinger DH, Hay DI (1977) Complete covalent structure of statherin, a tyrosine-rich
24 acidic peptide which inhibits calcium phosphate precipitation from human parotid saliva. *J*
25 *Biol Chem* 252, 1689-1695.
26
27

28
29 Siudikiene J, Machiulskiene V, Nyvad B, Tenovuo J, Nedzelskiene I (2008). Dental caries
30 increments and related factors in children with type 1 diabetes mellitus. *Caries Res* 42, 354-
31 362.
32
33

34
35 Szczepanski A, Mednieks MI, Hand AR (1998) Expression and distribution of parotid
36 secretory proteins in experimental diabetes. *Eur J Morphol* 36, 240-246.
37

38
39 Van Nieuw Amerongen A, Bolscher JG, Veerman EC (2004) Salivary proteins: protective
40 and diagnostic value in cariology? *Caries Res* 38, 247-253.
41

42
43 Venza M, Visalli M, Cucinotta M, Ciccì D, Passi P, Teti D (2006) Salivary histamine level
44 as a predictor of periodontal disease in type 2 diabetic and non-diabetic subjects. *J*
45 *Periodontol* 77, 1564-1571.
46
47

48
49 Vitorino R, Lobo MJ, Duarte JR, Ferrer-Correia AJ, Domingues PM, Amado FM (2005) The
50 role of salivary peptides in dental caries. *Biomed Chromatogr* 19, 214-222.
51

52
53 Yao Y, Berg EA, Costello CE, Troxler RF, Oppenheim FG (2003) Identification of protein
54 components in human acquired enamel pellicle and whole saliva using novel proteomics
55 approaches *J Biol Chem* 278, 5300-5308
56
57
58
59
60

Legends

Fig 1. Serous cells of submandibular gland of a diabetic patient, low (A) and high (B) magnification. Gold particles appear exclusively confined within secretory granules. The majority of them express a weak reactivity for statherin. Moreover, a large number of granules are completely negative.

Fig 2. Serous cell of submandibular gland of a non diabetic patient, low (A) and high (B) magnification. Most of secretory granules exhibit a strong reactivity for statherin, often associated with the dense core.

Fig 3. Statherin labeling densities of submandibular secretory granules in diabetics and in controls (gold particles / μm^2). *** = $p < 0.001$

Fig 4. Proportions of submandibular secretory granules with no labelling (group 0), with 1-11 (group 1-11), and with 12-46 (group 12-46) gold particles.

A: diabetic subjects; B: control subjects.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig 1. Serous cells of submandibular gland of a diabetic patient, low (A) and high (B) magnification. Gold particles appear exclusively confined within secretory granules. The majority of them express a weak reactivity for statherin. Moreover, a large number of granules are completely negative.
158x199mm (600 x 600 DPI)




Fig 2. Serous cell of submandibular gland of a non diabetic patient, low (A) and high (B) magnification. Most of secretory granules exhibit a strong reactivity for statherin, often associated with the dense core.

159x230mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig 3. Statherin labeling densities of submandibular secretory granules in diabetics and in controls (gold particles / μm^2). *** = $p < 0.001$
106x69mm (600 x 600 DPI)

Manuscript Copy




Fig 4. Proportions of submandibular secretory granules with no labelling (group 0), with 1-11 (group 1-11), and with 12-46 (group 12-46) gold particles.
92x108mm (600 x 600 DPI)