Search for 2β decay of cerium isotopes with CeCl_3 scintillator

P. Belli^a, R. Bernabei^{a,b,1}, F. Cappella^{c,d}, R. Cerulli^e, F.A. Danevich^f, A. d'Angelo^{c,d}, A. Di Marco^e, A. Incicchitti^c, F. Nozzoli^{a,b}, V.I. Tretyak^f

^aINFN, Sezione di Roma Tor Vergata, I-00133 Rome, Italy

^bDipartimento di Fisica, Università di Roma “Tor Vergata”, I-00133 Rome, Italy

^cINFN, Sezione di Roma, I-00185 Rome, Italy

^dDipartimento di Fisica, Università di Roma “La Sapienza”, I-00185 Rome, Italy

^eINFN, Laboratori Nazionali del Gran Sasso, 67010 Assergi (AQ), Italy

^fInstitute for Nuclear Research, MSP 03680 Kyiv, Ukraine

Abstract

Double beta processes in ^{136}Ce , ^{138}Ce and ^{142}Ce have been searched for by exploiting the active source approach with the help of a 6.9 g CeCl_3 crystal as a scintillator for the first time at the Gran Sasso National Laboratory (LNGS) of the INFN (Italy). The total measurement time is 1638 h; even such a small exposure has allowed the achievement of some improved half-life limits on 2β decay processes in these isotopes at the level of $10^{16} - 10^{18}$ yr. In particular (90% C.L.): $T_{1/2}^{0\nu\varepsilon\beta^+}(^{136}\text{Ce}) \geq 8.8 \times 10^{16}$ yr, $T_{1/2}^{2\nu\varepsilon\beta^+}(^{136}\text{Ce}) \geq 2.4 \times 10^{16}$ yr, $T_{1/2}^{2\nu2K}(^{136}\text{Ce}) \geq 3.2 \times 10^{16}$ yr, $T_{1/2}^{2\nu2K}(^{138}\text{Ce}) \geq 4.4 \times 10^{16}$ yr, and $T_{1/2}^{2\nu2\beta^-}(^{142}\text{Ce}) \geq 1.4 \times 10^{18}$ yr. Possible future perspectives are also briefly addressed.

PACS: 29.40.Mc, 23.40.-s

Keywords: CeCl_3 scintillator; double beta decay; low background.

1 INTRODUCTION

The neutrinoless (0ν) double beta (2β) decay of atomic nuclei $(A, Z) \rightarrow (A, Z \pm 2) + 2e^\mp$ is forbidden in the Standard Model (SM) since it violates the lepton number by two units [1]; however, it is predicted in many SM extensions. The double beta decay experiments are considered the best way to determine an absolute scale of the neutrino mass, to probe the nature of the neutrino, to establish the neutrino mass hierarchy, to search for the existence of right-handed admixtures in the weak interaction, and to test some other effects beyond the SM. During the last two decades the experimental sensitivity to the $0\nu2\beta^-$ decay mode reached levels up to $10^{23} - 10^{25}$ yr [1].

¹Corresponding author. E-mail address: rita.bernabei@roma2.infn.it (R. Bernabei)

In the SM the two-neutrino (2ν) double beta decay is an allowed rare process; to-date it is the rarest decay observed in direct laboratory experiments. It was detected only for 10 nuclides, and the corresponding half-lives are in the range of $10^{18} - 10^{24}$ yr [2, 3].

So far the experimental investigations in this field have been mainly focused on the $2\beta^-$ decays, while the results for double positron decay ($2\beta^+$), electron capture with positron emission ($\varepsilon\beta^+$), and capture of two electrons from the atomic shells (2ε) are much more modest (the most sensitive experiments gave limits on the 2ε , $\varepsilon\beta^+$ and $2\beta^+$ processes at level of $10^{17} - 10^{21}$ yr [2, 4, 5]). This is due to: i) the lower energy releases in the 2ε , $\varepsilon\beta^+$ and $2\beta^+$ processes with respect to those in the $2\beta^-$ decay (which imply lower probabilities for the processes and make difficult the background suppression); ii) the natural abundances of $2\beta^+$ isotopes (typically lower than 1% with only few exceptions). Nevertheless, efforts are in progress in this direction since the studies of the neutrinoless 2ε and $\varepsilon\beta^+$ decays could help to investigate a contribution of right-handed admixtures in weak interaction to the neutrinoless $2\beta^-$ decay [6].

An interesting case is the Cerium; in fact, it offers three double beta decaying isotopes: ^{136}Ce , ^{138}Ce and ^{142}Ce (see Table 1), and the recent development of new scintillating materials containing Ce also allows the exploitation of the efficient “source=detector” approach. In particular, the ^{136}Ce is a very interesting isotope since the high energy release allows the $2\beta^+$ decay mode, which is energetically possible only for 6 candidate-nuclei [2]. Moreover, some resonant neutrinoless 2ε captures in ^{136}Ce to the excited states of the ^{136}Ba are also energetically allowed (see e.g. [7]).

Up to now, the possible 2β decays of Ce isotopes were searched for only in 4 experiments: three of them exploited the active source approach with CeF_3 [8, 9] and $\text{GSO}(\text{Ce})$ crystals [10], and one the passive source approach with HPGe detector using the same CeCl_3 as in the present work [11]. The processes were not detected and only limits on half lives were set; for ^{136}Ce and ^{138}Ce they were in the range of $10^{15} - 10^{17}$ yr; for ^{142}Ce they were in the range of $10^{17} - 10^{19}$ yr.

In this paper we report the results obtained in the investigation of the 2β processes in Ce isotopes using for the first time the CeCl_3 crystal as a scintillator to realize the “source=detector” approach.

Table 1 summarizes the main information about the possible 2β decays in the CeCl_3 crystal.

Table 1: Potentially 2β active nuclides present in CeCl_3 crystals.

Transition	Energy release (keV) [12]	Isotopic abundance (%) [13]	Decay channels	Number of nuclei in 100 g of CeCl_3 crystal
$^{136}\text{Ce} \rightarrow ^{136}\text{Ba}$	2419(13)	0.185(0.002)	$2\varepsilon, \varepsilon\beta^+, 2\beta^+$	4.52×10^{20}
$^{138}\text{Ce} \rightarrow ^{138}\text{Ba}$	693(10)	0.251(0.002)	2ε	6.13×10^{20}
$^{142}\text{Ce} \rightarrow ^{142}\text{Nd}$	1416.7(2.1)	11.114(0.051)	$2\beta^-$	2.72×10^{22}

2 EXPERIMENTAL SET-UP AND MEASUREMENTS

2.1 CeCl_3 crystal scintillator

In recent years scintillating materials containing Cerium have attracted much attention in the developments of new fast scintillators with high density, large light yield, perfect linearity and good energy resolution for applications in fundamental and applied nuclear and particle physics, and in several other fields. In fact – due to the electric dipole allowed 5d-4f transitions in the atomic shell – the Ce^{3+} ions yield fast (10–100 ns) scintillation in the 300–500 nm wavelength range [14, 15]. Scintillating crystals, where Ce is not a dopant but a main constituent, were also studied such as CeF_3 , CeBr_3 , and CeCl_3 .

In particular, the detector used in the present measurements is a cylindrical $\varnothing 13 \times 13$ mm CeCl_3 crystal (6.9 g mass), housed – because of its hygroscopicity – in a combined plastic and copper container ($\varnothing 16 \times 20$ mm external size). Previous measurements were carried out on the detector radiopurity by using a HPGe detector deep underground at LNGS [11], and main scintillation features were studied in ref. [16].

The present data taking has been performed by equipping the crystal with a $\varnothing 7.6 \times 10$ cm Tetrasil-B light guide to further reduce the contribution to the background of the used 3" low background photomultiplier (PMT). The detector has been installed deep underground ($\simeq 3600$ m w.e.) in the low background DAMA/R&D set-up [5, 17] at LNGS. It was surrounded by Cu bricks and sealed in a low radioactive air-tight Cu box continuously flushed with high purity nitrogen gas (stored deeply underground for a long time) to avoid presence of residual environmental Radon. The copper box was surrounded by a passive shield made of 10 cm of high purity Cu, 15 cm of low radioactive lead, 1.5 mm of cadmium and 4/10 cm polyethylene/paraffin to reduce the external background. The whole shield has been closed inside a Plexiglas box, also continuously flushed by high purity nitrogen gas.

The signals from the PMT have been recorded – over a time window of 4000 ns and a sampling time of 1 GSample/s – by a 8 bit DC270 Acqiris transient digitizer (TD).

2.2 Calibration

The energy scale and the energy resolution of the CeCl_3 scintillator detector have been measured by using the following γ sources: i) ^{133}Ba (81 keV and 356 keV); ii) ^{22}Na (511 keV and 1275 keV); iii) ^{137}Cs (662 keV); iv) ^{60}Co (1173 keV and 1333 keV); see Fig. 1. The linearity of the energy calibration is shown in Fig. 2.

The dependence of the energy resolution of the CeCl_3 detector on the energy can be fitted by the function: $\sigma_\gamma/E_\gamma = a + b/\sqrt{E_\gamma}$, where E_γ is the energy of the γ quanta in keV; in particular, for the present assembling: $\sigma_\gamma/E_\gamma = 0.0121 + 0.237/\sqrt{E_\gamma}$.

Figure 1: Energy distributions measured by the CeCl₃ detector in the assembling with light guide when using the following γ sources: (top left) ^{133}Ba ; (top right) ^{22}Na ; (bottom left) ^{137}Cs ; (bottom right) ^{60}Co . See text.

Figure 2: Linearity of the energy response of the CeCl₃ scintillator as measured with the γ sources.

2.3 Pulse shape discrimination capability

The pulse shape discrimination (PSD) capability between α and $\gamma(\beta)$ particles has been investigated in ref. [16] with the crystal directly coupled to the PMT; the pulses, recorded by a 2.5 GSample/s TD over a time window of 500 ns, were analysed and a good discrimination

capability was found for the CeCl_3 crystal.

In the present work the possibility to apply the PSD to the production data collected with the CeCl_3 crystal in the present assembling (light guide and a 1 GSample/s TD) has been investigated. In particular, we use the Optimal Digital Filter Method [18]. For each signal $f(t)$, the numerical characteristic of its shape (shape indicator, SI) was defined as $SI = \sum f(t_k) \times P(t_k) / \sum f(t_k)$, where the sum is over the time channels k , starting from the origin of signal and averaging up to 70 ns. The $f(t_k)$ is the digitized amplitude (at the time t_k) of a given signal. The weight function $P(t)$ was defined as: $P(t) = \{f_\alpha(t) - f_\gamma(t)\} / \{f_\alpha(t) + f_\gamma(t)\}$, where $f_\alpha(t)$ and $f_\gamma(t)$ are the reference pulse shapes for α particles and γ quanta, respectively, obtained by summing up shapes of a few thousand γ or α events. To build the γ reference pulse we considered the data collected with the ^{22}Na and ^{60}Co sources, while for the α we considered the data collected in the production runs in the 1.5 – 2.3 MeV energy range², where the α contribution to the experimental energy distribution is dominant (see later). The distributions of the shape indicator for the two classes of events (energies above 1 MeV in γ scale) are depicted in Fig. 3; they are well described by Gaussian functions. Although there is still a statistical

Figure 3: (*Color online*) - Shape Indicator distributions of the events collected above 1 MeV: α particles (red, lower SI), γ particles (blue, higher SI).

separation between the two populations in the present assembling of the CeCl_3 crystal, it is poorer than in ref. [16]; this can be ascribed to the presence of the light guide, which reduces the far UV component of the scintillation light, and to the smaller sampling frequency. Thus, cautiously in the following analyses we will not profit of it.

2.4 Internal contaminations of the CeCl_3

In ref. [16] the residual contaminations in the CeCl_3 crystal were investigated and discussed in details; moreover, the activity of the CeCl_3 detector (including housing) was also previously

²The α energies are given in terms of calibrations with γ sources, that is in MeV electron equivalent. See also later in the figures.

investigated in ref. [11] on the basis of HPGe γ spectrometry. The present larger exposure and the used approach improve such an investigation in some aspects.

The energy distribution, measured deep underground in the present low background set-up during the 1638 h of data taking, is shown in Fig. 4.

Figure 4: Energy distribution measured deep underground during 1638 h live time inside the multi-component low background passive shield; the superimposed curve (color online) is the overall background fit estimated as in Figs. 9, 11 of ref. [16], where also the main components were shown. The background fit is well in agreement with the experimental data with exception just of the lowest energy bins; for this reason in the determination of the results on the double electron capture decay channels of ^{136}Ce and ^{138}Ce isotopes this fit has not been applied.

The measured background has a similar behaviour as in [16] and is dominated by the internal residual contaminants; in particular, the events below $\simeq 1400$ keV are mainly due to the radioactive decays of ^{138}La and to the β decays of radioactive isotopes in the ^{235}U chain, to which the α particles observed above $\simeq 1400$ keV belong. This has also allowed the determination of the relative light yield for α particles as compared with that for γ quanta – the so called α/β ratio³: $\alpha/\beta = 0.070 + 0.050 \times E_\alpha - 0.0026 \times E_\alpha^2$, where E_α is the energy of the α particles in MeV (E_α between 5.5 and 7.5 MeV).

In particular, the energy distribution of Fig. 4 has been analysed in order to more precisely determine the ^{235}U and ^{138}La content in the crystal, following the same procedures as in ref. [16] where the energy spectrum was compared with Monte Carlo expectations; the values obtained here are given in Table 2. The contents of ^{238}U and of ^{232}Th have been determined by analysing time correlated events in the TD time window (4000 ns) as described in the following.

During the present live time 14 double pulses events have been identified; two examples of them have already been shown in Fig. 7 of ref. [16]. These double pulses events cannot be ascribed to random coincidence; in fact, considering the mean rate measured above the energy threshold of $\simeq 50$ keV and the effective TD time window ($\simeq 3650$ ns), the expected number of random coincidences during the present data taking is: 8.3×10^{-5} .

The measured events can mainly be ascribed to α - α time correlated events produced by α decay of ^{219}Rn into ^{215}Po followed by ^{215}Po α decay ($T_{1/2} \sim 1.781$ ms) into ^{211}Pb (isotopes of the ^{235}U chain). In particular, the first pulse in each event can be due to the ^{219}Rn α decay

³The α/β ratio is defined as the ratio of α peak position in the energy scale measured with γ sources to the energy of α particles (E_α). Because γ quanta interact with the detector by β particles, we use the more convenient term “ α/β ” ratio.

(79.4% α of 6.819 MeV, 12.9% α of 6.553 MeV and 7.5% α of 6.425 MeV), while the second one can be due to the ^{215}Po α decay ($\simeq 100\%$ α of 7.386 MeV). Moreover, the number of the measured time correlated events (14) is in agreement with the expected number of α - α events in the effective TD time window: 16 α - α events, which can be derived from the ^{235}U measured contamination in the detector (see Table 2). Thus, upper limits on Bi-Po events can be set.

A Bi-Po event from ^{232}Th chain is given by the β decay of ^{212}Bi into ^{212}Po followed by the ^{212}Po α decay ($T_{1/2}=299(2)$ ns; 100% α of 8.785 MeV), while a Bi-Po event from ^{238}U chain is given by the β decay of ^{214}Bi into ^{214}Po followed by the ^{214}Po α decay ($T_{1/2}=164.3(2.0)$ μs ; $\simeq 100\%$ α of 7.687 MeV). The detection efficiency in the effective TD time window is $\simeq 100\%$ for Bi-Po events of the ^{232}Th chain and 1.5% for those of the ^{238}U chain.

Considering the expected α - α events from the ^{235}U residual contamination as background with respect to the Bi-Po events, the Bayesian approach [19] allows the estimation of the upper limit on the number of Bi-Po events: 6.61 (90% C.L.) from both ^{238}U and ^{232}Th chains. Thus, the activities: $a_{212\text{Bi}-212\text{Po}} \leq 0.16$ mBq/kg and $a_{214\text{Bi}-214\text{Po}} \leq 11$ mBq/kg (90% C.L.) are derived; then, considering the chains in equilibrium, the upper limits on the ^{232}Th and on the ^{238}U contaminations are achieved (see Table 2).

Table 2: Radioactive contamination of the CeCl_3 crystal. The upper limits (90% C.L.) arise from the analysis of the Bi-Po events. The chains have been considered in equilibrium.

Decay Chain	Nuclide	Activity (mBq/kg)	Contamination from the chain
^{232}Th	^{228}Th	< 0.16	< 40 ppt
^{235}U	^{227}Ac	$=284(2)$	$= 3.52(0.03)$ ppb
^{238}U	^{226}Ra	< 11	< 0.9 ppb
	^{138}La	$=862(31)$	$=948(35)$ ppb

The obtained limits and values are compatible and more precise than those of ref. [11, 16] with the only exception of the ^{238}U residual contamination, which was significantly higher in the HPGe measurements [11]; this can be ascribed – as already discussed also in ref. [16] – to a few mBq ^{238}U residual contamination in the housing materials (copper, optical window, etc.) which are of commercial quality.

3 The investigation on the 2β decay modes

As discussed in the following, there are no clear peculiarities in the energy distribution measured by the CeCl_3 detector, which could be interpreted as double beta decay of Cerium isotopes. Therefore, only lower half-life limits can be set according to the formula:

$$\lim T_{1/2} = N \cdot \eta \cdot t \cdot \ln 2 / \lim S,$$

where N is the number of potentially 2β unstable nuclei, η is the detection efficiency, t is the measuring time, and $\lim S$ is the number of events of the effect searched for which can be excluded at a given confidence level. The values of η in the considered energy interval, of S/η (see later) and of $\lim S$ for each considered decay mode are summarized in Table 3.

The response functions of the CeCl_3 detector for the 2β processes have been simulated with the help of the EGSnrc code [20]; the initial kinematics of the particles emitted in the decays

Table 3: Values of the interesting quantities for the calculation of the $T_{1/2}$ limits on the various decay channels (see later). The overall uncertainty on each η value does not exceed 5%.

Transition	Decay channel	Decay mode	Energy range (keV)	Efficiency η	S/η	S_{lim}/η 90%(68%) C.L.
$^{136}\text{Ce} \rightarrow ^{136}\text{Ba}$	$2\beta^+$	0ν	366-384	48%	$-(99 \pm 81)$	55(16)
		2ν	596-840	17%	$-(197 \pm 393)$	465(221)
	$\epsilon\beta^+$	0ν	1328-1504	54%	$-(53 \pm 55)$	46(16)
		2ν	200-1000	86%	$-(102 \pm 160)$	171(75)
	$2K$	0ν	56-92	79%	(58 ± 48)	137(106)
		2ν	56-92	87%	(53 ± 44)	125(97)
$^{138}\text{Ce} \rightarrow ^{138}\text{Ba}$	$2K$	0ν	56-92	73%	(64 ± 44)	151(117)
		2ν	56-92	87%	(53 ± 44)	125(97)
$^{142}\text{Ce} \rightarrow ^{142}\text{Nd}$	$2\beta^-$	0ν	1320-1496	84%	$-(26 \pm 36)$	36(15)
		2ν	200-1000	85%	$-(92 \pm 158)$	176(80)

has been generated with the DECAY0 event generator [21]. Similar procedure has been used to evaluate the background contribution from the measured residual contaminations.

Preliminarily the experimental sensitivity to each considered double beta decay mode is estimated by using the so-called 1σ approach where the excluded number of the events due to the effect searched for is estimated simply as the square root of the number of background counts in the chosen energy window; if the expected signal is a peak the considered energy interval is $\pm 1\sigma$ around the expected signal peak. Notwithstanding its simplicity, this method gives the right scale of the experimental sensitivity. Then, a suitable refined determination of the experimental limit is performed.

3.1 Double β processes in ^{136}Ce

As already mentioned above, the ^{136}Ce can decay through $2\beta^+$, $\epsilon\beta^+$ and 2ϵ . The used CeCl_3 crystal contains $N_{136} = 3.12 \times 10^{19}$ ^{136}Ce nuclei. The physics for the ^{136}Ce is rather rich. Two positrons could be emitted in its $2\beta^+$ decay with energy up to 375 keV; thus, only the ground state of ^{136}Ba can be populated. The annihilation of these positrons will give rise to four γ 's of 511 keV. In the $\epsilon\beta^+$ decay, not only the ground state but also the first excited level of ^{136}Ba at 818.5 keV can be populated. Finally, in the 2ϵ capture, many excited 0^+ and 2^+ levels of ^{136}Ba can be populated with the subsequent emission of different γ quanta in the de-excitation process.

The experimental sensitivity to the $2\nu 2\beta^+$ decay mode has preliminarily been determined by the 1σ approach. In this case one gets: $S/\eta \leq 422$ events in the energy interval 596 – 840 keV, which is the energy interval of interest for the decay mode (see Fig. 5); this gives a sensitivity: $T_{1/2}^{2\nu 2\beta^+} \geq 9.6 \times 10^{15}$ yr (68% C.L.). Then, a refined analysis is made: the energy distribution is fitted in the energy range 596 – 840 keV by the background model, which takes into account the measured residual contaminations, and the expected signal behaviour as evaluated by EGSnrc code. A $\chi^2/n.d.f. = 60.0/60 = 1.00$ is obtained (see Fig. 5). This approach gives a total number of events, which could be ascribed to the $2\nu 2\beta^+$ decay: $S/\eta = -(197 \pm 393)$ counts. This corresponds (in accordance with the Feldman-Cousins procedure [22]) to $\lim S/\eta = 465(221)$ counts at 90%(68%) C.L., giving for the considered process the half-life limit:

$$T_{1/2}^{2\nu 2\beta^+}({}^{136}\text{Ce}) \geq 0.87(1.83) \times 10^{16} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

Figure 5: (Color online) (*Left*) Energy distribution around 698 keV where it is expected the signal from the $2\nu 2\beta^+$ decay mode of ${}^{136}\text{Ce}$ following the detection of the two positrons and of one of the four γ 's emitted in the annihilation process. The dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown (in dot-dashed blue) the obtained 90% C.L. limit multiplied by a factor twenty. (*Right*) Energy distribution around 375 keV where it is expected the signal from the $0\nu 2\beta^+$ decay mode of ${}^{136}\text{Ce}$ when the annihilation γ 's escape the detector. The dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown (in dot-dashed blue) the obtained 90% C.L. limit multiplied by a factor ten.

In the case of the $0\nu 2\beta^+$ decay mode the maximum sensitivity has been obtained in the energy region around 375 keV. The 1σ approach in the energy interval 366–384 keV gives $S/\eta \leq 93$ events which corresponds to: $T_{1/2}^{0\nu 2\beta^+} \geq 4.4 \times 10^{16} \text{ yr}$ (68% C.L.). Then, the experimental data in the 320–436 keV energy interval has been fitted following the same procedure described above, obtaining a $\chi^2/n.d.f. = 38.1/28 = 1.36$ (see Fig. 5). The total number of events which could be ascribed to the $0\nu 2\beta^+$ decay is: $S/\eta = -(99 \pm 81)$ counts which corresponds to $\lim S/\eta = 55(16)$ counts at 90%(68%) C.L. Thus, the following half-life limit is obtained:

$$T_{1/2}^{0\nu 2\beta^+}({}^{136}\text{Ce}) \geq 0.74(2.53) \times 10^{17} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

In the $\varepsilon\beta^+$ decay, various particles are emitted: positron, X rays and Auger electrons from de-excitations in atomic shells, and γ quanta and/or conversion electrons from de-excitation of a daughter nucleus and from positron annihilation.

The maximum sensitivity to study the $2\nu\varepsilon\beta^+$ decay is obtained in the energy interval 200 – 1000 keV (see Fig. 6). The 1σ approach in this energy interval gives: $S/\eta \leq 164$ events, and $T_{1/2}^{2\nu\varepsilon\beta^+} \geq 2.5 \times 10^{16} \text{ yr}$ (68% C.L.). Then, the energy spectrum has been fitted in the same energy range according to the aforementioned procedure, obtaining a $\chi^2/n.d.f. = 257/199 = 1.29$ (see Fig. 6). The total number of events which could be ascribed to the $2\nu\varepsilon\beta^+$ decay mode is: $S/\eta = -(102 \pm 160)$ counts which corresponds to $\lim S/\eta = 171(75)$ counts at 90%(68%) C.L. and, thus, to:

Figure 6: (Color online) (*Left*) Energy distribution in the 200 – 1000 keV energy interval where it is expected the signal from the $2\nu\varepsilon\beta^+$ decay mode of ^{136}Ce following the detection of one positron and of the X-rays and Auger electrons. The dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown (in dot-dashed blue) the obtained 90% C.L. limit multiplied by a factor fifty. (*Right*) Energy distribution around 1397 keV where it is expected the signal from the $0\nu\varepsilon\beta^+$ decay mode of ^{136}Ce when the annihilation photons escape the detector. The dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown (in dot-dashed blue) the obtained 90% C.L. limit multiplied by a factor three.

$$T_{1/2}^{2\nu\varepsilon\beta^+}(^{136}\text{Ce}) \geq 2.4(5.4) \times 10^{16} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

For the $0\nu\varepsilon\beta^+$ decay, the 1σ approach was applied in the energy interval 1371 – 1423 keV ($\pm 1\sigma$ around the expected signal peak at 1397 keV), however in this case the tail of the nearby composite background structure affect this estimate. Anyhow, in this way we obtain $S/\eta \leq 26$ events, which corresponds to $T_{1/2}^{0\nu\varepsilon\beta^+} \geq 1.6 \times 10^{17} \text{ yr}$ (68% C.L.). Then, the energy spectrum was fitted in energy range 1328 – 1504 keV following the aforementioned procedure, obtaining a $\chi^2/n.d.f. = 17.1/21 = 0.81$ (see Fig. 6). The total number of events which could be ascribed to the $0\nu\varepsilon\beta^+$ decay mode is $S/\eta = -(53 \pm 55)$ counts; this corresponds to $\lim S/\eta = 46(16)$ counts at 90%(68%) C.L., and to:

$$T_{1/2}^{0\nu\varepsilon\beta^+}(^{136}\text{Ce}) \geq 0.88(2.53) \times 10^{17} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

As regards the case of 2ε decay, we have studied just the case of the double electron capture from K shell (binding energy: 37.44 keV). In particular, the $2\nu 2K$ gives a structure around $\simeq 75$ keV (sum of the binding energies of the two electrons from K shell). The 1σ approach in the energy range 72 – 78 keV ($\pm 1\sigma$ around the expected signal peak) gives $S/\eta \leq 39$ events which corresponds to: $T_{1/2}^{2\nu 2K} \geq 1.0 \times 10^{17} \text{ yr}$ (68% C.L.). Then, the energy spectrum was fitted in the energy range 56 – 92 keV by the background model (here – due to the small energy window and the quite smooth energy spectrum – it is assumed as a straight line) and the signal simulated with EGSnrc code (see Fig. 7); a $\chi^2/n.d.f. = 7.4/6 = 1.2$ is obtained. The total number of events which could be ascribed to the $2\nu 2K$ decay mode is: $S/\eta = (53 \pm 44)$ counts which corresponds to $\lim S/\eta = 125(97)$ counts at 90%(68%) C.L., and to:

$$T_{1/2}^{2\nu 2K}({}^{136}\text{Ce}) \geq 3.2(4.2) \times 10^{16} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

Figure 7: (Color online) Energy distribution around 75 keV where it is expected the signal from the $2\nu 2K$ decay mode of ${}^{136}\text{Ce}$ following the detection of the X-rays and Auger electrons as well as the signal from the $0\nu 2K$ decay mode. The horizontal dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown in dot-dashed (blue) line the obtained 90% C.L. limit multiplied by 2 on the $2\nu 2K$ and on $0\nu 2K$ decay modes (both for ${}^{136}\text{Ce}$ and ${}^{138}\text{Ce}$).

In the $0\nu 2\varepsilon$ process, in addition to the particles described above some other particle(s) should be emitted to take away the energy released instead by the two neutrinos in the $2\nu 2\varepsilon$ process. Usually a (bremsstrahlung) gamma quantum is assumed, and in the following we also suppose deexcitation by one γ quantum. Its energy will be equal to $E_\gamma = Q - E_{b1} - E_{b2}$, where E_{b1} and E_{b2} are the binding energies of the first and of the second captured electron on the corresponding atomic shells. For the ${}^{136}\text{Ce}$ this bremsstrahlung gamma quantum will have energy at least of 2344 keV. In the present experimental set-up the low mass of the used CeCl_3 crystal gives a very low peak efficiency for the detection of this gamma quantum of the order of 0.01%, which makes its detection very difficult. However the $0\nu 2K$ can also be identified by the peak at $\simeq 75$ keV, taking into account the different detection efficiency with the respect to the $2\nu 2K$ process. In particular, rescaling by efficiencies the result of the previous fit we obtain for the $0\nu 2K$ decay mode: $S/\eta = (58 \pm 48)$ counts; this corresponds to $\lim S/\eta = 137(106)$ counts at 90%(68%) C.L. and to:

$$T_{1/2}^{0\nu 2K}({}^{136}\text{Ce}) \geq 3.0(3.8) \times 10^{16} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

3.2 Double electron capture in ${}^{138}\text{Ce}$

As already mentioned in Introduction, the ${}^{138}\text{Ce}$ isotope can decay only through 2ε with $Q = 693(10)$ keV. Its natural abundance is $\delta = 0.251\%$; thus, the used CeCl_3 crystal contains $N_{138} = 4.23 \times 10^{19}$ nuclei.

We have studied the 2ε decay of the ${}^{138}\text{Ce}$ analysing the case of the K shell electron capture. In fact, the process of relaxation in the atomic shell of the ${}^{138}\text{Ba}$ after the $2\nu 2K$ capture in ${}^{138}\text{Ce}$ is exactly the same as for the ${}^{136}\text{Ce}$. Thus, we can use our previous calculations for the ${}^{136}\text{Ce}$ just correcting for the different number of ${}^{138}\text{Ce}$ nuclei. It gives the following limit:

$$T_{1/2}^{2\nu 2K}(^{138}\text{Ce}) \geq 4.4(5.7) \times 10^{16} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

The $0\nu 2K$ capture in ^{138}Ce is also similar to the $0\nu 2K$ process in ^{136}Ce . However, the energy of the emitted additional γ quantum (618 keV) is lower for the ^{138}Ce than for the ^{136}Ce (2344 keV). This leads to slightly lower efficiency for the detection of the 75 keV peak. The final values for the $T_{1/2}$ limit is the following:

$$T_{1/2}^{0\nu 2K}(^{138}\text{Ce}) \geq 3.6(4.7) \times 10^{16} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

3.3 $2\beta^-$ decays of ^{142}Ce

As already mentioned in Introduction, the ^{142}Ce isotope can decay through $2\beta^-$ with $Q = 1416.7$ keV. The natural abundance is $\delta = 11.114\%$, thus the used CeCl_3 crystal contains $N_{142} = 1.87 \times 10^{21}$ nuclei.

In the $2\beta^-$ decay the particles emitted are electrons and neutrinos (in the mode 2ν); thus, to detect this process with a good efficiency it is needed that the decaying isotope is inside the detector itself as in the present case.

The signal expected for the $2\nu 2\beta^-$ decay mode is continuous with a maximum around 400 keV (see Fig. 8, left). The maximum sensitivity to study this decay mode is achieved in the

Figure 8: (Color online) *(Left)* Energy distribution in the 200 – 1000 keV energy interval where it is mainly expected the signal from the $2\nu 2\beta^-$ decay mode of ^{142}Ce following the detection of the electrons. The dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown (in dot-dashed blue) the obtained 90% C.L. limit multiplied by a factor fifty. *(Right)* Energy distribution around 1417 keV where the signal from the $0\nu 2\beta^-$ decay mode of ^{142}Ce is expected. The dashed line (red) shows the best-fit curve and the histogram presents the experimental data; there is also shown (in dot-dashed blue) the obtained 90% C.L. limit multiplied by a factor three.

energy interval 200 – 1000 keV (see Fig. 8, left). The 1σ approach in this energy interval gives: $S/\eta \leq 165$ events, and $T_{1/2}^{2\nu 2\beta^-} \geq 1.5 \times 10^{18} \text{ yr}$ (68% C.L.). Then, the energy spectrum has been fitted in the same energy range by the background model and by the $2\beta^-$ signal simulated with

EGSnrc code, obtaining a $\chi^2/n.d.f. = 257/199 = 1.28$ (see Fig. 8, left). The total number of events which could be ascribed to the $2\nu 2\beta^-$ decay mode is: $S/\eta = -(92 \pm 158)$ counts which corresponds to $\lim S/\eta = 176(80)$ counts at 90%(68%) C.L. and, thus, to:

$$T_{1/2}^{2\nu 2\beta^-} (^{142}\text{Ce}) \geq 1.4(3.0) \times 10^{18} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

For the $0\nu 2\beta^-$ decay, the 1σ approach was applied in the energy interval 1391 – 1443 keV ($\pm 1\sigma$ around the expected signal peak at 1417 keV), however in this case the tail of the nearby composite background structure affects this estimate. Anyhow, in this way we obtain $S/\eta \leq 19$ events, which corresponds to $T_{1/2}^{0\nu 2\beta^-} \geq 1.3 \times 10^{19} \text{ yr}$ (68% C.L.). Then, the energy spectrum was fitted in the energy range 1320 – 1496 keV by the background model and the signal simulated with EGSnrc code, obtaining a $\chi^2/n.d.f. = 15.3/21 = 0.73$ (see Fig. 8, right). The total number of events which could be ascribed to the $0\nu 2\beta^-$ decay mode is $S/\eta = -(26 \pm 36)$ counts; this corresponds to $\lim S/\eta = 36(15)$ counts at 90%(68%) C.L., and to:

$$T_{1/2}^{0\nu 2\beta^-} (^{142}\text{Ce}) \geq 0.67(1.62) \times 10^{19} \text{ yr} \quad 90\%(68\%) \text{ C.L.}$$

4 Conclusions

A search for double beta processes in Cerium isotopes has been realized by using for the first time a small CeCl_3 crystal scintillator to realize the "source=detector" approach. In spite of the small exposed mass (6.9 g) and of the limited time of measurements (1638 h) some new improved half-life limits have been obtained as summarized in Table 4, where also the results of the previous most sensitive experiments are given for comparison.

Table 4: Half-life limits on 2β processes (ground state to ground state transitions) in Cerium isotopes at 90%(68%) C.L. Theoretical values for 0ν mode are given for the effective neutrino mass of 1 eV. More theoretical results can be found in compilations [2].

Transition	Decay channel	Decay mode	Exp. $T_{1/2}$ limit (yr)		Theoretical $T_{1/2}$ (yr)
			Present work	Previous results	
$^{136}\text{Ce} \rightarrow ^{136}\text{Ba}$	$2\beta^+$	0ν	$> 0.7(2.5) \cdot 10^{17}$	$> (6.9) \cdot 10^{17}$ [9]	$2.4 \cdot 10^{29}$ [6] $2.7 \cdot 10^{29}$ [23]
		2ν	$> 0.9(1.8) \cdot 10^{16}$	$> 1.8 \cdot 10^{16}$ [10]	$5.2 \cdot 10^{31}$ [6]
	$\varepsilon\beta^+$	0ν	$> 0.9(2.5) \cdot 10^{17}$	$> 3.8 \cdot 10^{16}$ [10]	$4.7 \cdot 10^{26}$ [6] $4.0 \cdot 10^{26}$ [23]
		2ν	$> 2.4(5.4) \cdot 10^{16}$	$> 2.6 \cdot 10^{15}$ [11]	$9.2 \cdot 10^{23}$ [6] $6.0 \cdot 10^{23}$ [24]
	$2K$	0ν	$> 3.0(3.8) \cdot 10^{16}$	$> 6.0 \cdot 10^{15}$ [10]	–
		2ν	$> 3.2(4.2) \cdot 10^{16}$	$> 2.7 \cdot 10^{16}$ [8]	$3.2 \cdot 10^{18} - 6.4 \cdot 10^{19}$ [25] $(3.2 - 5.1) \cdot 10^{21}$ [26] $9.6 \cdot 10^{21}$ [24]
	$2K$	0ν	$> 3.6(4.7) \cdot 10^{16}$	$> 1.9 \cdot 10^{15}$ [11]	–
		2ν	$> 4.4(5.7) \cdot 10^{16}$	$> 3.7 \cdot 10^{16}$ [8]	$\geq 2.1 \cdot 10^{26}$ [27]
$^{142}\text{Ce} \rightarrow ^{142}\text{Nd}$	$2\beta^-$	0ν	$> 0.7(1.6) \cdot 10^{19}$	$> (1.5) \cdot 10^{19}$ [9]	$2.8 \cdot 10^{24}$ [28]
		2ν	$> 1.4(3.0) \cdot 10^{18}$	$> 1.6 \cdot 10^{17}$ [10]	$2.2 \cdot 10^{20} - 4.2 \cdot 10^{21}$ [28] $2.3 \cdot 10^{23}$ [29]

This supports the interest in realizing larger mass and longer exposure deep underground with new CeCl_3 crystal scintillators exploiting the active source technique. A further relevant topic is the future preliminary selection of all the materials and of the growing/handling procedures as well as the development of chemical/physical purification techniques in order to improve the radiopurity of the crystal and housing materials. In particular, the ^{138}La contamination could be related mainly with the growing process; in fact, the crystal used here was realized just after the growth of a LaCl_3 scintillator. Moreover, the use of a larger TD sampling frequency and of different optical read-out should be considered in order to have the possibility to effectively apply the PSD for internal α background reduction when of interest. In addition, the $2\beta^+$ and $\varepsilon\beta^+$ processes can give strong signature when coincidences with other close detectors are investigated. It should be also stressed an advantage of much better energy resolution of CeCl_3 crystal scintillator ($\text{FWHM} \approx 5\%$ at 662 keV γ line of ^{137}Cs) in comparison with f.e. CeF_3 crystal scintillators ($\approx 18\%$ at the same energy) [8].

Further improvements in sensitivity can be reached by using enriched ^{136}Ce , increasing the detection efficiency by using larger CeCl_3 detector, and developing CeCl_3 scintillators with lower level of radioactive contamination. An experiment involving ≈ 100 kg of crystals enriched in ^{136}Ce to 20% (5×10^{25} nuclei of ^{136}Ce) could reach over 5 years of measurements the half-life sensitivity $T_{1/2} \approx 10^{25}$ yr (supposing zero background). Such a sensitivity could contribute to our understanding of the neutrino mass mechanism and right-handed currents in neutrinoless processes [6]. Particular interest is offered by the $2\nu 2\varepsilon$ decay mode of the ^{136}Ce , whose half-life is expected to be in the range $10^{18} - 10^{22}$ yr by theoretical estimates [2]; thus, this process should be observed in such an experiment.

5 Acknowledgments

The authors would like to express gratitude to the referees for their careful reading of the manuscript and their useful comments.

References

- [1] F.T. Avignone III, S.R. Elliott, J. Engel, Rev. Mod. Phys. 80 (2008) 481;
H.V. Klapdor-Kleingrothaus, Int. J. Mod. Phys. E 17 (2008) 505;
H. Ejiri, J. Phys. Soc. Japan 74 (2005) 2101;
F.T. Avignone III, G.S. King, Yu.G. Zdesenko, New J. Phys. 7 (2005) 6;
S.R. Elliot, J. Engel, J. Phys. G: Nucl. Part. Phys. 30 (2004) R183;
J.D. Vergados, Phys. Rep. 361 (2002) 1;
S.R. Elliot, P. Vogel, Ann. Rev. Nucl. Part. Sci. 52 (2002) 115;
Yu.G. Zdesenko, Rev. Mod. Phys. 74 (2002) 663.
- [2] V.I. Tretyak, Yu.G. Zdesenko, At. Data Nucl. Data Tables 61 (1995) 43; 80 (2002) 83.
- [3] A.S. Barabash, Phys. Rev. C 81 (2010) 035501.
- [4] Ju.M. Gavriljuk et al., Phys. At. Nucl. 69 (2006) 2124;
A.S. Barabash et al., J. Phys. G 34 (2007) 1721;
A.S. Barabash et al., Nucl. Phys. A 785 (2007) 371;

- H.J. Kim et al., Nucl. Phys. A 793 (2007) 171;
A.S. Barabash et al., Nucl. Phys. A 807 (2008) 269;
J. Dawson et al., Nucl. Phys. A 799 (2008) 167;
A.S. Barabash et al., Phys. Rev. C 80 (2009) 035501;
P. Belli et al., Eur. Phys. J. A 42 (2009) 171;
N.I. Rukhadze et al., J. Phys. Conf. Ser. 203 (2010) 012072.
- [5] P. Belli et al., Phys. Lett. B 658 (2008) 193;
P. Belli et al., Eur. Phys. J. A 36 (2008) 167;
P. Belli et al., Nucl. Phys. A 826 (2009) 256.
- [6] M. Hirsch et al., Z. Phys. A 347 (1994) 151.
- [7] A.S. Barabash, Phys. At. Nucl. 73 (2010) 162.
- [8] P. Belli et al., Nucl. Instrum. Meth. A 498 (2003) 352.
- [9] R. Bernabei et al., Il Nuovo Cim. A 110 (1997) 189.
- [10] F.A. Danevich et al., Nucl. Phys. A 694 (2001) 375.
- [11] P. Belli et al., Nucl. Phys. A 824 (2009) 101.
- [12] G. Audi, A.H. Wapstra, C. Thibault, Nucl. Phys. A 729 (2003) 337.
- [13] J.K. Bohlke et al., J. Phys. Chem. Ref. Data 34 (2005) 57.
- [14] C.W.E. van Eijk, Nucl. Instrum. Meth. A 460 (2001) 1.
- [15] K.W. Kramer, P. Dorenbos, H.U. Gudela, C.W.E. van Eijk, J. Mat. Chem. 16 (2006) 2773.
- [16] F. Cappella et al., Nucl. Instrum. Meth. A 618 (2010) 168.
- [17] P. Belli et al., Phys. Rev. C 76 (2007) 064603;
P. Belli et al., Nucl. Phys. A 789 (2007) 15;
R. Bernabei et al., Ukr. J. Phys. 51 (2006) 1037;
R. Bernabei et al., Nucl. Instrum. Meth. A 555 (2005) 270;
R. Cerulli et al., Nucl. Instr. Meth. A 525 (2004) 535;
P. Belli et al., Nucl. Instr. Meth. A 498 (2003) 352;
R. Bernabei et al., Nucl. Phys. A 705 (2002) 29;
P. Belli et al., Nucl. Phys. B 563 (1999) 97;
P. Belli et al., Astropart. Phys. 10 (1999) 115;
R. Bernabei et al., Il Nuovo Cim. A 110 (1997) 189;
R. Bernabei et al., Astropart. Phys. 7 (1997) 73.
- [18] E. Gatti, F. de Martini, Nuclear Electronics 2 (IAEA, Vienna, 1962) p. 265.
- [19] R.M. Barnett et al., Phys. Rev. D 54 (1996) 159.
- [20] I. Kawrakow, D. W. O. Rogers, The EGS Code System, NRCC Report PIRS-701 (2003).

- [21] O.A. Ponkratenko et al., Phys. At. Nucl. 63 (2000) 1282;
V.I. Tretyak, to be published.
- [22] G.J. Feldman, R.D. Cousins, Phys. Rev. D 57 (1998) 3873.
- [23] J. Suhonen, M. Aunola, Nucl. Phys. A 723 (2003) 271.
- [24] O.A. Rumyantsev, M.H. Urin, Phys. Lett. B 443 (1998) 51.
- [25] J. Suhonen, Phys. Rev. C 48 (1993) 574.
- [26] O. Civitarese, J. Suhonen, Phys. Rev. C 58 (1998) 1535.
- [27] J. Abad et al., J. de Physique 45 (1984) C3-147.
- [28] A. Staudt, K. Muto, H.V. Klapdor-Kleingrothaus, Europhys. Lett. 13 (1990) 31.
- [29] A. Bobyk et al., Eur. Phys. J. A 19 (2004) 327.