Lung function and overweight in school aged children after early childhood wheezing

Virpi H Sidoroff, Mari K Hyvärinen, Eija Piippo-Savolainen, Matti Korppi

► To cite this version:

Virpi H Sidoroff, Mari K Hyvärinen, Eija Piippo-Savolainen, Matti Korppi. Lung function and overweight in school aged children after early childhood wheezing. *Pediatric Pulmonology*, 2010, 10.1002/ppul.21386 . hal-00604878

HAL Id: hal-00604878

<https://hal.science/hal-00604878>

Submitted on 30 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lung function and overweight in school aged children after early childhood wheezing

Journal:	<i>Pediatric Pulmonology</i>
Manuscript ID:	PPUL-10-0168.R1
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	01-Nov-2010
Complete List of Authors:	Sidoroff, Virpi; University of Eastern Finland; Kuopio University Hospital, Department of Pediatrics Hyvärinen, Mari; Kuopio University Hospital, Department of Pediatrics Piippo-Savolainen, Eija; Kuopio University Hospital, Department of Pediatrics Korppe, Matti; Kuopio University Hospital, Department of Pediatrics
Keywords:	child, spirometry, obstruction, overweight, obesity, lung function, wheezing

SCHOLARONE™
Manuscripts

Lung function and overweight in school aged children after early childhood wheezing

Sidoroff Virpi¹ MD, Hyvärinen Mari K² MD, Piippo-Savolainen Eija² MD, PhD, Korppi Matti^{2,3} MD, PhD, Prof

¹ University of Eastern Finland, School of Medicine, Department of Pediatrics, Kuopio, Finland

² Department of Pediatrics, Kuopio University Hospital, Kuopio, Finland

³ Pediatric Research Center, Tampere University Hospital, Tampere, Finland

Corresponding author:

Sidoroff Virpi, Eteläkatu 10 b 7, 80110 Joensuu, Finland,

email: vpietika@hytti.uku.fi, tel +358-400015215

Abbreviated title: Lung function and overweight in wheezy children

Summary

Background and purpose. Recently, obesity has been connected with wheezing, asthma and reduced lung function. Most previous studies have been cross-sectional. The aim of the present follow-up study was to evaluate the association of preceding or current overweight or obesity with lung function at early and late school age after early childhood wheezing.

Material and methods. From the 100 children hospitalized for infection associated wheezing at <24 months of age, 83 attended the control visit at 4.0 years, 82 at 7.2 years and 81 at 12.3 years of age. Flow-volume spirometry was performed in 79 children at 7.2 years and in 80 children at 12.3 years of age. The weight status was assessed by calculating body mass index (BMI) at all visits. Age- and gender-specific BMI standard deviation scores (BMI-SDS) of >1.3 SD and > 2.0 SD were defined to mean overweight and obesity, respectively.

Results. Overweight at both 7.2 and 12.3 years of age was associated with decreased FEV_1/FVC (forced expiratory volume in one second/ forced vital capacity). Overweight and obesity at 7.2 years of age were associated with decreased FEV_1/FVC and MEF_{50} (maximal expiratory flow at 50 % of FVC) at 12.3 years of age. The results were similar by continuous and categorized analyses, being robust to adjustments for viral findings during early childhood wheezing and asthma maintenance medication at school age.

Conclusion. Overweight and obesity are significant risk factors for reduced lung function at school age after early childhood wheezing. Thus, early-life wheezers should avoid excessive weight gain during childhood.

Keywords: child, spirometry, obstruction, overweight, obesity, lung function, wheezing

Abbreviations:

BMI	body mass index
BMI-SDS	body mass index standard deviation score
TLC	total lung capacity
FRC	functional residual capacity
FVC	forced vital capacity
FEV ₁	forced expiratory volume in one second
FEV ₁ /FVC	forced expiratory volume in one second/forced vital capacity
MEF ₅₀	maximal expiratory flow at 50% of FVC
MEF ₂₅	maximal expiratory flow at 25% of FVC
aOR	adjusted odds ratios
CI	confidence intervals
RSV	respiratory syncytial virus

Introduction

Asthma and overweight or obesity have increased concomitantly among both children and adults during the last three decades in western countries¹⁻³. The studies have suggested an association between wheezing symptoms and obesity⁴⁻⁹, and also between asthma diagnoses and obesity¹⁰⁻¹³. When pulmonary function and bronchial reactivity have been studied objectively, the results have been mainly negative^{5, 14-18}.

In adults, obesity decreases both lung volume, as documented by reduced total lung capacity (TLC), forced vital capacity (FVC) and functional residual capacity (FRC), and airway flows, as documented by decreased forced expiratory volume in one second (FEV₁) and increased airway resistance^{19,20}. In addition, obesity leads to obstruction in peripheral airway²¹, as documented by decreased flows at 50% (MEF₅₀) or at 25% (MEF₂₅) of FVC. However, the largest impact seems to target at expiratory reserve volume, which means that restriction is the characteristic lung function disorder in obesity^{22,23}. In addition, weight reduction in obese adults elicited an improvement of lung function²⁴.

In children with asthma suggestive symptoms, high body mass index (BMI) was associated with subnormal lung function in one study¹⁸, but not in two others^{4,5}. In healthy children, lung function was either impaired²⁵, improved in boys and girls²⁶ or improved only in girls²⁷ by excessive weight gain.

We have prospectively followed-up a cohort of children hospitalized for infection associated wheezing at <24 months of age in 1992-1993. The weights and heights of the children were measured at follow-up visits at the median ages of 4.0, 7.2 and 12.3 years, and spirometry was performed at both visits at school age²⁸⁻³⁰.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The aim of the present study was to evaluate the association between overweight or obesity and lung function at school age in former early-life wheezers, with a special focus on the influence of an emerging overweight.

For Peer Review

Material and methods

Study design

One-hundred consecutive children, hospitalized for wheezing associated respiratory infection at age 1-23 months, were recruited into this follow-up study in 1992 -1993³¹.

Maintenance therapy for asthma was one of the exclusion criteria, and the wheezing episode was the first one in 87% of the included children³². The basic data were collected by interviewing the parents during hospitalization using a structured questionnaire. Seven respiratory viruses, including respiratory syncytial virus (RSV), were studied by antigen and antibody assays³². Later, viral studies were supplemented by polymerase chain reaction to RSV and rhinoviruses in frozen samples^{33,34}. In all, RSV was identified in 29/100 and rhinovirus in 27/81 cases^{33,35}.

Three follow-up visits were prospectively organized at the median ages of 4.0 years, 7.2 years and 12.3 years²⁸⁻³⁰. Asthma was present in 33/82 (40.2%) children at the 7.2-year visit²⁹ and in 32/81 (39.5%) children at the 12.3-year visit³⁰. Data on medical history, including the use of asthma medication during the preceding 12 months, were collected by interviewing the parents.

Assessment of weight status

The weights and heights of the children, measured by an experienced nurse using standard methods, were registered in all attending children at all controls visits. Body mass index (BMI) was calculated by using the equation: weight (kg) / [height (m)]². Due to the lack of Finnish references, we applied British gender- and age-specific population-based references. BMI standard deviation score (BMI-SDS), also called Z-score, was

assessed by an obesity calculator (NIBHI 2007)³⁶. This calculator applies British growth references from 1990, produced by the LMS method in 1995^{37,38}, and revised in 1996³⁹. The LMS method summarizes the distribution of BMI at each age by its median (M), coefficient of variation (S), and skewness expressed as a Box-Cox power (L) required to transform the data to normal⁴⁰.

In the present study, we used the same cut-off points for overweight and obesity as proposed by the British study evaluating the limits of obesity and overweight in different populations⁴¹. The cut-off point for overweight was BMI >91 percentile (approximating BMI-SDS >1.3) and for obesity BMI >98 percentile (approximating BMI-SDS >2.0)^{19,41}. In the British population, these cut-off points corresponds BMI 25 kg/m² and BMI 30 kg/m² levels, respectively, in the 18 years old young adults⁴¹.

Measurement of lung function

Lung function was studied by flow-volume spirometer (FVS) (Medikro, Kuopio, Finland) at the 7.2-year and 12.3-year study visits. The parameters were expressed as percentages of the gender-specific height-related reference values (% of predicted) for Finnish children^{42,44}. The parameters registered were FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FVC, MEF₅₀ (maximal expiratory flow at 50 % of FVC) and MEF₂₅ (maximal expiratory flow at 25 % of FVC). Either the measurement with the highest FEV₁ (at the 7.2-year study)²⁹ or the composite maximal curve (at the 12.3-year study)³⁰ was in each child recorded for further analyses. The cut-off limits for abnormal values were 80 % for FEV₁, 88 % for FEV₁/FVC, 62 % for MEF₅₀ and 48 % for MEF₂₅^{17,30,43}.

Lung function data were available from 79 children at the 7.2-year and from 81 children at the 12.3-year study visits. The numbers of children with anti-inflammatory maintenance medication for asthma during the 12 preceding months were 29/79 (36.77%) at age of 7.2 years and 18/80 (22.5%) at age of 12.3 years. At the 4.0-year visit, weight status was assessed for 74 children with lung function data available at the 7.2-year visit.

Statistics

The data were analysed using SPSS 17.0 software (SPSS Inc. Chicago, IL, USA). Mann-Whitney U-test was used in univariate analyses and analysis of variance (ANOVA) in multivariate analyses for continuous FVS parameters. In the previous studies in this cohort, RSV etiology of wheezing in early childhood was associated with a non-reversible restrictive pattern of lung function and non-RSV etiology (mostly rhinovirus) with an increased asthma risk and a reversible obstructive pattern of lung function at school age^{43, 44}. Therefore, analyses by ANOVA were adjusted also for RSV etiology of wheezing in early childhood, in addition to adjustment with anti-inflammatory asthma medication during the preceding 12 months before the study visit at school age. Logistic regression analysis, adjusted for RSV etiology of early childhood wheezing and anti-inflammatory asthma medication during the 12 preceding months before the study visit, was applied for categorized (by the limits of abnormal values in % predicted) FVS parameters, and the results were expressed as adjusted odds ratios (aOR) and their 95 % confidence intervals (95% CI).

Ethics

The study was approved by the joint Research Ethics Committee of Kuopio University and Kuopio University Hospital. Informed written consent was obtained from the parents of the children.

For Peer Review

Results

Overweight was present in 6.0% of children at 4.0 years, in 22.2% at 7.3 years and in 33.3% at 12.3 years of age. Less than 4% of children were obese at 4.0 years of age. The proportion of obese children was 8.6 % at 7.3 years and 19.8% at 12.3 years of age. Three of 5 overweight children at age 4.0 years were overweight at age 7.2 years, and all 3 obese children were obese, respectively. Fifteen of the 16 overweight children at age 7.2 years were overweight at age 12.3 years, and 4 of the 6 obese children were obese, respectively.

In univariate analyses at 7.2 years of age, the only significant association was between current overweight and lower FEV_1/FVC (Table 1). There were no significant associations between previous (4.0 years of age) overweight or obesity and any FVS parameter at age 7.2 years (Data not shown). In multivariate analyses, the association between obesity and lower FEV_1/FVC changed to significant (Table 1). All other associations between previous or current overweight or obesity and FVS remained as unchanged.

In univariate analyses at 12.3 years of age, previous (7.2 years of age) overweight was associated with lower FEV_1/FVC , and current (12.3 years of age) overweight was associated with lower FEV_1/FVC , MEF_{50} and MEF_{25} (Tables 2). Similarly, previous obesity was associated with lower FEV_1/FVC , and current obesity was associated with lower FEV_1/FVC , MEF_{50} and MEF_{25} (Table 3). In multivariate analyses, the association between current overweight and MEF_{50} changed to non-significant (Table 2). All other associations between previous or current overweight or obesity and FVS remained as unchanged.

In the logistic regression analyses, both previous and current overweight were significant risk factors for $FEV_1/FVC < 88\%$ at both 7.2 years (Table 4) and at 12.3 years of age (Table 5). Since overweight was highly permanent from 4.0 to 7.3 years of age and from 7.2 to 12.3 years of age, respectively, previous and current overweight could not be incorporated as variables in the same model. Therefore, the design of the study did not allow to assess their influences separately. Current obesity was a significant risk factor for $FEV_1/FVC < 88\%$ (aOR 8.07, 95% CI 2.10-31.05) and for $MEF_{25} < 48\%$ (aOR 4.34, 95% CI 1.06-17.79) at 12.3 years of age.

Discussion

There are three main results in the present study. First, the number of overweight and obese children increased constantly by age during the follow-up after early childhood wheezing until the median age of 12.3 years. Finally, a third was overweight, and about 20% were obese. The figures are higher than those calculated from self-reported weights and heights in 12-18 years old Finnish children in 1999, when 16.7% of the boys and 9.8% of the girls were overweight, and 1.4% and 0.4% were obese, assessed by using age- and gender-related international reference values⁴⁵. Second, current overweight was associated with decreased FEV₁/FVC at both 7.2 and 12.3 years of age. Overweight children had also reduced flows in small airways by MEF₂₅ and obese children by MEF₂₅ and MEF₅₀ at age 12.3 years. FEV₁/FVC is considered as the most sensitive and reliable measure of obstruction, but the reliability of MEF₂₅ has been questioned. Third, overweight and obesity at early school age preceded reduced FEV₁/FVC at age 12.3 years. But, since overweight was highly permanent from early to late school age, the influences of previous and current overweight could not be assessed separately.

The association between excessive weight gain and reduced lung function has been well documented in adults^{14-16,20,22}. In a Finnish study, weight reduction in obese adults even elicited an improvement of lung function²⁴. The effect of excessive weight gain to lung function seems to depend on the degree of obesity and on the presence and severity of asthma. When lung function was studied in 52 non-asthmatic non-obese, in 53 asthmatic non-obese, in 52 obese non-asthmatic and in 53 asthmatic obese adults, lung function was reduced in patients with asthma and obesity, and in addition, FVC and FRC were reduced

in obese adults also with no asthma¹⁶. In a birth cohort from New Zealand, an association was found between increasing BMI and decreasing FEV₁/FVC in 9-26 years old females, but not in males¹⁵. In a study on difficult-to-treat asthma, FEV₁ was increased and FRC and total lung capacity were decreased in obese patients⁴⁶. Thus, controlled studies in adults with asthma^{16,46}, as well as population-based studies in healthy adults^{15,22,23}, highlight the connection between obesity and decreased expiratory reserve volumes, which means that restriction is the characteristic lung function disorder in obesity.

In a population study from China, overweight had a beneficial effect on lung function in 7 to 20 years old healthy girls²⁷. In a population study from Mexico, lung function improved when weight increased in 8-11 years old healthy children, but after that age, lung function first reached a plateau and then decreased among the obese youths⁴⁷. Lung function by spirometry was compared between 657 overweight and 196 normal weight Greek children, and FVC, FEV₁, MEF_{25-75%} and FEV₁/FVC were reduced in overweight children¹⁸. In a population study from Australia, spirometry was performed in 2464 school children⁴⁸. Height-related FVC and FEV₁ increased by growth, but both FVC and FEV₁ decreased in relation to increasing overweight. The result was opposite in an other Australian population study in 5993 children; all variables in the FVS were, irrespective of BMI, within normal limits, but FEV₁ and FVC in the lowest BMI quintile were significantly lower than in other quintiles⁴. All these studies are cross-sectional with no follow-up available. The results seem confusing, but there may be following consistency; mild overweight may be beneficial but when substantial overweight or obesity develops, the effect turns to non-beneficial.

The results of the present follow-up study after early childhood wheezing showed that overweight was associated with reduced FEV_1/FVC at 7.2 and 12.3 years of age and reduced MEF_{50} and MEF_{25} at 12.3 years of age, in disagreement with most population studies in healthy children^{4,5,27,46,47}. However, the associations may be different in children with asthma; about 40% of our patients had asthma and all had wheezed in early childhood. Lung function and weight status were studied as a part of the childhood asthma management program (CAMP) in 1041 American children aged 5-12 years⁴⁹, and increasing BMI was associated with increasing FEV_1 and FVC, but decreasing FEV_1/FVC . The increase in FVC and decrease in FEV_1/FVC are in accordance with the results of our present study. In the CAMP study, weight status and lung function were studied by a cross-sectional design at enrolment of the patients, and no follow-up data have been, at least thus far, published. We found no evidence for a restrictive pattern of lung function disorder in overweight or obese children, though suggested to be present in obese asthmatic adults^{15,16,22,23,46}, and though was demonstrable after early childhood wheezing caused by RSV in 12.3 years old children of the present study^{43,44}.

The mechanisms leading to lung function disorders in association with overweight or obesity may be narrowing of the small peripheral airways²¹, in line with decreased MEF_{50} and MEF_{25} values in the overweight and obese children at the 12.3-year visit of the present study. In addition, decreased compliance of either the lungs due to narrow peripheral airways or the chest wall due to body adiposity may lead to subnormal lung function, including both decreased airway flows which mean an obstructive lung function disorder and decreased lung volumes which means a restrictive lung function disorder⁴⁸. The effect of excessive weight gain on lung function may be age-dependent, which

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

means that different mechanisms may influence in children and in adults and in children at different ages. In a case-control study from Denmark, FEV₁ and FVC were reduced in 193 obese men, compared with normal weight 215 controls, at 27 years of age, but after adjustments with current BMI, childhood BMI at 7 years of age was positively associated with both FEV₁ and FVC ⁵⁰.

The strengths of the present study are the long, over 10 years prospective follow-up time and the fact that weights and heights of the children were measured at all control visits by experienced nurses using calibrated scales, thus allowing a reliable follow-up of the weight status of the children. Prospectively collected data on potential confounding factors, like data on viral etiology of early childhood wheezing and use of anti-inflammatory medication at school age, were available for adjusted analyses. The small number of the patients is a remarkable limitation, and with no doubt, the study was underpowered to reveal all presenting associations. Our selected material, children hospitalized for wheezing at <24 months of age, can be considered as a weakness of the study lessening the generalizability of the results. On the other hand, early childhood wheezers are at a high risk for later pulmonary disorders, and knowing the risk factors relevant in this specific group of children, like overweight and obesity according to the present results, opens new ways for preventive measures.

In conclusion, overweight and obesity increased constantly by age during follow-up after wheezing in early childhood until the median age of 12.3 years. Overweight at school age was associated with bronchial obstruction, as manifested by decreased FEV₁/FVC at 7.2 years and 12.3 years of age, and also with decreased flows in small airways at 12.3 years

of age. Our results stress the importance of the prevention of excessive weight gain at pre-school and school age in children with early childhood wheezing.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements: We thank Vesa Kiviniemi, Ph.Lic., for statistical advice, Kuopio University Hospital, The National Foundation for Pediatric Research in Finland and Tampere Tuberculosis Foundation for financial support.

The authors do not have a financial relationship with any commercial entity that has an interest in the subjects of this manuscript.

For Peer Review

References:

1. Ford ES. The epidemiology of obesity and asthma. *J Allergy Clin Immunol.* 2005;115:897–909.
2. Akinbami LJ, Schoendorf KC. Trends in childhood asthma: prevalence, health care utilization, and mortality. *Pediatrics.* 2002; 110: 315-322
3. Lobstein T, Frelut ML. Prevalence of overweight among children in Europe. *Obes Rev.* 2003; 4: 195-200.
4. Schachter LM, Peat JK, Salome CM. Asthma and atopy in overweight children. *Thorax.* 2003; 58: 1031–1035.
5. Bibi H, Shoseyov D, Feigenbaum D, Genis M, Friger M, Peled R, Sharff S. The relationship between asthma and obesity in children: is it real or a case of over diagnosis? *J Asthma.* 2004; 41: 403–410.
6. Wickens K, Barry D, Friezema A, Rhodius R, Bone N, Purdie G, Crane J. Obesity and asthma in 11–12 year old New Zealand children in 1989 and 2000. *Thorax.* 2005; 60: 7–12.
7. Mamun AA, Lawlor DA, Alati R, O’Callaghan MJ, Williams GM, Najman JM. Increasing body mass index from age 5 to 14 years predicts asthma among adolescents: evidence from a birth cohort study: *Intern J Obes.* 2007; 31; 578-583
8. Joseph CLM, Havstad SL, Ownby DR, Zoratti E, Peterson EL, Stringer S, Johnson CC. Gender differences in the association of overweight and asthma morbidity among urban adolescents with asthma. *Pediatr Allergy Immunol.* 2009; 20: 362-369.

- 1
2
3 9. Kusunoki T, Morimoto T, Nishikomori R, Heike T, Ito M, Hosoi S, Nakahata T.
4
5 Obesity and the prevalence of allergic diseases in schoolchildren. *Pediatr Allergy*
6
7 *Immunol.* 2008; 19:527-534.
8
9
10
11 10. Gilliland FD, Berhane K, Islam T, McConnell R, Gauderman WJ, Gilliland SS, Avol
12
13 E, Peters J. Obesity and the risk of newly diagnosed asthma in school-age children. *Am J*
14
15 *Epidemiol.* 2003; 158: 406–415.
16
17
18 11. Gold DR, Damokosh AI, Dockery DW, Berkey CS. Body-mass index
19
20 as a predictor of incident asthma in a prospective cohort of children.
21
22 *Pediatr Pulmonol.* 2003; 36: 514–521.
23
24
25
26 12. Mannino DM, Mott J, Ferdinands JM, Camargo CA, Friedman M, Greves HM Redd
27
28 SC. Boys with high body masses have an increased risk of developing asthma: findings
29
30 from the National Longitudinal Survey of Youth (NLSY). *Intern J Obes (Lond).* 2006;
31
32 30: 6–13.
33
34
35
36 13. Jartti T, Saarikoski L, Jartti L, LIsinen I, Jula A, Huupponen R, Viikari J, Raitakari
37
38 OT. Obesity, adipokines and asthma. *Allergy.* 2009; 64: 770-777.
39
40
41 14. Schachter LM, Salome CM, Peat JK, Woolcock AJ. Obesity is a risk for asthma and
42
43 wheeze but not airway hyper-responsiveness. *Thorax.* 2001; 56: 4–8.
44
45
46 15. Hancox RJ, Milne BJ, Poulton R, Taylor DR, Greene JM, McLachlan CR, Cowan JO,
47
48 Flannery EM, Herbison GP, Sears MR. Sex differences in the relation between body
49
50 mass index and asthma and atopy in a birth cohort. *Am J Respir Crit Care Med.* 2005;
51
52 171: 440–445.
53
54
55
56
57
58
59
60

16. Nicolacakis K, Skowronski ME, Coreno AJ, West E, Nader NZ, Smith RL, McFadden ER. Observation on the physiological interactions between obesity and asthma. *J Appl Physiol*. 2008; 105: 1533-1541.
17. Viljanen A, Halttunen P, Kreus K, Viljanen BC: Spirometric studies in non-smoking healthy adults. *Scand J Clin Lab Invest*. 1982; 159 (Suppl): 5-20.
18. Spathopoulos D, Paraskakis E, Trypsianis G, Tsalkidis A, Arvanitidou V, Emporiadou M, Bouros D, Chatzimichael A. The Effect of Obesity on Pulmonary Lung Function of School Aged Children in Greece. *Pediatr Pulmon*. 2009; 44: 273-280.
19. Deane S, Thomson A. Obesity and pulmonologist. Review. *Arch Dis Child*. 2006; 91: 188-191.
20. Sin DD, Sutherland ER. Obesity and the lung: 4 obesity and asthma. *Thorax* 2008; 63: 1018-1023.
21. Rubinstein I, Zamel N, DuBarry L, Hoffstein MD. Airflow limitation in morbidly obese, nonsmoking men. *Ann Intern Med*. 1990; 112: 828-832.
22. Jones RL, Nzekwu MMU. The effects of body mass index on lung volumes. *Chest*. 2006; 130: 827-833.
23. Sin DD, Jones RL, Man SFP. Obesity is a risk factor for dyspnea but not for airflow obstruction. *Arch Intern Med*. 2002; 162: 1477-1481.
24. Stenius-Aarniala B, Pousa T, Kvarnström J, Grönlund E-L, Ylikahri M, Mustajoki P. Immediate and long term effects of weight reduction in obese people with asthma: randomized controlled study. *BMJ*. 2000; 320: 827-832

- 1
2
3 25. Li AM, Chan D, Wong E, Yin J, Nelson EAS, Fok TF. The effect of obesity on
4 pulmonary function. *Arch Dis Child* 2003; 88: 361-363.
5
6
7
8 26. He Q, Qong T, Du L, Jiang Z, Qui H, Gao Y, Liu J, Wu J, Yu IT . Respiratory Health
9 in overweight and obese chinese children. *Pediatr Pulmonol*. 2009; 44: 997-1002
10
11
12
13 27. Fung KP, Lau SP, Chow OKW, Lee J, Wong TW. Effect of overweight on lung
14 function. *Arch Dis Child*. 1990; 65: 512-515.
15
16
17
18 28. Reijonen TM, Kotaniemi-Syrjanen A, Korhonen K, Korppi M. Predictors of asthma
19 three years after hospital admission for wheezing in infancy. *Pediatrics*. 2000; 106: 1406-
20
21
22
23 1412
24
25
26 29. Kotaniemi-Syrjänen A, Reijonen TM, Korhonen K, Korppi M. Wheezing requiring
27 hospitalization in early childhood: predictive factors for asthma in a six-year follow-up.
28
29
30
31
32
33
34 30. Hyvärinen MK, Kotaniemi-Syrjänen A, Reijonen TM, Korhonen K, Korppi MO.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
31. Reijonen T, Korppi M, Kuikka L, Remes K. Anti-inflammatory therapy reduces wheezing after bronchiolitis. *Arch Pediatr Adolesc Med*. 1996; 150: 512-517.
32. Reijonen TM, Korppi M. One-year follow-up of young children hospitalized for wheezing: the influence of early anti-inflammatory therapy and risk factors for subsequent wheezing and asthma. *Pediatr Pulmonol*. 1998; 26: 113-119.

33. Kotaniemi-Syrjänen A, Vainionpää R, Reijonen TM, Waris M, Korhonen K, Korppi M. Rhinovirus-induced wheezing in infancy--the first sign of childhood asthma? *J Allergy Clin Immunol*. 2003; 11: 66-71.
34. Korppi M, Kotaniemi-Syrjänen A, Waris M, Vainionpää R, Reijonen TM. Rhinovirus-associated wheezing in infancy: comparison with respiratory syncytial virus bronchiolitis. *Pediatr Infect Dis J*. 2004; 23: 995-999.
35. Kotaniemi-Syrjänen, Laatikainen A, Waris M, Reijonen T, Vainionpää R, Korppi M. Respiratory syncytial virus infection in children hospitalized for wheezing: Virus-specific studies from infancy to preschool years. *Acta Paediatr*. 2005; 94: 159-164.
36. Northwest Institute for Bio-Health Informatics. Child obesity calculator using British 1990 growth reference data. Available at: <http://www.phsim.man.ac.uk/>. Accessed January 18, 2007.
37. Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Dis Child*. 1995; 73: 25-29.
38. Freeman JV, Cole TJ, Chinn S, Jones PR, White EM, Preece MA. Cross sectional stature and weight reference curves for the UK, 1990. *Arch Dis Child* 1995; 73: 17-24.
39. Cole TJ, Freeman JV, Preece MA. British 1990 growth reference centiles for weight, height, body mass index and head circumference fitted by maximum penalized likelihood. *Stat Med* 1998; 17: 407-429
40. Cole TJ. The LMS method for constructing normalized growth standards. *Eur J Clin Nutr* 1990; 44: 45-60.

- 1
2
3 41. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for
4
5 child overweight and obesity worldwide: international survey. *Brit Med J* 2000; 320:
6
7 1240-1246.
8
9
10
11 42. Koillinen H, Wanne O, Niemi V, Laakkonen E. Reference values for flow-volume
12
13 spirometry and peak expiratory flow rates in healthy Finnish children. *Finnish Medical*
14
15 *Journal* 1998; 5: 395-402. (in Finnish).
16
17
18 43. Kotaniemi-Syrjänen A, Reijonen TM, Korhonen K, Waris M, Vainionpää R, Korppi
19
20 M. Wheezing due to rhinovirus infection in infancy: Bronchial hyperresponsiveness at
21
22 school age. *Pediatr Internat.* 2008; 50: 506-510.
23
24
25 44. Hyvärinen MK, Kotaniemi-Syrjänen A, Reijonen T, Korhonen K, Korppi M. Lung
26
27 function and bronchial hyper-responsiveness 11 years after hospitalization for
28
29 chonchiolitis. *Acta Paeditr.* 2007; 96: 1464-1469.
30
31
32
33 45. Kautiainen S, Rimpelä A, Vikat A, Virtanen SM. Secular trends in overweight and
34
35 obesity among Finnish adolescents in 1977-1999. *Int J Obes Relat Metab Disord.* 2002;
36
37 26: 544-52.
38
39
40
41 46. van Veen IH, Ten Brinke A, Sterk PJ, Rabe KF, Bel EH. Airway inflammation in
42
43 obese and non-obese patients with difficult-to-treat asthma. *Allergy.* 2008; 63: 570-574.
44
45
46
47 47. Peres-Padilla R, Rojas R, Torres V, Borja-Aburto V, Olaiz G. Obesity among
48
49 Children Residing in Mexico City and its Impact on Lung Function: A Comparison with
50
51 Mexican-Americans. *Arch Med Res.* 2006; 156-171.
52
53
54
55
56
57
58
59
60

48. Lazarus R, Colditz G, Berkey C, Speizer F. Effects of body fat on ventilatory function in children and adolescents: cross-sectional findings from a random population sample of school children. *Pediatr Pulmon.* 1997; 24: 187-194.

49. Tantisira KG, Litonjua AA, Weiss ST, Fuhlbrigge AL (Childhood asthma management program research group). Association of body mass with pulmonary function in the childhood asthma management program (CAMP). *Thorax.* 2003; 58: 1036-1041.

50. Bua J, Prescott E, Schack-Nielsen L, Petersen L, Godtfredsen NS, Sørensen TI, Osler M. Weight history from birth through childhood and youth in relation to adult lung function, in Danish juvenile obese and non-obese men. *Int J Obes (Lond).* 2005; 29: 1055-1062.

Lung function and overweight in school aged children after early childhood wheezing

Sidoroff Virpi¹ MD, Hyvärinen Mari K² MD, Piippo-Savolainen Eija² MD, PhD, Korppi Matti^{2,3} MD, PhD, Prof

¹ University of Eastern Finland, School of Medicine, Department of Pediatrics, Kuopio, Finland

² Department of Pediatrics, Kuopio University Hospital, Kuopio, Finland

³ Pediatric Research Center, Tampere University Hospital, Tampere, Finland

Corresponding author:

Sidoroff Virpi, Eteläkatu 10 b 7, 80110 Joensuu, Finland,

email: vpietika@hytti.uku.fi, tel +358-400015215

Abbreviated title: Lung function and overweight in wheezy children

Summary

Background and purpose. Recently, obesity has been connected with wheezing, asthma and reduced lung function. Most previous studies have been cross-sectional. The aim of the present follow-up study was to evaluate the association of preceding or current overweight or obesity with lung function at early and late school age after early childhood wheezing.

Material and methods. From the 100 children hospitalized for infection associated wheezing at <24 months of age, 83 attended the control visit at 4.0 years, 82 at 7.2 years and 81 at 12.3 years of age. Flow-volume spirometry was performed in 79 children at 7.2 years and in 80 children at 12.3 years of age. The weight status was assessed by calculating body mass index (BMI) at all visits. Age- and gender-specific BMI standard deviation scores (BMI-SDS) of >1.3 SD and > 2.0 SD were defined to mean overweight and obesity, respectively.

Results. Overweight at both 7.2 and 12.3 years of age was associated with decreased FEV₁/FVC (forced expiratory volume in one second/ forced vital capacity). Overweight and obesity at 7.2 years of age were associated with decreased FEV₁/FVC and MEF₅₀ (maximal expiratory flow at 50 % of FVC) at 12.3 years of age. The results were similar by continuous and categorized analyses, being robust to adjustments for viral findings during early childhood wheezing and asthma maintenance medication at school age.

Conclusion. Overweight and obesity are significant risk factors for reduced lung function at school age after early childhood wheezing. Thus, early-life wheezers should avoid excessive weight gain during childhood.

Keywords: child, spirometry, obstruction, overweight, obesity, lung function, wheezing

Abbreviations:

BMI	body mass index
BMI-SDS	body mass index standard deviation score
TLC	total lung capacity
FRC	functional residual capacity
FVC	forced vital capacity
FEV ₁	forced expiratory volume in one second
FEV ₁ /FVC	forced expiratory volume in one second/forced vital capacity
MEF ₅₀	maximal expiratory flow at 50% of FVC
MEF ₂₅	maximal expiratory flow at 25% of FVC
aOR	adjusted odds ratios
CI	confidence intervals
RSV	respiratory syncytial virus

Introduction

Asthma and overweight or obesity have increased concomitantly among both children and adults during the last three decades in western countries¹⁻³. The studies have suggested an association between wheezing symptoms and obesity⁴⁻⁹, and also between asthma diagnoses and obesity¹⁰⁻¹³. When pulmonary function and bronchial reactivity have been studied objectively, the results have been mainly negative^{5, 14-18}.

In adults, obesity decreases both lung volume, as documented by reduced total lung capacity (TLC), forced vital capacity (FVC) and functional residual capacity (FRC), and airway flows, as documented by decreased forced expiratory volume in one second (FEV₁) and increased airway resistance^{19,20}. In addition, obesity leads to obstruction in peripheral airway²¹, as documented by decreased flows at 50% (MEF₅₀) or at 25% (MEF₂₅) of FVC. However, the largest impact seems to target at expiratory reserve volume, which means that restriction is the characteristic lung function disorder in obesity^{22,23}. In addition, weight reduction in obese adults elicited an improvement of lung function²⁴.

In children with asthma suggestive symptoms, high body mass index (BMI) was associated with subnormal lung function in one study¹⁸, but not in two others^{4,5}. In healthy children, lung function was either impaired²⁵, improved in boys and girls²⁶ or improved only in girls²⁷ by excessive weight gain.

We have prospectively followed-up a cohort of children hospitalized for infection associated wheezing at <24 months of age in 1992-1993. The weights and heights of the children were measured at follow-up visits at the median ages of 4.0, 7.2 and 12.3 years, and spirometry was performed at both visits at school age²⁸⁻³⁰.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The aim of the present study was to evaluate the association between overweight or obesity and lung function at school age in former early-life wheezers, with a special focus on the influence of an emerging overweight.

For Peer Review

Material and methods

Study design

One-hundred consecutive children, hospitalized for wheezing associated respiratory infection at age 1-23 months, were recruited into this follow-up study in 1992 -1993³¹. Maintenance therapy for asthma was one of the exclusion criteria, and the wheezing episode was the first one in 87% of the included children³². The basic data were collected by interviewing the parents during hospitalization using a structured questionnaire. Seven respiratory viruses, including respiratory syncytial virus (RSV), were studied by antigen and antibody assays³². Later, viral studies were supplemented by polymerase chain reaction to RSV and rhinoviruses in frozen samples^{33,34}. In all, RSV was identified in 29/100 and rhinovirus in 27/81 cases^{33,35}.

Three follow-up visits were prospectively organized at the median ages of 4.0 years, 7.2 years and 12.3 years²⁸⁻³⁰. Asthma was present in 33/82 (40.2%) children at the 7.2-year visit²⁹ and in 32/81 (39.5%) children at the 12.3-year visit³⁰. Data on medical history, including the use of asthma medication during the preceding 12 months, were collected by interviewing the parents.

Assessment of weight status

The weights and heights of the children, measured by an experienced nurse using standard methods, were registered in all attending children at all controls visits. Body mass index (BMI) was calculated by using the equation: weight (kg) / [height (m)]². Due to the lack of Finnish references, we applied British gender- and age-specific population-based references. BMI standard deviation score (BMI-SDS), also called Z-score, was

assessed by an obesity calculator (NIBHI 2007)³⁶. This calculator applies British growth references from 1990, produced by the LMS method in 1995^{37,38}, and revised in 1996³⁹. The LMS method summarizes the distribution of BMI at each age by its median (M), coefficient of variation (S), and skewness expressed as a Box-Cox power (L) required to transform the data to normal⁴⁰.

In the present study, we used the same cut-off points for overweight and obesity as proposed by the British study evaluating the limits of obesity and overweight in different populations⁴¹. The cut-off point for overweight was BMI >91 percentile (approximating BMI-SDS >1.3) and for obesity BMI >98 percentile (approximating BMI-SDS >2.0)^{19,41}. In the British population, these cut-off points corresponds BMI 25 kg/m² and BMI 30 kg/m² levels, respectively, in the 18 years old young adults⁴¹.

Measurement of lung function

Lung function was studied by flow-volume spirometer (FVS) (Medikro, Kuopio, Finland) at the 7.2-year and 12.3-year study visits. The parameters were expressed as percentages of the gender-specific height-related reference values (% of predicted) for Finnish children^{42,44}. The parameters registered were FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FVC, MEF₅₀ (maximal expiratory flow at 50 % of FVC) and MEF₂₅ (maximal expiratory flow at 25 % of FVC). Either the measurement with the highest FEV₁ (at the 7.2-year study)²⁹ or the composite maximal curve (at the 12.3-year study)³⁰ was in each child recorded for further analyses. The cut-off limits for abnormal values were 80 % for FEV₁, 88 % for FEV₁/FVC, 62 % for MEF₅₀ and 48 % for MEF₂₅^{17,30,43}.

Lung function data were available from 79 children at the 7.2-year and from 81 children at the 12.3-year study visits. The numbers of children with anti-inflammatory maintenance medication for asthma during the 12 preceding months were 29/79 (36.77%) at age of 7.2 years and 18/80 (22.5%) at age of 12.3 years. At the 4.0-year visit, weight status was assessed for 74 children with lung function data available at the 7.2-year visit.

Statistics

The data were analysed using SPSS 17.0 software (SPSS Inc. Chicago, IL, USA). Mann-Whitney U-test was used in univariate analyses and analysis of variance (ANOVA) in multivariate analyses for continuous FVS parameters. In the previous studies in this cohort, RSV etiology of wheezing in early childhood was associated with a non-reversible restrictive pattern of lung function and non-RSV etiology (mostly rhinovirus) with an increased asthma risk and a reversible obstructive pattern of lung function at school age^{43, 44}. Therefore, analyses by ANOVA were adjusted also for RSV etiology of wheezing in early childhood, in addition to adjustment with anti-inflammatory asthma medication during the preceding 12 months before the study visit at school age. Logistic regression analysis, adjusted for RSV etiology of early childhood wheezing and anti-inflammatory asthma medication during the 12 preceding months before the study visit, was applied for categorized (by the limits of abnormal values in % predicted) FVS parameters, and the results were expressed as adjusted odds ratios (aOR) and their 95 % confidence intervals (95% CI).

Ethics

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The study was approved by the joint Research Ethics Committee of Kuopio University and Kuopio University Hospital. Informed written consent was obtained from the parents of the children.

For Peer Review

Results

Overweight was present in 6.0% of children at 4.0 years, in 22.2% at 7.3 years and in 33.3% at 12.3 years of age. Less than 4% of children were obese at 4.0 years of age. The proportion of obese children was 8.6 % at 7.3 years and 19.8% at 12.3 years of age. Three of 5 overweight children at age 4.0 years were overweight at age 7.2 years, and all 3 obese children were obese, respectively. Fifteen of the 16 overweight children at age 7.2 years were overweight at age 12.3 years, and 4 of the 6 obese children were obese, respectively.

In univariate analyses at 7.2 years of age, the only significant association was between current overweight and lower FEV₁/FVC (Table 1). There were no significant associations between previous (4.0 years of age) overweight or obesity and any FVS parameter at age 7.2 years (Data not shown). In multivariate analyses, the association between obesity and lower FEV₁/FVC changed to significant (Table 1). All other associations between previous or current overweight or obesity and FVS remained as unchanged.

In univariate analyses at 12.3 years of age, previous (7.2 years of age) overweight was associated with lower FEV₁/FVC, and current (12.3 years of age) overweight was associated with lower FEV₁/FVC, MEF₅₀ and MEF₂₅ (Tables 2). Similarly, previous obesity was associated with lower FEV₁/FVC, and current obesity was associated with lower FEV₁/FVC, MEF₅₀ and MEF₂₅ (Table 3). In multivariate analyses, the association between current overweight and MEF₅₀ changed to non-significant (Table 2). All other associations between previous or current overweight or obesity and FVS remained as unchanged.

In the logistic regression analyses, both previous and current overweight were significant risk factors for FEV₁/FVC <88% at both 7.2 years (Table 4) and at 12.3 years of age (Table 5). Since overweight was highly permanent from 4.0 to 7.3 years of age and from 7.2 to 12.3 years of age, respectively, previous and current overweight could not be incorporated as variables in the same model. Therefore, the design of the study did not allow to assess their influences separately. Current obesity was a significant risk factor for FEV₁/FVC <88% (aOR 8.07, 95% CI 2.10-31.05) and for MEF₂₅ <48% (aOR 4.34, 95% CI 1.06-17.79) at 12.3 years of age.

Discussion

There are three main results in the present study. First, the number of overweight and obese children increased constantly by age during the follow-up after early childhood wheezing until the median age of 12.3 years. Finally, a third was overweight, and about 20% were obese. The figures are higher than those calculated from self-reported weights and heights in 12-18 years old Finnish children in 1999, when 16.7% of the boys and 9.8% of the girls were overweight, and 1.4% and 0.4% were obese, assessed by using age- and gender-related international reference values⁴⁵. Second, current overweight was associated with decreased FEV₁/FVC at both 7.2 and 12.3 years of age. Overweight children had also reduced flows in small airways by MEF₂₅ and obese children by MEF₂₅ and MEF₅₀ at age 12.3 years. FEV₁/FVC is considered as the most sensitive and reliable measure of obstruction, but the reliability of MEF₂₅ has been questioned. Third, overweight and obesity at early school age preceded reduced FEV₁/FVC at age 12.3 years. But, since overweight was highly permanent from early to late school age, the influences of previous and current overweight could not be assessed separately.

The association between excessive weight gain and reduced lung function has been well documented in adults^{14-16,20,22}. In a Finnish study, weight reduction in obese adults even elicited an improvement of lung function²⁴. The effect of excessive weight gain to lung function seems to depend on the degree of obesity and on the presence and severity of asthma. When lung function was studied in 52 non-asthmatic non-obese, in 53 asthmatic non-obese, in 52 obese non-asthmatic and in 53 asthmatic obese adults, lung function was reduced in patients with asthma and obesity, and in addition, FVC and FRC were reduced

in obese adults also with no asthma¹⁶. In a birth cohort from New Zealand, an association was found between increasing BMI and decreasing FEV₁/FVC in 9-26 years old females, but not in males¹⁵. In a study on difficult-to-treat asthma, FEV₁ was increased and FRC and total lung capacity were decreased in obese patients⁴⁶. Thus, controlled studies in adults with asthma^{16,46}, as well as population-based studies in healthy adults^{15,22,23}, highlight the connection between obesity and decreased expiratory reserve volumes, which means that restriction is the characteristic lung function disorder in obesity.

In a population study from China, overweight had a beneficial effect on lung function in 7 to 20 years old healthy girls²⁷. In a population study from Mexico, lung function improved when weight increased in 8-11 years old healthy children, but after that age, lung function first reached a plateau and then decreased among the obese youths⁴⁷. Lung function by spirometry was compared between 657 overweight and 196 normal weight Greek children, and FVC, FEV₁, MEF_{25-75%} and FEV₁/FVC were reduced in overweight children¹⁸. In a population study from Australia, spirometry was performed in 2464 school children⁴⁸. Height-related FVC and FEV₁ increased by growth, but both FVC and FEV₁ decreased in relation to increasing overweight. The result was opposite in an other Australian population study in 5993 children; all variables in the FVS were, irrespective of BMI, within normal limits, but FEV₁ and FVC in the lowest BMI quintile were significantly lower than in other quintiles⁴. All these studies are cross-sectional with no follow-up available. The results seem confusing, but there may be following consistency; mild overweight may be beneficial but when substantial overweight or obesity develops, the effect turns to non-beneficial.

The results of the present follow-up study after early childhood wheezing showed that overweight was associated with reduced FEV_1/FVC at 7.2 and 12.3 years of age and reduced MEF_{50} and MEF_{25} at 12.3 years of age, in disagreement with most population studies in healthy children^{4,5,27,46,47}. However, the associations may be different in children with asthma; about 40% of our patients had asthma and all had wheezed in early childhood. Lung function and weight status were studied as a part of the childhood asthma management program (CAMP) in 1041 American children aged 5-12 years⁴⁹, and increasing BMI was associated with increasing FEV_1 and FVC, but decreasing FEV_1/FVC . The increase in FVC and decrease in FEV_1/FVC are in accordance with the results of our present study. In the CAMP study, weight status and lung function were studied by a cross-sectional design at enrolment of the patients, and no follow-up data have been, at least thus far, published. We found no evidence for a restrictive pattern of lung function disorder in overweight or obese children, though suggested to be present in obese asthmatic adults^{15,16,22,23,46}, and though was demonstrable after early childhood wheezing caused by RSV in 12.3 years old children of the present study^{43,44}.

The mechanisms leading to lung function disorders in association with overweight or obesity may be narrowing of the small peripheral airways²¹, in line with decreased MEF_{50} and MEF_{25} values in the overweight and obese children at the 12.3-year visit of the present study. In addition, decreased compliance of either the lungs due to narrow peripheral airways or the chest wall due to body adiposity may lead to subnormal lung function, including both decreased airway flows which mean an obstructive lung function disorder and decreased lung volumes which means a restrictive lung function disorder⁴⁸.

The effect of excessive weight gain on lung function may be age-dependent, which

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

means that different mechanisms may influence in children and in adults and in children at different ages. In a case-control study from Denmark, FEV₁ and FVC were reduced in 193 obese men, compared with normal weight 215 controls, at 27 years of age, but after adjustments with current BMI, childhood BMI at 7 years of age was positively associated with both FEV₁ and FVC ⁵⁰.

The strengths of the present study are the long, over 10 years prospective follow-up time and the fact that weights and heights of the children were measured at all control visits by experienced nurses using calibrated scales, thus allowing a reliable follow-up of the weight status of the children. Prospectively collected data on potential confounding factors, like data on viral etiology of early childhood wheezing and use of anti-inflammatory medication at school age, were available for adjusted analyses. The small number of the patients is a remarkable limitation, and with no doubt, the study was underpowered to reveal all presenting associations. Our selected material, children hospitalized for wheezing at <24 months of age, can be considered as a weakness of the study lessening the generalizability of the results. On the other hand, early childhood wheezers are at a high risk for later pulmonary disorders, and knowing the risk factors relevant in this specific group of children, like overweight and obesity according to the present results, opens new ways for preventive measures.

In conclusion, overweight and obesity increased constantly by age during follow-up after wheezing in early childhood until the median age of 12.3 years. Overweight at school age was associated with bronchial obstruction, as manifested by decreased FEV₁/FVC at 7.2 years and 12.3 years of age, and also with decreased flows in small airways at 12.3 years

of age. Our results stress the importance of the prevention of excessive weight gain at pre-school and school age in children with early childhood wheezing.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements: We thank Vesa Kiviniemi, Ph.Lic., for statistical advice, Kuopio University Hospital, The National Foundation for Pediatric Research in Finland and Tampere Tuberculosis Foundation for financial support.

The authors do not have a financial relationship with any commercial entity that has an interest in the subjects of this manuscript.

For Peer Review

References:

1. Ford ES. The epidemiology of obesity and asthma. *J Allergy Clin Immunol.* 2005;115:897–909.
2. Akinbami LJ, Schoendorf KC. Trends in childhood asthma: prevalence, health care utilization, and mortality. *Pediatrics.* 2002; 110: 315-322
3. Lobstein T, Frelut ML. Prevalence of overweight among children in Europe. *Obes Rev.* 2003; 4: 195-200.
4. Schachter LM, Peat JK, Salome CM. Asthma and atopy in overweight children. *Thorax.* 2003; 58: 1031–1035.
5. Bibi H, Shoseyov D, Feigenbaum D, Genis M, Friger M, Peled R, Sharff S. The relationship between asthma and obesity in children: is it real or a case of over diagnosis? *J Asthma.* 2004; 41: 403–410.
6. Wickens K, Barry D, Friezema A, Rhodius R, Bone N, Purdie G, Crane J. Obesity and asthma in 11–12 year old New Zealand children in 1989 and 2000. *Thorax.* 2005; 60: 7–12.
7. Mamun AA, Lawlor DA, Alati R, O’Callaghan MJ, Williams GM, Najman JM. Increasing body mass index from age 5 to 14 years predicts asthma among adolescents: evidence from a birth cohort study: *Intern J Obes.* 2007; 31; 578-583
8. Joseph CLM, Havstad SL, Ownby DR, Zoratti E, Peterson EL, Stringer S, Johnson CC. Gender differences in the association of overweight and asthma morbidity among urban adolescents with asthma. *Pediatr Allergy Immunol.* 2009; 20: 362-369.

9. Kusunoki T, Morimoto T, Nishikomori R, Heike T, Ito M, Hosoi S, Nakahata T. Obesity and the prevalence of allergic diseases in schoolchildren. *Pediatr Allergy Immunol*. 2008; 19:527-534.
10. Gilliland FD, Berhane K, Islam T, McConnell R, Gauderman WJ, Gilliland SS, Avol E, Peters J. Obesity and the risk of newly diagnosed asthma in school-age children. *Am J Epidemiol*. 2003; 158: 406–415.
11. Gold DR, Damokosh AI, Dockery DW, Berkey CS. Body-mass index as a predictor of incident asthma in a prospective cohort of children. *Pediatr Pulmonol*. 2003; 36: 514–521.
12. Mannino DM, Mott J, Ferdinands JM, Camargo CA, Friedman M, Greves HM Redd SC. Boys with high body masses have an increased risk of developing asthma: findings from the National Longitudinal Survey of Youth (NLSY). *Intern J Obes (Lond)*. 2006; 30: 6–13.
13. Jartti T, Saarikoski L, Jartti L, LIsinen I, Jula A, Huupponen R, Viikari J, Raitakari OT. Obesity, adipokines and asthma. *Allergy*. 2009; 64: 770-777.
14. Schachter LM, Salome CM, Peat JK, Woolcock AJ. Obesity is a risk for asthma and wheeze but not airway hyper-responsiveness. *Thorax*. 2001; 56: 4–8.
15. Hancox RJ, Milne BJ, Poulton R, Taylor DR, Greene JM, McLachlan CR, Cowan JO, Flannery EM, Herbison GP, Sears MR. Sex differences in the relation between body mass index and asthma and atopy in a birth cohort. *Am J Respir Crit Care Med*. 2005; 171: 440–445.

16. Nicolacakis K, Skowronski ME, Coreno AJ, West E, Nader NZ, Smith RL, McFadden ER. Observation on the physiological interactions between obesity and asthma. *J Appl Physiol*. 2008; 105: 1533-1541.
17. Viljanen A, Halttunen P, Kreus K, Viljanen BC: Spirometric studies in non-smoking healthy adults. *Scand J Clin Lab Invest*. 1982; 159 (Suppl): 5-20.
18. Spathopoulos D, Paraskakis E, Trypsianis G, Tsalkidis A, Arvanitidou V, Emporiadou M, Bouros D, Chatzimichael A. The Effect of Obesity on Pulmonary Lung Function of School Aged Children in Greece. *Pediatr Pulmon*. 2009; 44: 273-280.
19. Deane S, Thomson A. Obesity and pulmonologist. Review. *Arch Dis Child*. 2006; 91: 188-191.
20. Sin DD, Sutherland ER. Obesity and the lung: 4 obesity and asthma. *Thorax* 2008; 63: 1018-1023.
21. Rubinstein I, Zamel N, DuBarry L, Hoffstein MD. Airflow limitation in morbidly obese, nonsmoking men. *Ann Intern Med*. 1990; 112: 828-832.
22. Jones RL, Nzekwu MMU. The effects of body mass index on lung volumes. *Chest*. 2006; 130: 827-833.
23. Sin DD, Jones RL, Man SFP. Obesity is a risk factor for dyspnea but not for airflow obstruction. *Arch Intern Med*. 2002; 162: 1477-1481.
24. Stenius-Aarniala B, Pousa T, Kvarnström J, Grönlund E-L, Ylikahri M, Mustajoki P. Immediate and long term effects of weight reduction in obese people with asthma: randomized controlled study. *BMJ*. 2000; 320: 827-832

25. Li AM, Chan D, Wong E, Yin J, Nelson EAS, Fok TF. The effect of obesity on pulmonary function. Arch Dis Child 2003; 88: 361-363.

26. He Q, Qong T, Du L, Jiang Z, Qui H, Gao Y, Liu J, Wu J, Yu IT . Respiratory Health in overweight and obese chinese children. Pediatr Pulmonol. 2009; 44: 997-1002

27. Fung KP, Lau SP, Chow OKW, Lee J, Wong TW. Effect of overweight on lung function. Arch Dis Child. 1990; 65: 512-515.

28. Reijonen TM, Kotaniemi-Syrjanen A, Korhonen K, Korppi M. Predictors of asthma three years after hospital admission for wheezing in infancy. Pediatrics. 2000; 106: 1406-1412

29. Kotaniemi-Syrjänen A, Reijonen TM, Korhonen K, Korppi M. Wheezing requiring hospitalization in early childhood: predictive factors for asthma in a six-year follow-up. Pediatr Allergy Immunol. 2002; 13: 418-425.

30. Hyvärinen MK, Kotaniemi-Syrjänen A, Reijonen TM, Korhonen K, Korppi MO. Teenage asthma after severe early childhood wheezing: an 11-year prospective follow-up. Pediatr Pulmonol. 2005; 40: 316-323.

31. Reijonen T, Korppi M, Kuikka L, Remes K. Anti-inflammatory therapy reduces wheezing after bronchiolitis. Arch Pediatr Adolesc Med. 1996; 150: 512-517.

32. Reijonen TM, Korppi M. One-year follow-up of young children hospitalized for wheezing: the influence of early anti-inflammatory therapy and risk factors for subsequent wheezing and asthma. Pediatr Pulmonol. 1998; 26: 113-119.

33. Kotaniemi-Syrjänen A, Vainionpää R, Reijonen TM, Waris M, Korhonen K, Korppi M. Rhinovirus-induced wheezing in infancy--the first sign of childhood asthma? *J Allergy Clin Immunol*. 2003; 11: 66-71.
34. Korppi M, Kotaniemi-Syrjänen A, Waris M, Vainionpää R, Reijonen TM. Rhinovirus-associated wheezing in infancy: comparison with respiratory syncytial virus bronchiolitis. *Pediatr Infect Dis J*. 2004; 23: 995-999.
35. Kotaniemi-Syrjänen, Laatikainen A, Waris M, Reijonen T, Vainionpää R, Korppi M. Respiratory syncytial virus infection in children hospitalized for wheezing: Virus-specific studies from infancy to preschool years. *Acta Paediatr*. 2005; 94: 159-164.
36. Northwest Institute for Bio-Health Informatics. Child obesity calculator using British 1990 growth reference data. Available at: <http://www.phsim.man.ac.uk/>. Accessed January 18, 2007.
37. Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Dis Child*. 1995; 73: 25-29.
38. Freeman JV, Cole TJ, Chinn S, Jones PR, White EM, Preece MA. Cross sectional stature and weight reference curves for the UK, 1990. *Arch Dis Child* 1995; 73: 17-24.
39. Cole TJ, Freeman JV, Preece MA. British 1990 growth reference centiles for weight, height, body mass index and head circumference fitted by maximum penalized likelihood. *Stat Med* 1998; 17: 407-429
40. Cole TJ. The LMS method for constructing normalized growth standards. *Eur J Clin Nutr* 1990; 44: 45-60.

- 1
2
3 41. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for
4
5 child overweight and obesity worldwide: international survey. *Brit Med J* 2000; 320:
6
7 1240-1246.
8
9
10
11 42. Koillinen H, Wanne O, Niemi V, Laakkonen E. Reference values for flow-volume
12
13 spirometry and peak expiratory flow rates in healthy Finnish children. *Finnish Medical*
14
15 *Journal* 1998; 5: 395-402. (in Finnish).
16
17
18 43. Kotaniemi-Syrjänen A, Reijonen TM, Korhonen K, Waris M, Vainionpää R, Korppi
19
20 M. Wheezing due to rhinovirus infection in infancy: Bronchial hyperresponsiveness at
21
22 school age. *Pediatr Internat.* 2008; 50: 506-510.
23
24
25 44. Hyvärinen MK, Kotaniemi-Syrjänen A, Reijonen T, Korhonen K, Korppi M. Lung
26
27 function and bronchial hyper-responsiveness 11 years after hospitalization for
28
29 chonchiolitis. *Acta Paediatr.* 2007; 96: 1464-1469.
30
31
32 45. Kautiainen S, Rimpelä A, Vikat A, Virtanen SM. Secular trends in overweight and
33
34 obesity among Finnish adolescents in 1977-1999. *Int J Obes Relat Metab Disord.* 2002;
35
36 26: 544-52.
37
38
39 46. van Veen IH, Ten Brinke A, Sterk PJ, Rabe KF, Bel EH. Airway inflammation in
40
41 obese and non-obese patients with difficult-to-treat asthma. *Allergy.* 2008; 63: 570-574.
42
43
44 47. Peres-Padilla R, Rojas R, Torres V, Borja-Aburto V, Olaiz G. Obesity among
45
46 Children Residing in Mexico City and its Impact on Lung Function: A Comparison with
47
48 Mexican-Americans. *Arch Med Res.* 2006; 156-171.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 48. Lazarus R, Colditz G, Berkey C, Speizer F. Effects of body fat on ventilatory function
4
5 in children and adolescents: cross-sectional findings from a random population sample of
6
7 school children. *Pediatr Pulmon*. 1997; 24: 187-194.
8
9
10
11 49. Tantisira KG, Litonjua AA, Weiss ST, Fuhlbrigge AL (Childhood asthma
12
13 management program research group). Association of body mass with pulmonary
14
15 function in the childhood asthma management program (CAMP). *Thorax*. 2003; 58:
16
17 1036-1041.
18
19
20
21 50. Bua J, Prescott E, Schack-Nielsen L, Petersen L, Godtfredsen NS, Sørensen TI, Osler
22
23 M. Weight history from birth through childhood and youth in relation to adult lung
24
25 function, in Danish juvenile obese and non-obese men. *Int J Obes (Lond)*. 2005; 29:
26
27 1055-1062.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Flow-volume spirometry (FVS, % of predicted, mean, SD) in 74 study subjects at 7.2 years of age, in relation to current overweight or obesity (at 7.2 years of age).

Parameters in FVS ^b	Current overweight ^a				Current obesity ^a			
	Yes (N=16) Mean, (SD)	No (N=58) Mean, (SD)	Univariate p	Multivariate p ^d	Yes (N=7) Mean, (SD)	No (N=72) Mean, (SD)	Univariate p ^c	Multivariate p ^d
FVC	108.71 (19.75)	102.06 (16.48)	0.252	0.219	106.73 (8.78)	103.27 (18.00)	0.448	0.467
FEV ₁	100.16 (14.16)	99.95 (13.94)	0.977	0.890	96.58 (4.78)	100.33 (14.46)	0.523	0.574
FEV ₁ /FVC	92.63 (8.95)	98.47 (8.93)	0.017	0.025	90.81 (7.62)	97.76 (9.17)	0.076	0.034
MEF ₅₀	84.00 (17.73)	85.10 (20.39)	0.700	0.814	81.30 (19.03)	85.19 (19.87)	0.546	0.615
MEF ₂₅	66.00 (19.85)	71.27 (22.76)	0.394	0.454	62.77 (17.30)	70.78 (22.50)	0.398	0.305

^a Overweight BMI-SDS >1.3 (obese children included); obesity BMI-SDS > 2,0

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FCV, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Mann-Whitney U -test

^d Analysis of variance, adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months).

Table 2. Flow-volume spirometry (FVS, % of predicted, mean, SD) in 80 study subjects at 12.3 years of age, in relation to previous (at 7.2 years of age) and current (at 12.3 years of age) overweight.

Parameters in FVS ^b	Previous overweight ^a		Univariate p ^c	Multivariate p ^d	Current overweight ^a		Univariate p ^c	Multivariate p ^d
	Yes (N=16) Mean, (SD)	No (N=58) Mean, (SD)			Yes (N=27) Mean, (SD)	No (N=53) Mean, (SD)		
FVC	101.03 (8.40)	97.21 (13.20)	0.131	0.306	100.56 (11.50)	96.38 (12.68)	0.162	0.164
FEV₁	90.21 (12.96)	93.03 (12.96)	0.684	0.420	90.68 (13.60)	93.02 (21.50)	0.439	0.500
FEV₁/FVC	89.15 (10.06)	96.01 (7.63)	0.013	0.003	90.09 (8.81)	96.81 (7.36)	0.001	0.001
MEF₅₀	74.88 (20.68)	83.91 (18.82)	0.121	0.095	75.63 (20.47)	84.39 (17.70)	0.037	0.076
MEF₂₅	64.49 (23.75)	73.18 (19.25)	0.138	0.139	62.96 (19.58)	76.13 (19.85)	0.006	0.009

^a BMI-SDS >1.3

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FVC, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Mann-Whitney U -test

^d Analysis of variance, adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Flow-volume spirometry (FVS, % of predicted, mean, SD) in 80 study subjects at 12.3 years of age, in relation to previous (at 7.2 years of age) and current (at 12.3 years of age) obesity.

Parameters in FVS ^b	Previous obesity ^a		Univariate p ^c	Multivariate p ^d	Current obesity ^a		Univariate p ^c	Multivariate p ^d
	Yes	No			Yes	No		
	(N=6) Mean, (SD)	(N=68) Mean, (SD)			(N=16) Mean, (SD)	(N=64) Mean, (SD)		
FVC	99.05 (9.47)	97.95 (12.65)	0.797	0.769	98.39 (9.12)	97.64 (13.12)	0.674	0.837
FEV ₁	87.28 (13.30)	92.87 (12.89)	0.434	0.318	87.44 (12.69)	93.43 (12.86)	0.136	0.116
FEV ₁ /FVC	87.76 (6.99)	95.12 (8.57)	0.022	0.026	88.85 (9.78)	95.97 (7.51)	0.010	0.002
MEF ₅₀	70.46 (15.72)	82.97 (19.52)	0.132	0.107	71.68 (18.81)	83.87 (18.41)	0.025	0.028
MEF ₂₅	60.03 (10.59)	72.34 (20.84)	0.113	0.130	61.02 (21.40)	74.35 (19.68)	0.021	0.024

^a BMI-SDS >2

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FCV, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Mann-Whitney U -test

^d Analysis of variance, adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months).

Table 4. Logistic regression: abnormal lung function in flow-volume spirometry (FVS) at 7.2 years of age, in relation to previous (at 4.0 years of age) and current (at 7.2 years of age) overweight.

Abnormal FVS ^b	Previous overweight ^a		aOR ^c (95% CI)	Current overweight ^a		aOR ^c (95% CI)
	Yes (N=5)	No (N=69)		Yes (N=17)	No (N=57)	
FEV₁<80%	0	4	-	1	3	1.01 (0.10-10.58)
FEV₁/FVC<88%	3	10	15.46 (1.65-145.26)	6	7	3.71 (1.01-13.60)
MEF₅₀<62%	2	11	6.46 (0.772-33.95)	3	11	0.89 (0.21-3.80)
MEF₂₅<48%	2	12	4.74 (0.61-36.76)	3	11	0.89 (0.22-3.70)
At least one abnormal parameter	3	16	6.63 (0.89-49.46)	6	14	1.63 (0.51-5.22)

^a BMI-SDS >1.3

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FVC, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Odds ratio (95% confidence interval) adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months)

Table 5. Logistic regression: abnormal lung function in flow-volume spirometry (FVS) at 12.3 years of age, in relation to previous (at 7.2 years of age) and current (at 12.3 years of age) overweight.

Abnormal FVS ^b	Previous overweight ^a		aOR ^c (95% CI)	Current overweight ^a		aOR ^c (95% CI)
	Yes (N=16)	No (N=58)		Yes (N=24)	No (N=50)	
FEV ₁ <80%	2	10	0.70 (0.13-3.60)	5	9	1.00 (0.29-3.46)
FEV ₁ /FVC<88%	6	6	5.64 (1.43-22.20)	8	5	4.02 (1.14-14.22)
MEF ₅₀ <62%	3	9	1.29 (0.30-5.54)	7	6	2.64 (0.78-8.97)
MEF ₂₅ <48%	4	6	3.51 (0.76-16.13)	6	5	2.47 (0.65-9.38)
At least one abnormal parameter	6	17	1.45 (0.45-4.67)	11	15	1.70 (0.63-4.55)

^a BMI-SDS >1.3

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FCV, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Odds ratio (95% confidence interval) adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months)

Table 1. Flow-volume spirometry (FVS, % of predicted, mean, SD) in 74 study subjects at 7.2 years of age, in relation to current overweight or obesity (at 7.2 years of age).

Parameters in FVS ^b	Current overweight ^a				Current obesity ^a			
	Yes (N=16) Mean, (SD)	No (N=58) Mean, (SD)	Univariate p	Multivariate p ^d	Yes (N=7) Mean, (SD)	No (N=72) Mean, (SD)	Univariate p ^c	Multivariate p ^d
FVC	108.71 (19.75)	102.06 (16.48)	0.252	0.219	106.73 (8.78)	103.27 (18.00)	0.448	0.467
FEV₁	100.16 (14.16)	99.95 (13.94)	0.977	0.890	96.58 (4.78)	100.33 (14.46)	0.523	0.574
FEV₁/FVC	92.63 (8.95)	98.47 (8.93)	0.017	0.025	90.81 (7.62)	97.76 (9.17)	0.076	0.034
MEF₅₀	84.00 (17.73)	85.10 (20.39)	0.700	0.814	81.30 (19.03)	85.19 (19.87)	0.546	0.615
MEF₂₅	66.00 (19.85)	71.27 (22.76)	0.394	0.454	62.77 (17.30)	70.78 (22.50)	0.398	0.305

^a Overweight BMI-SDS >1.3 (obese children included); obesity BMI-SDS > 2.0

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FVC, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Mann-Whitney U -test

^d Analysis of variance, adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Flow-volume spirometry (FVS, % of predicted, mean, SD) in 80 study subjects at 12.3 years of age, in relation to previous (at 7.2 years of age) and current (at 12.3 years of age) overweight.

Parameters in FVS ^b	Previous overweight ^a		Univariate p ^c	Multivariate p ^d	Current overweight ^a		Univariate p ^c	Multivariate p ^d
	Yes	No			Yes	No		
	(N=16) Mean, (SD)	(N=58) Mean, (SD)			(N=27) Mean, (SD)	(N=53) Mean, (SD)		
FVC	101.03 (8.40)	97.21 (13.20)	0.131	0.306	100.56 (11.50)	96.38 (12.68)	0.162	0.164
FEV ₁	90.21 (12.96)	93.03 (12.96)	0.684	0.420	90.68 (13.60)	93.02 (21.50)	0.439	0.500
FEV ₁ /FVC	89.15 (10.06)	96.01 (7.63)	0.013	0.003	90.09 (8.81)	96.81 (7.36)	0.001	0.001
MEF ₅₀	74.88 (20.68)	83.91 (18.82)	0.121	0.095	75.63 (20.47)	84.39 (17.70)	0.037	0.076
MEF ₂₅	64.49 (23.75)	73.18 (19.25)	0.138	0.139	62.96 (19.58)	76.13 (19.85)	0.006	0.009

^a BMI-SDS >1.3

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FCV, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Mann-Whitney U -test

^d Analysis of variance, adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months).

Table 3. Flow-volume spirometry (FVS, % of predicted, mean, SD) in 80 study subjects at 12.3 years of age, in relation to previous (at 7.2 years of age) and current (at 12.3 years of age) obesity.

Parameters in FVS ^b	Previous obesity ^a		Univariate p ^c	Multivariate p ^d	Current obesity ^a		Univariate p ^c	Multivariate p ^d
	Yes	No			Yes	No		
	(N=6) Mean, (SD)	(N=68) Mean, (SD)			(N=16) Mean, (SD)	(N=64) Mean, (SD)		
FVC	99.05 (9.47)	97.95 (12.65)	0.797	0.769	98.39 (9.12)	97.64 (13.12)	0.674	0.837
FEV₁	87.28 (13.30)	92.87 (12.89)	0.434	0.318	87.44 (12.69)	93.43 (12.86)	0.136	0.116
FEV₁/FVC	87.76 (6.99)	95.12 (8.57)	0.022	0.026	88.85 (9.78)	95.97 (7.51)	0.010	0.002
MEF₅₀	70.46 (15.72)	82.97 (19.52)	0.132	0.107	71.68 (18.81)	83.87 (18.41)	0.025	0.028
MEF₂₅	60.03 (10.59)	72.34 (20.84)	0.113	0.130	61.02 (21.40)	74.35 (19.68)	0.021	0.024

^a BMI-SDS >2

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FVC, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Mann-Whitney U -test

^d Analysis of variance, adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months).

Table 4. Logistic regression: abnormal lung function in flow-volume spirometry (FVS) at 7.2 years of age, in relation to previous (at 4.0 years of age) and current (at 7.2 years of age) overweight.

Abnormal FVS ^b	Previous overweight ^a		aOR ^c (95% CI)	Current overweight ^a		aOR ^c (95% CI)
	Yes (N=5)	No (N=69)		Yes (N=17)	No (N=57)	
FEV ₁ <80%	0	4	-	1	3	1.01 (0.10-10.58)
FEV ₁ /FVC<88%	3	10	15.46 (1.65-145.26)	6	7	3.71 (1.01-13.60)
MEF ₅₀ <62%	2	11	6.46 (0.772-33.95)	3	11	0.89 (0.21-3.80)
MEF ₂₅ <48%	2	12	4.74 (0.61-36.76)	3	11	0.89 (0.22-3.70)
At least one abnormal parameter	3	16	6.63 (0.89-49.46)	6	14	1.63 (0.51-5.22)

^a BMI-SDS >1.3

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FCV, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Odds ratio (95% confidence interval) adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months)

Table 5. Logistic regression: abnormal lung function in flow-volume spirometry (FVS) at 12.3 years of age, in relation to previous (at 7.2 years of age) and current (at 12.3 years of age) overweight.

Abnormal FVS ^b	Previous overweight ^a		aOR ^c (95% CI)	Current overweight ^a		aOR ^c (95% CI)
	Yes (N=16)	No (N=58)		Yes (N=24)	No (N=50)	
FEV₁<80%	2	10	0.70 (0.13-3.60)	5	9	1.00 (0.29-3.46)
FEV₁/FVC<88%	6	6	5.64 (1.43-22.20)	8	5	4.02 (1.14-14.22)
MEF₅₀<62%	3	9	1.29 (0.30-5.54)	7	6	2.64 (0.78-8.97)
MEF₂₅<48%	4	6	3.51 (0.76-16.13)	6	5	2.47 (0.65-9.38)
At least one abnormal parameter	6	17	1.45 (0.45-4.67)	11	15	1.70 (0.63-4.55)

^a BMI-SDS >1.3

^b FVC (forced vital capacity), FEV₁ (forced expiratory volume in one second), FEV₁/FCV, MEF₅₀ (maximal expiratory flow at 50% of FVC) and MEF₂₅ (maximal expiratory flow at 25% of FVC), expressed as % of predicted.

^c Odds ratio (95% confidence interval) adjusted for RSV etiology of early wheezing and current anti-inflammatory asthma medication (during the preceding 12 months)