A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, south-west Indian Ocean

Alexandra Gross, Jeremy Kiszka, Olivier van Canneyt, Pierre Richard,
Vincent Ridoux

► To cite this version:

Alexandra Gross, Jeremy Kiszka, Olivier van Canneyt, Pierre Richard, Vincent Ridoux. A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, south-west Indian Ocean. *Estuarine, Coastal and Shelf Science*, 2009, pp.367-374. hal-00606243

HAL Id: hal-00606243

<https://hal.science/hal-00606243>

Submitted on 5 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**A preliminary study of habitat and resource partitioning among co-occurring
tropical dolphins around Mayotte, south-west Indian Ocean**

Gross Alexandra¹, Kiszka Jeremy^{1,2}, Van Canneyt Olivier³, Richard Pierre¹, Ridoux Vincent^{1,3*}

Running title: Tropical dolphin niche segregation

1: Littoral ENvironnement et Sociétés (LIENSs), UMR6250, Université de la Rochelle, 2 rue
Olympe de Gouges, 17000 La Rochelle, France.

2: Direction de l'Environnement et du Développement Durable, Collectivité Départementale de
Mayotte, BP 101, 97600 Mamoudzou, Mayotte, France. Email : jeremy.kiszka@wanadoo.fr

3: Centre de Recherche sur les Mammifères Marins, Université de la Rochelle, 23 avenue Albert
Einstein, F-17071 La Rochelle cedex, France.

* Corresponding author: vincent.ridoux@univ-lr.fr

Tel.: +33(0)5.4644.9910

Abstract

Mayotte in the southwest Indian Ocean is characterized by high dolphin diversity. They may coexist within a fairly small area around the island because they exploit neither the same preferential habitats nor the same resources. This preliminary study aimed to investigate ecological niche segregation among these delphinid communities: the Indo-Pacific bottlenose dolphin, *Tursiops aduncus*, the pantropical spotted dolphin, *Stenella attenuata*, the spinner dolphin, *Stenella longirostris*, and the melon-headed whale, *Peponocephala electra*. Two approaches were used. Habitat preferences were investigated by analysing dolphin sighting data and associated physiographical characteristics. Resource partitioning was explored by analysing C and N stable isotopes in skin and blubber biopsies. Only *T. aduncus*, which showed clear association with coastal habitats in the lagoon, differed from the others in terms of habitat preferences, characterised by shallow depth and slope, and proximity to the coast. All other species shared similar oceanic habitats immediately outside the lagoon, these being of higher depth and slope, greater distance from the coast and were not discernable by discriminant analysis. The two *Stenella* species and the melon-headed whale displayed very high overlap in habitat physiographic variables. The analysis of stable isotopes confirmed the ecological isolation of *T. aduncus* and revealed a clear segregation of *P. electra* compared to the two *Stenella* that was not apparent in the habitat analysis. This may reflect ecological differences that were not observable from diurnal surface observations.

Key-words: dolphins, ecological niche, habitats, stable isotopes, south-west Indian Ocean, Mayotte.

1. Introduction

In biological communities, each species has its own unique niche, which provides the conditions and resources needed for its survival. A shared resource in limited supply will bring about competition between members of the same species (intra-specific competition) or between individuals of different species (inter-specific competition). Competition can take two different forms: interference, which is a direct, often aggressive, interaction between individuals, or exploitation-competition, in which individuals interact with each other indirectly, by responding to a resource level which has been depressed by the activity of competitors (Begon et al., 1986).

A niche occupied by a species in the absence of competitors is termed its fundamental niche, whereas in the presence of competitors, species may be confined to a realized niche, which is shaped by the presence of competing sympatric species (Begon et al., 1986). Hence, the coexistence of potentially competing species is often made possible by the differentiation of their realized ecological niches. The first mechanism that allows for niche differentiation is resource partitioning. In this case, different species living in the same habitat exploit the resources differently. For example, predators of different size may feed on prey of different size, hence minimising the overlap between the various predators' prey size ranges. Prey specialization presumably allows for niche partitioning in areas of sympatry (Ballance, 2002). The second mechanism involves spatial (microhabitat differentiation) or temporal separation in the availability of the different resources (these become available at different times of the day or different seasons of the year; Huisman and Weissing, 2001).

Small delphinids belong to numerous species which have similar morphological characteristics. This situation suggests that fine-scale mechanisms allow for the partitioning of habitats and resources when and where the different species live in sympatry. A study of the cetacean community of Great Abaco, Bahamas, has shown that the ecological niches of the four species

that permanently live there do not overlap (MacLeod et al., 2004). These species capture prey at different depths of the water column. Other cetacean species are observed in the study area only during the season when prey abundance is sufficiently high to support their presence, while they are competitively excluded for the rest of the year (MacLeod et al., 2004). Indeed, the spatial distribution of marine predators is mainly determined by the distribution and availability of their prey, these in turn varying according to physical, chemical and biological characteristics of the water masses (Forcada, 2002).

The dietary ecology of marine mammals and their trophic level can be determined using different methods: traditional methods analyse faeces and regurgitated food of living animals, as well as the stomach contents of dead animals (from strandings or by-catch; e.g. Pusineri et al., 2007). A more recent method for studies of dietary ecology is stable isotope analysis of blubber, skin or muscle samples (Bearhop et al., 1993). The carbon and nitrogen isotope ratios ($^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$, expressed hereafter as $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of a consumer reflect those of its diet, with a slight retention of the heavier isotope and excretion of the lighter one (Das et al., 2003). As a consequence, tissues will be enriched with heavy isotopes at every trophic level. The minor stepwise trophic enrichment of the carbon-isotope ratio limits its use in assessing trophic levels, but enhances its use in tracking carbon sources through a food chain. The carbon isotope ratio of secondary and tertiary consumers should thus reflect the source of carbon at the base of their food chain (Kelly, 2000). The higher enrichment of the nitrogen isotope ratio of consumers compared to their prey makes it very useful for the determination of their trophic level (Kelly, 2000). Thus, both ratios help elucidate trophic relations and habitat use.

Mayotte, a volcanic island in the northern Mozambique Channel (southwest Indian Ocean), is characterized by the permanent presence of more than 20 species of cetaceans (Kiszka et al., 2007). Of these, the most important in coastal waters are the Indo-Pacific bottlenose dolphin,

Tursiops aduncus (Ehremberg, 1833), the pan-tropical spotted dolphin, *Stenella attenuata* (Gray, 1846), the spinner dolphin, *S. longirostris* (Gray, 1828), and the melon-headed whale, *Peponocephala electra* (Gray, 1846). The island has a great variety of marine ecosystems offering a large diversity of habitats: coasts, mangroves, an extended lagoon (1100 km²), different kinds of reefs (fringing, pinnacles, and barrier), a steep insular slope with many submarine canyons and seamounts, and the open ocean (Quod et al., 2000). At least twelve species of delphinids may coexist in a fairly small area around the island because they exploit neither the same habitats nor the same resources (Kiszka et al., 2007). The present study aimed to investigate ecological niche segregation among the resident dolphin community of Mayotte, especially the Indo-Pacific bottlenose dolphin, the pantropical spotted dolphin, the spinner dolphin and the melon-headed whale. We concentrated on these four species as they are of fairly similar size and can be found within the same proximity around Mayotte, in closely-related habitats within a small area and at all seasons (Kiszka et al., 2007). This is particularly so for the two *Stenella* species and the melon-headed whale which are all encountered immediately outside the barrier-reef and in the channels, whereas existing literature suggests that they would be more differentiated habitat-wise, with the spinner dolphin feeding offshore but resting inshore, the melon headed whale being an oceanic squid-eater and the Indo-Pacific bottlenose dolphin dwelling in nearshore coastal habitats (Norris et al., 1994; Silva et al., 2005; Perryman, 2002; Wells and Scott, 2002). Two approaches were used for the study: a comparison of habitat by the analysis of dolphin sighting data and associated behavioural and physiographic characteristics (group size, depth, slope, distance to the coast and proximity to the different kinds of reefs), and the study of habitat and resource partitioning by the analysis of C and N stable isotopes from skin and blubber biopsies.

2. Material and Methods

2.1 Study area

The volcanic island of Mayotte (45°10'E, 12°50'S), which is part of the Comoros archipelago, is located in the northern Mozambique Channel (Indian Ocean) between Madagascar and Southeast Africa. Its surface area is 376 km² and it is composed of two main islands and about 30 islets scattered within and around a lagoon. Mayotte is surrounded by a large lagoon-reef complex, whose width varies from 3 to 15 km. Fringing reefs surround the archipelago, an inner double-reef is present off the south-west end of Mayotte, and the barrier reef, which is interrupted by numerous channels, separates the lagoon itself (maximum depth 80 m) from the external slope and more oceanic habitats.

The four species of interest, the Indo-Pacific bottlenose dolphin, the pantropical spotted dolphin, the spinner dolphin, and the melon-headed whale, range in size from about 200 cm and 90 kg for the spinner dolphin to 250 cm and 250 kg for the melon-headed whale (Perrin, 2002a and b, Perryman, 2002, Wells and Scott, 2002).

2.2 Data and sample collection

Data were collected from 1997 to 2005, during small boat-based surveys dedicated to studying marine mammals (Figure 1). Several types of boats were used: a 7 m catamaran equipped with two, four-stroke, 60-hp outboard engines; a 7 m boat equipped with two, two-stroke, 40-hp outboard engines; a 6.4 m cabin boat equipped with an inboard four-stroke and 150-hp outboard engine. Surveys were conducted during daylight hours, i.e. between 0700 h and 1800 h, in sea conditions not exceeding Beaufort 3. The survey vessels did not follow pre-defined transects but sampling covered all habitats within the lagoon and over the external insular slope (Figure 1). Effort varied according to month (Figure 2), with more effort being applied in the austral summer

(November to January). When dolphins were encountered, preliminary information records included group size (maximum, minimum, best estimate), geographic position, activity (travelling, resting, foraging/feeding, socialising, milling, play), group classification on the basis of the relative size of individuals (adults, sub-adults, calves), research boat disturbance (bowride, approach, avoidance, no response) as well as group formation (tight, loose, dispersed, variable, convergent; Shane, 1990; Würsig et al., 1998). This study is still on-going and, therefore, only the sighting locations and associated physiographic variables are analysed here.

When conditions were optimal (good weather and sea state, dolphins closely approaching the boat), biopsies were collected using a cross-bow (*BARNETT Veloci-Speed® Class*) with Finn Larsen bolts and tips (20-mm). The dolphins were hit below the dorsal fin, when close (3-10 meters) to the research boat. Sampling periods spanned all seasons but sample sizes did not allow seasonal comparisons (January, August and December for *T. aduncus*, N=4; January, February, March and October for *S. attenuata*, N=4; February, March, May and October for *S. longirostris*, N=5; March, July and December for *P. electra*, N=4). It was not possible to determine sex, size or age of the individuals biopsied. Blubber and skin biopsy samples were preserved separately in 90° ethanol before shipping and subsequent analysis. Biopsy sampling was conducted under scientific permit #78/DAF/2004.

2.3 Database

We constituted a database in which every dolphin observation was associated with the physiographic characteristics (distance to the coast, to the different reefs and to the closest channel, as well as depth and slope of seafloor and the variance of these two parameters) corresponding to the GPS (Global Positioning System) fixes of the observation. The distance data were obtained using GIS (Geographic Information System) software ArcView (ArcGIS 8.2) by

ESRI (Environmental Systems Research Institute). Bathymetric data were obtained from *Service Hydrographique et Océanographique de la Marine* (SHOM). Interpolation of bathymetry data, needed to generate depth and slope data for each sighting, was undertaken with the extension Spatial Analyst by kriging transformation of the raster file into an interpolated data file. This was obtained by calculating the mean value of the twelve points closest to every bathymetric point in a 1 km² grid. This interpolated data file thus allowed us to generate depth and slope data over the whole study area and therefore to associate bathymetry data to any dolphin observation in the area.

2.4 Data analysis

The environmental data were first compared between species using basic methods (non-parametric analyses and ANOVA). Then we used multivariate statistical methods including discriminant linear and quadratic analyses to examine how well sightings were assigned to the correct species from the combination of associated physiographic variables. All methods were implemented with the software R (R-2.2.1, R Development Core Team, 2005).

2.5 Stable isotope analyses

Blubber and skin were separated for each biopsy. The ethanol they contained was evaporated at 45°C over 48 h and the samples were ground and freeze-dried (Hobson et al., 1997). Lipids were extracted using cyclohexane (C₆H₁₂) prior to analysis because they are depleted in ¹³C; if they were not extracted, this would cause a bias in the isotopic signature of ¹³C (De Niro and Epstein, 1978; Tieszen et al., 1983). Small sub-samples (0.35 to 0.45 mg ± 0.001 mg) were prepared for analysis. Stable isotope measurements were performed with a continuous-flow isotope-ratio mass spectrometer (Isoprime, Micromass) coupled to an elemental analyser (Eurovector EA 3024).

Results are expressed in δ notation relative to PeeDee Belemnite and atmospheric N_2 for $\delta^{13}C$ and $\delta^{15}N$, respectively, according to the equation $\delta X = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 10^3$, where X is ^{13}C or ^{15}N and R is the isotope ratio $^{13}C/^{12}C$ or $^{15}N/^{14}N$, respectively. Replicate measurements of internal laboratory standards (acetanilide) indicated that measurement errors were $<0.15\text{‰}$ and $<0.20\text{‰}$ for $\delta^{13}C$ and $\delta^{15}N$, respectively. Percent C and N elemental composition of tissues were obtained using the elemental analyzer and used to calculate the sample C:N ratio, indicating a good lipid removal efficiency when <4 .

3. Results

3.1 General

Our cetacean sightings comprised 394 positional data for *S. longirostris* ($n = 208$), *S. attenuata* ($n = 88$), *T. aduncus* ($n = 83$) and *P. electra* ($n = 15$) in all sectors around Mayotte (Figure 2) and all seasons (Figure 3). Indo-Pacific bottlenose were mostly found within the lagoon, very often in the immediate proximity of the fringing reef, and to a lesser extent over the North West bank located outside the main barrier reef. The two *Stenella* species were observed all along the barrier reef on its outer side and made only a few incursions into the lagoon. Finally, the melon-headed whale was only seen outside the barrier reef.

3.2 Habitat use and niche partitioning

A preliminary Principal Component Analysis (PCA plot not shown) allowed us to select group size, distance to coast, depth and slope as the set of physiographic variables with least redundancy. Distances to the various reefs and to the nearest channel were heavily redundant

relative to distance from coast and therefore not retained. A simple comparison of the distribution of these variables for the four focal species was performed using Kruskal-Wallis tests (Figure 4). *T. aduncus* is observed at short distances from the coast with shallow bathymetry and moderate slopes, whereas the other three species cannot be significantly differentiated in terms of habitat characteristics. In particular, the two *Stenella* species manifested similar characteristics in group size and environmental preferences. On the other hand, group size differed strongly between *T. aduncus* that lives in small groups (median = 6) and *P. electra* that is mostly found in groups of several hundred individuals (median = 300).

Linear discriminant analysis separated the species according to their environmental characteristics (Figure 5). The predictive power of the linear discriminant analysis was good only for *T. aduncus* which can be explained by its habitat preferences (Table 1, upper part). The other species' habitat characteristics were too similar to permit acceptable predictions. The good prediction for *S. longirostris* might be attributable only to the considerable number of sighting data for this species. Finally, the quadratic discriminant analysis provided slightly better predictive results, especially for *P. electra* (Table 1, lower part).

The different methods used to compare the preferred habitats of the four delphinid species under study all clearly showed that *T. aduncus* differs from the other species in its preferred environmental parameters. *S. attenuata* and *S. longirostris* share similar environmental characteristics. *P. electra* is characterised by a larger group size and deeper depth preference, but resembles *Stenella* in terms of the majority of its other environmental preferences (distance and slope).

3.3 Resource partitioning

Stable isotope ratios were lower in blubber than in skin but the pattern of differences observed between species was similar in both tissues (Figure 6). Skin and blubber of *T. aduncus* have the highest values of $\delta^{13}\text{C}$. The two *Stenella* species are not discernable from each other as shown by the extensive overlap in standard deviation for both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$, and have the lowest values for $\delta^{13}\text{C}$. *P. electra* has the highest $\delta^{15}\text{N}$ and a $\delta^{13}\text{C}$ values that are intermediate between those for the two *Stenella* species and *T. aduncus*. The intra-specific variance is represented by the standard deviation, which is more important for the $\delta^{13}\text{C}$ values than for the $\delta^{15}\text{N}$, except in the blubber samples of *T. aduncus* where intra-specific variance in the $\delta^{15}\text{N}$ is prevalent.

4. Discussion

The present study comprises a preliminary analysis of habitat and resource use among an assemblage of co-existing tropical delphinids living around Mayotte in the southwest Indian Ocean. The principal finding is that, among the four species of interest, the Indo-Pacific bottlenose dolphin is clearly differentiated from the other species in terms of both habitat preference and stable isotope analyses. From the variables tested here, the other three species can hardly be separated in terms of their preferred habitats but stable isotope analysis revealed a dietary segregation between the melon-headed whale and the two species of the genus *Stenella*. This was not initially evident in the sighting data analyses that described the dolphins' diurnal use of habitats. However, some limitations render these findings preliminary. The sightings constituted presence-only data as the observation effort could not be readily quantified and, hence, the data cannot provide significant information concerning dolphin absence. In this work, we tried to characterise the habitats where dolphins were found, not assess the overall distribution of each species around the island. Nevertheless, the field surveys comprehensively covered the

study area such that all habitats were visited and the sighting data of the four focal species were considered representative of their habitat preference.

The Indo-Pacific bottlenose dolphin, *Tursiops aduncus*, occupies an ecological niche that clearly differs from those occupied by the other study species. *T. aduncus* was observed only inside the lagoon except in the northern part of the island where the reef system is in the form of an open bank outside the lagoon. This species is generally found close to the fringing reef which constitutes the ecosystem where prey of this coastal dolphin may be concentrated. The high $\delta^{13}\text{C}$ value in its tissues indicates a benthic carbon source that dolphins more easily access in coastal habitats (Hobson, 1999). This species has indeed been observed feeding near mangroves, along the fringing reefs or over seagrass beds. Its isotopic signature of $\delta^{15}\text{N}$ is similar to that of the two *Stenella*, but this cannot be interpreted as an indication of similar trophic levels as the $\delta^{15}\text{N}$ values of the local primary producers *versus* those outside the lagoon have not been investigated at this stage.

T. aduncus lives and forages individually or in small groups inside the lagoon, where the water is shallow and large predators are absent, and probably feeds on prey that would not aggregate in large schools (Mann et al., 2000). Indo-Pacific bottlenose dolphins forage over reefs or soft bottom substrata and near the shore relatively close to the island of Mayotte and around Zanzibar (Tanzania) (Amir et al., 2005).

The pantropical spotted dolphin and the spinner dolphin share a barrier reef-associated habitat outside the lagoon. They are also found where the water depth rapidly attains a depth of 100 m. In comparison to the Indo-Pacific bottlenose dolphin, their tissues had a lower $\delta^{13}\text{C}$, suggesting it was derived from an oceanic carbon source; oceanic phytoplankton is reported to be ^{13}C -depleted

relative to marine phanerogams (Hobson, 1999). Their isotopic ratios suggested their diet comprised oceanic prey.

S. attenuata and *S. longirostris* can occasionally be observed inside the lagoon where they might take advantage of the safety it provides for resting. This kind of behaviour has been observed in Hawaiian spinner dolphin populations (Norris et al., 1994). *S. longirostris* generally lives in single-species groups from several dozens to several hundreds of individuals, e.g. around the main Hawai'i island (Norris et al., 1994). Their aggregation in large groups might offer some protection against predators but it might also facilitate feeding through communal hunting on large pelagic fish schools. In the present work, *S. attenuata* was rarely observed in single-species groups but was generally found in association with *S. longirostris*, forming important mixed-species groups. Both species might take advantage of a larger group size for safety and foraging. Nevertheless, a large group size increases the potential for intra-specific competition as well as inter-specific competition if the two species feed together. A study in the southwest Atlantic has shown that distributions of spotted dolphins and spinner dolphins may partially overlap (Moreno et al., 2005). Associations of spotted dolphins and spinner dolphins are frequently found in Hawai'i, where the two species travel together while showing many aggressive interactions, but they do not seem to feed together (Psarakos et al., 2003). Conversely, in the Azores, common dolphins, *Delphinus delphis*, and Atlantic spotted dolphins, *Stenella frontalis*, associate and forage together (Clua and Grosvalet, 2001). As *S. attenuata* and *S. longirostris* have similar habitat preferences and similar isotopic signatures, both for carbon and nitrogen, there is no evidence of ecological niche differentiation in these two species. Two options could be investigated in the future: either the two dolphins share the same resources in the same habitats, presumably because food is not limiting there, or they segregate when feeding on different

resources but their prey have similar isotopic signatures because they rely on the same carbon source at the same trophic level.

The melon-headed whale, *Peponocephala electra*, prefers steep slopes and slightly deeper water than the two *Stenella* species. *Peponocephala electra* forms large groups of several hundred individuals. As it was always observed in a habitat that overlaps with that of the spinner and spotted dolphins, analysis of its habitat characteristics failed to distinguish between it and the two *Stenella* species. This might be due to the fact that melon-headed whale sightings were only made on groups that occasionally approached the island, as the surveys were conducted in a limited area around Mayotte and groups living farther offshore in the open ocean would not have been observed. In this case, our understanding of the preferred habitat of this species remains marginal, being limited to its nearshore fringe. Stable isotope analysis, on the other hand, clearly showed that there is an ecological differentiation between *P. electra* and the two *Stenella* species. Values of $\delta^{13}\text{C}$ for *P. electra*'s were intermediate between those of the *Stenella* species and *T. aduncus*. This would suggest that carbon isotopic composition of *P. electra* is influenced more by benthic primary production than that of the two *Stenella* species. *Peponocephala electra* is known for its oceanic habitat (Perryman, 2002), and possible interpretations for its $\delta^{13}\text{C}$ include the possibility that it is a deep-diving species that feeds on prey derived from a detritus-based food web unlike the *Stenella* that feed on a phytoplankton-based food web. In addition, *P. electra* clearly differs in its $\delta^{15}\text{N}$, this being indicative of a higher trophic level, suggesting its diet includes more carnivorous fish and squid than the spotted and spinner dolphins. Again, as in *T. aduncus*, all sources of primary production should be investigated concerning their $\delta^{15}\text{N}$ to adequately interpret the trophic levels.

If there is high inter-individual variance in the $\delta^{15}\text{N}$ (*versus* $\delta^{13}\text{C}$) values within a species, the interpretation is that the species is composed of individuals that have varied feeding habits and prey on organisms at different trophic levels ($\delta^{15}\text{N}$) or are found in different habitats ($\delta^{13}\text{C}$). A low variance indicates that all the individuals are specialist feeders on similar prey or within the same habitat. Thus, stable isotope variance is sometimes considered a measure of niche width (Bearhop et al., 2004) or at least the inter-individual measure of niche width. Results of our stable isotope analysis indicate that *P. electra* is a fairly specialized feeder, whereas *S. attenuata*, *S. longirostris* and especially *T. aduncus* would be more eclectic feeders, which, in the latter case, is in line with previous studies (Mann et al., 2000).

Detectable differences between the standard deviations in the stable isotope composition in the skin and blubber samples might be attributable to their differential rates in tissue renewal: this takes a few days for epidermis but several months for the collagen matrix of the blubber (Abend and Smith, 1995). The residence time of elements in tissues depends on metabolic turnover rates (Rubenstein and Hobson, 2004). This enables tracking of an animal's foraging history as demonstrated by a study on pilot whales (*Globicephala melas*) in the North Atlantic (Abend and Smith, 1995). *T. aduncus* and the two *Stenella* species clearly differ in the respective isotopic carbon signatures in their skin and blubber, indicating that they feed on trophic webs with different carbon sources over the long term.

Each species' habitat preference is presumably based on their prey distributions (Baumgartner et al., 2000) which, in turn, are related to water depth (Hastie et al., 2005) and, indirectly, bathymetric features that influence currents and productivity (Fiedler, 2002). Foraging behaviour seems to be closely related to submarine habitat characteristics (Hastie et al., 2004). It has been demonstrated that a variable bathymetry contributes to global delphinid abundance, promoting

the regional abundance of different species (Gannier, 2005). The rich marine biodiversity of Mayotte is possibly related to its variety of habitats and these provide numerous ecological niches for delphinid prey.

5. Conclusions

The present study provides preliminary ecological indications of niche differentiation and resource partitioning within the Mayotte delphinid communities. The only species that differs from the others in terms of habitat characteristics is the coastal-dwelling Indo-Pacific bottlenose dolphin, whereas pantropical spotted and spinner dolphins and the melon-headed whale share similar oceanic habitats immediately outside the lagoon. Stable isotope analysis confirmed the ecological specialisation of *T. aduncus* and, in addition to this, revealed a clear segregation of *P. electra*, from the two *Stenella* species in terms of their feeding that was not apparent in the habitat analysis. This may reflect behavioural differences that were not detectable from diurnal surface observations. Further work should strengthen these conclusions through dedicated sampling of habitat preference and an expansion of the stable isotope studies. These should focus on seasonal changes in food partitioning and the isotopic composition of a series of putative prey species and primary producers characteristic of the inshore-offshore gradient of habitats found around Mayotte.

Acknowledgements

Sightings were collected from 1997 to 2004 during the austral winter (July to October) as part of the humpback whale (*Megaptera novaeangliae*) surveys conducted by the *Observatoire des Mammifères Marins de Mayotte* (*Direction de l'Agriculture et de la Forêt*, DAF) and the Cetacean Conservation and Research Program (American Museum of Natural History-Wildlife

Conservation Society). The other sightings were collected from November 2004 to August 2005 during a dolphin research project conducted by the *Office National de la Chasse et de la Faune Sauvage* (ONCFS) and the Agriculture and Forestry Office (*Direction de l'Agriculture et de la Forêt*). We thank Peter Ersts (American Museum of Natural History, Center for Biodiversity and Conservation) for providing the map showing tracks of the cetacean surveys conducted around Mayotte. We are particularly grateful to Gaël Guillou (*Université de la Rochelle*) for undertaking the mass spectrometer analyses. Funding for field work was provided by the *Ministère de l'Energie, l'Ecologie, le Développement Durable et de l'Aménagement du Territoire* (MEEDDAT) and the *Collectivité Départementale de Mayotte* (CDM). We thank Robin Rolland, Alban Jamon, Wilfrid Fousse, Ismaël Oussen (DAF), Sarah Caceres, Franck Charlier, Denis Girou (ONCFS), Didier Fray (CDM) and the personnel of *Brigade Nature* (CDM) for assistance in the field in Mayotte.

References

Abend, A.G., Smith, T.D., 1995. Differences in ratios of stable isotopes of nitrogen in long-finned pilot whales (*Globicephala melas*) in the western and eastern North Atlantic. *ICES Journal of Marine Science* 52: 837-841.

Amir, O.A., Berggren, P., Ndaro, S.G.M., Jiddawi, N.S., 2005. Feeding ecology of the Indo-Pacific bottlenose dolphin (*Tursiops aduncus*) incidentally caught in the gillnet fisheries off Zanzibar, Tanzania. *Estuarine Coastal and Shelf Science* 63: 429-437.

400 Ballance, L.T. 2002. Cetacean Ecology. *In*: Perrin, W.F., Würsig, B., Thewissen, J.G.M.
 401 (Editors), Encyclopedia of Marine Mammals, Academic Press, San Diego, pp 208-214.
 402

403 Baumgartner, M.F., Mullin, K.D., May, L.N., Leming, T.D., 2001. Cetacean habitats in the
 404 northern Gulf of Mexico. Fishery Bulletin 99: 219-239.
 405

406 Bearhop, S., Adams, C.E., Waldron, S., Fuller, R.A., Macleod, H., 2004. Determining trophic
 407 niche width: A novel approach using stable isotope analysis. Journal of Animal Ecology 73:
 408 1007-1012.
 409

410 Begon, M., Harper, J.L., Townsend, C.R., 1986. Ecology – Individuals, Populations and
 411 Communities. Blackwell Scientific Publications, Boston, 876 pp.
 412

413 Chapman, J.L., Reiss, M.J., 1999. Ecology – Principles and Applications. Cambridge University
 414 Press, Cambridge, 336 pp.
 415

416 Clua, E., Grosvalet, F., 2001. Mixed-species feeding aggregation of dolphins, large tunas and
 417 seabirds in the Azores. Aquatic Living Resources 14: 11-18.
 418

419 Das, K., Lepoint, G., Leroy, Y., Bouquegneau, J.M., 2003. Marine mammals from the southern
 420 North Sea: feeding ecology data from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements. Marine Ecology Progress
 421 Series 263: 287-298.
 422

423 De Niro, M.J., Epstein, S., 1978. Influence of diet on the distribution of carbon isotopes in
 424 animals. *Geochimica Cosmochimica Acta* 42: 495-506.

425

426 Estep, M.F., Dabrowski, H., 1980. Tracing food webs with stable hydrogen isotopes. *Science*
 427 209: 1537–1538.

428

429 Fiedler, P.C., 2002. Ocean environment. *In*: Perrin, W.F., Würsig, B., Thewissen, J.G.M.
 430 (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp 824-830.

431

432 Forcada, J., 2002. Distribution. *In*: Perrin, W.F., Würsig, B., Thewissen, J.G.M. (Editors),
 433 *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp 325-333.

434

435 Gannier, A., 2005. Summer distribution and relative abundance of delphinids in the
 436 Mediterranean Sea. *Revue d'Ecologie (Terre et Vie)* 60: 223-238.

437

438 Hastie, G.D., Swift, R.J., Slessor, G., Thompson, P.M., Turrell, W.R., 2005. Environmental
 439 models for predicting oceanic dolphin habitat in the Northeast Atlantic. *ICES Journal of Marine*
 440 *Science* 62: 760-770.

441

442 Hastie, G.D, Wilson, B., Wilson, L.J., Parsons, K.M., Thompson, P.M. 2004. Functional
 443 mechanisms underlying cetacean distribution patterns: hotspots for bottlenose dolphins are linked
 444 to foraging. *Marine Biology* 144: 397-403.

445

446 Hobson, K.A., 1999. Tracing origins and migration of wildlife using stable isotopes: a review.
 447 *Oecologia* 120: 314-326.
 448

449 Hobson, K.A., Gibbs, H.L., Gloutney, M.L. 1997. Preservation of blood and tissue samples for
 450 stable-carbon and stable-nitrogen isotope analysis. *Canadian Journal of Zoology* 75:1720-1723.
 451

452 Huisman, J., Weissing, F.J., 2001. Biological conditions for oscillations generated by
 453 multispecies oscillations. *Ecology* 82: 2682-2695.
 454

455 Kelly, J.F., 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian
 456 trophic ecology. *Canadian Journal of Zoology* 78: 1-27.
 457

458 Kiszka, J., Ersts, P.J., Ridoux, V., 2007. Cetacean diversity around the Mozambique Channel
 459 island of Mayotte (Comoros Archipelago). *Journal of Cetacean Research and Management* 9:
 460 105-109.
 461

462 Macleod, C.D., Hauser, N., Peckham, H., 2004. Diversity, relative density and structure of the
 463 cetacean community in summer months east of Great Abaco, Bahamas. *Journal of the Marine*
 464 *Biological Association of the U.K.* 84: 469-474.
 465

466 Mann, J., Connor, R.C., Tyack, P.L., Whitehead, H., 2000. *Cetacean Societies – Field Studies of*
 467 *Dolphins and Whales.* University of Chicago Press, Chicago, 433 pp.
 468

469 Moreno, I.B., Zerbini, A.N., Danilewicz, D., De Oliveira Santos, M.C., Simoes-Lopes, P.C.,
 470 Lailson-Brito, Jr. J., Azevedo, A.F., 2005. Distribution and habitat characteristics of dolphins of
 471 the genus *Stenella* (Cetacea: Delphinidae) in the southwest Atlantic Ocean. *Marine Ecology*
 472 *Progress Series* 300: 229–240.
 473
 474 Norris, K.S., Würsig, B., Wells, R. S., Würsig, M., 1994. *The Hawaiian Spinner Dolphin*,
 475 University of California Press, Berkeley, 408 pp.
 476
 477 Perrin, W.F., 2002a. Pantropical spotted dolphin *Stenella attenuata*. *In*: Perrin, W.F., Würsig, B.,
 478 Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp
 479 865-867.
 480
 481 Perrin, W.F., 2002b. Spinner dolphin *Stenella longirostris*. *In*: Perrin, W.F., Würsig, B.,
 482 Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp
 483 1174-1178.
 484
 485 Perrin, W.F., Würsig, B., Thewissen, J.G.M., 2002. *Encyclopedia of Marine Mammals*,
 486 Academic Press, San Diego, 1414 pp.
 487
 488 Perryman, W.L., 2002. Melon-headed whale *Peponocephala electra*. *In*: Perrin, W.F., Würsig,
 489 B., Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego,
 490 pp 733-735.
 491

492 Psarakos, S., Herzing, D.L., Marten, K., 2003. Mixed-species associations between Pantropical
 493 spotted dolphins (*Stenella attenuata*) and Hawaiian spinner dolphins (*Stenella longirostris*) off
 494 Oahu, Hawaii. *Aquatic Mammals* 29: 390–395.
 495
 496 Quod, J.P., Naim, O., Abdourazi, F., 2000. The Comoros archipelago. *In*: Sheppard, C. (Editors),
 497 Seas at the Millennium: An Environmental Evaluation. Pergamon Press, Oxford, pp 243-52.
 498
 499 R Development Core Team, 2005. R: A language and environment for statistical computing. R
 500 Foundation for Statistical Computing. Vienna, Austria. (<http://www.Rproject.org>)
 501
 502 Pusineri, C., Magnin, V., Meynier, L., Spitz, J., Hassani, S., Ridoux, V., 2007. Food and feeding
 503 ecology of the common dolphin (*Delphinus delphis*) in the northeast Atlantic and comparison
 504 with its diet in neritic areas. *Marine Mammal Science* 23: 30-47.
 505
 506 Rubinstein, D.R., Hobson, K.A., 2004. From birds to butterflies: animal movement patterns and
 507 stable isotopes. *Trends in Ecology and Evolution* 19: 256-263.
 508
 509 Silva, J.M., Silva, F.J.L., Sazima, I., 2005. Rest, nurture, sex, release, and play: diurnal
 510 underwater behaviour of the spinner dolphin at Fernando de Noronha Archipelago, SW Atlantic.
 511 *Journal of Ichthyology and Aquatic Biology* 9: 161–176.
 512
 513 Tieszen, L.L., Boutton, T.W., Tesdahl, K.G., Slade, N.A., 1983. Fractionation and turnover of
 514 stable carbon isotopes in animal tissues: implications for $\delta^{13}\text{C}$ analysis of diet. *Oecologia* 57: 32-
 515 37.

516

517 Shane, S.H., 1990. Behaviour and ecology of the bottlenose dolphin at Sanibel Island, Florida. In:
518 S. Leatherwood, R.R. Reeves (Editors.). The Bottlenose Dolphin. Academic Press, San Diego,
519 pp. 256-266.

520

521 Wells, R.S., Scott, M.D., 2002. Bottlenose dolphins *Tursiops truncatus* and *T. aduncus*. In:
522 Perrin, W.F., Würsig, B., Thewissen, J.G.M. (Editors), Encyclopedia of Marine Mammals,
523 Academic Press, San Diego, pp 122-128.

524

525 Würsig, B., Lynn, S.K., Jefferson, T.A., Mullin, K.D., 1998. Behaviour of cetaceans in the
526 Northern Gulf of Mexico relative to survey ships and aircraft. Aquatic Mammals 24: 41-50.

527

528

Figure captions

Figure 1: Location of the study area and observation routes from July 2004 to August 2005. Isobath (left) and GPS tracks of sighting surveys (right) are shown around Mayotte Island with barrier reef.

Figure 2: General locations of delphinid sightings around Mayotte and its reefs.

Figure 3: Temporal distribution of effort expressed as number of sightings per three-month period.

Figure 4: Delphinid sighting median values, 50, 75 percentiles and outliers of (a) group size, (b) distance to the coast, (c) depth and (d) seafloor slope. Kruskal Wallis tests showed that *T. aduncus* differed significantly in all cases with p -values $<10^{-6}$.

Figure 5: Sighting density for melon-headed whale (dotted line), pan-tropical spotted (black line), spinner (dashed line) and Indo-Pacific bottlenose dolphin (dashed and dotted line) along the first discriminant axis.

Figure 6: Stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in ‰) values in delphinid blubber (right-hand graph) and skin (left-hand graph). Graphs show average values and standard deviations. Black squares represent *S. longirostris* (N=5), white squares *S. attenuata* (N=4), black diamonds *T. aduncus* (N=4) and white diamonds *P. electra* (N=4).

Table 1: Discriminant analysis: linear and quadratic predictions for the four species. Each sighting was assigned to one of the four study species on the basis of its associated physiographic characteristics. Only *T. aduncus* data were correctly assigned to the right species whereas all others were mostly assigned to *S. longirostris* as they all share similar habitat characteristics.

Linear discriminant analysis

	<i>Peponocephala electra</i>	<i>Stenella attenuata</i>	<i>Stenella longirostris</i>	<i>Tursiops aduncus</i>
<i>P. electra</i>	2	3	5	0
<i>S. attenuata</i>	1	3	1	2
<i>S. longirostris</i>	12	79	194	19
<i>T. aduncus</i>	0	2	8	62
Good prediction	14%	3%	93%	77%

Quadratic discriminant analysis

	<i>Peponocephala electra</i>	<i>Stenella attenuata</i>	<i>Stenella longirostris</i>	<i>Tursiops aduncus</i>
<i>P. electra</i>	11	4	4	0
<i>S. attenuata</i>	1	18	18	1
<i>S. longirostris</i>	3	46	131	4
<i>T. aduncus</i>	0	19	55	78
Good prediction	79%	20%	63%	96%

Fig.1

597

598 Fig.2

599

600

601 Fig.3

602

603

604

605

606

607 Fig.4

608

609 Fig. 5

610

611

612

613

Fig. 6