Epidemiology of borderline oxacillin-resistant Staphylococcus aureus in pediatric cystic fibrosis.

Timothy Ronan Leahy, Yvonne Yau, Eshetu Atenafu, Mary Corey, Felix Ratjen, Valerie Waters

► To cite this version:

Timothy Ronan Leahy, Yvonne Yau, Eshetu Atenafu, Mary Corey, Felix Ratjen, et al.. Epidemiology of borderline oxacillin-resistant Staphylococcus aureus in pediatric cystic fibrosis.. Pediatric Pulmonology, 2011, 10.1002/ppul.21383 . hal-00614855

HAL Id: hal-00614855

<https://hal.science/hal-00614855>

Submitted on 17 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Epidemiology of borderline oxacillin-resistant
Staphylococcus aureus in pediatric cystic fibrosis.**

Journal:	<i>Pediatric Pulmonology</i>
Manuscript ID:	PPUL-10-0257.R1
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	19-Oct-2010
Complete List of Authors:	leahy, timothy; Institute of Molecular Medicine, Genomic Research Laboratory Yau, Yvonne; The Hospital for Sick Children, Paediatric Laboratory Medicine Atenafu, Eshetu; The Hospital for Sick Children, Child Health Evaluative Sciences Corey, Mary; Hospital for Sick Children Ratjen, Felix; Hospital for Sick Children, Division of Resp. Medicine Waters, Valerie; Hospital for Sick Children
Keywords:	methicillin resistant Staphylococcus aureus , mecA, pulsed-field gel electrophoresis

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Epidemiology of borderline oxacillin-resistant *Staphylococcus aureus* in pediatric cystic fibrosis.

Timothy Ronan Leahy*, MB MRCPI, Institute of Molecular Medicine, Trinity College Dublin, St James’s Hospital, James’s St, Dublin 8, Ireland; formerly of Division of Infectious Diseases, Department of Pediatrics, The Hospital for Sick Children, Toronto.

Yvonne C.W. Yau, MD, FRCPC, Division of Microbiology, Department of Pediatric Laboratory Medicine, The Hospital for Sick Children, Toronto.

Eshetu Atenafu, MSc, Child Evaluative Health Sciences, Research Institute, The Hospital for Sick Children, Toronto.

Mary Corey, PhD, Child Evaluative Health Sciences, Research Institute, The Hospital for Sick Children, Toronto.

Felix Ratjen, MD, PhD, FRCPC, Division of Respiratory Medicine, Department of Pediatrics, The Hospital for Sick Children, Toronto.

Valerie Waters, MD, MSc, FRCPC Division of Infectious Diseases, Department of Pediatrics, The Hospital for Sick Children, 555 University Avenue, Toronto, M5G 1X8, Canada.

*Corresponding author: telephone number +353 1 8961000, fax number +353 1 8963503, email address leahyt@tcd.ie

Keywords: methicillin resistant *Staphylococcus aureus* (MRSA), *mecA*, pulsed-field gel electrophoresis.

Abbreviated title. BORSA in pediatric cystic fibrosis

24 Summary

25 A single-centre retrospective study was undertaken in children with cystic fibrosis (CF)
26 to evaluate 1) risk factors for acquisition; 2) molecular epidemiology; and 3) impact on
27 disease progression of borderline oxacillin-resistant *Staphylococcus aureus* (BORSA)
28 versus *mecA*-positive methicillin-resistant *Staphylococcus aureus* (MRSA). The study
29 comprised of 1) identification of all children with at least one respiratory specimen
30 positive for either BORSA or MRSA during the study period; 2) compilation of relevant
31 clinical and epidemiological data from 12-month period leading up to first positive
32 (index) culture; 3) microbiological and molecular characterisation of index isolates and 4)
33 measurement of subsequent clinical outcome. Thirty-eight children were identified with
34 at least one positive isolate; 4 were excluded due to insufficient clinical or laboratory
35 data. Eighteen children (53%) grew BORSA in their index culture. Children who
36 acquired BORSA only (n=16) were more likely to have had prior MSSA colonization
37 (p<0.0001). Usage of oral cephalexin (p<0.01) and inhaled tobramycin (p<0.03) prior to
38 index culture was significantly and independently associated with acquisition of BORSA.
39 The majority of BORSA isolates were hyper β -lactamase producers and susceptible to a
40 greater range of antibiotics. Strain relatedness was not evident within the BORSA group.
41 There was no difference in disease progression between the two groups. This is the first
42 study to demonstrate that a significant proportion of *S. aureus* isolates with methicillin
43 resistance in the CF population are BORSAs that lack *mecA*. Antibiotic pressure may
44 lead to the development of BORSA in CF patients. Prospective studies are needed to
45 assess its clinical impact.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

46 **Introduction**

48 Methicillin-resistant *Staphylococcus aureus* (MRSA) has emerged as an important
49 respiratory pathogen in the cystic fibrosis (CF) population. The US CF registry reported a
50 significant rise in MRSA infection among CF patients, from 2.1% in 1996 to 21.2% in
51 2007¹. MRSA infection has been associated with accelerated deterioration in pulmonary
52 function ², increased hospitalization ³, impaired growth, increased antibiotic usage ⁴ and
53 increased mortality in CF patients ⁵.

54 According to the Clinical Laboratory Standards Institute (CLSI), methicillin
55 resistance in *S. aureus* is defined as having an oxacillin minimum inhibitory
56 concentration (MIC) > 4 µg/ml⁶. The primary mechanism of resistance for MRSA is the
57 production of an altered penicillin binding protein (PBP2a) with reduced affinity for
58 methicillin that is encoded by the *mecA* gene ⁷. We have identified a group of CF patients
59 infected with borderline resistant *S. aureus* (BORSA) isolates that are characterized by
60 oxacillin MICs at or just above the susceptibility breakpoint of 4 µg/ml and lack the
61 *mecA* gene ⁷. The implications of isolating BORSA in a respiratory culture from CF
62 patients have not previously been explored. The aim of this study was therefore to
63 compare the risk factors for acquisition, the molecular epidemiology and subsequent
64 clinical impact on disease progression of BORSA infection in comparison with *mecA*-
65 positive MRSA infection in pediatric CF patients.

68 **Materials and Methods**

69

70 Study design and patient population

71 This was a single-centre retrospective comparative study of pediatric CF patients
72 whose care was primarily based at the CF clinic at the Hospital for Sick Children,
73 Toronto between 1992 and 2007. The clinic currently cares for approximately 280
74 children up to the age of eighteen. Respiratory samples (expectorated sputum or
75 oropharyngeal swabs) were sent from all children attending the clinic to the microbiology
76 laboratory for culture at every 3-month visit and during acute pulmonary exacerbations
77 (APEs). Patients with one or more respiratory specimens positive for *S. aureus* with
78 methicillin resistance (either BORSA or MRSA) were included in the analysis. Cases
79 with three or more specimens positive for BORSA or MRSA during the study period
80 were considered persistently colonized². The study was approved by the Hospital for
81 Sick Children's Research Ethics Board.

83 Data collection

84 Patients with BORSA or MRSA isolated from their respiratory specimens were
85 identified using the microbiology database, and cross-referenced with the CF clinical
86 database. Date of the first BORSA or MRSA-positive respiratory culture (index culture)
87 for each patient was identified. The medical and laboratory records for each patient were
88 examined using a standardized data collection form. Baseline patient characteristics were
89 assessed at the date of index culture, and risk factors for acquisition of BORSA or MRSA
90 infection were assessed over the twelve-month period prior to the index culture. Impact

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

91 on clinical progression was assessed by examination of the time period from index
92 culture until discharge from the clinic or until December 31st 2008, whichever date was
93 later. Data collected included information on patients’ demographics, pulmonary function
94 tests (PFTs), respiratory cultures, hospital admissions, acute pulmonary exacerbations
95 and antibiotic use. An **acute pulmonary exacerbation** was defined as a change in
96 pulmonary status considered by the attending physician to require antibiotic treatment,
97 either oral or intravenous. An “antibiotic day” was defined as one day for each antibiotic
98 used, for example a 14-day course of two antibiotics was defined as 28 “antibiotic days”.
99 The purpose of this distinction was to capture more complex antibiotic regimens that
100 might be employed to treat MRSA infected patients. Patients were classified as being on
101 oral cephalexin if they were prescribed cephalexin at the time of index culture of at least
102 one month’s duration (in all instances, this occurred in the absence of signs or symptoms
103 of a pulmonary exacerbation). Patients were classified as being on inhaled tobramycin if
104 they were prescribed inhaled tobramycin at the time of index culture of at least one
105 month’s duration (**for previous *P. aeruginosa* infection**). Patients were classified as being
106 on oral ciprofloxacin if they were prescribed at least one course of ciprofloxacin (in all
107 instances, this occurred in the presence of signs or symptoms of a pulmonary
108 exacerbation). Patients were categorized as co-colonized with other CF pathogens using
109 previously defined criteria ⁸. Only patients with at least 6 months of follow-up after the
110 index culture were included in the analysis of outcome measures.

114 Microbiology

115 **Specimen Processing**

116 Respiratory specimens from CF patients were screened for MRSA using media
117 containing 2 µg/ml of oxacillin: an in-house mannitol salt agar (MRSA agar base,
118 Acumedia, Lansing, MI, and 1% D-mannitol, Difco, Detroit, MI) supplemented with 1%
119 thymidine and 2 µg/ml of oxacillin (both from Sigma, St. Louis, MO) was used from
120 1992 to 2001, and Oxacillin Resistance Screening Agar Base (ORSAB, Oxoid, Nepean,
121 Ontario, Canada) was used from 2002 onward. Plates were incubated at 35°C aerobically
122 and observed for the presence of growth at 24 hours and at 48 hours. Suspicious colonies
123 were identified as *S. aureus* by Gram stain, catalase, slide coagulase, and tube coagulase
124 tests.

126 **Antimicrobial Susceptibility**

127 Antimicrobial susceptibility of *S. aureus* isolates was tested by a combination of
128 oxacillin 6µg/ml screen plate, disk diffusion^{6,9} and an automated system (Vitek, bio-
129 Merieux, St. Laurent, Quebec, from 1992 to Nov 2002, and the BD Automated Phoenix
130 System, BD Diagnostic Systems, Sparks, MD from Nov 2002 onward). Inducible
131 resistance to clindamycin was detected by disk diffusion using the D-zone test^{6,9}. In
132 addition, each strain was examined for the presence of PBP2a by latex agglutination
133 (PBP2' test kit, Oxoid, Hants, UK). Organisms that were negative for PBP2a by latex
134 agglutination but grew on the oxacillin screen plate were sent to the reference laboratory
135 (Central Provincial Health Laboratory, Ontario) for minimum inhibitory concentration
136 (MIC) by agar dilution and polymerase chain reaction (PCR) for the detection of the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

mecA gene. An isolate was deemed as BORSA if the oxacillin MIC was $\geq 4 \mu\text{g/ml}$, PBP2a was absent and the *mecA* PCR was negative⁷. An isolate was deemed as MRSA if the oxacillin MIC was $\geq 4\mu\text{g/ml}$, the latex agglutination for PBP2a was positive and/or the *mecA* PCR was positive⁶.

β-Lactamase Production

All BORSAs were tested for the production of β-lactamase as a potential mechanism of oxacillin resistance by disk diffusion with ampicillin (10 μg) and amoxicillin-clavulinic (20 μg/ 10 μg) disks as previously described (13). An isolate was deemed as hyper β-lactamase producing if the ampicillin zone size was resistant and the amoxicillin-clavulanic zone size was susceptible, suggesting borderline resistance was secondary to large amounts of β-lactamase⁹(13).

Molecular Characterization

The *mecA* and *nuc* gene were co-amplified with a multiplex real time PCR as described previously¹⁰. The presence of Panton-Valentine leukocidin was examined by amplification of the *lukF-PV* and *lukS-PV* genes using primer and protocols previously described¹¹. Pulsed-field gel electrophoresis (PFGE) was conducted using *SmaI* as per manufacturer protocol (Bio-rad Laboratories, Hercules, CA) on all newly identified MRSA strains. Electrophoretically-generated DNA profiles were analyzed using the criteria proposed by Tenover et al to determine strain relatedness¹².

1
2
3 160 Statistical analysis
4

5
6 161 Descriptive statistics, frequency distributions and percentages were calculated for
7
8 162 the outcome variables and other covariates of interest. Normality assumption test was
9
10 163 carried out for the continuous variables using the Kolmogorov-Smirnov test. Student's t-
11
12 164 test or Kruskal-Wallis test (if normality assumption failed) was conducted for continuous
13
14 165 variables within the two groups. The Chi-square or Fisher's exact test, as appropriate,
15
16 166 was carried out for the dichotomous variables to assess the association with the two
17
18 167 groups. To analyze independent risk factors, a multivariate analysis using a logistic
19
20 168 regression model was applied. A p-value of <0.05 was considered statistically significant.
21
22 169 The statistical programme SAS (version 9.1; SAS Institute Inc., Cary, NC) was used.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Patient population

Between 1992 and 2007, 38 patients had at least one respiratory specimen positive for either BORSA or MRSA. Of these, three were excluded from our analysis because their care was not primarily based at our hospital, thus we had no clinical data for these patients. In addition, *mecA* PCR could not be performed on one isolate retrospectively therefore 34 patients were included in the study. Of these 34 patients, two patients were initially colonized with *mecA*- negative and subsequently *mecA*-positive strains and were excluded from the final statistical analysis as it was unclear how to categorize them in terms of patient characteristics and outcomes. Ultimately 16 patients with *mecA* positive MRSA and 16 with *mecA* negative BORSA on the index culture were compared for baseline characteristics. For the longitudinal assessment, 15 patients in each group were compared since one patient in each group did not have follow-up data for at least 6 months.

Patient characteristics and risk factors for BORSA and MRSA

The two groups were similar in terms of baseline characteristics (Table 1). There was no difference in pulmonary function or rate of decline of pulmonary function in the year prior to the index culture. In addition, there was no difference in the number of acute pulmonary exacerbations, hospital admissions or overall antibiotic use for the treatment of acute pulmonary exacerbations. A significantly greater proportion of BORSA infected patients were colonized with MSSA in the year prior (94% vs 25%, $p<0.0001$). Likewise,

1
2
3 194 a higher proportion of BORSA infected patients received oral antibiotics (cephalexin)
4
5 195 (75% vs 25%, $p=0.005$).
6
7
8 196 Mean duration of therapy among those patients on cephalexin was 337 days (range 30-
9
10 197 365 days) in the BORSA group and 311 days (range 150-365 days) in the MRSA group;
11
12 198 cephalexin treatment was instituted not for pulmonary exacerbation but as treatment for
13
14 199 MSSA identified on routine sputum cultures. Furthermore, a significantly greater
15
16 200 number of BORSA infected patients had received a course of oral ciprofloxacin (31% vs
17
18 201 0%, $p=0.043$) or were on inhaled antibiotics (tobramycin) (88% vs 44%, $p=0.009$) in the
19
20 202 year prior to the index culture than patients with MRSA. No inter-group difference was
21
22 203 detected in any of the other co-colonizing respiratory pathogens. Patients with BORSA
23
24 204 strains had significantly more clinic visits in the year prior to first isolate ($n=7.94$) than
25
26 205 patients with *mecA*-positive strains ($n=4.69$, $p=0.049$) although this was not a significant
27
28 206 risk factor in the logistic regression model
29
30
31
32
33

34 207 A logistic regression model was applied to determine the independence of
35
36 208 epidemiological and clinical variables that were correlated with BORSA infection.
37
38 209 Colonization with MSSA and use of oral cephalexin were significantly associated as 68%
39
40 210 of patients with MSSA were treated with cephalexin and 81% of patients treated with
41
42 211 cephalexin were colonized with MSSA ($p<0.05$). The use of oral cephalexin (regression
43
44 212 coefficient (β)=1.15, SD=0.47, p value=0.015) and inhaled tobramycin (regression
45
46 213 coefficient (β)=1.16, SD=0.53, p value=0.028) in the year prior to index culture were
47
48 214 significantly and independently associated with BORSA infection.
49
50
51
52

53 215

54
55 216 Microbiological characteristics of BORSA and MRSA isolates
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

217 The microbiological characteristics of the BORSA isolates are illustrated in Table
218 2. Fourteen of the 16 isolates had oxacillin MICs at or just above the breakpoint (4 to 8
219 µg/ml) but 2 isolates had MICs of 16 µg/ml. All of the BORSA isolates were negative for
220 PBP2a by latex agglutination and were *mecA* negative by PCR. Twelve of the 16 BORSA
221 isolates were hyper β-lactamase producers. This was shown by resistance to ampicillin by
222 disk diffusion that was reverted to susceptible with the addition of clavulanic acid, a β-
223 lactamase inhibitor. The remaining 4 isolates had ampicillin zone size that were
224 unchanged by addition of clavulin by disk diffusion.

225 In comparison to the MRSA isolates, a significantly greater proportion of the
226 BORSA isolates were susceptible to both erythromycin (81% vs 13%, $p < 0.0001$) and
227 clindamycin (94% vs 31%, $p < 0.001$). None of the BORSA isolates were D-test positive
228 but 5 (45%) of the MRSA isolates were D-test positive ($p = 0.043$). All of the BORSA
229 and MRSA isolates were susceptible to trimethoprim-sulphamethoxazole. None of the
230 BORSA isolates tested produced PVL. Three of the 16 MRSA isolates were identified as
231 USA300 epitypes, and produced PVL.

232
233 Relatedness of BORSA and MRSA isolates

234 Pulsed field gel electrophoresis (PFGE) revealed that none of the BORSA strains
235 were related by the Tenover criteria¹³ (Figure 1). Within the MRSA group, we found
236 four separate instances of strain relatedness during the course of the study: a pair of
237 siblings shared an indistinguishable strain, four patients shared possibly related strains,
238 two patients shared a closely related strain with a third patient sharing a possibly related

239 strain and, finally, three patients (including a pair of siblings) shared an indistinguishable
240 strain.

241

242 Clinical outcomes of patients

243 To compare the clinical impact of BORSA versus MRSA infection on disease
244 progression, we assessed 99.7 patient-years of follow-up after the index culture with a
245 comparable mean duration of follow up of 3.37 years in the MRSA group and 3.27 years
246 in the BORSA group ($p=0.507$) (Table 3). Patients infected with MRSA were more
247 likely to have persistent infection at the end of follow up than patients with BORSA
248 infection (80% vs 53%) but this was not statistically significant ($p=0.121$). The annual
249 decline in percent predicted forced expiratory volume in 1 second (FEV_1), was greater
250 among the MRSA infected patients (5.8%) than those infected with BORSA (2.8%)
251 although this difference was not significant ($p = 0.77$). Neither was there a significant
252 difference in the number of acute pulmonary exacerbations, hospital admissions, inpatient
253 days or clinic visits. One patient in the MRSA group died, and another progressed to lung
254 transplant. Three patients in the BORSA group had a lung transplant during the study
255 period.

256

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

This is the first study to show that a significant proportion (50%) of *S. aureus* isolates with methicillin resistance infecting CF patients are BORSA which lack the *mecA* gene, the main genetic determinant of methicillin resistance in *S. aureus*⁷. The present study demonstrates that antibiotic use in CF patients colonized with MSSA is a risk factor specifically for infection with BORSA. Previous studies have shown that ciprofloxacin and cephalosporin use is a risk factor for acquisition of MRSA in CF patients¹⁴ but, to our knowledge, no other study has examined the differential risk factors for BORSA compared to MRSA acquisition.

Transmission of traditional MRSA between patients^{15,16}, within the community¹⁷ and even within households¹⁸ has been well described in the CF population. Evidence of person to person transmission of *mecA*-positive MRSA was also present in this study with shared strains among CF patients and siblings. The mechanism of resistance is relevant to how CF patients acquire MRSA as the *mecA* gene is absent in its entirety from MSSA^{19 7}. It is believed to have originated from another staphylococcus species, such as *Staphylococcus sciuri*, and been transposed into MSSA²⁰. Although horizontal transfer of *mecA* between a *S. epidermidis* and MSSA isolate within the same patient has been described²¹, transmission of *mecA*-positive MRSA isolates between individuals is likely more common as there are a well defined number of *mecA*-positive MRSA clones worldwide²².

Our data would suggest that CF patients may acquire *mecA*-negative BORSA isolates in a different way. In contrast to MRSA isolates, BORSA isolates had unique

1
2
3 280 PFGE patterns and appeared not to be shared between patients in our study. CF patients
4
5 281 colonized with MSSA and exposed to oral cephalexin or inhaled tobramycin (which can
6
7 282 act synergistically with β -lactam antibiotics against *S. aureus*) were at significantly
8
9 283 increased risk of acquiring BORSA infection. This suggests that BORSA isolates may
10
11 284 arise from endogenous MSSA isolates that develop other mechanisms of resistance to
12
13 285 methicillin in response to antibiotic pressure. Previous studies have shown that *S. aureus*
14
15 286 strains with borderline susceptibility to oxacillin have a significant drop in the penicillin
16
17 287 MICs to the susceptible range with the addition of a β -lactamase inhibitor²³. This
18
19 288 predicts the response of oxacillin MICs to the addition of a β -lactamase inhibitor and
20
21 289 suggests that a large amount of β -lactamase production is the mechanism of resistance to
22
23 290 oxacillin in the majority of BORSA, as observed in 12 of 16 isolates in our study.
24
25 291 However, in the 4 remaining BORSA strains where the ampicillin zone size did not
26
27 292 change significantly with the addition of clavulanic acid, the mechanism of methicillin
28
29 293 resistance is unknown. In the aforementioned study, the authors described *S. aureus*
30
31 294 strains with intermediate MICs to oxacillin that remained the same in the presence of
32
33 295 clavulanic acid. The authors suggested that oxacillin resistance was due to factors other
34
35 296 than β -lactamase inactivation²³. It is possible that it is due to a modification of “normal”
36
37 297 penicillin-binding proteins (PBPs) such as PBP 1, PBP 2 and PBP 4 as was previously
38
39 298 shown in clinical BORSA isolates by Tomasz et al²⁴. Exposure of susceptible
40
41 299 staphylococci to increasing concentrations of β -lactam antibiotics in the laboratory is
42
43 300 known to cause PBP alterations associated with increased MICs²⁵. In this study,
44
45 301 persistent antibiotic pressure may have similarly modified PBPs in endogenous MSSA
46
47 302 isolates colonizing CF patients and resulting in the development of *mecA*-negative, non-
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

hyper β lactamase producing BORSA isolates. Thus, in the prevention of the emergence of BORSA in CF patients, other infection control measures, such as patient segregation, may not be as important as the judicious use of antibiotics.

In our study, the majority of patients treated with oral cephalexin were colonized with MSSA. Although there were no signs and symptoms of a pulmonary exacerbation at the time, we are limited in a retrospective study to the information that is recorded in the medical charts. Their antibiotic treatment was thus likely driven by their MSSA positive respiratory cultures and/or more subtle symptoms. Primary prophylaxis for *S. aureus* (the treatment of all newly diagnosed CF patients with antistaphylococcal agents irrespective of respiratory culture results) is not routinely practiced in North America due to concerns of *P. aeruginosa* acquisition^{26,27}. A systematic review of prophylactic antibiotics in cystic fibrosis, however, suggests that *P. aeruginosa* acquisition in this context is associated less frequently with narrow spectrum anti-staphylococcal antibiotics such as flucloxacillin compared with broader spectrum anti-staphylococcal antibiotics such as cephalexin²⁸. In our study, although patients treated with oral cephalexin were not receiving prophylaxis against *S. aureus*, they were receiving prolonged courses of cephalexin for MSSA detected on routine culture of the respiratory tract. This prolonged cephalexin use was identified as an independent risk factor for the acquisition of BORSA isolates and occurred recently, primarily in the last 5 years of the study.

The clinical significance of BORSA infection in CF patients is unknown. Our study lacked a sufficient sample size and is not powered to clearly delineate the clinical impact of BORSA infection. Previous studies have reported accelerated deterioration in pulmonary function², increased hospitalization³, impaired growth, increased antibiotic

usage⁴ and increased mortality⁵ in CF patients infected with MRSA. However, these studies are limited by a lack of information on the laboratory methodology of identifying MRSA and thus it is not known to what degree these effects are attributable to BORSA or MRSA.

Detection of BORSA isolates may also influence the choice of antibiotics in the treatment of these infections in the setting of a pulmonary exacerbation. In our study, BORSA isolates were more susceptible to erythromycin and clindamycin than the MRSA isolates. This finding is not surprising as the *mecA* gene is located on a mobile genetic element (the Staphylococcal Cassette Chromosome *mec*: SCC*mec*) which can contain additional drug resistance genes²². Although earlier studies suggested that infections due to *mecA* negative BORSA isolates could be treated with β -lactam antibiotics^{29,30}, more recent cases report failure of cloxacillin and even vancomycin in the treatment of endocarditis caused by BORSA, depending on the mechanism of methicillin resistance^{31,32}. This has implications for how respiratory specimens from CF patients should be screened for MRSA. Some of the newer MRSA selective media such as chromogenic media for MRSA that use cefoxitin as the selective agent will not detect *mecA*-negative BORSA isolates as we have observed in our own clinical laboratory. Thus these isolates could be missed from CF respiratory specimens unless full antimicrobial susceptibilities were performed on all *S. aureus* isolates. Hence, CF patients could potentially be treated with suboptimal antibiotics during a pulmonary exacerbation.

There were several limitations to this study. It is unclear whether CF patients infected with BORSA isolates behave more like CF patients infected with MSSA or CF patients infected with MRSA. Future studies assessing the clinical impact of BORSA in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CF should compare BORSA, MRSA and MSSA colonized patients in a longitudinal fashion, controlling for potential confounding factors and include more extensive PFGE comparisons to determine the evolution of BORSA. In addition, the small number of patients limited the number of variables that could be included in the logistic regression model of risk factors. This study was also not powered to detect differences in clinical outcomes between the groups. It was a single centre study and may not reflect findings in other CF populations. Finally, in a quarter of patients with *mecA*-negative BORSA, the mechanism of resistance to methicillin remains unknown.

In conclusion, CF patients with *mecA*-positive MRSA most commonly acquire their infection through person to person transmission whereas BORSA is likely preferentially selected out from endogenous MSSA in CF patients due to persistent antibiotic pressure. Prospective studies are needed to examine the prevalence, clinical impact and appropriate management of BORSA in the CF population, as well as the underlying mechanism of resistance.

1
2
3 368 **Acknowledgements**
4

5
6 369 The authors wish to acknowledge the assistance of Dr ML Mulvey at the National
7
8 370 Reference Laboratory for his assistance in performing multiplex PCR for the MRSA
9
10 371 isolates, Dana Kovach, Division of Microbiology in performing the pulsed field gel
11
12 372 electrophoresis and Derek Stephens of the Child Health Evaluative Service for his
13
14 373 assistance in the data analysis.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

References

1. Cystic. Patient Registry 2007 Annual Report. Cystic Fibrosis Foundation.

2. Dasenbrook EC, Merlo CA, Diener-West M, Lechtzin N, Boyle MP. Persistent methicillin-resistant *Staphylococcus aureus* and rate of FEV1 decline in cystic fibrosis. *Am J Respir Crit Care Med* 2008;178(8):814-821.

3. Ren CL, Morgan WJ, Konstan MW, Schechter MS, Wagener JS, Fisher KA, Regelman WE. Presence of methicillin resistant *Staphylococcus aureus* in respiratory cultures from cystic fibrosis patients is associated with lower lung function. *Pediatr Pulmonol* 2007;42(6):513-518.

4. Miall LS, McGinley NT, Brownlee KG, Conway SP. Methicillin resistant *Staphylococcus aureus* (MRSA) infection in cystic fibrosis. *Arch Dis Child* 2001;84(2):160-162.

5. Dasenbrook E, Checkley, W, Merlo, CA, Lechtzin, N, Boyle, MP. MRSA is associated with increased mortality: a 10 year cohort study of National Patient Registry. *Pediatr Pulmonol* 2009;Suppl. 32:310-311.

6. Clinical. Performance standards for antimicrobial susceptibility testing. Wayne, Pa: Clinical Laboratory Standards Institute; 2007.

7. Chambers HF. Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. *Clin Microbiol Rev* 1997;10(4):781-791.

8. Lee TW, Brownlee KG, Conway SP, Denton M, Littlewood JM. Evaluation of a new definition for chronic *Pseudomonas aeruginosa* infection in cystic fibrosis patients. *J Cyst Fibros* 2003;2(1):29-34.

9. Clinical. performance standards for antimicrobial disk susceptibility tests. Wayne, Pa: Clinical Laboratory Standards Institute; 2006.

10. Mulvey MR, MacDougall L, Cholin B, Horsman G, Fidyk M, Woods S. Community-associated methicillin-resistant *Staphylococcus aureus*, Canada. *Emerg Infect Dis* 2005;11(6):844-850.

11. Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, Vandenesch F, Etienne J. Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 1999;29(5):1128-1132.

12. Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH, Swaminathan B. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995;33(9):2233-2239.

13. Tenover F, Arbeit R, Goering R, Mickelsen P, Murray B, Persing D, Swaminathan B. Interpreting Chromosomal DNA Restriction Patterns Produced by Pulsed-Field Gel Electrophoresis: Criteria for Bacterial Strain Typing. *Journal of Clinical Microbiology* 1995;33(9):2233-2239.

14. Nadesalingam K, Conway SP, Denton M. Risk factors for acquisition of methicillin-resistant *Staphylococcus aureus* (MRSA) by patients with cystic fibrosis. *J Cyst Fibros* 2005;4(1):49-52.

15. Schlichting C, Branger C, Fournier JM, Witte W, Boutonnier A, Wolz C, Gouillet P, Doring G. Typing of *Staphylococcus aureus* by pulsed-field gel

- electrophoresis, zymotyping, capsular typing, and phage typing: resolution of clonal relationships. *J Clin Microbiol* 1993;31(2):227-232.
16. Givney R, Vickery A, Holliday A, Pegler M, Benn R. Methicillin-resistant *Staphylococcus aureus* in a cystic fibrosis unit. *J Hosp Infect* 1997;35(1):27-36.
17. Goodrich JS, Sutton-Shields TN, Kerr A, Wedd JP, Miller MB, Gilligan PH. Prevalence of community-associated methicillin-resistant *Staphylococcus aureus* in patients with cystic fibrosis. *J Clin Microbiol* 2009;47(4):1231-1233.
18. Stone A, Quittell L, Zhou J, Alba L, Bhat M, Decelie-Germana J, Rajan S, Bonitz L, Welter JJ, Dozor AJ, Gherson I, Lowy FD, Saiman L. *Staphylococcus aureus* Nasal Colonization Among Pediatric CF Patients and Their Household Contacts. *Pediatr Infect Dis J* 2009.
19. Kreiswirth B, Kornblum J, Arbeit RD, Eisner W, Maslow JN, McGeer A, Low DE, Novick RP. Evidence for a clonal origin of methicillin resistance in *Staphylococcus aureus*. *Science* 1993;259(5092):227-230.
20. Wu S, Piscitelli C, de Lencastre H, Tomasz A. Tracking the evolutionary origin of the methicillin resistance gene: cloning and sequencing of a homologue of *mecA* from a methicillin susceptible strain of *Staphylococcus sciuri*. *Microb Drug Resist* 1996;2(4):435-441.
21. Wielders CL, Vriens MR, Brisse S, de Graaf-Miltenburg LA, Troelstra A, Fleer A, Schmitz FJ, Verhoef J, Fluit AC. In-vivo transfer of *mecA* DNA to *Staphylococcus aureus* [corrected]. *Lancet* 2001;357(9269):1674-1675.
22. Deurenberg RH, Vink C, Kalenic S, Friedrich AW, Bruggeman CA, Stobberingh EE. The molecular evolution of methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect* 2007;13(3):222-235.
23. McDougal LK, Thornsberry C. The role of beta-lactamase in staphylococcal resistance to penicillinase-resistant penicillins and cephalosporins. *J Clin Microbiol* 1986;23(5):832-839.
24. Tomasz A, Drugeon HB, de Lencastre HM, Jabes D, McDougall L, Bille J. New mechanism for methicillin resistance in *Staphylococcus aureus*: clinical isolates that lack the PBP 2a gene and contain normal penicillin-binding proteins with modified penicillin-binding capacity. *Antimicrob Agents Chemother* 1989;33(11):1869-1874.
25. Tonin E, Tomasz A. Beta-lactam-specific resistant mutants of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1986;30(4):577-583.
26. Stutman HR, Lieberman JM, Nussbaum E, Marks MI. Antibiotic prophylaxis in infants and young children with cystic fibrosis: a randomized controlled trial. *J Pediatr* 2002;140(3):299-305.
27. Ratjen F, Comes G, Paul K, Posselt HG, Wagner TO, Harms K. Effect of continuous antistaphylococcal therapy on the rate of *P. aeruginosa* acquisition in patients with cystic fibrosis. *Pediatr Pulmonol* 2001;31(1):13-16.
28. Smyth A, Walters S. Prophylactic antibiotics for cystic fibrosis. *Cochrane Database Syst Rev* 2003(3):CD001912. DOI 10.1002/14651858.CD001912.
29. Chambers HF, Archer G, Matsuhashi M. Low-level methicillin resistance in strains of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1989;33(4):424-428.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

30. Massanari RM, Pfaller MA, Wakefield DS, Hammons GT, McNutt LA, Woolson RF, Helms CM. Implications of acquired oxacillin resistance in the management and control of Staphylococcus aureus infections. J Infect Dis 1988;158(4):702-709.

31. Skinner S, Murray M, Walus T, Karlowsky JA. Failure of cloxacillin in treatment of a patient with borderline oxacillin-resistant Staphylococcus aureus endocarditis. J Clin Microbiol 2009;47(3):859-861.

32. Nelson L, Cockram CS, Lui G, Lam R, Lam E, Lai R, Ip M. Community case of methicillin-resistant Staphylococcus aureus infection. Emerg Infect Dis 2006;12(1):172-174.

For Peer Review

Figure legends

Figure 1.

A. PFGE of *Sma*I restriction fragments of representative MRSA isolates (lanes 2-6).

Strains in lanes 3, 4 are indistinguishable and strains in lanes 2, 6 are closely related.

Lanes 1,7: lambda ladder.

B. Representative BORSA isolates (lanes 1-5) that are unrelated. Lane 6: lambda ladder.

For Peer Review

Epidemiology of borderline oxacillin-resistant *Staphylococcus aureus* in pediatric cystic fibrosis.

Timothy Ronan Leahy*, MB MRCPI, Institute of Molecular Medicine, Trinity College
Dublin, St James's Hospital, James's St, Dublin 8, Ireland; formerly of Division of
Infectious Diseases, Department of Pediatrics, The Hospital for Sick Children, Toronto.

Yvonne C.W. Yau, MD, FRCPC, Division of Microbiology, Department of Pediatric Laboratory Medicine, The Hospital for Sick Children, Toronto.

Eshetu Atenafu, MSc, Child Evaluative Health Sciences, Research Institute, The Hospital
for Sick Children, Toronto.

Mary Corey, PhD, Child Evaluative Health Sciences, Research Institute, The Hospital for Sick Children, Toronto.

Felix Ratjen, MD, PhD, FRCPC, Division of Respiratory Medicine, Department of Pediatrics, The Hospital for Sick Children, Toronto.

Valerie Waters, MD, MSc, FRCPC Division of Infectious Diseases, Department of Pediatrics, The Hospital for Sick Children, 555 University Avenue, Toronto, M5G 1X8, Canada.

*Corresponding author: telephone number +353 1 8961000, fax number +353 1 8963503, email address leahyt@tcd.ie

Keywords: methicillin resistant *Staphylococcus aureus* (MRSA), *mecA*, pulsed-field gel electrophoresis.

Abbreviated title. BORSA in pediatric cystic fibrosis

24 Summary

25 A single-centre retrospective study was undertaken in children with cystic fibrosis (CF)
26 to evaluate 1) risk factors for acquisition; 2) molecular epidemiology; and 3) impact on
27 disease progression of borderline oxacillin-resistant *Staphylococcus aureus* (BORSA)
28 versus *mecA*-positive methicillin-resistant *Staphylococcus aureus* (MRSA). The study
29 comprised of 1) identification of all children with at least one respiratory specimen
30 positive for either BORSA or MRSA during the study period; 2) compilation of relevant
31 clinical and epidemiological data from 12-month period leading up to first positive
32 (index) culture; 3) microbiological and molecular characterisation of index isolates and 4)
33 measurement of subsequent clinical outcome. Thirty-eight children were identified with
34 at least one positive isolate; 4 were excluded due to insufficient clinical or laboratory
35 data. Eighteen children (53%) grew BORSA in their index culture. Children who
36 acquired BORSA only (n=16) were more likely to have had prior MSSA colonization
37 (p<0.0001). Usage of oral cephalexin (p<0.01) and inhaled tobramycin (p<0.03) prior to
38 index culture was significantly and independently associated with acquisition of BORSA.
39 The majority of BORSA isolates were hyper β -lactamase producers and susceptible to a
40 greater range of antibiotics. Strain relatedness was not evident within the BORSA group.
41 There was no difference in disease progression between the two groups. This is the first
42 study to demonstrate that a significant proportion of *S. aureus* isolates with methicillin
43 resistance in the CF population are BORSAs that lack *mecA*. Antibiotic pressure may
44 lead to the development of BORSA in CF patients. Prospective studies are needed to
45 assess its clinical impact.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Methicillin-resistant *Staphylococcus aureus* (MRSA) has emerged as an important respiratory pathogen in the cystic fibrosis (CF) population. The US CF registry reported a significant rise in MRSA infection among CF patients, from 2.1% in 1996 to 21.2% in 2007¹. MRSA infection has been associated with accelerated deterioration in pulmonary function², increased hospitalization³, impaired growth, increased antibiotic usage⁴ and increased mortality in CF patients⁵.

According to the Clinical Laboratory Standards Institute (CLSI), methicillin resistance in *S. aureus* is defined as having an oxacillin minimum inhibitory concentration (MIC) > 4 µg/ml⁶. The primary mechanism of resistance for MRSA is the production of an altered penicillin binding protein (PBP2a) with reduced affinity for methicillin that is encoded by the *mecA* gene⁷. We have identified a group of CF patients infected with borderline resistant *S. aureus* (BORSA) isolates that are characterized by oxacillin MICs at or just above the susceptibility breakpoint of 4 µg/ml and lack the *mecA* gene⁷. The implications of isolating BORSA in a respiratory culture from CF patients have not previously been explored. The aim of this study was therefore to compare the risk factors for acquisition, the molecular epidemiology and subsequent clinical impact on disease progression of BORSA infection in comparison with *mecA*-positive MRSA infection in pediatric CF patients.

68 **Materials and Methods**

69

70 Study design and patient population

71 This was a single-centre retrospective comparative study of pediatric CF patients
72 whose care was primarily based at the CF clinic at the Hospital for Sick Children,
73 Toronto between 1992 and 2007. The clinic currently cares for approximately 280
74 children up to the age of eighteen. Respiratory samples (expectorated sputum or
75 oropharyngeal swabs) were sent from all children attending the clinic to the microbiology
76 laboratory for culture at every 3-month visit and during acute pulmonary exacerbations
77 (APEs). Patients with one or more respiratory specimens positive for *S. aureus* with
78 methicillin resistance (either BORSA or MRSA) were included in the analysis. Cases
79 with three or more specimens positive for BORSA or MRSA during the study period
80 were considered persistently colonized². The study was approved by the Hospital for
81 Sick Children's Research Ethics Board.

83 Data collection

84 Patients with BORSA or MRSA isolated from their respiratory specimens were
85 identified using the microbiology database, and cross-referenced with the CF clinical
86 database. Date of the first BORSA or MRSA-positive respiratory culture (index culture)
87 for each patient was identified. The medical and laboratory records for each patient were
88 examined using a standardized data collection form. Baseline patient characteristics were
89 assessed at the date of index culture, and risk factors for acquisition of BORSA or MRSA
90 infection were assessed over the twelve-month period prior to the index culture. Impact

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

91 on clinical progression was assessed by examination of the time period from index
92 culture until discharge from the clinic or until December 31st 2008, whichever date was
93 later. Data collected included information on patients’ demographics, pulmonary function
94 tests (PFTs), respiratory cultures, hospital admissions, acute pulmonary exacerbations
95 and antibiotic use. An acute pulmonary exacerbation was defined as a change in
96 pulmonary status considered by the attending physician to require antibiotic treatment,
97 either oral or intravenous. An “antibiotic day” was defined as one day for each antibiotic
98 used, for example a 14-day course of two antibiotics was defined as 28 “antibiotic days”.
99 The purpose of this distinction was to capture more complex antibiotic regimens that
100 might be employed to treat MRSA infected patients. Patients were classified as being on
101 oral cephalexin if they were prescribed cephalexin at the time of index culture of at least
102 one month’s duration (in all instances, this occurred in the absence of signs or symptoms
103 of a pulmonary exacerbation). Patients were classified as being on inhaled tobramycin if
104 they were prescribed inhaled tobramycin at the time of index culture of at least one
105 month’s duration (for previous *P. aeruginosa* infection). Patients were classified as being
106 on oral ciprofloxacin if they were prescribed at least one course of ciprofloxacin (in all
107 instances, this occurred in the presence of signs or symptoms of a pulmonary
108 exacerbation). Patients were categorized as co-colonized with other CF pathogens using
109 previously defined criteria ⁸. Only patients with at least 6 months of follow-up after the
110 index culture were included in the analysis of outcome measures.

114 Microbiology

115 **Specimen Processing**

116 Respiratory specimens from CF patients were screened for MRSA using media
117 containing 2 µg/ml of oxacillin: an in-house mannitol salt agar (MRSA agar base,
118 Acumedia, Lansing, MI, and 1% D-mannitol, Difco, Detroit, MI) supplemented with 1%
119 thymidine and 2 µg/ml of oxacillin (both from Sigma, St. Louis, MO) was used from
120 1992 to 2001, and Oxacillin Resistance Screening Agar Base (ORSAB, Oxoid, Nepean,
121 Ontario, Canada) was used from 2002 onward. Plates were incubated at 35°C aerobically
122 and observed for the presence of growth at 24 hours and at 48 hours. Suspicious colonies
123 were identified as *S. aureus* by Gram stain, catalase, slide coagulase, and tube coagulase
124 tests.

126 **Antimicrobial Susceptibility**

127 Antimicrobial susceptibility of *S. aureus* isolates was tested by a combination of
128 oxacillin 6µg/ml screen plate, disk diffusion^{6,9} and an automated system (Vitek, bio-
129 Merieux, St. Laurent, Quebec, from 1992 to Nov 2002, and the BD Automated Phoenix
130 System, BD Diagnostic Systems, Sparks, MD from Nov 2002 onward). Inducible
131 resistance to clindamycin was detected by disk diffusion using the D-zone test^{6,9}. In
132 addition, each strain was examined for the presence of PBP2a by latex agglutination
133 (PBP2' test kit, Oxoid, Hants, UK). Organisms that were negative for PBP2a by latex
134 agglutination but grew on the oxacillin screen plate were sent to the reference laboratory
135 (Central Provincial Health Laboratory, Ontario) for minimum inhibitory concentration
136 (MIC) by agar dilution and polymerase chain reaction (PCR) for the detection of the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

mecA gene. An isolate was deemed as BORSA if the oxacillin MIC was $\geq 4 \mu\text{g/ml}$, PBP2a was absent and the *mecA* PCR was negative⁷. An isolate was deemed as MRSA if the oxacillin MIC was $\geq 4\mu\text{g/ml}$, the latex agglutination for PBP2a was positive and/or the *mecA* PCR was positive⁶.

β-Lactamase Production

All BORSAs were tested for the production of β-lactamase as a potential mechanism of oxacillin resistance by disk diffusion with ampicillin (10 μg) and amoxicillin-clavulinic (20 μg/ 10 μg) disks as previously described (13). An isolate was deemed as hyper β-lactamase producing if the ampicillin zone size was resistant and the amoxicillin-clavulanic zone size was susceptible, suggesting borderline resistance was secondary to large amounts of β-lactamase⁹(13).

Molecular Characterization

The *mecA* and *nuc* gene were co-amplified with a multiplex real time PCR as described previously¹⁰. The presence of Panton-Valentine leukocidin was examined by amplification of the *lukF-PV* and *lukS-PV* genes using primer and protocols previously described¹¹. Pulsed-field gel electrophoresis (PFGE) was conducted using *SmaI* as per manufacturer protocol (Bio-rad Laboratories, Hercules, CA) on all newly identified MRSA strains. Electrophoretically-generated DNA profiles were analyzed using the criteria proposed by Tenover et al to determine strain relatedness¹².

1
2
3 160 Statistical analysis
4

5
6 161 Descriptive statistics, frequency distributions and percentages were calculated for
7
8 162 the outcome variables and other covariates of interest. Normality assumption test was
9
10 163 carried out for the continuous variables using the Kolmogorov-Smirnov test. Student's t-
11
12 164 test or Kruskal-Wallis test (if normality assumption failed) was conducted for continuous
13
14 165 variables within the two groups. The Chi-square or Fisher's exact test, as appropriate,
15
16 166 was carried out for the dichotomous variables to assess the association with the two
17
18 167 groups. To analyze independent risk factors, a multivariate analysis using a logistic
19
20 168 regression model was applied. A p-value of <0.05 was considered statistically significant.
21
22 169 The statistical programme SAS (version 9.1; SAS Institute Inc., Cary, NC) was used.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Patient population

Between 1992 and 2007, 38 patients had at least one respiratory specimen positive for either BORSA or MRSA. Of these, three were excluded from our analysis because their care was not primarily based at our hospital, thus we had no clinical data for these patients. In addition, *mecA* PCR could not be performed on one isolate retrospectively therefore 34 patients were included in the study. Of these 34 patients, two patients were initially colonized with *mecA*- negative and subsequently *mecA*-positive strains and were excluded from the final statistical analysis as it was unclear how to categorize them in terms of patient characteristics and outcomes. Ultimately 16 patients with *mecA* positive MRSA and 16 with *mecA* negative BORSA on the index culture were compared for baseline characteristics. For the longitudinal assessment, 15 patients in each group were compared since one patient in each group did not have follow-up data for at least 6 months.

Patient characteristics and risk factors for BORSA and MRSA

The two groups were similar in terms of baseline characteristics (Table 1). There was no difference in pulmonary function or rate of decline of pulmonary function in the year prior to the index culture. In addition, there was no difference in the number of acute pulmonary exacerbations, hospital admissions or overall antibiotic use for the treatment of acute pulmonary exacerbations. A significantly greater proportion of BORSA infected patients were colonized with MSSA in the year prior (94% vs 25%, $p<0.0001$). Likewise,

194 a higher proportion of BORSA infected patients received oral antibiotics (cephalexin)
195 (75% vs 25%, $p=0.005$).
196 Mean duration of therapy among those patients on cephalexin was 337 days (range 30-
197 365 days) in the BORSA group and 311 days (range 150-365 days) in the MRSA group;
198 cephalexin treatment was instituted not for pulmonary exacerbation but as treatment for
199 MSSA identified on routine sputum cultures. Furthermore, a significantly greater
200 number of BORSA infected patients had received a course of oral ciprofloxacin (31% vs
201 0%, $p=0.043$) or were on inhaled antibiotics (tobramycin) (88% vs 44%, $p=0.009$) in the
202 year prior to the index culture than patients with MRSA. No inter-group difference was
203 detected in any of the other co-colonizing respiratory pathogens. Patients with BORSA
204 strains had significantly more clinic visits in the year prior to first isolate ($n=7.94$) than
205 patients with *mecA*-positive strains ($n=4.69$, $p=0.049$) although this was not a significant
206 risk factor in the logistic regression model

207 A logistic regression model was applied to determine the independence of
208 epidemiological and clinical variables that were correlated with BORSA infection.
209 Colonization with MSSA and use of oral cephalexin were significantly associated as 68%
210 of patients with MSSA were treated with cephalexin and 81% of patients treated with
211 cephalexin were colonized with MSSA ($p<0.05$). The use of oral cephalexin (regression
212 coefficient (β)=1.15, SD=0.47, p value=0.015) and inhaled tobramycin (regression
213 coefficient (β)=1.16, SD=0.53, p value=0.028) in the year prior to index culture were
214 significantly and independently associated with BORSA infection.

215

216 Microbiological characteristics of BORSA and MRSA isolates

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

217 The microbiological characteristics of the BORSA isolates are illustrated in Table
218 2. Fourteen of the 16 isolates had oxacillin MICs at or just above the breakpoint (4 to 8
219 μg/ml) but 2 isolates had MICs of 16 μg/ml. All of the BORSA isolates were negative for
220 PBP2a by latex agglutination and were *mecA* negative by PCR. Twelve of the 16 BORSA
221 isolates were hyper β-lactamase producers. This was shown by resistance to ampicillin by
222 disk diffusion that was reverted to susceptible with the addition of clavulanic acid, a β-
223 lactamase inhibitor. The remaining 4 isolates had ampicillin zone size that were
224 unchanged by addition of clavulin by disk diffusion.

225 In comparison to the MRSA isolates, a significantly greater proportion of the
226 BORSA isolates were susceptible to both erythromycin (81% vs 13%, $p < 0.0001$) and
227 clindamycin (94% vs 31%, $p < 0.001$). None of the BORSA isolates were D-test positive
228 but 5 (45%) of the MRSA isolates were D-test positive ($p = 0.043$). All of the BORSA
229 and MRSA isolates were susceptible to trimethoprim-sulphamethoxazole. None of the
230 BORSA isolates tested produced PVL. Three of the 16 MRSA isolates were identified as
231 USA300 epitypes, and produced PVL.

232
233 Relatedness of BORSA and MRSA isolates

234 Pulsed field gel electrophoresis (PFGE) revealed that none of the BORSA strains
235 were related by the Tenover criteria¹³ (Figure 1). Within the MRSA group, we found
236 four separate instances of strain relatedness during the course of the study: a pair of
237 siblings shared an indistinguishable strain, four patients shared possibly related strains,
238 two patients shared a closely related strain with a third patient sharing a possibly related

239 strain and, finally, three patients (including a pair of siblings) shared an indistinguishable
240 strain.

241

242 Clinical outcomes of patients

243 To compare the clinical impact of BORSA versus MRSA infection on disease
244 progression, we assessed 99.7 patient-years of follow-up after the index culture with a
245 comparable mean duration of follow up of 3.37 years in the MRSA group and 3.27 years
246 in the BORSA group ($p=0.507$) (Table 3). Patients infected with MRSA were more
247 likely to have persistent infection at the end of follow up than patients with BORSA
248 infection (80% vs 53%) but this was not statistically significant ($p=0.121$). The annual
249 decline in percent predicted forced expiratory volume in 1 second (FEV_1), was greater
250 among the MRSA infected patients (5.8%) than those infected with BORSA (2.8%)
251 although this difference was not significant ($p = 0.77$). Neither was there a significant
252 difference in the number of acute pulmonary exacerbations, hospital admissions, inpatient
253 days or clinic visits. One patient in the MRSA group died, and another progressed to lung
254 transplant. Three patients in the BORSA group had a lung transplant during the study
255 period.

256

Discussion

This is the first study to show that a significant proportion (50%) of *S. aureus* isolates with methicillin resistance infecting CF patients are BORSA which lack the *mecA* gene, the main genetic determinant of methicillin resistance in *S. aureus*⁷. The present study demonstrates that antibiotic use in CF patients colonized with MSSA is a risk factor specifically for infection with BORSA. Previous studies have shown that ciprofloxacin and cephalosporin use is a risk factor for acquisition of MRSA in CF patients¹⁴ but, to our knowledge, no other study has examined the differential risk factors for BORSA compared to MRSA acquisition.

Transmission of traditional MRSA between patients^{15,16}, within the community¹⁷ and even within households¹⁸ has been well described in the CF population. Evidence of person to person transmission of *mecA*-positive MRSA was also present in this study with shared strains among CF patients and siblings. The mechanism of resistance is relevant to how CF patients acquire MRSA as the *mecA* gene is absent in its entirety from MSSA^{19 7}. It is believed to have originated from another staphylococcus species, such as *Staphylococcus sciuri*, and been transposed into MSSA²⁰. Although horizontal transfer of *mecA* between a *S. epidermidis* and MSSA isolate within the same patient has been described²¹, transmission of *mecA*-positive MRSA isolates between individuals is likely more common as there are a well defined number of *mecA*-positive MRSA clones worldwide²².

Our data would suggest that CF patients may acquire *mecA*-negative BORSA isolates in a different way. In contrast to MRSA isolates, BORSA isolates had unique

1
2
3 280 PFGE patterns and appeared not to be shared between patients in our study. CF patients
4
5 281 colonized with MSSA and exposed to oral cephalexin or inhaled tobramycin (which can
6
7 282 act synergistically with β -lactam antibiotics against *S. aureus*) were at significantly
8
9 283 increased risk of acquiring BORSA infection. This suggests that BORSA isolates may
10
11 284 arise from endogenous MSSA isolates that develop other mechanisms of resistance to
12
13 285 methicillin in response to antibiotic pressure. Previous studies have shown that *S. aureus*
14
15 286 strains with borderline susceptibility to oxacillin have a significant drop in the penicillin
16
17 287 MICs to the susceptible range with the addition of a β -lactamase inhibitor²³. This
18
19 288 predicts the response of oxacillin MICs to the addition of a β -lactamase inhibitor and
20
21 289 suggests that a large amount of β -lactamase production is the mechanism of resistance to
22
23 290 oxacillin in the majority of BORSA, as observed in 12 of 16 isolates in our study.
24
25 291 However, in the 4 remaining BORSA strains where the ampicillin zone size did not
26
27 292 change significantly with the addition of clavulanic acid, the mechanism of methicillin
28
29 293 resistance is unknown. In the aforementioned study, the authors described *S. aureus*
30
31 294 strains with intermediate MICs to oxacillin that remained the same in the presence of
32
33 295 clavulanic acid. The authors suggested that oxacillin resistance was due to factors other
34
35 296 than β -lactamase inactivation²³. It is possible that it is due to a modification of “normal”
36
37 297 penicillin-binding proteins (PBPs) such as PBP 1, PBP 2 and PBP 4 as was previously
38
39 298 shown in clinical BORSA isolates by Tomasz et al²⁴. Exposure of susceptible
40
41 299 staphylococci to increasing concentrations of β -lactam antibiotics in the laboratory is
42
43 300 known to cause PBP alterations associated with increased MICs²⁵. In this study,
44
45 301 persistent antibiotic pressure may have similarly modified PBPs in endogenous MSSA
46
47 302 isolates colonizing CF patients and resulting in the development of *mecA*-negative, non-
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

hyper β lactamase producing BORSA isolates. Thus, in the prevention of the emergence of BORSA in CF patients, other infection control measures, such as patient segregation, may not be as important as the judicious use of antibiotics.

In our study, the majority of patients treated with oral cephalexin were colonized with MSSA. Although there were no signs and symptoms of a pulmonary exacerbation at the time, we are limited in a retrospective study to the information that is recorded in the medical charts. Their antibiotic treatment was thus likely driven by their MSSA positive respiratory cultures and/or more subtle symptoms. Primary prophylaxis for *S. aureus* (the treatment of all newly diagnosed CF patients with antistaphylococcal agents irrespective of respiratory culture results) is not routinely practiced in North America due to concerns of *P. aeruginosa* acquisition^{26,27}. A systematic review of prophylactic antibiotics in cystic fibrosis, however, suggests that *P. aeruginosa* acquisition in this context is associated less frequently with narrow spectrum anti-staphylococcal antibiotics such as flucloxacillin compared with broader spectrum anti-staphylococcal antibiotics such as cephalexin²⁸. In our study, although patients treated with oral cephalexin were not receiving prophylaxis against *S. aureus*, they were receiving prolonged courses of cephalexin for MSSA detected on routine culture of the respiratory tract. This prolonged cephalexin use was identified as an independent risk factor for the acquisition of BORSA isolates and occurred recently, primarily in the last 5 years of the study.

The clinical significance of BORSA infection in CF patients is unknown. Our study lacked a sufficient sample size and is not powered to clearly delineate the clinical impact of BORSA infection. Previous studies have reported accelerated deterioration in pulmonary function², increased hospitalization³, impaired growth, increased antibiotic

usage⁴ and increased mortality⁵ in CF patients infected with MRSA. However, these studies are limited by a lack of information on the laboratory methodology of identifying MRSA and thus it is not known to what degree these effects are attributable to BORSA or MRSA.

Detection of BORSA isolates may also influence the choice of antibiotics in the treatment of these infections in the setting of a pulmonary exacerbation. In our study, BORSA isolates were more susceptible to erythromycin and clindamycin than the MRSA isolates. This finding is not surprising as the *mecA* gene is located on a mobile genetic element (the Staphylococcal Cassette Chromosome *mec*: SCC*mec*) which can contain additional drug resistance genes²². Although earlier studies suggested that infections due to *mecA* negative BORSA isolates could be treated with β -lactam antibiotics^{29,30}, more recent cases report failure of cloxacillin and even vancomycin in the treatment of endocarditis caused by BORSA, depending on the mechanism of methicillin resistance^{31,32}. This has implications for how respiratory specimens from CF patients should be screened for MRSA. Some of the newer MRSA selective media such as chromogenic media for MRSA that use cefoxitin as the selective agent will not detect *mecA*-negative BORSA isolates as we have observed in our own clinical laboratory. Thus these isolates could be missed from CF respiratory specimens unless full antimicrobial susceptibilities were performed on all *S. aureus* isolates. Hence, CF patients could potentially be treated with suboptimal antibiotics during a pulmonary exacerbation.

There were several limitations to this study. It is unclear whether CF patients infected with BORSA isolates behave more like CF patients infected with MSSA or CF patients infected with MRSA. Future studies assessing the clinical impact of BORSA in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

349 CF should compare BORSA, MRSA and MSSA colonized patients in a longitudinal
350 fashion, controlling for potential confounding factors and include more extensive PFGE
351 comparisons to determine the evolution of BORSA. In addition, the small number of
352 patients limited the number of variables that could be included in the logistic regression
353 model of risk factors. This study was also not powered to detect differences in clinical
354 outcomes between the groups. It was a single centre study and may not reflect findings in
355 other CF populations. Finally, in a quarter of patients with *mecA*-negative BORSA, the
356 mechanism of resistance to methicillin remains unknown.

357 In conclusion, CF patients with *mecA*-positive MRSA most commonly acquire
358 their infection through person to person transmission whereas BORSA is likely
359 preferentially selected out from endogenous MSSA in CF patients due to persistent
360 antibiotic pressure. Prospective studies are needed to examine the prevalence, clinical
361 impact and appropriate management of BORSA in the CF population, as well as the
362 underlying mechanism of resistance.

363
364
365
366
367

1
2
3 368 **Acknowledgements**
4

5
6 369 The authors wish to acknowledge the assistance of Dr ML Mulvey at the National
7
8 370 Reference Laboratory for his assistance in performing multiplex PCR for the MRSA
9
10 371 isolates, Dana Kovach, Division of Microbiology in performing the pulsed field gel
11
12 372 electrophoresis and Derek Stephens of the Child Health Evaluative Service for his
13
14 373 assistance in the data analysis.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

References

1. Cystic. Patient Registry 2007 Annual Report. Cystic Fibrosis Foundation.
2. Dasenbrook EC, Merlo CA, Diener-West M, Lechtzin N, Boyle MP. Persistent methicillin-resistant *Staphylococcus aureus* and rate of FEV1 decline in cystic fibrosis. *Am J Respir Crit Care Med* 2008;178(8):814-821.
3. Ren CL, Morgan WJ, Konstan MW, Schechter MS, Wagener JS, Fisher KA, Regelman WE. Presence of methicillin resistant *Staphylococcus aureus* in respiratory cultures from cystic fibrosis patients is associated with lower lung function. *Pediatr Pulmonol* 2007;42(6):513-518.
4. Miall LS, McGinley NT, Brownlee KG, Conway SP. Methicillin resistant *Staphylococcus aureus* (MRSA) infection in cystic fibrosis. *Arch Dis Child* 2001;84(2):160-162.
5. Dasenbrook E, Checkley, W, Merlo, CA, Lechtzin, N, Boyle, MP. MRSA is associated with increased mortality: a 10 year cohort study of National Patient Registry. *Pediatr Pulmonol* 2009;Suppl. 32:310-311.
6. Clinical. Performance standards for antimicrobial susceptibility testing. Wayne, Pa: Clinical Laboratory Standards Institute; 2007.
7. Chambers HF. Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. *Clin Microbiol Rev* 1997;10(4):781-791.
8. Lee TW, Brownlee KG, Conway SP, Denton M, Littlewood JM. Evaluation of a new definition for chronic *Pseudomonas aeruginosa* infection in cystic fibrosis patients. *J Cyst Fibros* 2003;2(1):29-34.
9. Clinical. performance standards for antimicrobial disk susceptibility tests. Wayne, Pa: Clinical Laboratory Standards Institute; 2006.
10. Mulvey MR, MacDougall L, Cholin B, Horsman G, Fidyk M, Woods S. Community-associated methicillin-resistant *Staphylococcus aureus*, Canada. *Emerg Infect Dis* 2005;11(6):844-850.
11. Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, Vandenesch F, Etienne J. Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 1999;29(5):1128-1132.
12. Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH, Swaminathan B. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995;33(9):2233-2239.
13. Tenover F, Arbeit R, Goering R, Mickelsen P, Murray B, Persing D, Swaminathan B. Interpreting Chromosomal DNA Restriction Patterns Produced by Pulsed-Field Gel Electrophoresis: Criteria for Bacterial Strain Typing. *Journal of Clinical Microbiology* 1995;33(9):2233-2239.
14. Nadesalingam K, Conway SP, Denton M. Risk factors for acquisition of methicillin-resistant *Staphylococcus aureus* (MRSA) by patients with cystic fibrosis. *J Cyst Fibros* 2005;4(1):49-52.
15. Schlichting C, Branger C, Fournier JM, Witte W, Boutonnier A, Wolz C, Gouillet P, Doring G. Typing of *Staphylococcus aureus* by pulsed-field gel

- electrophoresis, zymotyping, capsular typing, and phage typing: resolution of clonal relationships. *J Clin Microbiol* 1993;31(2):227-232.
16. Givney R, Vickery A, Holliday A, Pegler M, Benn R. Methicillin-resistant *Staphylococcus aureus* in a cystic fibrosis unit. *J Hosp Infect* 1997;35(1):27-36.
17. Goodrich JS, Sutton-Shields TN, Kerr A, Wedd JP, Miller MB, Gilligan PH. Prevalence of community-associated methicillin-resistant *Staphylococcus aureus* in patients with cystic fibrosis. *J Clin Microbiol* 2009;47(4):1231-1233.
18. Stone A, Quittell L, Zhou J, Alba L, Bhat M, Decelie-Germana J, Rajan S, Bonitz L, Welter JJ, Dozor AJ, Gherson I, Lowy FD, Saiman L. *Staphylococcus aureus* Nasal Colonization Among Pediatric CF Patients and Their Household Contacts. *Pediatr Infect Dis J* 2009.
19. Kreiswirth B, Kornblum J, Arbeit RD, Eisner W, Maslow JN, McGeer A, Low DE, Novick RP. Evidence for a clonal origin of methicillin resistance in *Staphylococcus aureus*. *Science* 1993;259(5092):227-230.
20. Wu S, Piscitelli C, de Lencastre H, Tomasz A. Tracking the evolutionary origin of the methicillin resistance gene: cloning and sequencing of a homologue of *mecA* from a methicillin susceptible strain of *Staphylococcus sciuri*. *Microb Drug Resist* 1996;2(4):435-441.
21. Wielders CL, Vriens MR, Brisse S, de Graaf-Miltenburg LA, Troelstra A, Fleer A, Schmitz FJ, Verhoef J, Fluit AC. In-vivo transfer of *mecA* DNA to *Staphylococcus aureus* [corrected]. *Lancet* 2001;357(9269):1674-1675.
22. Deurenberg RH, Vink C, Kalenic S, Friedrich AW, Bruggeman CA, Stobberingh EE. The molecular evolution of methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect* 2007;13(3):222-235.
23. McDougal LK, Thornsberry C. The role of beta-lactamase in staphylococcal resistance to penicillinase-resistant penicillins and cephalosporins. *J Clin Microbiol* 1986;23(5):832-839.
24. Tomasz A, Drugeon HB, de Lencastre HM, Jabes D, McDougall L, Bille J. New mechanism for methicillin resistance in *Staphylococcus aureus*: clinical isolates that lack the PBP 2a gene and contain normal penicillin-binding proteins with modified penicillin-binding capacity. *Antimicrob Agents Chemother* 1989;33(11):1869-1874.
25. Tonin E, Tomasz A. Beta-lactam-specific resistant mutants of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1986;30(4):577-583.
26. Stutman HR, Lieberman JM, Nussbaum E, Marks MI. Antibiotic prophylaxis in infants and young children with cystic fibrosis: a randomized controlled trial. *J Pediatr* 2002;140(3):299-305.
27. Ratjen F, Comes G, Paul K, Posselt HG, Wagner TO, Harms K. Effect of continuous antistaphylococcal therapy on the rate of *P. aeruginosa* acquisition in patients with cystic fibrosis. *Pediatr Pulmonol* 2001;31(1):13-16.
28. Smyth A, Walters S. Prophylactic antibiotics for cystic fibrosis. *Cochrane Database Syst Rev* 2003(3):CD001912. DOI 10.1002/14651858.CD001912.
29. Chambers HF, Archer G, Matsuhashi M. Low-level methicillin resistance in strains of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1989;33(4):424-428.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

30. Massanari RM, Pfaller MA, Wakefield DS, Hammons GT, McNutt LA, Woolson RF, Helms CM. Implications of acquired oxacillin resistance in the management and control of Staphylococcus aureus infections. J Infect Dis 1988;158(4):702-709.

31. Skinner S, Murray M, Walus T, Karlowsky JA. Failure of cloxacillin in treatment of a patient with borderline oxacillin-resistant Staphylococcus aureus endocarditis. J Clin Microbiol 2009;47(3):859-861.

32. Nelson L, Cockram CS, Lui G, Lam R, Lam E, Lai R, Ip M. Community case of methicillin-resistant Staphylococcus aureus infection. Emerg Infect Dis 2006;12(1):172-174.

For Peer Review

Figure legends

Figure 1.

A. PFGE of *Sma*I restriction fragments of representative MRSA isolates (lanes 2-6).

Strains in lanes 3, 4 are indistinguishable and strains in lanes 2, 6 are closely related.

Lanes 1,7: lambda ladder.

B. Representative BORSA isolates (lanes 1-5) that are unrelated. Lane 6: lambda ladder.

For Peer Review

Table 1. Characteristics of patients with MRSA and BORSA

	MRSA (n=16)	BORSA (n=16)	P-value
Patient characteristics at index culture			
Mean age (range) in yr	10.4 (1.8-17.5)	13.1 (8.5-17.3)	0.078
No. (%) male	10 (63)	8 (50)	0.476
Mean BMI z-score	-0.18	-0.40	>0.5
No. (%) homozygous DF508	7 (44)	7 (44)	>0.5
No. (%) pancreatic insufficient	15 (94)	14 (88)	>0.5
No. (%) CF related diabetes	1 (6)	2 (13)	>0.5
Mean % predicted FEV ₁ (range)	68.5 (41-99)	69.4 (30-113)	>0.5
Respiratory flora in 12-month period prior to index culture			
No. (%) MSSA colonized	4 (25)	15 (94)	<0.0001
No. (%) <i>P. aeruginosa</i> colonized	3 (19)	6 (38)	0.433
No. (%) <i>A. fumigates</i> colonized	1 (6)	5 (31)	0.171
No. (%) <i>H. influenza</i> colonized	1 (6)	1 (6)	>0.5
No. (%) <i>B. cepacia</i> colonized	1 (6)	0 (0)	>0.5
No. (%) <i>S. maltophilia</i> colonized	1 (6)	1 (6)	>0.5
Antibiotic usage in 12-month period prior to index culture			
No. (%) of patients on oral cephalexin	4 (25)	12 (75)	0.005
No. (%) of patients on inhaled tobramycin	7 (44)	14 (88)	0.009
No. (%) of patients on oral ciprofloxacin	0(0)	5(31)	0.043
Disease activity during 12-month period prior to index culture			
Mean change (range) in % predicted FEV ₁	-2.5 (-11.6 to +10.7)	-3.8 (-28.9 to +10.2)	>0.5
Mean no. (range) of exacerbations	0.8 (0-4)	1.4 (0-8)	>0.5
Mean no. (range) of hospital admissions	0.4 (0-4)	0.7 (0-6)	>0.5
Mean no. (range) of inpatient days	3.4 (0-37)	7.6 (0-61)	>0.5
Mean no. (range) of IV antibiotic days	6.1 (0-70)	14.2 (0-130)	>0.5
Mean no. (range) of clinic visits	4.7 (3-10)	7.9 (3-35)	0.049

Abbreviations: BMI – body mass index, MSSA- methicillin sensitive *Staphylococcus aureus*, FEV₁-
Forced expiratory volume in 1 second.

Table 1. Characteristics of patients with MRSA and BORSA

	MRSA (n=16)	BORSA (n=16)	P-value
Patient characteristics at index culture			
Mean age (range) in yr	10.4 (1.8-17.5)	13.1 (8.5-17.3)	0.078
No. (%) male	10 (63)	8 (50)	0.476
Mean BMI z-score	-0.18	-0.40	>0.5
No. (%) homozygous DF508	7 (44)	7 (44)	>0.5
No. (%) pancreatic insufficient	15 (94)	14 (88)	>0.5
No. (%) CF related diabetes	1 (6)	2 (13)	>0.5
Mean % predicted FEV ₁ (range)	68.5 (41-99)	69.4 (30-113)	>0.5
Respiratory flora in 12-month period prior to index culture			
No. (%) MSSA colonized	4 (25)	15 (94)	<0.0001
No. (%) <i>P. aeruginosa</i> colonized	3 (19)	6 (38)	0.433
No. (%) <i>A. fumigates</i> colonized	1 (6)	5 (31)	0.171
No. (%) <i>H. influenza</i> colonized	1 (6)	1 (6)	>0.5
No. (%) <i>B. cepacia</i> colonized	1 (6)	0 (0)	>0.5
No. (%) <i>S. maltophilia</i> colonized	1 (6)	1 (6)	>0.5
Antibiotic usage in 12-month period prior to index culture			
No. (%) of patients on oral cephalexin	4 (25)	12 (75)	0.005
No. (%) of patients on inhaled tobramycin	7 (44)	14 (88)	0.009
No. (%) of patients on oral ciprofloxacin	0(0)	5(31)	0.043
Disease activity during 12-month period prior to index culture			
Mean change (range) in % predicted FEV ₁	-2.5 (-11.6 to +10.7)	-3.8 (-28.9 to +10.2)	>0.5
Mean no. (range) of exacerbations	0.8 (0-4)	1.4 (0-8)	>0.5
Mean no. (range) of hospital admissions	0.4 (0-4)	0.7 (0-6)	>0.5
Mean no. (range) of inpatient days	3.4 (0-37)	7.6 (0-61)	>0.5
Mean no. (range) of IV antibiotic days	6.1 (0-70)	14.2 (0-130)	>0.5
Mean no. (range) of clinic visits	4.7 (3-10)	7.9 (3-35)	0.049

Abbreviations: BMI – body mass index, MSSA- methicillin sensitive *Staphylococcus aureus*, FEV₁-

Forced expiratory volume in 1 second.

Table 2. Microbiological characteristics of BORSA isolates

Isolate	Oxacillin MIC (µg/ml)	PBP2a latex agglutination	<i>mecA</i> PCR	AMP/AMC zone size (mm)†	Hyper β- lactamase production	ERY	CLI	SXT
SA 002	8	neg	neg	13/23	Yes	S	S	S
SA 003	16	neg	neg	12/26	Yes	I	S*	S
SA 004	4	neg	neg	38/40	No	S	S	S
SA 006	8	neg	neg	35/36	No	S	S	S
SA 011	8	neg	neg	18/29	Yes	S	S	S
SA 016	4	neg	neg	15/26	Yes	S	S	S
SA 018	8	neg	neg	14/28	Yes	S	S	S
SA 020	4	neg	neg	17/28	Yes	S	S	S
SA 021	4	neg	neg	16/26	Yes	S	S	S
SA 022	8	neg	neg	30/29	No	S	S	S
SA 023	8	neg	neg	16/21	Yes	R	S*	S
SA 029	16	neg	neg	10/27	Yes	S	S	S
SA 034	8	neg	neg	12/21	Yes	S	S	S
SA 035	4	neg	neg	29/31	No	S	S	S
SA 036	4	neg	neg	16/23	Yes	S	S	S
SA 038	8	neg	neg	11/24	Yes	R	S*	S

†Ampicillin (AMP) susceptible zone size ≥ 29 mm, resistant ≤ 28 mm
Amoxicillin clavulanic acid (AMC) susceptible zone size ≥ 20 mm, resistant ≤ 19 mm⁹
* D-test negative , AMP-ampicillin, AMC-amoxicillin clavulanic acid
ERY-erythromycin, CLI-clindamycin, SXT-trimethoprim-sulphamethoxazole
S-susceptible I-intermediate R-resistant

Table 3. Clinical outcomes of patients with MRSA and BORSA

	MRSA	BORSA	P-
Clinical outcomes	(n = 15)	(n = 15)	value
Mean duration (range) of follow up in years	3.37 (0.76-14.8)	3.27 (0.66-5.57)	0.507
Mean annual change (range) in % predicted FEV1	-5.8 (-33.1 to +3.2)	-2.8 (-10.2 to +7.7)	0.770
Mean no. (range) of exacerbations per year	1.8 (0-5.3)	2.4 (0.3-7.6)	0.443
Mean no. (range) of hospital admissions per year	0.9 (0-6.6)	1.3 (0-4.7)	0.242
Mean no. (range) of inpatient days per year	11.5 (0-109.2)	16.1 (0-91.7)	0.226
Mean no. (range) of IV antibiotic days per year	26.1 (0-224.1)	33.3 (0-200.6)	0.361
Mean no. (range) of clinic visits per year	7.7 (3.7-37.4)	11.7 (2.8-80.9)	0.272
No. (%) of patients with persistent infection	12 (80)	8 (53.3)	0.121

83x176mm (150 x 150 DPI)