Local adaptation and the evolution of species ranges under climate change

K.E. Atkins, J.M.J. Travis

► To cite this version:

K.E. Atkins, J.M.J. Travis. Local adaptation and the evolution of species ranges under climate change. Journal of Theoretical Biology, 2010, 266 (3), pp.449. 10.1016/j.jtbi.2010.07.014 . hal-00616258

HAL Id: hal-00616258

<https://hal.science/hal-00616258>

Submitted on 21 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript

Local adaptation and the evolution of species ranges
under climate change

K.E. Atkins, J.M.J. Travis

PII: S0022-5193(10)00363-2
DOI: doi:10.1016/j.jtbi.2010.07.014
Reference: YJTBI6076

To appear in: *Journal of Theoretical Biology*

Received date: 5 August 2009
Revised date: 13 July 2010
Accepted date: 14 July 2010

Cite this article as: K.E. Atkins and J.M.J. Travis, Local adaptation and the evolution of species ranges under climate change, *Journal of Theoretical Biology*, doi:[10.1016/j.jtbi.2010.07.014](https://doi.org/10.1016/j.jtbi.2010.07.014)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

www.elsevier.com/locate/jtbi

LOCAL ADAPTATION AND THE EVOLUTION OF SPECIES RANGES UNDER CLIMATE CHANGE

Atkins, K.E. ^{*†‡} Travis J.M.J. [§]

Corresponding Author: katherine.atkins@yale.edu

+1 (203) 737 4874 (Tel)

*Present Address: Department of Epidemiology and Public Health, Yale University School of Medicine, New Haven, CT 06520

[†]Centre for Infectious Diseases, University Of Edinburgh, West Mains Road, EH9 3JT, UK

[‡]Depts. of Biology and Mathematics, University of York, Heslington, York, YO10 5DD, UK

[§]Institute of Biological Sciences, Zoology Building, Tillydrone Avenue, University of Aberdeen, Aberdeen, AB24 2TZ, UK

Abstract

The potential impact of climate change on biodiversity is well documented. A well developed range of statistical methods currently exists that projects the possible future habitat of a species directly from the current climate and a species distribution. However, studies incorporating ecological and evolutionary processes remain limited. Here, we focus on the potential role that local adaptation to climate may play in driving the range dynamics of sessile organisms. Incorporating environmental adaptation into a stochastic simulation yields several new insights. Counter-intuitively, our simulation results suggest that species with broader ranges are not necessarily more robust to climate change. Instead, species with broader ranges can be more susceptible to extinction as locally adapted genotypes are often blocked from range shifting by the presence of cooler adapted genotypes that persist even when their optimum climate has left them behind. Interestingly, our results also suggest that it will not always be the cold-adapted phenotypes that drive polewards range expansion. Instead, range shifts may be driven by phenotypes conferring adaptation to conditions prevalent towards the centre of a species' equilibrium distribution. This may have important consequences for the conservation method termed predictive provenancing. These initial results highlight the potential importance of local adaptation in determining how species will respond to climate change and we argue that this is an area requiring urgent theoretical and empirical attention.

Keywords: local density dependence; environmental gradient; environmental niche; extinction, species' range.

Introduction

The threat that climate change poses to biodiversity is well documented (Parmesan and Yohe, 2003; Pearson et al., 2002; Peterson et al., 2002; Thomas et al., 2004; Walther et al., 2002). Many species have already exhibited responses to the changing environment; in a meta-analysis Parmesan and Yohe (2003) found that 434 species out of 893 had shifted in distribution and/or abundance. A key question is whether climate change increases the risk that species will become extinct, and initial results from a study by Thomas et al. (2004) are alarming. Even for a best-case scenario, using the minimum expected climate change and ignoring the potential for dispersal limitation, the authors predict a 9-13% loss in species abundance. Using the worst climate scenario and with extreme dispersal limitation this figure rises to a staggering 38-52%. Whilst this study represents a useful exercise, not least in terms of focussing minds on the problem at hand, it simultaneously highlights some important deficiencies that it shares with the many bioclimate envelope models (BEMs), widely used to predict future ranges (Hampe, 2004). There is an urgent need for the development of a new generation of models to make this type of prediction. However, this goal is hindered by a basic lack of understanding of the likely consequences of key ecological and evolutionary processes for range dynamics under climate change. Local adaptation is a good example of one of these processes: many species comprise collections of populations each adapted to local climatic conditions. However, there has been little formal consideration of the impact the degree of local adaptation present within a species might have on range dynamics (Polechová et al., 2009), and no attempt to

incorporate these processes into predictive modelling. It is likely that two aspects of
 50 a species' adaptation will be important. First, the steepness with which an individual's
 fitness declines as it is moved from its optimal climate and second, the total
 52 range of environmental conditions under which a species, as a whole, can survive.
 In this contribution, we take a first step and develop a relatively simple simulation
 54 model to explore the issue of how local adaptation influences a species ability to track
 changing climate.

A plethora of studies have used bioclimate envelope modelling to predict the
 future biogeographic ranges of species (Bakkenes et al., 2002; Broennimann et al.,
 58 2006; Peterson et al., 2002; Schwarz et al., 2006; Thuiller et al., 2005). Refined
 methods are being developed that entail the use of increasingly sophisticated statistics
 60 (e.g. Gavin and Hu (2005)), genetic algorithms (e.g. Tarmansen et al. (2006)) and
 neural networks (e.g. Pearson and Dawson (2003)), and it has even been argued
 62 that the optimal approach should use a combination of these approaches in order
 to look for concordance between them (Araujo et al., 2005). Intense activity in this
 64 field has provided a suite of methods that obtain a relationship between the climate
 and a species range prior to environmental change: thus defining the 'climate space'
 66 occupied by the species. Utilising future climate scenarios, these methods predict the
 future climate space for the species. However, there is a growing realisation that this
 68 correlational approach alone is ill-equipped to predict future species ranges (Davis
 et al., 2005; Hampe, 2004; Lawton, 2000; Woodward and Beerling, 1997).

70 The dynamics of a species' range during a period of climate change will be dependent
 upon the existence and interplay of various ecological and evolutionary pro-

cesses. Dispersal is perhaps the most obvious of these and it is anticipated that species with greater dispersal ability will be better able to track a changing climate (Travis and Dytham, 2002; Travis et al., 2009). Midgley et al. (2002) incorporate simple dispersal functions into a model predicting future ranges of Cape Proteaceae. The description of the spatial population dynamics used in this model is relatively crude but, in linking a model of ecological dynamics with a BEM, Midgley et al. (2002) take an important step in the right direction.

Theoretical studies, often focussing on invasion biology rather than range dynamics under climate change, have considered a range of evolutionary dynamics during expansions: Garcia-Ramos and Rodriguez (2002) demonstrate the interacting roles played by local adaptation and habitat heterogeneity in determining spread rate; several theoretical studies have now demonstrated that range expansion might be accelerated by the evolution of increased dispersal propensity at expanding fronts (e.g. Travis and Dytham, 2002; Burton et al., 2010) and recent empirical evidence confirms that this is an important effect (Phillips et al., 2006); Klopstein et al. (2006) demonstrate that neutral mutations arising on the edge of a range expansion sometimes ‘surf’ on the wave of advance and can thus reach a wider spatial distribution and higher frequency than would be expected in a stationary population. Klopstein et al. (2006) suggest that this surfing phenomenon may increase the rate of evolution of spatially expanding populations. More recent extensions to this work have demonstrated that this surfing dynamic can be important for non-neutral mutations. Even deleterious mutations can sometimes attain high abundance at an expanding front (Burton and Travis, 2008a; Excoffier and Ray, 2008; Travis et al., 2007) and this effect

can substantially modify evolutionary dynamics where fitness landscapes are rugged
(Burton and Travis, 2008b). In this paper we concentrate on local adaptation, an
important evolutionary process that, to date, has rather surprisingly received little
attention in terms of its potential role in range expansions and determining a species
dynamics during an episode of climate change.

Considerable empirical evidence indicates that locally adapted phenotypes are
commonplace within many species (Lambrechts et al., 1996; McNeilly and Antonovic,
1968; Riihimäki et al., 2005; Santamaria et al., 2003). For example, it has long
been known that many plant species show genetic differentiation in the timing of
flowering (McNeilly and Antonovic (1968); this variation is frequently clinal with
northern populations consistently different from southern counterparts (Riihimäki
et al., 2005). Occurrence of a genotype outside of its usual range may result in
suboptimal flowering time and a corresponding reduction in fitness. Similar patterns
have been described for a bird species: across Europe, blue tits exhibit adaptive
differences in photoresponsiveness that results in laying dates coinciding with local
caterpillar availability (Lambrechts et al., 1996). These two examples both involve the
timing of key events being optimally aligned to seasonal environmental conditions,
but there are other ways in which local adaptation may be manifest. It is likely
that within many species there is variability in thermal tolerances with individuals
towards warmer parts of the range adapted to function better in the heat while
directional selection will have resulted in individuals found in cooler regions having
a physiology that equips them to function efficiently in the cold (e.g. fish (Feminella
and Matthews, 1984), amphibians (Olsson and Uller, 2003) and insects (Ayres and

Scriber, 1994)).

There is now recognition that local adaptation is likely to be an important determinant of species' responses to climate change and an interdisciplinary treatment of this issue, linking evolution and ecology, has already been called for (Pertoldi and Bach, 2007). We establish how local adaptation drives the dynamics of biogeographic ranges during and following a period of climate change. We utilise a spatially explicit individual based simulation to model the impact of climate change on a single species. This model uses both stochastic demographic parameters and allows inclusion of local density dependent rates for survival of individuals. In our discussion we emphasise the potential implications of our results for conservation biology and highlight where we believe future effort is required to develop improved theory related to the evolutionary ecology of range shifting.

The Model

There has been considerable recent interest in modelling the ecological and evolutionary dynamics of spatially structured populations that live along environmental gradients (e.g. Alleaume-Benharira et al., 2006; Travis et al., 2006; Dytham, 2009) and some studies have extended this approach to consider the range shifting of populations structured in this way (e.g. Brooker et al., 2007; Mustin et al., 2009). We adopt a similar approach in this contribution and below we describe, in turn, the structure of the landscape, the ecological and evolutionary features of the model, and the simulation experiments that we conduct.

The Environment

Individuals of the single species inhabit a two-dimensional landscape that, unless specified otherwise, has dimensions 200 rows (x value) columns by 200 columns (y value), which are ordered from the bottom left hand corner (i.e. increasing an individual's x location would move the individual up the grid, whilst increasing its y location would move it to the right of the grid). A value of (x, y) determines an individual's location. We assume that environmental conditions vary latitudinally (i.e. different rows have different conditions, for example due to variation in climate) and the parameter, $\theta(x)$, is used to denote this condition. We assume a linear gradient, b , in environmental conditions (i.e. for every value of y , the condition at x , $\theta(x)$ is bx). An optimal environmental location, J , is defined as the climate under which the highest potential population growth rates are obtained. Potential growth rate

declines with distance from this optimum location. J is defined as a row value, x , on the grid. In all our simulations, we allow populations to first establish under stable environmental conditions and in these periods the environmental conditions within each row remains constant through time. To simulate environmental change (e.g. climate warming), we increase $\theta(x)$ for each row, x and increase J by ν each year. The period of climate change lasts for T years. An example of an environmental condition changing latitudinally could be a warming climate, where the local temperature increases in all locations over time. *Local* adaptation to this condition would allow individuals to reside in the climate in which they are adapted. An optimal location, J , corresponds to a position where it is always favourable to be near, e.g. Pease et al. (1989) give the example of an advancing glacier, from which there exists an optimal place to reside. In the case of a warming climate, this optimal location could be any number of things e.g. a moving front of adverse weather conditions, the moving range of a quick-to-adapt parasite vector or food stuff in response to the changing climate. *Global* adaptation would allow individuals to reside near to this optimal location.

The Ecology

We assume that each cell on the lattice can support no more than one individual (i.e. we are using a patch occupancy model). In each year we simulate, in turn, reproduction, dispersal of offspring and mortality. All individuals first have the opportunity to reproduce. To avoid potential biases towards particular locations or individuals, the order in which individuals are given the opportunity for reproduction is randomised. The probability that an individual reproduces depends upon its fitness (see

Local Adaptation and Evolution, below). A single offspring is born to a reproducing adult and this offspring immediately disperses. This once-in-a-lifetime dispersal event occurs via a uniformly distributed Moore neighbourhood (nearest eight cells). Offspring that disperse to an unoccupied cell automatically establish. In most simulations, dispersal to an occupied cell results in automatic death and there is thus a strong priority effect in relation to space occupancy. However, in some simulations we relax this assumption and allow offspring to displace established individuals with probability, p . We always assume reflective boundary conditions as a way of maintaining the current population size if the individuals are able to find space. After all individuals have had an opportunity to reproduce, and offspring have dispersed, we simulate mortality. All individuals are subject to stochastic mortality with probability, μ . Mortality is independent of both the condition of the environment and an individual's phenotype.

Local Adaptation and Evolution

Each individual carries two alleles. These are represented as integer values and their mean determines the environmental condition to which the individual is adapted. For example, an individual possessing two alleles with values 24 and 28, has phenotype, $z=26$ (and is optimally adapted to the conditions in a row x such that $\theta(x) = 26$). The probability that an individual reproduces in any given year depends upon both where the individual is located in relation to the environmental location with highest potential productivity, J , and the degree to which the individual is adapted to its local environment, $\theta(x)$. Formally, the probability of reproducing is given by the

following expression:

$$\exp [((\theta(x) - z)^2/2V_s) + ((J - x)^2/2W_s)]$$

where, the first term accounts for the degree to which the individual is locally adapted and the second accounts for the proximity of the individual to the location with highest potential productivity. V_s determines the steepness of decline in an individual's fitness as it is moved further from the conditions to which it is best adapted, while W_s determines the rate at which maximum potential reproduction declines with distance from J . Note that W_s is akin to the notion of environmental tolerance as described in Pease et al. (1989).

It is worth emphasising the distinction between W_s and V_s and the need for both parameters in our model using a simple illustrative example. Consider first two individuals, one of which is perfectly adapted to the conditions at location J while the other is perfectly adapted to conditions found 10 rows higher on the lattice than J . Initially we will assume both are located in the environments to which they are adapted. Despite the fact that both are optimally adapted, their reproductive fitness will not be the same as the first individual (at location J) is in the inherently more productive environment. Our parameter W_s determines just how much lower the reproductive fitness is of the individual found 10 rows away from J . Next consider the case that these same two individuals are both located at the position 10 rows away from J . Now the first individual will have the lower reproductive fitness because, while both are in the same environment, the first is not optimally adapted to that

environment. The amount that the first individual's fitness is reduced due to it being
 216 less well-adapted is determined by V_s .

When a birth event occurs the offspring inherits two alleles: a copy of one of its
 218 parent's alleles (chosen randomly from the set of 2 with equal probability) and a copy
 from an individual, selected at random, in the vicinity of its parent. This vicinity can
 220 be considered as the pollen dispersal neighbourhood and we term it h . If the vicinity
 contains no individuals then the offspring inherits both alleles from its parent (this
 222 is akin to 'selfing'). During the copying of alleles that ensues, mutation occurs at
 each locus with a probability of β . When a mutation occurs, the value of the allele
 224 is modified by an amount randomly drawn from the discrete uniform distribution,
 $(-10, 10)$.

226 Simulation Experiments

We have conducted a number of simulation experiments to explore the dynamics
 228 of the model. In the first set, we run the model under a stable climate (i.e. main-
 taining a constant position of maximum potential reproduction, J). We run the
 230 model for a range of parameter values and, in particular, vary both V_s and W_s . We
 output population abundance and mean phenotype across the environmental gradi-
 232 ent. Subsequently, we run a large number of simulations where we initially allow the
 population to obtain quasiequilibrium under stable environmental conditions before
 234 simulating a period of climate change. In these simulations we are most interested
 in the probability that a population survives an episode of climate change. We run
 236 sensitivity analyses to ask, in turn, how this depends upon V_s and W_s , the strength

of the priority effect (p), the rate (ν) and duration of climate change (T), and the
rate and size of mutations. In the final set of simulations we vary the probability of
mortality (β), the landscape width (by altering the number of columns on the grid),
and the dispersal neighbourhood for pollen (h). In all the sensitivity analyses, we
run 50 replicate simulations for each parameter combination.

Results

Under stable environmental conditions, population distributions establish as one would expect. For smaller W_s (corresponding to a steeper decline in maximum potential reproductive rate with distance from J), smaller ranges emerge (Figures 1a,1b and 1c). The corresponding mean phenotype is shown in Figures 1d, 1e and 1f. For smaller W_s , there is a greater mismatch between the local environment and the mean phenotype away from J (in these figures $J=140$).

Most of our results relate to range dynamics under climate change. Figure 2 illustrates typical dynamics showing snapshots at various stages in the climate change period. Initially, we allow the population to obtain quasiequilibrium under stable climate and this is what is shown in Figure 2a. At this stage, we define a ‘climate envelope’ for our simulated species. We establish the range of environmental conditions within which 99% of the individuals are found. We show the location of this climate envelope during the period of environmental change to illustrate how well the population’s range responds. In the example given in Figure 2, as the environment changes, the population’s range lags behind its climate envelope. This is most marked at the expanding front, with the leading edge of the range advance falling a substantial distance behind the shifting climate window. However, it is also apparent that at the trailing edge some individuals are found well behind what we defined as the species’ climate envelope.

Running the simulation model for a wide range of parameter combinations revealed a rich range of behaviours. We illustrate some of the most important features

in Figure 3a which shows mean phenotypes across the environmental gradient on one axis, and how this distribution of phenotypes changes during and subsequent to a period of climate change (see Figure 3). For most runs of the model where the population survives the episode of climate change we observe that the range width following climate change is narrower (Figures 3a, 3b and 3d). This is a consequence of the loss of phenotypes adapted to the warmest and coolest extremes that were occupied by the population prior to the onset environmental change. Ultimately phenotypic variation is recovered and the range expands to reoccupy its original niche but this can take many thousands of generations. A common feature of the results is the initial expansion of the cold-adapted phenotypes (shown in green in Figure 3a) after the onset of change, but this is followed by a rapid collapse of these phenotypes (Figures 3b, 3c and 3d) and, in some cases, a rapid collapse of the whole population (Figure 3c). Figure 3c illustrates an important result: even when there is a considerable overlap between the range of a species prior to climate change and the climate envelope of the species following change (shown by the white lines in Figure 3) extinction can still occur. Finally, we sometimes observe dynamics where the population splits into distinct subpopulations comprising individuals with quite distinct phenotypes. Because they are adapted to quite different climates, these subpopulations can remain spatially disjunct for quite lengthy periods following climate change (see Figure 3d).

Figures 4 to 6 show the results of our sensitivity analyses, conducted to establish to roles of the different model parameters in determining the likelihood that a population is able to survive a period of climate change. Unsurprisingly, steeper declines in

individuals' reproductive fitness as they are moved away from their own optima
 (i.e. small V_s), lead to reduced probabilities of populations surviving climate change
 (Figure 4a). However, for the sensitivity analysis around W_s , the results are a little
 less expected. Higher probabilities of extinction are observed when the fundamental
 environment niche of the species is broader (i.e. the chance of extinction is greater
 for higher W_s - see Figure 4b).

Perhaps our most important result is illustrated in Figure 5. This figure shows
 the results of simulations conducted to establish the role of p in determining the
 likelihood that species survive an episode of climate change. Recall that p determines
 the probability that an offspring is able to displace an already established individual
 from the site to which it disperses. The counterintuitive result shown in Figure 4b
 was obtained for $p = 0.0$ (i.e. there is an absolute priority effect). When the opposite
 is assumed, and offspring always displace existing occupants (i.e. $p = 1.0$), we find
 that species that have broader fundamental niches (higher W_s) are now more likely to
 survive than those with narrower niches (lower W_s). Interestingly, when we assume
 an offspring has a 50% chance of displacing an existing occupant ($p = 0.5$), we find
 that the initial result holds and extinction risk is higher for higher W_s .

We find that the probability that a species' range survives a period of climate
 change is far more sensitive to some of the model's parameters than others. In
 general we find that the results are largely insensitive to the mutation parameters,
 neither the mutation rate nor the effect size of a single mutation have noticeable
 impacts (results not shown). However, the scale of pollen dispersal, the length of the
 period of climate change, the extrinsic probability of mortality and the width of the

310 landscape all have major effects. In general, extinction risk is higher when pollen
disperses shorter distances (Figure 6a), the period of climate change is longer (Figure
312 6b), the probability of extrinsic mortality is lower (Figure 6c) and the landscape is
broader (Figure 6d).

Discussion

In this contribution, we have presented results from a simulation model that integrates key features from similar models exploring species' dynamics in response to environmental change (Brooker et al., 2007; Dytham, 2009; Mustin et al., 2009, e.g.) and from previous analytical theory exploring local adaptation on environmental gradients (Kirkpatrick and Barton, 1997; Pease et al., 1989). We establish that local adaptation can play a major role in determining the dynamics of range shifting under climate change, and suggest that this is likely to have important implications for predictions of the impact of environmental change on both future species ranges and the rates of extinction.

Two results from our work clearly highlight how local adaptation can alter the nature of range shifting. First, a species may fail to survive a period of climate change, even when there is an overlap between its range prior to climate change and the area where climate is predicted to be suitable following climate change. Second, we find that often a range begins to expand in the direction of climate change before rapidly collapsing to extinction. Both these results have important applied implications. The first suggests that any predictions based upon treating the species as a single homogenous unit may be flawed and that, for this reason, simple extensions of the bioclimate envelope approach are likely to be inadequate. The second implies that we need to be extremely cautious before inferring that species that are already expanding their ranges in the direction of climate change will continue to do so.

In our initial simulations, we found that species occupying broader niches more

at risk of extinction due to climate change. This counterintuitive result required further investigation, and the key assumption driving this difference turned out to be local density dependence in the form of space occupancy. In the initial simulation model, individuals occupy a patch until they die and are never displaced by a newborn individual. When we alter the priority effect within the simulations, such that established individuals are no longer invulnerable to displacement by newborn individuals, we find that species with broader ranges in a stable climate are better able to survive a period of environmental change. Pease et al. (1989) concluded that without local density dependence, a species with a broader range was less likely to become extinct during an episode of climate change. This is agreement with our findings.

Examining the original distributions of the population before climate change in the simulation, it becomes clear why the result has been obtained. The individuals at the periphery of the range (those lying on the lines at the extrema of the current climate) are lower in density and will necessarily have a lower birth rate than those towards the centre of the distribution (given the same local adaptation). The offspring of these individuals will however be able to establish more readily since the density of individuals decreases away from the globally optimal location. Thus, we note that the absolute gradient of the density of individuals away from this global optimum increases with decreasing W_s . As a result, we see that decreasing W_s increases the availability of patches surrounding the optimum location, into which offspring can be born. Therefore as the climate starts to shift and the optimal location moves upwards, the high frequency of better adapted individuals with higher birth rates around the

centre of the distribution have more space to move forward in the range and the individuals at the front of the distribution are soon optimally located, making them more fecund (despite being less locally adapted) but with fewer density dependent effects limiting their birth rate. This then creates a faster turnover of new individuals, on which selection can act. When W_s is large, this movement and selection of phenotypes is much impaired leading to an overall reduction in the adaptability of the population and an increased chance of extinction. The local density-dependence effects have a much more striking effect on extinction probability than mutation rate or mutation size, which do not ameliorate extinction probability if density dependent effects are present.

The important role played by local density dependence, mediated through space occupancy effects, has recently been demonstrated for species assemblages and our results clearly demonstrate that it can also be important within a species. Takenaka (2004) uses simulation modelling to explore the distributions of coexisting tree species as they shift their ranges due to climate change. Takenaka's work shows that the replacement of one species by another better adapted to the new climatic conditions can be dramatically slowed if mature individuals continue to block space for lengthy periods even though the climate no longer suits them. Brooker et al. (2007) also described a similar space blocking effect in a simulation model describing the range dynamics of an assemblage comprising species arranged along an environmental stress gradient. The authors highlighted that the longevity of an organism would determine the strength of the effect. Our work highlights that these effects are not limited to interspecific interactions; prolonged space occupancy beyond the point at which an

individual is well adapted to local conditions can similarly limit the range shifting of locally adapted phenotypes within a single species. An important determinant of species' extinction, both within and between species, will be the relative rates at which fecundity and mortality are impacted as the climate shifts. If fecundity declines relatively rapidly, space occupancy effects will be greater, while if mortality increases more rapidly, when environmental conditions are displaced from those to which an individual is adapted, the effect will be weaker. Additionally, higher rates of fitness-independent mortality will reduce the strength of the space blocking effect and allow more successful range shifting (see Figure 6c for an example). In future work, it will be interesting to explore how the partitioning of local adaptation within and between species may influence the spatial dynamics of a species guild under climate change scenarios.

An interesting feature of our results is that a species can become extinct even when the climate envelopes before and after climate change overlap. The realisation of the simulation model illustrated in Figure 3c shows this effect very clearly. Representation of a species as a single unit, as is the case in the statistical pattern matching methods used in bioclimate envelope modelling (Bakkenes et al., 2002; Broennimann et al., 2006; Schwarz et al., 2006; Thuiller et al., 2005) neglects this potentially crucial effect. There is an urgent need for a careful consideration of the taxonomic resolution at which we should be modelling the response of biodiversity to climate change. As further information becomes available on the extent to which populations within species are differentially adapted to local conditions, a critical appraisal of the importance of this evolutionary process will be required.

This question of the organisational level at which species are represented also
 406 needs addressing within the community using dynamic vegetation models to pre-
 dict both changes in ecological and biogeochemical processes during climate change
 408 (Cramer et al., 2001; Morales et al., 2005). Many of these models do not even work
 at the resolution of the species, instead they group species into functional types and
 410 assume that all members of a functional type share an identical environmental niche.
 In reality, species (or populations) towards the cooler end of the environmental space
 412 that can be occupied by the functional type will not function in warmer conditions as
 effectively as those species (or populations) that have adapted to the heat, and vice
 414 versa. This may have important consequences not only for the projections of future
 patterns of biodiversity made by this type of model but also for their projections of
 416 future biogeochemical cycles and balances.

A new generation of models is required that takes the best parts from bioclimate
 418 modelling and incorporates ecological and evolutionary dynamics. The results pre-
 sented in this paper make a strong case for the inclusion of local adaptation as a
 420 matter of urgency. One conservation management option to mitigate some of the ef-
 fects described in this paper would be the translocation of genotypes within a species
 422 range. Our simulations tend to show that the phenotypes adapted to the local condi-
 tions found in the core of the range tend to be the most important for range shifting,
 424 and from this we would suggest that a potential strategy would be to translocate
 individuals (for plants, pollen might suffice) from core areas to populations located
 426 towards the expanding front. A model parameterised with species-specific informa-
 tion on dispersal distance and the genetics of local adaptation would be an invaluable

aid in developing this type of strategy, and could be used to provide recommendations on the optimal amount and frequency of translocation.

The model considered in this paper has much in common with a model used to explore the dynamics of speciation along environmental gradients (Doebeli and Dieckmann, 2003). Interestingly, we have observed that in some regions of parameter space we find that disruptive selection can result in two (or more) spatially distinct species arising in our simulation model. This is not the main topic of this paper, however we note that both during and following a period of climate change a species' range tends to become fragmented (see Figures 2 and 3d). In some cases this spatial separation can last for a considerable period following the stabilisation of climate (Figure 3d) and it is interesting to speculate that in some cases this may act to promote speciation. It would certainly be worthwhile extending recent speciation models (Artzy-Randrup and Kondrashov, 2006; Bolnick, 2006; Yukilevich and True, 2006) to consider the potential impacts of environmental change.

Understanding the response of biodiversity at different organisational levels under a period of rapid environmental change represents the greatest challenge faced by ecologists today. Here, we have considered how local adaptation within a sessile species' range can impact on that species dynamics during climate change. By applying ideas and methods that are already well developed in the ecological literature we have obtained several new insights that we believe have important consequences for how we should go about both predicting and also managing changes in biodiversity under climate change. We hope that the results from theoretical studies such as this one make a persuasive case for a closer integration between the statistical modelling

that currently dominates the climate impacts literature and dynamic spatial models
incorporating biological processes.

Acknowledgements

Thanks to Zaq Coelho and Gustav Delius for continued support. This research was
funded by NERC and ALTERNET.

References

- Alleaume-Benharira, M., Pen, I. R., Ronce, O., 2006. Geographical patterns of adaptation within a species' range: Interactions between drift and gene flow. *Journal of Evolutionary Biology* 19, 203–215.
- Araujo, M. B., Whittaker, R. J., Ladle, R. J., Erhard, M., 2005. Reducing uncertainty in projections of extinction risk from climate change. *Global Ecology and Biogeography* 14, 529–538.
- Artzy-Randrup, Y., Kondrashov, A. S., 2006. Sympatric speciation under incompatibility selection. *Proceedings of the National Academy of Sciences of the United States of America* 103, 11619–11624.
- Ayres, M. P., Scriber, J. M., 1994. Local adaptation to regional climates in *Papilio canadensis* (Lepidoptera: Papilionidae). *Ecological Monographs* 64, 465–482.
- Bakkenes, M., Alkemade, J. R. M., Ihle, F., 2002. Assessing effects of forecasted climate change on the diversity and distribution of European higher plants for 2050. *Global Change Biology* 8, 390–407.
- Bolnick, D. I., 2006. Multi-species outcomes in a common model of sympatric speciation. *Journal of Theoretical Biology* 241, 734–744.
- Broennimann, O., Thuiller, W., Hughes, G., Midgley, G. F., Alkemade, J. M. R., Guisan, A., 2006. Do geographic distribution, niche property and life form explain plants' vulnerability to global change? *Global Change Biology* 12, 1079–1093.

- Brooker, R. W., Travis, J. M. J., Clark, E. J., Dytham, C., 2007. Modelling species' range shifts in a changing climate: The impacts of biotic interactions, dispersal distance and the rate of climate change. *Journal of Theoretical Biology* 245, 59–65.
- Burton, O., Phillips, B., Travis, J., 2010. Trade-offs and the evolution of life-histories during range expansion. *Ecology Letters*, *In press*.
- Burton, O. J., Travis, J. M. J., 2008a. The frequency of fitness peak shifts is increased at expanding range margins due to mutation surfing. *Genetics* 179, 941–950.
- Burton, O. J., Travis, J. M. J., 2008b. Landscape structure and boundary effects determine the fate of mutations occurring during range expansions. *Heredity* 101, 329–340.
- Cramer, W., Bondeau, A., Woodward, F. I., Prentice, I. C., Betts, R. A., Brovkin, V., Cox, P. M., Fisher, V., Foley, J. A., Friend, A. D., Kucharik, C., Lomas, M. R., Ramankutty, N., Sitch, S., Smith, B., White, A., Young-Molling, C., 2001. Global response of terrestrial ecosystem structure and function to CO_2 and climate change: results from six dynamic global vegetation models. *Global Change Biology* 7, 357–373.
- Davis, M. B., Shaw, R. G., Etterson, J. R., 2005. Evolutionary responses to changing climate. *Ecology* 86, 1704–1714.
- Dytham, C., 2009. Evolved dispersal strategies at range margins. *Proceedings of the Royal Society B: Biological Sciences* 276, 1407–1413.

- Excoffier, L., Ray, N., 2008. Surfing during population expansions promotes genetic revolutions and structuration. *Trends in Ecology and Evolution* 23, 347–351.
- Feminella, J. W., Matthews, W. J., 1984. Intraspecific differences in thermal tolerance of *Etheostoma spectabile* (agassiz) in constant versus fluctuating environments. *Journal of Fish Biology* 25, 455–461.
- Garcia-Ramos, G., Rodriguez, D., 2002. Evolutionary speed of species' invasions. *Evolution* 56, 661–668.
- Gavin, D. G., Hu, F. S., 2005. Bioclimatic modelling using gaussian mixture distributions and multiscale segmentation. *Global Ecology and Biogeography* 14, 491–501.
- Hampe, A., 2004. Bioclimate envelope models: what they detect and what they hide. *Global Ecology and Biogeography* 13, 469–471.
- Kirkpatrick, M., Barton, N. H., 1997. Evolution of a species' range. *American Naturalist* 150, 1–23.
- Klopfstein, S., Currat, M., Excoffier, L., 2006. The fate of mutations surfing on the wave of a range expansion. *Molecular Biology and Evolution* 23, 482–490.
- Lambrechts, M. M., Perret, P., Blondel, J., 1996. Adaptive differences in the timing of egg laying between different populations of birds result from variation in photoresponsiveness. *Proceedings of the Royal Society of London Series B - Biological Sciences* 263, 19–22.
- Lawton, J., 2000. Ecological Consequences of Heterogeneity (ed. M. J. Hutchings, E.

- 516 *John and A. J. A. Stewart*). Cambridge University Press, concluding Remarks: a
Review of Some Open Questions.
- 518 McNeilly, T., Antonovic, J., 1968. Evolution in closely adjacent plant populations 4:
barriers to gene flow. *Heredity* 23, 205.
- 520 Midgley, G. F., Hannah, L., Millar, D., Rutherford, M. C., Powrie, L., 2002. As-
sessing the vulnerability of species richness to anthropogenic climate change in a
522 biodiversity hotspot. *Global Ecology and Biogeography* 11, 445–451.
- 524 Morales, P., Sykes, M. T., Prentice, I. C., Smith, P., Smith, B., Bugmann, H., Zierl,
B., Friedlingstein, P., Viovy, N., Sabate, S., Sanchez, A., Pla, E., Gracia, C. A.,
Sitch, S., Arneth, A., Ogee, J., 2005. Comparing and evaluating process-based
526 ecosystem model predictions of carbon and water fluxes in major european forest
biomes. *Global Change Biology* 11, 2211–2233.
- 528 Mustin, K., Benton, T., Dytham, C., Travis, J., 2009. The dynamics of climate-
induced range shifting; perspectives from simulation modelling. *Oikos* 118, 131–
530 137.
- 532 Olsson, M., Uller, T., 2003. Thermal environment, survival and local adaptation in
the common frog, *Rana temporaria*. *Evolutionary Ecology Research* 5, 431–437.
- 534 Parmesan, C., Yohe, G., 2003. A globally coherent fingerprint of climate change
impacts across natural systems. *Nature* 421, 37–42.
- 536 Pearson, R. G., Dawson, T. P., 2003. Predicting the impact of climate change on

- 536 the distribution of species: are bioclimate envelope models useful? *Global Ecology*
and *Biogeography* 12, 361–371.
- 538 Pearson, R. G., Dawson, T. P., Berry, P. M., 2002. A spatial evaluation of climate
impact on the envelope of species. *Ecological Modelling* 154, 289–300.
- 540 Pease, C. M., Lande, R., Bull, J. J., 1989. A model of population growth, dispersal
and evolution in a changing environment. *Ecology* 70, 1657–1664.
- 542 Pertoldi, C., Bach, L. A., 2007. Evolutionary aspects of climate-induced changes and
the need for multidisciplinary. *Journal of Thermal Biology* 32, 118–124.
- 544 Peterson, A. T., Ortega-Huerta, M. A., Bartley, J., Sanchez-Cordero, V., Soberon,
J., Buddemeier, R. H., Stockwell, D. R. B., 2002. Future projections for Mexican
546 faunas under global climate change scenarios. *Nature* 416, 626–629.
- Phillips, B. L., Brown, G. P., Webb, J. K., Shine, R., 2006. Invasion and the evolution
548 of speed in toads. *Nature* 439, 7078.
- Polechová, J., Barton, N., Marion, G., 2009. Species' range: Adaptation in space and
550 time. *American Naturalist* 174, E186–E204.
- Riihimäki, M., Podolsky, R., Kuittinen, H., Koelewijn, H., Savolainen, O., 2005. Study-
552 ing genetics of adaptive variation in model organisms: flowering time variation in
Arabidopsis lyrata. *Genetica* 123, 63–74.
- 554 Santamaria, L., Figuerola, J., Pilon, J. J., Mjelde, M., Green, A. J., Boer, T. D.,

King, R. A., Gornall, R. J., 2003. Plant performance across latitude: The role of
 556 plasticity and local adaptation in an aquatic plant. *Ecology* 84, 2454–2461.

Schwarz, M. W., Iverson, L. R., Prasad, A. M., Matthews, S. N., O'Connor, R. J.,
 558 2006. Predicting extinctions as a result of climate change. *Ecology* 87, 1611–1615.

Takenaka, A., 2004. Local coexistence of tree species and the dynamics of global dis-
 560 tribution pattern along an environmental gradient: a simulation study. *Ecological
 Research* 20, 297–304.

Termansen, M., McClean, C. G., Preston, C. D., 2006. The use of genetic algorithms
 562 and bayesian classification to model species distributions. *Ecological Modelling*
 564 192, 410–424.

Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Colling-
 566 ham, Y. C., Erasmus, B. F. N., de Siqueira, M. F., Grainger, A., Hannah, L.,
 Hughes, L., Huntley, B., van Jaarsveld, A. S., Midgley, G. F., Miles, L., Ortega-
 568 Huerta, M. A., Peterson, A. T., Phillips, O. L., Williams, S. E., 2004. Extinction
 risk from climate change. *Nature* 427, 145–147.

Thuiller, W., Lavorel, S., Araujo, M. B., 2005. Climate change threats to plant
 570 diversity in Europe. *Proceedings of the National Academy of Sciences of the USA*
 572 102, 8245–8250.

Travis, J., Brooker, R., Clark, E., Dytham, C., 2006. The distribution of positive
 574 and negative species interactions across environmental gradients on a dual-lattice
 model. *Journal of Theoretical Biology* 241, 896–902.

- 576 Travis, J. M. J., Dytham, C., 2002. Dispersal evolution during invasions. *Evolutionary Ecology Research* 4, 1119–1129.
- 578 Travis, J. M. J., Munkemuller, T., Burton, O. J., Best, A. S., Dytham, C., Johst, K., 2007. Deleterious mutations can surf to high densities on the wave front of an
580 expanding population. *Molecular Biology and Evolution* 24, 2334–2343.
- Travis, J. M. J., Mustin, K., Benton, T. G., Dytham, C., 2009. Accelerating invasion rates result from the evolution of density-dependent dispersal. *Journal of Theoretical Biology* 259, 151158.
- 582
- 584 Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.-M., Hoegh-Guldberg, O., Bairlein, F., 2002. Ecological responses to
586 recent climate change. *Nature* 416, 389–395.
- Woodward, F. I., Beerling, D. J., 1997. The dynamics of vegetation change: Health
588 warnings for equilibrium ‘dodo’ models. *Global Ecology and Biogeography Letters* 6, 413–418.
- 590 Yukilevich, R., True, J. R., 2006. Divergent outcomes of reinforcement speciation: The relative importance of assortative mating and migration modification. *American Naturalist* 167, 638–654.
592

Figure 1: Frequency distributions for population size and trait value for the equilibrium in a constant environment, where the environment changes with x . The results were averaged over 50 realisations of the model. A set of simulations were run for a different values of $2\ln W_s$ given on the graph by different colours (red-2, blue-3, green-4, yellow-5, black-6). An increasing value of W_s relates to an increased environmental niche range, seen in Figures (a), (b) and (c). In Figures (d), (e) and (f) the mean phenotype is an average of all allele values (2 per individual) existing on the environmentally invariant line $y = \text{const}$. In the case of zero density, the mean phenotype is given a value 0.

Figure 2: Snapshots of one realisation of the model showing population inhabiting the 2d lattice landscape. The grid corresponds to the individuals on the grid with the invariant environment of each row and changing environmental gradient through each column. The colour grid shows the optimal trait value for each location, $\theta(x)$, running from hot (red) to cool (blue). Individual patch colour shows trait value to which the individual is adapted, z . The climate envelope (shown by black lines) indicates the area holding approximately 98% of the population at pre-climate change equilibrium, with the median at the optimal. (a) shows the equilibrium in a constant environment before any climate change has occurred; (b), (c), (d) and (e) show the population at a specified time after change onset and (f) shows final distribution after climate change. Illustrative parameters are chosen as $W_s = \exp(5)/2$ and $V_s = 25$, under a climate change rate of $\nu = 0.3$ over $T = 200$ years to emphasise the range shift. During the climate shift this envelope tracks the environmental optimal for the population. The population splits into clusters before combining once the shift ends.

Figure 3: Four realisations of the model through a period of climate change. The colours show the population and its local adaptation, z (blue through to red). (a) shows population persistence with the majority of individuals tracking the climate; $T=300, V_s=100, W_s=500, \nu=25$ leading to the subpopulation at the rear and very front of the range being lost but population persisting due to the middle third of the pre shift population moving. (b) shows population persistence through climate change with only a small subpopulation tracking the climate; $T=300, V_s=10, W_s=500, \nu=23$ allowing population movement due to the middle of the front half of the range shifting, although front margin does show signs of initial movement. (c) shows population extinction during climate change; $T=100, V_s=50, W_s=250, \nu=40$ produces a climate shift too fast for adaptation. (d) shows population persistence with two small subpopulations tracking the climate; $T=300, V_s=7.5, W_s=500, \nu=23$ produces similar subpopulation shifting its range as seen in (b) but shift occurs in two geographically distinct subpopulations, which eventually rejoin for good on reaching the new equilibrium.

Figure 4: Sensitivity analysis for variance revealing both population variances (V_s and W_s) having opposite effects on extinction probability in the model with (a) showing the effect of variance around local optimal adaptation and (b) the effect of variance around the optimal location. The extinction probability for any population can be noted as a non-decreasing function of the rate of climate change, ν . Notice that an increase in the environmental niche (an increase in W_s) leads to a increase, or no change, in the extinction probability.

Figure 5: The effect of decreased density dependent effects on proportion of populations extinct under climate change. Red to Blue scale corresponds as before to 100% to 0% of populations extinct after climate shift. Note the qualitative difference in the two trends: in (a) where $p=0.5$, a large range width corresponds to a higher extinction probability, whereas in (b) a large range width allows better population persistence.

Figure 6: The effect of different parameters on species extinction. The plots show the proportion of realisations (out of a total of 50) where the species became extinct with varying rates of climate change. In all cases, $W_s = 150$ and unless otherwise varied, parameter values are kept constant at $N = 200$, $T = 200$, $V_s = 50$, $\mu = 0.005$, $\sigma = 1.207$ and maximum mutation size is set at 10.