Fast control cost for heat-like semigroups: Lebeau-Robbiano strategy and Hautus test

Luc Miller

► To cite this version:

Luc Miller. Fast control cost for heat-like semigroups: Lebeau-Robbiano strategy and Hautus test. Wellposedness and controllability of evolution equations, Dec 2010, Oberwolfach, Germany. pp.3269-3309, 10.4171/OWR/2010/56 . hal-00620867

HAL Id: hal-00620867

<https://hal.science/hal-00620867>

Submitted on 8 Sep 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MATHEMATISCHES FORSCHUNGSMINISTRIUM OBERWOLFACH

Report No. 56/2010

DOI: 10.4171/OWR/2010/56

**Mini-Workshop: Wellposedness and Controllability of
Evolution Equations**

Organised by
Birgit Jacob, Wuppertal
Jonathan Partington, Leeds
Sandra Pott, Paderborn/Glasgow
Hans Zwart, Enschede

December 12th – December 18th, 2010

Mathematics Subject Classification (2000): Primary: 47D06, 93B05, 93C20. Secondary: 93B28, 47A57, 47B35, 42B37.

Preprint of the extended abstract of the talk by Luc Miller.

The full report can be downloaded from the website of the MFO via
“Scientific Programme/Annual Schedules/Mini-Workshops in 2010/Mini-Workshop:
Wellposedness and Controllability of Evolution Equations/OWR_2010_56.pdf”.

Abstracts

Fast control cost for heat-like semigroups: Lebeau-Robbiano strategy and Hautus test

LUC MILLER

(joint work with Thomas Duykaerts)

Since the seminal work of Russell and Weiss in [7], resolvent conditions for various notions of admissibility, observability and controllability, and for various notions of linear evolution equations have been studied intensively, sometimes under the name of infinite-dimensional Hautus test, cf. [8, 2]. This talk based on [1] investigates resolvent conditions for null-controllability in arbitrary time: necessary conditions for general semigroups, and sufficient conditions for analytic normal semigroups and semigroups with negative self-adjoint generators.

1. INTRODUCTION

Let $-A$ be the generator of a strongly continuous semigroup on a Hilbert space $\mathcal{E}$. Let C be a bounded operator from the domain $D(A)$ with the graph norm to another Hilbert space $\mathcal{F}$. The norms in $\mathcal{E}$ and $\mathcal{F}$ are denoted $\|\cdot\|$. We refer to the monograph [8] for a full account of the control theory of semigroups.

Recall the usual *admissibility* condition (for some time $T > 0$ hence all $T > 0$),

$$(1) \quad \exists K_T > 0, \forall v \in D(A), \quad \int_0^T \|Ce^{-tA}v\|^2 dt \leq K_T \|v\|^2.$$

If C is admissible for A then null-controllability at time T is equivalent to *final-observability* at time T (cf. [8], i.e.

$$(2) \quad \exists \kappa_T > 0, \forall v \in \mathcal{E}, \quad \|e^{-TA}v\|^2 \leq \kappa_T \int_0^T \|Ce^{-tA}v\|^2 dt.$$

The control property investigated here is (2) for all $T > 0$.

1.1. Control cost. The coefficient κ_T in (2) is the *control cost*: it is the ratio of the size of the input over the size of the initial state which the input steers to the zero final state in a lapse of time T . It blows up as $T \rightarrow 0$. E.g. for the heat semigroup on a compact manifold M with Dirichlet boundary conditions observed from a subset Ω : $\kappa_T \leq c_0 \exp(2c/T)$, $T \in (0, 1)$, where c_0 is a positive constant, implies $c \geq d^2/4$ where d is the furthest a point of M can be from Ω , and is implied by $c > 3L^2/4$ where L is the length of the longest generalized geodesic in M which does not intersect Ω ($L < +\infty$ is known as the condition of Bardos-Lebeau-Rauch).

For many evolutions of parabolic type, κ_T is bounded by $c_0 \exp(2c/T^\beta)$ where c , c_0 and β are positive constants. E.g. thermoelastic plates without rotatory inertia, the plate equation with square root damping, diffusions in discontinuous media or in a potential well, diffusions generated by the fractional Laplacian or non-selfadjoint elliptic generators, cf. references in [5].

1.2. Resolvent conditions. The resolvent condition: $\exists M > 0$,

$$(3) \quad \|v\|^2 \leq \frac{M}{(\operatorname{Re} \lambda)^2} \|(A - \lambda)v\|^2 + \frac{M}{\operatorname{Re} \lambda} \|Cv\|^2, \quad v \in D(A), \quad \operatorname{Re} \lambda > 0.$$

was introduced in [7] as a necessary condition for exact observability in infinite time of exponentially stable semigroups.

When A is skew-adjoint (equivalently when the semigroup is a unitary group), it was proved in [3] that the following resolvent condition is necessary and sufficient for final-observability (hence exact observability) in some time $T > 0$: $\exists M > 0$,

$$\|v\|^2 \leq M \|(iA - \lambda)v\|^2 + M \|Cv\|^2, \quad v \in D(A), \quad \lambda \in \mathbb{R}.$$

We refer to [6] for more background and references. This result was extended to some more general groups in [2, theorem 1.2].

When $-A$ generates an exponentially stable normal semigroup, [2, theorem 1.3] proves that the resolvent condition (3) is sufficient for the weaker notion:

$$(4) \quad \exists T > 0, \exists \kappa_T > 0, \forall v \in \mathcal{E}, \|e^{-TA}v\|^2 \leq \kappa_T \int_0^\infty \|Ce^{-tA}v\|^2 dt.$$

In this framework (4) implies (2) for *some* time T .

But it seems that resolvent conditions for final-observability for *any* $T > 0$ in (2) has not been investigated yet, although it is quite natural for heat-like semigroups.

2. RESULTS

2.1. Necessary resolvent conditions for semigroups. The proof mainly consists in changing i into -1 in [3, lemma 5.2]. Cf. also the proof of [7, theorem 1.2].

Theorem 1. *Let $B_T = \sup_{t \in [0, T]} \|e^{-tA}\|$ be the semigroup bound up to time T .*

If (1) and (2) hold then : $\forall v \in D(A)$, $\lambda \in \mathbb{C}$, $\operatorname{Re} \lambda > 0$,

$$\|v\|^2 \leq 2e^{2T \operatorname{Re} \lambda} \left((B_T^2 + 2\kappa_T K_T) \frac{\|(A - \lambda)v\|^2}{(\operatorname{Re} \lambda)^2} + \kappa_T \frac{\|Cv\|^2}{\operatorname{Re} \lambda} \right),$$

Theorem 2. *If final-observability (2) holds for all $T \in (0, T_0]$ with the control cost $\kappa_T = c_0 e^{\frac{2c}{T^\beta}}$ for some positive β , c and c_0 then the resolvent condition*

$$\|v\|^2 \leq a_0 e^{2a(\operatorname{Re} \lambda)^\alpha} (\|(A - \lambda)v\|^2 + \|Cv\|^2), \quad v \in D(A), \quad \operatorname{Re} \lambda > 0,$$

holds with power $\alpha = \frac{\beta}{\beta+1}$ and rate $a = c^{\frac{1}{\beta+1}} \frac{\beta+1}{\beta^\alpha}$.

It still holds for $\lambda \in \mathbb{C}$ if $\operatorname{Re} \lambda$ is replaced by $\operatorname{Re}_+ \lambda := \max\{\operatorname{Re} \lambda, 0\}$.

2.2. Sufficient resolvent conditions for an analytic normal semigroup. The proof is based on the Lebeau-Robbiano strategy of [5]. N.b. (1) is not assumed.

Theorem 3. *Assume that $-A$ generates an analytic normal semigroup, hence there exists $\omega \in \mathbb{R}$ and $\theta \in [0, \frac{\pi}{2})$ such that $\sigma(A) \subset \{z \in \mathbb{C} : \arg(z - \omega) \leq \theta\}$.*

The resolvent condition with $\alpha \in (0, 1)$, $\omega_0 < \omega$, $\lambda_0 > \omega_0$, positive a_0 and a ,

$$\|v\|^2 \leq \frac{\cos^2 \theta}{(\lambda - \omega_0)^2} \|(A - \lambda)v\|^2 + a_0 e^{2a\lambda^\alpha} \|Cv\|^2, \quad v \in D(A), \quad \lambda \geq \lambda_0,$$

implies final-observability (2) for all time $T > 0$ with the control cost estimate

$$\limsup_{T \rightarrow 0} T^\beta \ln \kappa_T \leq 2a^{\beta+1}(\beta+1)^{\beta(\beta+1)}\beta^{-\beta^2}, \quad \text{where } \beta = \frac{\alpha}{1-\alpha}.$$

2.3. Sufficient resolvent condition for a negative self-adjoint generator.

The proof combines the Lebeau-Robbiano strategy of [5], the control transmutation method of [4] (which deduces the final-observability of the heat-like equation $\dot{v} + Av = 0$ from the exact observability of the wave-like equation $\ddot{w} + Aw = 0$) and results on resolvent conditions from [6].

Theorem 4. Assume that the positive self-adjoint operator A and the operator C bounded from $D(\sqrt{A})$ with the graph norm to $\mathcal{F}$ satisfy the admissibility and observability conditions with nonnegative powers γ and positive δ , L_* and M_* :

$$\begin{aligned} \|Cv\|^2 &\leq L_* \lambda^\gamma \left(\frac{1}{\lambda} \|(A - \lambda)v\|^2 + \|v\|^2 \right), \quad v \in D(A), \quad \lambda \geq \inf A, \\ \|v\|^2 &\leq M_* \lambda^\delta \left(\frac{1}{\lambda} \|(A - \lambda)v\|^2 + \|Cv\|^2 \right), \quad v \in D(A), \quad \lambda \geq \inf A. \end{aligned}$$

If $\gamma + \delta < 1$ then final-observability (2) holds for all $T > 0$ with the cost estimate

$$\limsup_{T \rightarrow 0} T^\beta \ln \kappa_T < +\infty, \quad \text{where } \beta = \frac{1 + \gamma + \delta}{1 - \gamma - \delta}.$$

The assumption of the control transmutation method corresponds to $\gamma = \delta = 0$. The Russell-Weiss condition (3) corresponds to $\delta = -1$.

A logarithmic improvement of this theorem is also proved in [1] thanks to this new variant of the Lebeau-Robbiano strategy of [5]:

Theorem 5. Assume the admissibility condition (1) and that $-A$ generates a normal semigroup. If the logarithmic observability condition on spectral subspaces

$$\|v\|^2 \leq a_0 e^{2a\lambda/((\log(\log \lambda))^\alpha \log \lambda)} \|Cv\|^2, \quad v \in \mathbf{1}_{\operatorname{Re} A < \lambda} \mathcal{E}, \quad \lambda \geq \lambda_0.$$

holds with $\alpha > 2$, λ_0 , a_0 , a positive then final-observability (2) holds for all $T > 0$.

REFERENCES

- [1] T. Duyckaerts and L. Miller, *Resolvent conditions for the control of parabolic equations*, manuscript, 2010, soon available online at <http://hal.archives-ouvertes.fr>.
- [2] B. Jacob and H. Zwart, *On the Hautus test for exponentially stable C_0 -groups*, SIAM J. Control Optim. **48** (2009), no. 3, 1275–1288.
- [3] L. Miller, *Controllability cost of conservative systems: resolvent condition and transmutation*, J. Funct. Anal. **218** (2005), no. 2, 425–444.
- [4] L. Miller, *The control transmutation method and the cost of fast controls*, SIAM J. Control Optim. **45** (2006), no. 2, 762–772.
- [5] L. Miller, *A direct Lebeau-Robbiano strategy for the observability of heat-like semigroups*, Discrete Contin. Dyn. Syst. B **14** (2010), no. 4, 1465–1485.
- [6] L. Miller, *Resolvent conditions for the control of unitary groups*, manuscript, 2010.
- [7] D. L. Russell and G. Weiss, *A general necessary condition for exact observability*, SIAM J. Control Optim. **32** (1994), no. 1, 1–23.
- [8] M. Tucsnak and G. Weiss, *Observation and control for operator semigroups*, Birkhäuser Advanced Texts: Basel Textbooks, Birkhäuser Verlag, Basel, 2009.