Thrombosis preventive potential of chicory coffee consumption: a clinical study

Edit Schumacher, Éva Vigh, Valéria Molnár, Péter Kenyeres, Gergely Fehér,
Gábor Késmárky, Kálmán Tóth, János Garai

► To cite this version:

Edit Schumacher, Éva Vigh, Valéria Molnár, Péter Kenyeres, Gergely Fehér, et al.. Thrombosis preventive potential of chicory coffee consumption: a clinical study. *Phytotherapy Research*, 2011, 10.1002/ptr.3481 . hal-00625153

HAL Id: hal-00625153

<https://hal.science/hal-00625153>

Submitted on 21 Sep 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Thrombosis preventive potential of chicory coffee consumption: a clinical study

Journal:	<i>Phytotherapy Research</i>
Manuscript ID:	PTR-10-0363.R2
Wiley - Manuscript type:	Full Paper
Date Submitted by the Author:	19-Feb-2011
Complete List of Authors:	Schumacher, Edit; University of Pécs, Medical School, Department of Pathophysiology and Gerontology; University of Pécs, Medical School, 1st Department of Internal Medicine Vigh, Éva; University of Pécs, Medical School, Department of Pathophysiology and Gerontology Molnár, Valéria; University of Pécs, Medical School, Department of Pathophysiology and Gerontology Kenyeres, Péter; University of Pécs, Medical School, 1st Department of Internal Medicine Fehér, Gergely; University of Pécs, Medical School, 1st Department of Internal Medicine Késmárky, Gábor; University of Pécs, Medical School, 1st Department of Internal Medicine Tóth, Kálmán; University of Pécs, Medical School, 1st Department of Internal Medicine Garai, János; University of Pécs, Medical School, Department of Pathophysiology and Gerontology
Keyword:	caffeic acid, chicory coffee, hemorheological parameters, macrophage migration inhibitor factor, platelet aggregation

SCHOLARONE™
Manuscripts

Thrombosis preventive potential of chicory coffee consumption: a clinical study

Running title: Chicory coffee: a thrombosis preventive beverage?

Edit Schumacher^{1,2}, Éva Vigh^{1,*}, Valéria Molnár¹, Péter Kenyeres², Gergely Fehér²,
Gábor Késmárky², Kálmán Tóth² and János Garai¹

¹ Department of Pathophysiology and Gerontology, Medical School, University of Pécs,
H-7624 Pécs, Hungary

² 1st Department of Internal Medicine, Medical School, University of Pécs, H-7624
Pécs, Hungary

* Correspondence to:

Éva Vigh

Department of Pathophysiology and Gerontology, Medical School, University of Pécs,
H-7624 Pécs, Hungary. Tel.: +36 72 536246; Fax: +36 72 526247

E-mail: eva.vigh@aok.pte.hu

Keywords: caffeic acid, chicory coffee, hemorheological parameters, macrophage
migration inhibitor factor, platelet aggregation

ABSTRACT

Protective effects of plant polyphenol intake on cardiovascular morbidity and mortality are widely acknowledged. Caffeine-free chicory coffee is a rich source of plant phenolics, including caffeic acid, which inhibits in vitro platelet aggregation, and also phenylpyruvate tautomerase enzymatic activity of the proinflammatory cytokine, macrophage migration inhibitory factor (MIF). To assess whether chicory coffee consumption might confer cardiovascular benefits we performed a clinical intervention study with 27 healthy volunteers, who consumed 300 mL chicory coffee every day for one week. The dietary intervention produced variable effects on platelet aggregation, depending on the inducer used for the aggregation test. Whole blood and plasma viscosity were both significantly decreased, along with serum MIF levels, after one week of chicory coffee consumption. Moreover, significant improvements were seen in red blood cell deformability. No changes in hematocrit, fibrinogen level or red blood cell counts were detected.

The full spectrum of these effects is unlikely to be attributable to a single compound present in chicory coffee, nevertheless, the phenolics, including caffeic acid, are expected to play a substantial role. In conclusion, our study offers an encouraging starting-point to delineate the antithrombotic and antiinflammatory effects of phenolic compounds found in chicory coffee.

INTRODUCTION

Diverse biological effects of plant polyphenols have been revealed in countless papers published throughout the last decades. Epidemiological studies suggest that consumption of polyphenol-rich food might impede the onset and the progression of certain human diseases (Kris-Etherton *et al.*, 2002). Phenolic compounds are abundant in fruits, vegetables, coffee, wine and olive oil (Huang *et al.*, 1986). The favourable effects of moderate red wine consumption on certain cardiovascular risk factors have been documented (Fuhrman *et al.*, 1995). The possible benefits of coffee, the other widely popular polyphenol-rich beverage, still await clarification. The risk of acute coronary syndromes was found to follow a J-shaped trend dependent on the dose of coffee consumed per day (Panagiotakos *et al.*, 2003). The upsurge with higher doses might be attributable to elevated blood pressure caused by the caffeine content of coffee brewed from *Coffea arabica* beans. Nevertheless, moderate consumption conferred decreased risk compared with non-consumers. The antithrombotic effect of coffee has been reported to be independent from its caffeine, but rather to be attributable to its phenolic acids able to incorporate into platelets (Natella *et al.*, 2008). The influence of coffee consumption on lifetime cardiovascular risk has been analyzed recently in two large cohorts. A dose dependent protective effect has been found that seems to be more pronounced in female consumers (Lopez-Garcia *et al.*, 2008; Lopez-Garcia *et al.*, 2009). These reports also found the protective effect independent from caffeine.

A cup of coffee is an abundant source of chlorogenic acid, i.e. caffeic acid esterified with quinic acid. Caffeic acid becomes available for absorption following hydrolysis. Known to be absorbed almost totally (~95%) from the small intestine caffeic

acid reaches its peak blood level within 2 hours after oral intake (Olthof *et al.*, 2001; Nardini *et al.*, 2002).

Macrophage migration inhibitory factor (MIF) has been the first non-immunoglobulin immune-mediator described (Bloom and Bennett, 1966; David, 1966). The basal serum concentration of this proinflammatory cytokine is 3-5 ng/mL (Metz and Bucala, 1997). Its level rises in infectious or non-infectious inflammatory conditions such as sepsis (Bernhagen *et al.*, 1993) or rheumatoid arthritis (Onodera *et al.*, 1999). Enzymatic tautomerase (Rosengren *et al.*, 1996; Rosengren *et al.*, 1997) and thiol-protein oxidoreductase (Kleemann *et al.*, 1998) activities of MIF have been revealed recently. Certain antiinflammatory phytochemicals inhibit the tautomerase activity in a concentration dependent manner (Molnar and Garai, 2005). Among these molecules caffeic acid exhibits one of the best inhibitory potential concerning either phenylpyruvate- or dopachrome- tautomerase activities (Molnar and Garai, 2005; Senter *et al.*, 2002). Although the exact role of the enzymatic activity of this cytokine remains to be delineated MIF tautomerase has already attained a reputation as a promising pharmacologic target in inflammatory conditions (Lubetsky *et al.*, 2002). Caffeic acid and its derivatives have been reported to inhibit platelet aggregation *in vitro* and *in vivo* (Hung *et al.*, 2005; Hsiao *et al.*, 2007) – an effect confirmed in our laboratory as well.

Chicory (*Cichorium intybus*) is one of the richest dietary sources of caffeic acid and its derivatives e.g. chlorogenic acid. An alcoholic (40%) extract (1:5) of the whole plant has been reported to contain 12.98 ± 0.06 g% of caffeic acid derivatives (Kocsis *et al.*, 2003). Of note, coffee brewed from ground roasted chicory roots does not contain caffeine and has a long history being used as coffee substitute or admixture (Galasko *et al.*, 1989). In our present study we have addressed the clinical effects of chicory coffee

consumption on platelet aggregation, on hemorheological factors and on serum MIF levels.

MATERIALS AND METHODS

Study population

Our clinical study has been approved by the local Regional Ethical Committee according to the actual amended version of the World Medical Association Declaration of Helsinki. Written informed consent has been obtained from all participants.

In our self-controlled study 27 healthy volunteers (13 women, 14 men) were recruited (mostly university students). The average age was 23 ± 0.4 years. Five of them were smokers and 17 of them were habitual coffee drinkers. The volunteers were asked to refrain from consuming (Arabic) coffee or tea one week before and during the whole period of the study, because these also contain caffeic acid. If it became unavoidable caffeine pills were provided. No vitamins or drugs were allowed to be taken one week before and during the study. The volunteers were also asked to refrain from alcohol consumption.

Intervention and sample collection

The volunteers consumed 300 mL coffee in the morning each day prepared from 20g (3 tablespoons) ground chicory coffee throughout the one week study period. We aimed to observe the effects of both, a single dose and a week long daily consumption of chicory coffee on the following parameters: platelet aggregation, hemorheological factors, fibrinogen and MIF concentration. Blood samples were collected from 27

volunteers on the first day just before and 2 hours after the first coffee consumption in fasting conditions. 24 volunteers also donated blood samples on the eighth day following one week daily chicory coffee consumption. The remaining 3 participants were dropouts. They were unable to consume daily the requested amount of coffee because of taste aversion.

Statistical analysis

Results are expressed as mean $\pm$ SEM. All parameters were normally distributed. Statistical comparisons between 2 time points were performed by one-way Student's *t*-test. The ANOVA Repeated Measures analysis was applied if the parameter was measured at 3 different time points during the study. *p*-values < 0.05 were accepted as statistically significant.

Platelet aggregation

Blood samples were collected from cubital veins in tubes containing 3.8% sodium citrate, and then incubated at 37°C for 20 minutes. Platelet-rich plasma was prepared from venous blood by centrifugation at 900g for 10 minutes. After carefully removing platelet-rich plasma, the remaining specimen was further centrifuged at 3600g for 10 minutes to obtain platelet-poor plasma. Platelet aggregation was measured in Carat TX-4 platelet aggregometer (Carat Ltd., Hungary). Samples were incubated at 37°C and continuously stirred at 1000 rpm during the measurement. 50 μ L of adenosine diphosphate (ADP) (5 and 10 μ M), collagen (2 μ g/mL) or epinephrine (10 μ M) was added to the platelet-rich plasma to induce platelet aggregation. All measurements were carried out within 2 hours after vein puncture.

Whole blood and plasma viscosity

Blood samples were collected in tubes containing lithium heparin as anticoagulant. Plasma and whole blood viscosity values were determined with Hevimet 40 capillary viscometer (Hemorex Ltd; Hungary) at 37°C within 2 hours after blood sampling. In this viscosimeter the flow of whole blood (or plasma) is detected optoelectronically along the capillary tube. Apparent whole blood viscosity values were obtained at 90 1/s shear rate (Toth *et al.*, 1994). Plasma fibrinogen concentration was determined by von Clauss’s method.

Red blood cell deformability

Red blood cell (RBC) deformability was determined at various shear stresses by laser diffraction analysis, using an ektacytometer, LORCA (Laser-assisted Optical Rotational Cell Analyzer, RR Mechatronics, Hoorn, Netherland). The system has been described elsewhere in details (Hardeman *et al.*, 1994). Twenty five µl of EDTA-anticoagulated blood was suspended in a PBS solution of PVP (Poly-Vinyl-Pyrrolidone 360 kD, Sigma-Aldrich, Budapest, Hungary). The viscosity of the PVP solution was 30 ± 2 mPa x s, all measurements were performed at 37°C (Johnson, 1989). Elongation indexes were calculated from the diffraction patterns for shear stresses between 0.3-30 Pa: a higher elongation index indicates greater red blood cell deformation.

Serum MIF levels

Blood samples were collected in tubes without anticoagulant. Serum MIF levels were assessed by ELISA with a Duo set ELISA Development System from R&D

Systems, Minneapolis, MN, US according to the manufacturer's instructions. The amount of MIF was expressed as pmol/mL.

RESULTS

Caffeic acid inhibits platelet aggregation in vitro (Hung *et al.*, 2005). This finding has been confirmed by us as well: 1.2 mM caffeic acid decreased collagen induced platelet aggregation in vitro by 50% (data not shown). 10 μ M ADP induced platelet aggregation has increased after chicory coffee consumption at 2 hours and on the 8th day as well (Fig. 1A). 5 μ M ADP induced platelet aggregation did not change significantly (Fig. 1B). Collagen-induced platelet aggregation decreased from baseline at 2 hours (Fig. 1C), but epinephrine-induced aggregation increased (Fig. 1D).

No changes in basic hematologic parameters like hematocrit or red blood cell counts were detected (Table 1). Whole blood viscosity significantly decreased after one week daily chicory coffee consumption (Fig. 2A). At all fluid shear stresses lower than 30 Pa significant improvements were seen there in RBC deformability, an acknowledged determinant of whole blood viscosity (Table 2). A significant decrease in plasma viscosity was evident after one week of chicory coffee consumption (Fig. 2B), while no changes were detected in fibrinogen levels, one of the main determinants of plasma viscosity (Table 1).

Daily consumption of chicory coffee for one week significantly decreased serum MIF levels (Fig. 3).

DISCUSSION

The participants observed no side effects during the study. Changes in hemorheological parameters detected in the course of this dietary intervention are unlikely to be attributable to only one single compound, but rather to an array of agents absorbed from chicory coffee. The inhibitory effect of chicory consumption on platelet aggregation seems to vary with the inductor used. This might be attributable to a particular mechanism by which agents in chicory coffee interfere with platelet aggregation. Nevertheless, in vivo antiplatelet activity was found much weaker than expected from preceding in vitro studies with caffeic acid, one of the most abundant plant phenolic of this beverage. Consumption of chicory coffee for one week, however, significantly improved hemorheological parameters, like plasma- and whole blood viscosity and red blood cell deformability. These changes indicate that daily chicory coffee consumption might be preventive against certain microcirculatory pathologies. The decrease in whole blood viscosity might largely be attributable to the increased red blood cell deformability, however, the modest but significant decrease in plasma viscosity in the face of unchanged fibrinogen levels appears perplexing. Further studies are warranted to clarify the exact mechanisms and to identify the compounds behind these effects of chicory coffee.

Baskurt *et al.* (1998) found significantly decreased RBC deformability in septic states at fluid shear stresses less than 5 Pa in humans and rats. Elevated MIF levels are acknowledged indicators of septic state (Bernhagen *et al.*, 1993). Potato peel extract rich in caffeic acid has been reported to protect erythrocytes against oxidative damage, a known cause of impaired cell deformability (Singh and Rajini, 2008). Daily

consumption of chicory coffee rich in antiinflammatory phenols (including caffeic acid) for one week duration substantially decreased serum MIF levels of healthy volunteers in our study in parallel with the improved red blood cell deformability. To decide whether these phenomena are related or are merely coincidental further studies are needed. Notably MIF upregulates adhesion molecules on endothelial cells (Amin *et al.*, 2006), hence suppression of MIF levels could be regarded preventive against monocyte adhesion and emigration, a key element in initiation of the atherosclerotic process.

In conclusion, our study constitutes an encouraging starting point to investigate further the antithrombotic, antiinflammatory and beneficial hemorheologic effects of phenolic compounds from chicory coffee.

Acknowledgement

We are indebted to the volunteers participating in the study. The complimentary sample of ground roasted chicory provided to this study by Multi-Cikória Kft. Hungary is gratefully acknowledged. The authors have no financial or commercial conflict of interest.

REFERENCES

- Amin MA, Haas CS, Zhu K *et al.* 2006. Migration inhibitory factor up-regulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 via Src, PI3 kinase, and NFkappaB. *Blood* **107**: 2252–2261.
- Baskurt OK, Gelmont D, Meiselman HJ. 1998. Red blood cell deformability in sepsis. *Am J Respir Crit Care Med* **157**: 421–427.

- Bernhagen J, Calandra T, Mitchell RA *et al.* 1993. MIF is a pituitary-derived cytokine that potentiates lethal endotoxaemia. *Nature* **365**: 756–759.
- Bloom BR, Bennett B. 1966. Mechanism of a reaction in vitro associated with delayed-type hypersensitivity. *Science* **153**: 80–82.
- David JR. 1966. Delayed hypersensitivity in vitro: its mediation by cell-free substances formed by lymphoid cell-antigen interaction. *Proc Natl Acad Sci USA* **56**: 72–77.
- Fuhrman B, Lavy M, Aviram M. 1995. Consumption of red wine with meals reduces the susceptibility of human plasma and low-density lipoprotein to lipid peroxidation. *Am J Clin Nutr* **61**: 549–554.
- Galasko GT, Furman KI, Alberts E. 1989. The caffeine contents of non-alcoholic beverages. *Food Chem Toxicol* **27**: 49–51.
- Hardeman MR, Goedhart PT, Dobbe JGG, Lettinga KP. 1994. Laser-assisted optical rotational cell analyzer (LORCA). 1. A new instrument for measurement of various structural hemorheological parameters. *Clin Hemorheol* **14**: 605–618.
- Hsiao G, Lee JJ, Lin KH *et al.* 2007. Characterization of a novel and potent collagen antagonist, caffeic acid phenethyl ester, in human platelets: In vitro and in vivo studies. *Cardiovasc Res* **75**: 782–792.
- Huang HM, Johanning GL, O'Dell BL. 1986. Phenolic acid content of food plants and possible nutritional implications. *J Agric Food Chem* **34**: 48–51.
- Hung CC, Tsai WJ, Kuo LM, Kuo YH. 2005. Evaluation of caffeic acid amide analogues as anti-platelet aggregation and anti-oxidative agents. *Bioorg Med Chem* **13**: 1791–1797.
- Johnson RM. 1989. Ektacytometry of red blood cells. *Methods Enzymol* **173**: 35–54.

- Kleemann R, Kapurniotu A, Frank RW *et al.* 1998. Disulfide analysis reveals a role for macrophage migration inhibitory factor (MIF) as thiol-protein oxidoreductase. *J Mol Biol* **280**: 85–102.
- Kocsis I, Hagymasi K, Kery A, Szoke E, Blazovics A. 2003. Effects of chicory on pancreas status of rats in experimental dislipidemia. *Acta Biol Szeged* **47**: 143–146.
- Kris-Etherton PM, Hecker KD, Bonanome A *et al.* 2002. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Am J Med* **113**: 71S–88S.
- Lopez-Garcia E, van Dam RM, Li TY, Hu FB. 2008. The relationship of coffee consumption with mortality. *Ann Intern Med* **148**: 904–914.
- Lopez-Garcia E, Rodriguez-Artalejo F, Rexrode KM, Logroscino G, Hu FB, van Dam RM. 2009. Coffee consumption and risk of stroke in women. *Circulation* **119**: 1116–1123.
- Lubetsky JB, Dios A, Han J *et al.* 2002. The tautomerase active site of macrophage migration inhibitory factor is a potential target for discovery of novel anti-inflammatory agents. *J Biol Chem* **277**: 24976–24982.
- Metz CN, Bucala R. 1997. Role of macrophage migration inhibitory factor in the regulation of the immune response. *Adv Immunol* **66**: 197–223.
- Molnar V, Garai J. 2005. Plant-derived anti-inflammatory compounds affect MIF tautomerase activity. *Int Immunopharmacol* **5**: 849–856.
- Nardini M, Cirillo E, Natella F, Scaccini C. 2002. Absorption of phenolic acids in human after coffee consumption. *J Agric Food Chem* **50**: 5735–5741.
- Natella F, Nardini M, Belelli F *et al.* 2008. Effect of coffee drinking on platelets: inhibition of aggregation and phenols incorporation. *Brit J Nutr* **100**: 1276–1282.

- 1
2
3
4
5 Olthof MR, Hollman PC, Katan MB. 2001. Chlorogenic acid and caffeic acid are
6
7 absorbed in humans. *J Nutr* **131**: 66–71.
8
9 Onodera S, Tanji H, Suzuki K *et al.* 1999. High expression of macrophage migration
10
11 inhibitory factor in the synovial tissues of rheumatoid joints. *Cytokine* **11**: 163–167.
12
13 Panagiotakos DB, Pitsavos C, Chryschoou C, Kokkinos P, Toutouzas P, Stefanadis C.
14
15 2003. The J-shaped effect of coffee consumption on the risk of developing acute
16
17 coronary syndromes: The CARDIO2000 case-control study. *J Nutr* **133**: 3228–
18
19 3232.
20
21
22
23 Rosengren E, Bucala R, Aman P *et al.* 1996. The immunoregulatory mediator
24
25 macrophage migration inhibitory factor (MIF) catalyzes a tautomerization reaction.
26
27 *Mol Med* **2**: 143–149.
28
29
30 Rosengren E, Aman P, Thelin S *et al.* 1997. The macrophage migration inhibitory factor
31
32 MIF is a phenylpyruvate tautomerase. *FEBS Lett* **417**: 85–88.
33
34
35 Senter PD, Al-Abed Y, Metz CN *et al.* 2002. Inhibition of macrophage migration
36
37 inhibitory factor (MIF) tautomerase and biological activities by acetaminophen
38
39 metabolites. *Proc Natl Acad Sci USA* **99**: 144–149.
40
41
42 Singh N, Rajini PS. 2008. Antioxidant-mediated protective effect of potato peel extract
43
44 in erythrocytes against oxidative damage. *Chem Biol Interact* **173**: 97–104.
45
46
47 Toth K, Habon T, Horvath I, Mezey B, Juricskay I, Mozsik G. 1994. Hemorheological
48
49 and hemodynamical parameters in patients with ischemic heart disease at rest and
50
51 at peak exercise. *Clin Hemorheol* **14**: 329–338.
52
53
54
55
56
57
58
59
60

Table 1. Effect of chicory coffee consumption on basic hematologic parameters.Mean values $\pm$ SEM.

	Control (n=27)	1 week (n=24)
Red blood cell count (T/L)	4.79 $\pm$ 0.09	4.74 $\pm$ 0.09
Hemoglobin (g/L)	143.81 $\pm$ 2.57	142.33 $\pm$ 2.84
Hematocrit (%)	41.82 $\pm$ 0.59	41.38 $\pm$ 0.65
Fibrinogen (g/L)	2.63 $\pm$ 0.09	2.65 $\pm$ 0.11
Platelet count(G/L)	241.93 $\pm$ 8.04	241.67 $\pm$ 9.25

Table 2. Effect of chicory coffee consumption on red blood cell deformability. The reported elongation indexes represent RBC deformability at different shear stresses.

Mean values $\pm$ SEM.

Shear stress (Pa)	Control (n=27)	2 hours (n=27)	1 week (n=24)
0.3	0.0545 $\pm$ 0.0018	0.0532 $\pm$ 0.0017	0.0594 $\pm$ 0.0018** ##
0.53	0.1094 $\pm$ 0.0027	0.1089 $\pm$ 0.0024	0.1141 $\pm$ 0.0030* #
0.95	0.2067 $\pm$ 0.0027	0.2065 $\pm$ 0.0029	0.2120 $\pm$ 0.0033** #
1.69	0.3176 $\pm$ 0.0027	0.3177 $\pm$ 0.0027	0.3235 $\pm$ 0.0028* #
3	0.4194 $\pm$ 0.0019	0.4186 $\pm$ 0.0023	0.4240 $\pm$ 0.0024* #
5.33	0.4974 $\pm$ 0.0014	0.4966 $\pm$ 0.0020	0.5015 $\pm$ 0.0019** ##
9.49	0.5497 $\pm$ 0.0013	0.5493 $\pm$ 0.0018	0.5533 $\pm$ 0.0016** ##
16.87	0.5961 $\pm$ 0.0013	0.5957 $\pm$ 0.0016	0.5985 $\pm$ 0.0015** ##
30	0.6276 $\pm$ 0.0008	0.6277 $\pm$ 0.0011	0.6290 $\pm$ 0.0011

*difference from baseline $p < 0.05$, **difference from baseline $p < 0.01$

#difference from 2 hours $p < 0.05$, ##difference from 2 hours $p < 0.01$

Schumacher 1 ↑

Figure 1.

Effect of chicory coffee consumption on platelet aggregation.

A: Platelet aggregation induced by 10 μ M ADP. B: Platelet aggregation induced by 5 μ M ADP. C: Platelet aggregation induced by 2 μ g/ml collagen. D: Platelet aggregation induced by 10 μ M epinephrine.

Mean values $\pm$ SEM. *p < 0.05, ** p < 0.01
1134x815mm (150 x 150 DPI)

Schumacher 2 ↑

Figure 2.
Effect of chicory coffee consumption on blood and plasma viscosity.
A: Changes in whole blood viscosity. *p < 0.05, **p < 0.01; B: Changes in plasma viscosity. *p < 0.05
Mean values ± SEM.

845x1368mm (150 x 150 DPI)

Schumacher 3 ↑

Figure 3.
Effect of chicory coffee consumption on serum MIF levels.
** $p < 0.01$, + $p < 0.06$
Mean values $\pm$ SEM
805x690mm (150 x 150 DPI)