Effect of tidal flooding on metal distribution in pore waters of marsh sediments and its transport to water column (Tagus estuary, Portugal)

Juan Santos-Echeandía, Carlos Vale, Miguel Caetano, Patrícia Pereira,
Ricardo Prego

► To cite this version:

Juan Santos-Echeandía, Carlos Vale, Miguel Caetano, Patrícia Pereira, Ricardo Prego. Effect of tidal flooding on metal distribution in pore waters of marsh sediments and its transport to water column (Tagus estuary, Portugal). *Marine Environmental Research*, 2010, 70 (5), pp.358. 10.1016/j.marenvres.2010.07.003 . hal-00631692

HAL Id: hal-00631692

<https://hal.science/hal-00631692>

Submitted on 13 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Effect of tidal flooding on metal distribution in pore waters of marsh sediments and its transport to water column (Tagus estuary, Portugal)

Authors: Juan Santos-Echeandía, Carlos Vale, Miguel Caetano, Patrícia Pereira, Ricardo Prego

PII: S0141-1136(10)00112-1

DOI: [10.1016/j.marenvres.2010.07.003](https://doi.org/10.1016/j.marenvres.2010.07.003)

Reference: MERE 3465

To appear in: *Marine Environmental Research*

Received Date: 1 June 2010

Revised Date: 16 July 2010

Accepted Date: 18 July 2010

Please cite this article as: Santos-Echeandía, J., Vale, C., Caetano, M., Pereira, P., Prego, R. Effect of tidal flooding on metal distribution in pore waters of marsh sediments and its transport to water column (Tagus estuary, Portugal), *Marine Environmental Research* (2010), doi: 10.1016/j.marenvres.2010.07.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Effect of tidal flooding on metal distribution in pore waters of marsh sediments and its transport to water column (Tagus estuary, Portugal)

Juan Santos-Echeandía^{a,b,*}, Carlos Vale^b, Miguel Caetano^b, Patrícia Pereira^b and Ricardo Prego^a

^aMarine Research Institute (CSIC), Department of Oceanography, Eduardo Cabello 6, 36208, Vigo, Spain

^bIPIMAR, National Institute of Biological Resources, Avenida Brasília 1446-006, Lisboa, Portugal

Carlos Vale: cvale@ipimar.pt

Miguel Caetano: mcaetano@ipimar.pt

Patrícia Pereira: patbio@ipimar.pt

Ricardo Prego: prego@iim.csic.es

***Corresponding author:** Juan Santos Echeandía

INRB/IPIMAR – National Institute of Biological Resources

Department of Environment and Biodiversity

Av. Brasília, 1449-006 Lisboa, Portugal

jecheandia@ipimar.pt

TLF: +351.21.3027073

FAX: +351.21.3015948

Abstract

Sediment cores and flooding water were collected at 0, 5, 10 and 50 minutes of tidal inundation in two sites of the Rosário salt marsh located in the proximity of a heavy industrialised zone of Tagus estuary colonised by pure stands of *Spartina maritima* (low marsh) and *Sarcocornia fruticosa* (high marsh). The cores were sliced in 5-cm layers and sediment solids, pore water, and belowground biomass were separated in order to measure Fe, Mn, Zn, Cu, Pb and Cd. The pore waters and sediments colonised by *S. fruticosa*, as well as belowground biomass presented high concentrations of Zn, Cu, Pb and Cd. Belowground biomass exceeded in one order of magnitude the metal levels in sediments. Abundant belowground biomass and small dimension of *S. fruticosa* roots facilitates the root-sediment interactions and presumably the metal retention in the higher marsh. The novelty of this work is the result of tidal inundation on pore water concentrations of metals in salt marsh sediments and their exportation to the water column. Concentrations in pore waters varied at minute scales, but 50 minutes after inundation levels were comparable to the initial values. The metal levels in flooding water increased abruptly during the first 10 to 20 minutes of inundation. The concentration peaks (Fe=60 µM, Mn=7.5 µM, Zn=1.7 µM, Cu=550 nM, Pb=100 nM, Cd=1.7 nM) reached one to two orders of magnitude above the values found in subsequent periods of inundation. The advective transports during the 50-min inundation during two daily pulses of inundation were: Fe (9520 and 1640), Mn (24), Zn (220 and 82), Cu (74 and 16), Pb (13 and 15) and Cd (0.3 and 0.08) for *S. maritima* and *S. fruticosa*, respectively. These quantities exceeded 3 to 4 orders of magnitude of the corresponding predicted diffusive fluxes (Fick 1st law) on a daily basis. This work emphasizes the importance of tidal flooding over salt marsh sediments to the metal exportation to the water column.

Keywords: Salt marsh; metals; pore-waters; Tagus estuary; advective; diffusive; fluxes; tidal flooding.

1. Introduction

The post-mobilisation of deposited iron, manganese and trace elements has been extensively studied in deep and coastal sediments (e.g., review Sundby, 2006). On the basis of vertical properties of sediment pore water (pH, redox potential, dissolved oxygen, sulphur compounds, nutrients and metals) and of metal concentrations in solid sediments, it was proposed a sequence of reactions for the oxidation of the labile sedimentary organic matter that reaches the sediment surface (Aller, 1977; Froelich et al., 1979). Oxygen diffused from the water column into the upper sediment layer is the most favourable oxidant, and as concentration decreases with the depth other oxidants are consumed to degrade the organic matter. Reactions of similar free energy like nitrate and manganese oxide may overlap, (Burdige, 1993; van Capellen and Yang, 1996). The chemical zonation in the upper sediments is characterised by gradients with the depth, concentrations in pore waters exceeding largely the levels in overlying water. These differences result in diffusive fluxes of solutes from sediments to water column (Berner, 1980). The presence and activity of macroinvertebrates, causing perturbation and irrigation in the upper sediments, enhances the sediment-water exchanges of solutes (Brune et al., 2000). The pore waters of inter-tidal permeable sediments are renewed as tidal water floods the area causing exports of solutes by pulses (Falcão and Vale, 1990; Caetano et al., 2007).

The salt marsh sediments have specificities with relevance to early diagenesis and sediment-water exchanges such as: deliver of oxygen into subsurface layers of sediments by plant roots (Clothier and Green, 1997; Caetano and Vale, 2002); supply of organic matter as roots dye (Caçador et al., 2004; Pereira et al., 2007); presence of a mosaic of small channels in the upper sediments where tidal water excurses during flooding. Works have reported two situations: anaerobic conditions in the rooting sediments, meaning that delivered oxygen was insufficient to oxidise the large amount of degradable organic matter (Haines and Dunn, 1985; Mendelssohn et al., 1995; Sundby et al., 2003); oxic to suboxic conditions suggesting an excess of oxygen supplied to the sediment for the requirements. Various studies have been focused on the influence of root-sediment interactions on sediment chemistry (Sundby et al., 2003; 2005; Taillefert et al., 2007; Koretsky et al., 2008a,b) and response of plants (Windham et al., 2001; Jacob and Otte, 2003; Weis and Weis, 2004; Caetano et al., 2007; 2008). However, sediment-water exchanges in salt marsh are poorly documented (Taillefert et al., 2007).

Several studies showed that Tagus salt marshes, including Rosário, incorporate large quantities of anthropogenic metals into the colonised sediments (e.g. Caçador et al., 1993; 1996; 2000; Sundby et al., 1998; Caetano et al., 2007). Retention of contaminated particles from the anthropogenic sources (Vale, 1990) and root-sediment interactions appear to contribute to the metal enrichment of colonised sediments and belowground biomass (Caçador et al., 1993; 1996). Recent studies in Rosário have shown that as root dye the accumulated and sorbed metals are redistributed in the sediments (Pereira et al., 2007). Few works have been focused on Tagus marshes pore waters. Reboreda and Caçador (2007) have reported high concentrations of trace metals in bulk pore waters of rooting sediment. By using an amalgam voltammetric microelectrode Sundby et al. (2003; 2005) proved that dissolved oxygen, Fe, Mn and Pb concentrations in pore waters of Rosário salt marsh sediments vary seasonally.

This paper reports the changes of Fe, Mn, Zn, Cu, Pb and Cd concentrations in pore water and flooding water during the first 50 minutes of the tidal inundation over sediments of the Tagus salt marsh (Rosário) colonised by *Spartina maritima* and *Sarcocornia fruticosa*. On the basis of the registered concentrations it was estimated the diffusive fluxes between sediments and overlying water during submerged periods. In addition, it was calculated the amounts exported from the colonised sediments during the first 50 minutes of tidal inundation. The magnitude of the two contributions to the water column is discussed.

2. Material and methods

2.1. Study area

The 340 km² of the Tagus estuary represent one of the largest transitional systems in Europe. Almost 40% of the estuary is composed by inter-tidal mudflats and its southern and eastern shores contain extensive areas of salt marshes colonised by *Sarcocornia fruticosa*, *Sarcocornia perennis*, *Halimione portulacoides* and *Spartina maritima*. The marsh selected for this study (Rosário) is located in the southern shoreline of the estuary (Figure 1). It covers an area of 200 ha (Crespo, 1993), being characterized by homogeneous stands of *S. maritima* as a pioneer species in the lower part, pure stands of *H. portulacoides* across the 20-50 cm elevation transect, and *S. fruticosa* and *S. perennis* in the higher salt marsh. The marsh is fully inundated twice a day by tidal action (2-4m of tidal amplitude) through a highly branched system of channels that cross the elevation transect. The channels have 0.5 to 1.5 m depth promoting the inundation of the higher marsh even at low amplitude tides. The study area is located in the proximity of a heavy industrialised zone that includes several chemical plants. During the past four decades these industries discharged effluents enriched in various contaminants into the estuary that were transported towards the Rosário marsh by the semi-diurnal tidal flooding.

2.2. Sampling strategy

Sediment cores of 30-cm long were collected, as well as flooding water over the short period of tidal inundation in the two sites of the Rosário salt marsh colonised by pure stands of *S. fruticosa* (high marsh) and *S. maritima* (low marsh) and separated by less than 5 meters. Sampling took place in March 2008. At low tide, four sediment cores were collected at each site, when sediment was exposed to the atmosphere, and water retained in the main channel of the marsh was sampled by hand into acid pre-cleaned syringes. One of the cores was sampled at each site to measure temperature, dissolved oxygen and salinity all along its depth. When tidal water starts to flood each site, two additional sediment cores and flooding water were collected at each time of inundation: 5, 10 and 50 minutes. Flooding water was sampled 1 cm above the sediment surface directly into acid pre-cleaned syringes. All water samples were acidified with suprapure HCl (pH < 2). The cores were sliced immediately after sampling in 5-cm layers, prepared composite samples for each layer of the cores collected at each site, and material stored in acid precleaned HDPE vials avoiding air presence inside. Sampling took place in less than three minutes. Core material consisted of dense rooting sediments with no evidence of burrowing worms, crabs or bivalves. The water and sediment samples were kept in refrigerated boxes and immediately transported to the laboratory. A more detailed description of these sampling procedures can be found in previous works (Caetano et al, 2007).

2.3. Sample treatment

Pore waters were separated from the sediment layers by centrifugation at 10160 rcf x g for 30 minutes at +4° C and filtered through 0.45 µm cellulose acetate membranes. Filtration and subsequent manipulation

of the samples were carried out in a glove box under argon atmosphere in order to avoid alteration of the initial conditions (Caetano et al., 2007; Santos-Echeandia et al., 2009). Pore-water samples were acidified with suprapure HCl (pH < 2). The remaining solid material of each sediment layer was washed with Milli-Q water (18.2 MΩ.cm) smoothly using a teflon sieve of 212-μm mesh size and tweezers to separate belowground biomass from sediment particles. Sediment particles and belowground biomass of each layer was oven dried at 40 °C and homogenised with an agate mortar. Procedures were done in order to subsequent determinations of total Fe, Mn, Zn, Cu, Pb and Cd concentrations in water, sediment and biological samples.

2.4. Methods

Dissolved oxygen was measured in sediment layers using a Diamond Electro-Tech needle electrode following the method described in Brotas et al. (1990). Temperature was measured with a Pt100 sensor. The salinity of flooding and pore waters was measured using a conductivity electrode. Each composite sample was divided in three subsamples and analyzed independently for trace metals. Total dissolved Mn and Fe in overlying, flooding and pore waters were determined by graphite furnace atomic absorption spectrometry (Varian SpectrAA Zeeman 220). Concentrations were determined using the interpolation method in a previously designed calibration curve. Detection limits (defined as three times the standard deviation) were 2.6 nM for Fe and 1.9 nM for Mn. The reproducibility, measured as the RSD was always lower than 5%. The rest of total dissolved metals (Zn, Cu, Pb and Cd) were determined by means of stripping voltammetry using a Metrohm VA-694 equipped with a hanging mercury drop electrode (HMDE) as the working electrode, Ag/AgCl as the reference electrode, and a Pt wire as the counter-electrode. Prior to determination, samples were diluted (when necessary) and UV-digested for 1 h using UV-Digester equipped with a high-pressure mercury lamp of 200 W (Achterberg and Van Den Berg., 1994). The simultaneous determination of Zn, Cu, Pb and Cd in the dissolved phase was carried out using the method of standard additions by means of differential pulse anodic stripping voltammetry (DPASV). One blank was run every five samples and results were blank-corrected. The accuracy of the analytical procedure was assessed by the analysis of three different certified reference materials (CASS-4, SLEW-3 and SLRS-4), obtaining good agreement with the certified concentrations (Table 1). The detection limits of the analytical procedure, defined as three times the standard deviation of the blanks, were 0.5 nM for Zn, 0.7 nM for Cu, 22 pM for Pb and 20 pM for Cd. The precision of the analysis measured as relative standard deviation was: 5–10% for Zn (at 20 nM), 5–10% for Cu (at 10 nM), 5–10% for Pb (at 300 pM) and 25–40% for Cd (at 60 pM).

Approximately 100 mg of sediment was mineralized completely with 6 cm³ of HF (40%) and 1 cm³ of Aqua-Regia (HCl-36%:HNO₃-60%; 3:1) in closed Teflon bombs at 100°C during 1 hour. The bomb contents were evaporated to near dryness in Teflon vials, redissolved with HNO₃, heated for 20 min at 75 °C and diluted to 50 cm³ with Milli-Q water (Caetano et al., 2008). Plant material (≈200 mg) was digested with a mixture of HNO₃ (60%) and H₂O₂ (30%) in Teflon bombs at 60 °C for 12 hours and at 100 °C for 1 hour. Two procedural blanks were prepared using the same analytical procedure and reagents, and included within each batch of 20 samples. Iron, Mn and Zn were analysed by flame atomic absorption spectrometry (Perkin Elmer AA100) with an air-acetylene flame. Metal concentrations were determined with the standard additions method. Concentrations of Cu, Pb and Cd were determined using a quadrupole ICP-MS (Thermo Elemental, X-Series) equipped with a Peltier Impact bead spray chamber and a concentric Meinhard nebulizer. Coefficients of variation for metal counts (n=5) varied between 0.5 and 2%. The precision and accuracy of each metal concentration measurements, determined through repeated analysis of certified references materials (MESS2, PACS2 and MAG1 for sediment and BCR60

and BCR61 for plant material) (Table 1), using In as internal standard, were 1-4% and 2-5%, respectively (data not shown). Procedural blanks always accounted for less than 1% of the total metal concentrations in samples.

2.5. Estimation of sediment-water exchange

Different levels of trace metals between pore waters of upper sediment layers and overlying water imply their export/import to/from the water column. Since inter-tidal sediments are alternatively covered with water and exposed to the atmosphere, two types of fluxes should be considered: diffusive fluxes during the submerged period and advective transport associated with the tidal flooding.

Diffusive flux. On the basis of the measured concentration profiles, the benthic fluxes of trace metals are predicted by Fick's first law of diffusion (Berner, 1980):

$$F_{dif} = \phi D_s (C_0 - C_p) / \Delta x \quad (1)$$

where ϕ is the porosity, D_s is the diffusion coefficient, $C_0 = [Me]$ in pore water of the topmost sediment layer, $C_p = [Me]$ in overlying water and x is the space co-ordinate, negative into sediment and origin at sediment-water interface. Sediment porosity (ϕ) was calculated from the following equation:

$$\phi = m_{PW} / (m_{PW} + m_s/2.65 + m_R/1.2) \quad (2)$$

where m_{PW} represents the pore water mass, m_s the sediment solids mass and m_R the roots mass. The 2.65 value is the density of the sediment considering to be formed by pure mud (Berner, 1980) and the 1.2 is the density of the roots. The diffusion coefficient (D_s) was calculated from the following equation:

$$D_s = \phi^2 D_0 \quad (3)$$

where D_0 represents the diffusion coefficient of a metal at a given temperature (calculated from Li and Gregory, 1974).

Advective flux. As tidal water floods over inter-tidal sediments the pore water solutes tends to be exported. The associated pressure difference and tidal water movement are the driven mechanism for this export. By using the temporal variation of trace metals concentrations in the flooding water the advective transport (T) to the water column are calculated using the following expression (Caetano et al., 2007):

$$T = \sum [(C_{t+1} - C_t) / 2 - C_i] (h_{t+1} - h_t) \quad (4)$$

where C_{t+1} and C_t are $[Me]$ in the flooding water at times $t+1$ and t and h_{t+1} , C_i is the residual concentration (lowest value obtained for the flooding water) and h_t are the water depth at the same times. It was observed that water depth during the field measurements increased on the average 0.5 and 1 cm each minute of inundation in *S. fruticosa* and *S. maritima* areas, respectively. The transport of trace metals is calculated for the first 50 minutes of inundation. Since inter-tidal sediments are inundated twice a day, the advective daily flux (T) was multiplied by a factor of two.

2.6. Statistics

Statistical software (Statistica 7.0) was used for statistical analyses. Considering the number of samples, analysis was performed by the non-parametric Kruskal-Wallis test in order to detect metal concentrations differences in sediment solids, belowground biomass and pore water. Correlations between parameters were searched by Spearman rank correlations ($p < 0.05$).

3. Results

3.1. Metal concentrations in rooting sediments

The rooting sediments from the two sampling sites contained between 10.2 and 12.3% of Al, a proxy for the preponderance fine grain particles (Windom et al, 1989). The sediments colonised by *S. fruticosa* presented significantly higher levels of Pb ($p<0.05$), Zn, Cu and Cd ($p<0.1$) than by *S. maritima* (table 2). Maximum concentrations reached 290, 2408, 83 and $4.1 \mu\text{g g}^{-1}$, respectively. The depth variation of the metal concentration differed in the two sampling sites. Whereas the first 15-cm depth of sediments colonised by *S. maritima* exhibited elevated concentrations of Cu and Pb, the 15-25 cm layers of sediments colonised by *S. fruticosa* showed enhanced levels of Cd and Zn.

3.2. Plant belowground biomass

Belowground biomass of the two plants was concentrated in the upper 15-cm depth, decreasing uniformly with the depth (table 3). The belowground biomass of *S. fruticosa* 3873 g m^{-2} , calculated as the amount present the upper 25-cm, exceeded 1292 g m^{-2} found for *S. maritima*. The roots of *S. maritima* were thicker (maximum diameter of 3 mm) than the roots of *S. fruticosa* that were usually $<1 \text{ mm}$. These observations combined with biomass point to a higher specific surface area of the belowground biomass of *S. fruticosa*.

3.3. Metal concentrations in belowground biomass

The concentrations of Cu, Cd and Pb differed significantly ($p<0.05$ and $p<0.1$) in the belowground biomass of the two plants (Table 3). The metal concentration varied with the depth, although levels were not correlated to the quantity of roots presented at each layer. In fact, roots in the first 5-cm sediment layer contained generally lower metal concentrations than in deeper layers with elevated biomass. The most noticeable aspect is the accumulated levels of metals (Cu, Pb and Cd) in roots being one order of magnitude above the concentrations found in the corresponding rooting sediments (tables 2 and 3). The maximum concentrations ($\mu\text{g g}^{-1}$) in belowground biomass of *S. maritima* and *S. fruticosa* were: 3571 and 11017 (Zn), 212 and 334 (Cu), 1528 and 2739 (Pb) and 19 and 30 (Cd), respectively.

3.4. Pore water characteristics of sediments under air-exposed conditions

Temperature of air-exposed sediments in the sampling period (early morning) ranged within a narrow interval ($14\text{--}15^\circ\text{C}$) and showed no substantial differences between higher and lower marsh, and with the depth. Table 4 presents the maximum and minimum values of parameters in pore waters of sediments colonised by the two plants under air-exposed conditions. Salinity varied from lower ($32.4\text{--}33.6\%$) to higher marsh ($28.5\text{--}31.8\%$), reflecting freshwater drainage to the upper area colonised by *S. fruticosa*. In both cores dissolved oxygen remained $< 2 \mu\text{M}$ at all depths. Maximum values of dissolved Fe and Mn in sediments colonised by *S. maritima* and *S. fruticosa* were in the same order of magnitude (12.7 and $65.3 \mu\text{M}$ of Fe and 34.4 and $14.5 \mu\text{M}$ of Mn, respectively). On the contrary, dissolved levels of the analysed trace elements were one order of magnitude higher in the area colonised by *S. fruticosa*. The differences had higher significance for Zn, Cu and Pb ($p<0.001$) than for Cd ($p<0.05$). Concentrations in pore waters of *S. maritima* and *S. fruticosa* reached: 1.2 and $20.6 \mu\text{M}$ (Zn), 28 and 201 nM (Cu), 15 and 268 nM (Pb), 1.4 and 75 nM (Cd), respectively.

3.5. Time variation of metal concentrations in pore waters

Figures 2 and 3 compare the depth metal concentrations in pore waters of sediments at air-exposed conditions ($t=0$), 5, 10 and 50 minutes after the tidal flooding of the marsh. Vertical concentrations at these

periods of time varied with the analysed elements at the two surveyed sites. It is remarkable that pore waters at $t=5$ min. of the surface sediment layers in the marsh colonised by *S. maritima* exhibited lower metal concentrations than at $t=0$ (Figure 2). The differences were less clear for the higher marsh covered with *S. fruticosa* (Figure 3), meaning that tidal inundation had a lower effect on the modification of pore water chemistry. The profiles at 10 and 50 minutes inundation pointed to establishing conditions towards those registered at air-exposed period. This tendency was better observed in the lower marsh. The shape of profiles differs in the two sites. Whereas pore waters from the upper layers of the lower marsh showed higher metal concentrations, sub-surface maxima were found in the pore waters of sediments covered by *S. fruticosa*.

3.6. Metal concentrations in flooding water

The levels of Fe, Mn, Zn, Cu, Pb and Cd in the dissolved fraction of flooding water over the two sites of the marsh increased abruptly in the first 2 to 5 minutes of inundation (figure 4). The sharp concentration peaks (Fe=60 μ M, Mn=7.5 μ M, Zn=2.5 nM, Cu=550 nM, Pb=100 nM, Cd=1.7 nM) reached one to two orders of magnitude above the values found in subsequent periods of inundation. Furthermore, these maximum values are within the range of levels registered in pore waters (table 4). The maximum of Fe, Mn, Cu and Cd concentrations in the area colonised by *S. fruticosa* exceeded the values observed in lower marsh (*S. maritima*).

4. Discussion

A noticeable remark of the present work is the consistent differences of belowground biomass and Zn, Cu, Pb and Cd concentrations in solid fine sediments, roots and pore waters between sites colonised by *S. maritima* and *S. fruticosa*. Higher values were registered in the sediment column of the upper marsh covered by *S. fruticosa*. Despite the daily flooding of the full marsh by the tide, differences of metal concentrations in pore waters stand.

4.1. The influence of belowground biomass on metal retention

The sediments of the Rosário marsh and adjacent inter-tidal flats consist of fine particles contaminated by discharges of the industrial complex localised in the southern part of the Tagus estuary. Various works have pointed to the retention of these contaminants in inter-tidal flats (Vale, 1990) and in the adjoining salt marshes (Caçador and Vale, 2001; Caetano et al., 2007). However, the levels of anthropogenic metals in marsh sediments exceed largely the values found in the non-vegetated sediments (Caçador et al., 1993, 1996a,b, 2000). The enrichment is confined to the rooting sediments and varies with the plant species that cover the marsh sediment. The differences found on metal concentrations between sediments colonised by *S. maritima* and *S. fruticosa* corroborate the previous findings.

The iron-oxides plaque found on roots of plants in northern European and US marshes (Otte et al., 1989; Mendelssohn et al., 1995; St-Cyr and Campbell, 1996; Cambrollé et al., 2008) increases the thickness in Tagus marshes to form hollow cylinders around the roots (Vale et al., 1990). The growth of these structures results from the reaction between O_2 , which is delivered by roots during most time of the year, and Fe (II) in pore waters that diffuses towards the roots (Sundby et al., 1998)). As observed in other marshes (St-Cyr and Crowder 1990, Otte 1991, St-Cyr and Campbell 1996), trace elements are incorporated in the iron-oxides (Vale et al., 1990; Caetano and Vale, 2002; Vale et al., 2003). However, significant relationships between trace elements and Fe concentration in the bulk rooting sediments of the two sites were not found ($p \geq 0.05$), suggesting that Fe precipitates are not the major driver to retain trace

elements in sediments colonised by *S. fruticosa* and *S. maritima*. Enhanced levels of total Fe and Mn in the 0-5 cm layer may indicate the precipitation of oxides by the O_2 diffused from the surface rather than delivered by roots. Since these plants have a short or absence of dormancy in winter (Catarino and Caçador, 1981) it is unlikely a stop on the O_2 delivery. Therefore the lack of oxygen observed in the sediment column may indicate that compounds, like ammonium and sulphide, have consumed the supplied O_2 . Reduced species of Fe and Mn are also oxidised by root-derived oxygen, although the higher background concentration do not allow differentiating until oxygen is in excess. Nevertheless, it appears that oxygen delivered by the roots was not used to oxidise substantial amounts of Fe(II) in Fe(III) and of Mn(II) in Mn(IV) and enhance the Fe and Mn in solid sediments, since sediment layers of higher belowground biomass did not present increases of these elements.

Previous works on the Tagus marshes indicated a correlation between metal concentration in rooting sediments and loss on ignition (Caçador et al., 1999). These relationships pointed to the importance of organic matter, which at these layers derives mainly from the decomposing roots, on retention of trace elements (Jones, 1998; Weis and Weis, 2004). This effect may be emphasised in marshes with abundant belowground biomass, like the Rosário site colonised by *S. fruticosa*. Because belowground biomass of this plant was found to have an elevated specific area (Sundby et al., 2005), sediment-pore water interaction is expected to be higher than in the sediments covered with *S. maritima*. The enhancement of this interaction, among others related to the different strategies of the two studied plants (Groenendijk and Vink-Lieavaart, 1987; Caetano et al., 2008), may induce higher levels of trace elements found in the roots of *S. fruticosa* in comparison to the *S. maritima*. A study of metal release from decomposing root litter in Tagus marshes over a 7-month period pointed to complex behaviour of the surveyed metals. Litter in spring may act as a cation exchanger adsorbing Cu and Pb from sediments, while Zn and Cd stocks decreased progressively in plants (Pereira et al, 2007; Caçador et al., 2009). As metals exchange with the decomposing litter the metal concentrations in pore waters may enhance. The litter decomposition and the release of exudates from live roots are a continuous source of organic matter that has direct impact on metals availability (Sundby et al., 2005; Pereira et al, 2007). In a previous study over the annual growth cycle dissolved sulphide was absent in rooting sediments, even in periods of lack of oxygen (Sundby et al., 2003). Assuming similar conditions in the sediment, no substantial amounts of metals in pore waters are expected to precipitate as metal sulphides. In the absence of this effective removal, the peak concentrations of dissolved Cu, Cd, Pb and Zn in pore waters are presumably linked to organic complexes stabilized in solution.

4.2. Metal exchange between sediment and overlying water at submerged period

Vertical pore water profiles differed in the two studied sites. At 50-min after inundation of the marsh and at air-exposed conditions, *S. maritima* sediments exhibited in general higher levels of Fe, Mn, Zn, Cu, Pb and Cd in pore waters of the upper layers. Pore waters of *S. fruticosa* sediments showed a sub-surface maximum of Zn, Pb and Cd concentrations below the layers of higher root biomass (table 3) and of Cu levels at layers of higher root density. The different metal concentration between pore water of 0-5 cm layer and overlying water induce a diffusive flux estimated by the Fick's 1st law (equation 1). The similarity of concentrations at t=50 min. of inundation and at air-exposed conditions (t=0) suggests that no exchanges are foreseen by molecular diffusion after that period of inundation. The best estimation for pore water concentrations under submerged period is, thus, the average between concentration at t=50 min and t=0 (assumed as representative of the final submerged period). It was considered that sediments covered by *S. maritima* and *S. fruticosa* are submerged 6 and 4 hours per tide, respectively. Table 5 shows the predicted amounts ($\text{nmol m}^{-2} \text{d}^{-1}$) of metals exchanged daily with the overlying water. Sediments covered

by *S. maritima* imported Cu and Zn (-16 and $-137 \text{ nmol m}^{-2} \text{ d}^{-1}$) and exported Fe, Mn, Pb and Cd (105 , 4800 , 0.4 and $0.04 \text{ nmol m}^{-2} \text{ d}^{-1}$, respectively). The sediments colonised by *S. fruticosa* exported larger amounts of Zn, Mn, Cu and Cd (570 , 290 , 80 and 8 , respectively) and imported Fe and Pb (-510 and -4 , respectively). In these calculations it was considered the value of porosity (ϕ) in equation (2) commonly used for fine sediments. However, the presence of micro-channels around the belowground plant parts near the sediment-water interface may facilitate the exchange of solutes by molecular diffusion. Therefore exchanges of metals between sediment-water interfaces driven by diffusive gradients are underestimated.

4.3. Effect of tidal flooding on metal concentration in pore waters and overlying water

The decrease of metal concentrations in pore water during the first 5 and 10 minutes of inundation (figures 2 and 3) clearly indicates that tidal flooding modifies the chemical equilibrium in pore water of the upper sediment layers. These results are in line with studies performed in Ria Formosa (SW Europe) showing the modification of nutrients, Cu and Cd profiles in pore water as permeable inter-tidal sediments are flooded (Falcão and Vale, 1990; Caetano et al, 2007). During inundation overlying water containing lower metal concentrations mixes with pore waters of the upper sediment layers. Alterations associated with precipitation/dissolution and sorption/desorption during this period should not also be excluded (Caetano et al., 1997; Heuttel et al., 1998). In general, profiles registered 50 minutes after the inundation were closer to the initial conditions suggesting that equilibrium have been nearly achieved. In the first minutes of tidal flooding all determined elements increased considerably their concentrations in water that inundates the sediments from both sites. The abrupt increases are in line with the decreases of metal concentrations in pore water pointing to a substantial pulse of solutes exported to the water column. Although different trends on pore water concentration were found between both plants during the inundation the overall process induced by tide was similar in *S. maritima* and *S. fruticosa*.

4.4. Advective transport of metal associated to tidal flooding

The increase of metal concentrations in overlying waters during the first minutes of tidal flooding is coherent with the decrease on the pore waters and points to the escape of metals from pore water of salt marsh sediments (figure 3). On the basis of these measurements one may calculate the advective transport to the water column using the equation (4). Table 4 presents the amounts of Fe, Mn, Zn, Cu, Pb and Cd on a daily basis, which correspond to the sum of two pulses associated with the two tidal flooding periods. The most remarkable aspect is that quantities are three to four orders of magnitude above those predicted by Fick's 1st law. For example, the diffusive flux of Pb was between -4 and $0.4 \text{ nmol m}^{-2} \text{ d}^{-1}$ while advective transport reached $13\text{-}15 \times 10^3 \text{ nmol m}^{-2} \text{ d}^{-1}$. Previous works on permeable inter-tidal sediments have shown the importance of pulse mechanisms associated with the tidal flushing on sediment-water exchanges (Falcão and Vale, 1990; Caetano et al, 2007). The current work permits to extend the relevance of this mechanism to salt marsh sediments covered with *S. maritima* and *S. fruticosa*, two plants commonly found in south European salt marshes.

Although sediments of Rosário marsh consist of fine-grained sediments, the micro-channels formed by the root mosaic have probably a major role on water and solutes circulation driven by the increase of pressure on sediments as tide inundates the area. The amounts exported by this mechanism were higher in *S. maritima* than in *S. fruticosa* area, although the metal concentration in pore waters of sediments covered by this plant exceeded largely that found in *S. maritima*. The most plausible explanation for this apparent discrepancy is the smaller size of the *S. fruticosa* roots and their random and intricate distribution which create more obstacles to pore water flushing during the first moments of tidal flooding. Otherwise, *S. maritima* has thicker tillers and roots promoting increased exchanges between pore water and flooding

waters. These results are important from an ecological point-of-view since marshes subjected to tidal inundations also act as a source of dissolved metals from sediment to the water column being available for macrobenthic organisms and small fishes. These findings also support a previous works that showed the transfer of trace metals to marine environment across the detritus generated by salt marsh plants (Pereira et al, 2007; Caçador et al., 2009). Considering the magnitude and the periodicity of advective export, successive tidal inundation would deplete the pool of metals in the salt marsh sediment if not balanced by equivalent inputs. Conservation of mass requires an import during the submerged period by settling of suspended particles, advection of water, deposition of particles during ebb tide and exchanges of metals with belowground biomass.

5. Conclusions

This work emphasizes the importance of tidal flooding over salt marsh sediments to the metal exportation to the water column. The accumulation of metals in the sediments close to the plant roots, together with the arriving of the tide results in an important exportation of metals from the pore water of salt marsh areas to the overlying waters. More than the diffusive flux, the advective transport has been demonstrated to be an important contribution to the overall flux. This type of fluxes should be taken into account when assessing trace metal contamination problems and remediation in salt marsh areas. Furthermore, one may speculate that inundation of coastal areas associated with the climatic changes, either to increase of water levels or floods will result in additional fluxes of trace elements from intertidal areas to the water column.

Acknowledgments

The authors wish to thank the colleagues Rute Cesário (IPIMAR) and Ana García-Blanco (IIM-CSIC) for the field work and technical assistance. This article is a contribution to the Spanish-Portuguese Actions of references 2007PT0021 and FCT/CSIC9/CSIC/08. Dr. Santos-Echeandía would like to thank the Basque Government for the financial support (post-doctoral contract).

References

- Achterberg, E. P., Van Den Berg C., 1994. In-line ultraviolet-digestion of natural water samples for trace metal determination using an automated voltammetric system. *Analytica Chimica Acta* 291, 213-232.
- Aller, R., 1977. The influence of macrobenthos on chemical diagenesis of marine sediments. PhD Thesis, Yale University, USA.
- Berner, R., 1980. Early Diagenesis. A Theoretical Approach. Princeton University Press, USA.
- Brotas, V., Ferreira, A., Vale, C., Catarino, F., 1990. Oxygen profiles in intertidal sediments of Ria Formosa (S. Portugal). *Hydrobiologia* 207, 123-129.
- Brune, A., Frenzel, P., Cypionka, H., 2000. Life at the oxic-anoxic interface: microbial activities and adaptations. *FEMS Microbiology Reviews* 24, 691-710.
- Burdige, D., 1993. The biogeochemistry of manganese and iron reduction in marine sediments. *Earth Science Reviews* 35, 249-284.
- Caçador, I., Vale C., 2001. Salt marshes, in: Prasad, M., editor. *Metals in the Environment*. Hyderabad: Marcel Dekker Inc. p. 95-115.

- Caçador, I., Vale, C., Catarino, F., 1993. Effects of plants on the accumulation of Zn, Pb, Cu and Cd in sediments of the Tagus estuary salt marshes, Portugal. J-P. (Vernet ed.). Environmental Contamination. Elsevier. Amsterdam Studies in Environmental Science 55, 355–365.
- Caçador, I., Vale, C., Catarino, F., 1996a. Accumulation of Zn, Pb, Cu and Ni in sediments between roots of the Tagus estuary salt marshes, Portugal. Estuarine Coastal and Shelf Sciences 42, 393–403.
- Caçador, I., Vale, C., Catarino, F., 1996b. The influence of plants on concentration and fractionation of Zn, Pb, and Cu in salt marsh sediments (Tagus Estuary, Portugal). Journal of Aquatic Ecosystems Health 5, 193–198.
- Caçador, I., Vale, C., Catarino, F., 1999. Relationship between metal concentrations and organic matter content in the Tagus estuary salt marsh sediments, in: Halophyte use in different climates Backhuys Publishers, Leiden, The Netherlands.
- Caçador, I., Vale, C., Catarino, F., 2000. Seasonal variation of Zn, Pb, Cu and Cd concentrations in the root-sediment system of *Spartina maritima* and *Halimione portulacoides* from Tagus estuary salt marshes. Marine Environmental Research 49, 279–290.
- Caçador, I., Costa, A., Vale, C., 2004. Carbon storage in Tagus salt marsh sediments. Water Air Soil Pollution: Focus 4, 701–714.
- Caçador, I., Caetano, M., Duarte, B., Vale, C., 2009. Stock and losses of trace metals from salt marsh plants. Marine Environmental Research 67, 75–82.
- Caetano, M., Vale, C., 2002. Retention of arsenic phosphorus in iron-rich concretions of Tagus salt marshes. Marine Chemistry 79, 261–271.
- Caetano, M., Fonseca, N., Cesário, R., Vale, C., 2007. Mobility of Pb in salt marshes recorded by total content and stable isotopic signature. Science Total Environment 380, 84–92.
- Caetano, M., Falcão, M., Vale, C., Bebianno, M., 1997. Tidal flushing of ammonium, iron and manganese from inter-tidal sediment pore waters. Marine Chemistry 58, 203–211.
- Caetano, M., C. Vale, R. Cesário e N. Fonseca, 2008. Evidence for preferential depths of metal retention in roots of salt marsh plants. Science of the Total Environment 390, 466–474.
- Cambrollé, J., Redondo-Gómez, S., Mateos-Naranjo, E., Figueroa, M.E., 2008. Comparison of the role of two *Spartina* species in terms of phytostabilization and bioaccumulation of metals in the estuarine sediment. Marine Pollution Bulletin 56, 2037–2042.
- Catarino, F., Caçador, I., 1981. Produção de biomassa estratégia de desenvolvimento em *Spartina maritima* e outros elementos da vegetação dos sapais do estuário do Tejo. Boletim Sociedade Broteriana 2, 384–403.
- Clothier, B., Green, S., 1997. Roots: the big movers of water and chemicals in soil. Soil Science 162, 534–543.
- Crespo, R., 1993. Cartografia do habitat potencial de Passeriformes no Estuário do Tejo por processamento digital de imagem. Degree Thesis FC – University of Lisbon, Lisbon.
- Froelich, P., G. Klinkhammer, M. Bender, N. Luedtke, G. Heath, D. Cullen, P. Dauphin, D. Hammond, B. Hartman and V. Maynard, 1979. Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis. Geochimica et Cosmochimica Acta 43, 1075–1091.
- Falcão, M., Vale, C. 1990. Study of the Ria Formosa ecosystem: benthic nutrient remineralization and tidal variability of nutrients in the water. Hydrobiologia 207, 137–146.
- Groenendijk, A., Vink-Lieavaart, M., 1987. Primary production and biomass on a Dutch salt marsh: Emphasis on the belowground component. Vegetatio 70, 21–27.
- Haines, B., Dunn, E., 1985. Coastal marshes, in: Chabot B, Mooney H, editors. Physiological Ecology of North American Plant Communities. London: Chapman and Hall. p. 323–47.

- Heuttel, M., Ziebis, W., Forster, S., Luther, G., 1998. Advective transport affecting metal and nutrient distributions and interfacial fluxes in permeable sediments. *Geochimica et Cosmochimica Acta* 62, 613-631.
- Li, Y., Gregory, S., 1974. Diffusion of ions in sea water and in deep-sea sediments. *Geochimica et Cosmochimica Acta* 38, 703-714.
- Jacob, D., Otte, M., 2003. Conflicting processes in the wetland plant rhizosphere: metal retention or mobilization? *Water Air Soil Pollution* 3, 91-104.
- Jones, D., 1998. Organic acids in the rhizosphere—a critical review. *Plant and Soil* 205, 24-44.
- Koretsky, C., Cuellar, A., Haveman, M., Beuving, L., Shattuck, T., Wagner, M., 2008a. Influence of *Spartina* and *Juncus* on saltmarsh sediments. II. Trace element geochemistry. *Chemical Geology* 255, 100-113.
- Koretsky, C., Haveman, M., Cuellar, A., Beuving, L., Shattuck, T., Wagner, M., 2008b. Influence of *Spartina* and *Juncus* on saltmarsh sediments. I. PoreWater Geochemistry. *Chemical Geology* 255, 87-99.
- Mendelssohn, I., Kleiss, B., Wakeley, J., 1995. Factors controlling the formation of oxidising root channels: a review. *Wetlands* 15, 37-46.
- Otte, M.L., Rozema, J., Koster, L., Haarsma, M.S., Broekman, R.A., 1989. Iron plaque on roots of *Aster tripolium* L.: interaction with zinc uptake. *New Phytologist* 111, 309-317.
- Otte, R., 1991. Heavy metals and arsenic in vegetation of salt marshes and floodplains. PhD Thesis, Vrije Universiteit, Amsterdam.
- Pereira, P., Caçador, I., Vale, C., Caetano, M., Costa, A., 2007. Decomposition of belowground biomass litter and metal dynamics in salt marshes (Tagus Portugal). *Science Total Environment* 380, 93-101.
- Reboreda, R., Caçador, I., 2007. Copper, zinc and lead speciation in salt marsh sediments colonised by *Halimione portulacoides* and *Spartina maritima*. *Chemosphere* 69, 1655-1661.
- Santos-Echeandia, J., Prego, R., Cobelo-García, A., Millward, G.E., 2009. Porewater geochemistry in a Galician Ria (NW Iberian Peninsula): Implications for benthic fluxes of dissolved trace elements (Co, Cu, Ni, Pb, V, Zn). *Marine Chemistry* 117, 77-87.
- St-Cyr, L., Campbell, P., 1996. Metals (Fe, Mn, Zn) in root plaque of submerged aquatic plant collected in situ: Relations with metal concentrations in adjacent sediments and in the root tissue. *Biogeochemistry* 33, 45-76.
- St-Cyr, L., Crowder, A., 1990. Manganese and copper in the root plaque of *Phragmites australis*. *Soil Science* 149, 191-198.
- Sundby, B., Vale, C., Caçador, I., Catarino, F., Madureira, M., Caetano, M., 1998. Metal-rich concretions on the roots of the salt marsh plants: mechanism and rate of formation. *Limnology and Oceanography* 43, 245-252.
- Sundby, B., Vale, C., Caetano, M., Luther, G., 2003. Redox chemistry in the root zone of a salt marsh sediment in the Tagus estuary, Portugal. *Aquatic Geochemistry* 9, 257-271.
- Sundby, B., Caetano, M., Vale, C., Gobeil, C., Luther, G., Nuzzio, D., 2005. Root-induced cycling of lead in salt marsh sediments. *Environmental Science and Technology* 39, 2080-2086.
- Sundby, B., 2006. Transient state diagenesis in continental margin muds. *Marine Chemistry* 102, 2-12.
- Taillefert, M., Neuhuber, S., Bristow, G.W., 2007. The effect of tidal forcing on biogeochemical processes in intertidal salt marsh sediments. *Geochemical Transactions* 8, 6.

- 532 Vale, C., Catarino, F., Cortesão, C., Caçador, I., 1990. Presence of metal-rich rhizoconcretions on the
533 roots of *Spartina maritima* from the salt marshes of the Tagus Estuary, Portugal. Science of the
534 Total Environment 97/98, 617–626.
- 535 Vale, C., Caetano, M., Raimundo, J., 2003. Incorporation of trace elements in iron-rich concretions around
536 plant roots of Tagus estuary salt marshes (Portugal). Journal of Soils and Sediments 3, 208–212.
- 537 Vale, C., 1990. Temporal variations of particulate metals in the Tagus river estuary. Science of the Total
538 Environment 97, 137–154.
- 539 Van Cappellen, P., Y. Wang, 1996. Cycling of iron and manganese in surface sediments: a general theory
540 for the coupled transport and reaction of carbon, oxygen, nitrogen, sulphur, iron and manganese.
541 American Journal Science 296, 197-243.
- 542 Weis, J., Weis, P., 2004. Metal uptake, transport and release by wetland plants: implications for
543 phytoremediation and restoration. Environmental International 30, 685–700.
- 544 Windham, L., Weis, J., Weis, P., 2001. Lead uptake, distribution, and effects in two dominant salt marsh
545 macrophytes, *Spartina alterniflora* (Cordgrass) and *Phragmites australis* (Common reed). Marine
546 Pollution Bulletin 42, 811–816.
- 547 Windom, H., Schropp, S., Calder, F., Ryan, J., Smith, R., Burney, L., Lewis, F.G., Rawlison, C.H., 1989.
548 Natural trace metal concentrations in estuarine and coastal marine sediments of the southeastern
549 United States. Environmental Science and Technology 23, 314–320.

Table 1. Certified values and measured average concentrations of Fe (μM), Mn, Zn, Cu, Pb and Cd (nM) in the reference materials in SLEW-3 (estuarine water), CASS-4 (nearshore seawater) and SLRS-4 (river water); number of replicates in brackets.

	Values	Fe (μM)	Mn (nM)	Zn (nM)	Cu (nM)	Pb (nM)	Cd (nM)
SLEW-3	Obtained			2.72 $\pm$ 1.18 (n=3)	27.0 $\pm$ 0.07 (n=3)		0.386 $\pm$ 0.07 (n=3)
	Certified			3.07 $\pm$ 0.57	24.39 $\pm$ 1.89		0.427 $\pm$ 0.036
CASS-4	Obtained			5.8 $\pm$ 1.0 (n=13)	9.4 $\pm$ 0.5 (n=6)	0.068 $\pm$ 0.012 (n=13)	0.236 $\pm$ 0.033 (n=13)
	Certified			5.8 $\pm$ 0.9	9.3 $\pm$ 0.8	0.047 $\pm$ 0.017	0.231 $\pm$ 0.027
SLRS-4	Obtained	0.23 $\pm$ 0.02 (n=5)	63.5 $\pm$ 2.3 (n=5)				
	Certified	0.18 $\pm$ 0.09	61.3 $\pm$ 3.3				

Table 2 – Concentrations and standard deviation of Fe (%), Mn, Zn, Cu, Pb and Cd ($\mu\text{g g}^{-1}$) in sediment solids colonised by *Spartina maritima* and *Sarcocornia fruticosa* from Rosario salt marsh. Probability (p) was provided by Kruskal-Wallis test considering the data of all depths.

Colonised sediment			
	Depth (cm)	<i>S. maritima</i> (lower marsh)	<i>S. fruticosa</i> (higher marsh)
Fe ° (%)	0 - 5	4.3±0.3	4.7±0.4
	5 - 10	3.9±0.1	3.8±0.3
	10 - 15	4.0±0.1	4.8±0.2
	15 - 20	4.0±0.2	6.8±1.5
	20 - 25	4.7±0.1	5.6±0.5
Mn ° ($\mu\text{g g}^{-1}$)	0 - 5	311±26	234±44
	5 - 10	178±11	190±22
	10 - 15	172±1	189±28
	15 - 20	170±17	210±28
	20 - 25	181±14	247±14
Zn * ($\mu\text{g g}^{-1}$)	0 - 5	482±24	333±22
	5 - 10	501±36	465±61
	10 - 15	513±13	890±183
	15 - 20	596±101	1075±304
	20 - 25	728±2	2408±131
Cu * ($\mu\text{g g}^{-1}$)	0 - 5	65±18	71±12
	5 - 10	71±4	50±22
	10 - 15	65±1	79±10
	15 - 20	27±3	83±33
	20 - 25	41±2	83±36
Pb ** ($\mu\text{g g}^{-1}$)	0 - 5	187±31	290±89
	5 - 10	166±13	168±98
	10 - 15	136±19	266±13
	15 - 20	24±4	257±13
	20 - 25	56±1	264±127
Cd * ($\mu\text{g g}^{-1}$)	0 - 5	1.2±0.4	0.80±0.2
	5 - 10	0.30±0.1	1.4±0.9
	10 - 15	0.50±0.1	2.5±1.9
	15 - 20	1.1±0.2	3.1±2.8
	20 - 25	1.6±0.7	4.1±2.8

* - $p < 0.1$; ** - $p < 0.05$; ° not significant

Table 3 – Dry weight belowground biomass (g m^{-2}), concentrations and standard deviation of Fe (%), Mn, Zn, Cu, Pb and Cd ($\mu\text{g g}^{-1}$) in belowground biomass of *Spartina maritima* and *Sarcocornia fruticosa* from Rosario salt marsh. Probability (p) was provided by Kruskal-Wallis test considering the data of all depths. Roots were inexistent in 20-25-cm layer.

	Depth (cm)	Belowground biomass	
		<i>S. maritima</i> (lower marsh)	<i>S. fruticosa</i> (higher marsh)
Biomass ** (g m^{-2})	0 – 5	282	1001
	5 -10	425	1058
	10 – 15	346	641
	15 – 20	238	829
	20 – 25	-	344
Fe ° (%)	0 – 5	2.8±0.5	3.3±0.3
	5 -10	9.8±4.2	1.7±0.1
	10 – 15	4.4±0.7	4.4±0.3
	15 – 20	4.5±0.2	6.0±1.0
	20 – 25	-	5.2±0.6
Mn ° ($\mu\text{g g}^{-1}$)	0 – 5	211±235	47±18
	5 -10	210±80	44±5
	10 – 15	99±10	89±44
	15 – 20	90±34	176±40
	20 – 25	-	190±9
Zn ° ($\mu\text{g g}^{-1}$)	0 – 5	2923±1888	576±156
	5 -10	2468±1439	3313±1355
	10 - 15	2538±764	5116±585
	15 - 20	3571±740	5875±585
	20 - 25	-	11017±1693
Cu ** ($\mu\text{g g}^{-1}$)	0 - 5	104±59	227±123
	5 -10	149±73	305±41
	10 - 15	190±49	334±22
	15 - 20	212±106	293±3
	20 - 25	-	236±4
Pb * ($\mu\text{g g}^{-1}$)	0 - 5	522±218	599±221
	5 -10	1528±243	1196±398
	10 - 15	1123±222	2739±186
	15 - 20	453±88	2106±186
	20 - 25	-	2113±74
Cd ** ($\mu\text{g g}^{-1}$)	0 - 5	19±14	13±7
	5 -10	9±9	27±5
	10 - 15	11±3	23±2
	15 - 20	17±3	23±2
	20 - 25	-	30±4

* - $p < 0.1$; ** - $p < 0.05$; ° not significant

Table 4 – Ranges of temperature, salinity and Fe, Mn, Zn (μM), Cu, Pb and Cd (nM) pore water concentrations in sediments (0-25 cm) at the air exposure situation ($t=0$), colonised *Spartina maritima* and *Sarcocornia fruticosa* from Rosario salt marsh. Probability (p) was provided by Kruskal-Wallis test considering the data of all depths.

		Pore waters	
		<i>S. maritima</i> (lower marsh)	<i>S. fruticosa</i> (higher marsh)
Temperature	$^{\circ}\text{C}$	14-15	14-15
Salinity		32.4 – 33.6	28.5 – 31.8
Fe $^{\circ}$	μM	0.32 – 13	0.17 – 65
Mn $^{\circ}$		5.1 – 34	5.8 – 14
Zn **		0.33 – 1.2	1.2 – 14
Cu **	nM	0.99 – 28	27 – 201
Pb **		1.0 – 15	29 – 268
Cd **		0.10 – 1.4	0.27 - 75

** - $p < 0.05$; $^{\circ}$ not significant

Table 5 – Estimated diffusive fluxes and advective transport of Fe, Mn, Zn, Cu, Pb and Cd in Rosário sediments colonised *Spartina maritima* and *Sarcocornia fruticosa* from Rosário salt marsh. Positive fluxes indicate net exportation of metals from the sediment to the overlying waters.

	Diffusive flux (nmol m ⁻² d ⁻¹)		Advective flux (10 ³ x nmol m ⁻² d ⁻¹)	
	<i>S. maritima</i>	<i>S. fruticosa</i>	<i>S. maritima</i>	<i>S. fruticosa</i>
	(lower marsh)	(upper marsh)	(lower marsh)	(upper marsh)
Fe	105	-510	9520	1640
Mn	4800	290	24	-17
Zn	-140	570	220	82
Cu	-17	27	74	16
Pb	0.4	-4.0	13	15
Cd	0.04	3.0	0.3	0.08

FIGURE CAPTIONS

Figure 1 – Location of the Rosario salt marsh in the Tagus estuary.

Figure 2 - Depth metal concentrations in pore waters of sediments at air-exposed conditions ($t=0$), 5, 10 and 50 minutes after the tidal flooding of the lower marsh colonised by *S. maritima*. Standard deviation bars of the three replicates are included.

Figure 3 - Depth metal concentrations in pore waters of sediments at air-exposed conditions ($t=0$), 5, 10 and 50 minutes after the tidal flooding of the higher marsh colonised by *S. fruticosa*. Standard deviation bars of the three replicates are included.

Figure 4 – Time course evolution of Fe, Mn, Zn, Cu, Pb and Cd concentrations in the dissolved fraction of flooding water in the areas colonised by *S. fruticosa* and *S. maritima* of the Rosario marsh. Standard deviation bars of the three replicates are included.

Figure 1

Figure 2

Figure 3

Figure 4