Development of Atlantic cod () exposed to produced water during early life stages: Effects on embryos, larvae, and juvenile fish

Sonnich Meier, H. Craig Morton, Gunnar Nyhammer, Bjørn Einar Grøsvik, Valeri Makhotin, Audrey Geffen, Stepan Boitsov, Karen Anita Kvestad, Anneli Bohne-Kjersem, Anders Goksøyr, et al.

► To cite this version:

Sonnich Meier, H. Craig Morton, Gunnar Nyhammer, Bjørn Einar Grøsvik, Valeri Makhotin, et al.. Development of Atlantic cod () exposed to produced water during early life stages: Effects on embryos, larvae, and juvenile fish. Marine Environmental Research, 2010, 70 (5), pp.383. 10.1016/j.marenvres.2010.08.002 . hal-00631697

HAL Id: hal-00631697

<https://hal.science/hal-00631697>

Submitted on 13 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Development of Atlantic cod (*Gadus morhua*) exposed to produced water during early life stages: Effects on embryos, larvae, and juvenile fish

Authors: Sonnich Meier, H. Craig Morton, Gunnar Nyhammer, Bjørn Einar Grøsvik, Valeri Makhotin, Audrey Geffen, Stepan Boitsov, Karen Anita Kvestad, Anneli Bohne-Kjersem, Anders Goksøyr, Arild Folkvord, Jarle Klungsøyr, Asbjørn Svoldal

PII: S0141-1136(10)00117-0

DOI: [10.1016/j.marenvres.2010.08.002](https://doi.org/10.1016/j.marenvres.2010.08.002)

Reference: MERE 3468

To appear in: *Marine Environmental Research*

Received Date: 8 December 2009

Revised Date: 10 August 2010

Accepted Date: 13 August 2010

Please cite this article as: Meier, S., Morton, H.C., Nyhammer, G., Grøsvik, B.E., Makhotin, V., Geffen, A., Boitsov, S., Kvestad, K.A., Bohne-Kjersem, A., Goksøyr, A., Folkvord, A., Klungsøyr, J., Svoldal, A. Development of Atlantic cod (*Gadus morhua*) exposed to produced water during early life stages: Effects on embryos, larvae, and juvenile fish, *Marine Environmental Research* (2010), doi: 10.1016/j.marenvres.2010.08.002

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Development of Atlantic cod (*Gadus morhua*) exposed to produced water
during early life stages: Effects on embryos, larvae, and juvenile fish.**

Sonnich Meier^{a*}, H. Craig Morton^a, Gunnar Nyhammer^b, Bjørn Einar Grøsvik^{ab}, Valeri
Makhotin^c, Audrey Geffen^b, Stepan Boitsov^a, Karen Anita Kvestad^b, Anneli Bohne-Kjersem^b,
Anders Goksøyr^b, Arild Folkvord^b, Jarle Klungsøyr^a, Asbjørn Svardal^{ad}

^aInstitute of Marine Research, Bergen, Norway

^b Department of Molecular Biology/Biology, University of Bergen, Norway

^cDepartment of Ichthyology, Moscow State University, Russia.

^dDepartment of Medicine, Section for Pharmacology, University of Bergen, Norway

*Corresponding author:

Sonnich Meier

Institute of Marine Research

P.O.Box 1870, N-5817 Nordnes, Bergen, Norway

Fax: +47 55238555

E-mail: Sonnich.meier@imr.no

Abstract

Produced water (PW) contains numerous toxic compounds of natural origin, such as dispersed oil, metals, alkylphenols (APs), and polycyclic aromatic hydrocarbons (PAHs). In addition, PW also contains many different chemicals which have been added during the oil production process. In the study described here, cod were exposed to real PW collected from an oil production platform in the North Sea. This was done in order to best recreate the most realistic field-exposure regime in which fish will be affected by a wide range of chemicals. The biological effects found in this study therefore cannot be assigned to one group of chemicals alone, but are the result of exposure to the complex chemical mixture found in real PW. Since APs are well known to cause endocrine disruption in marine organisms, we focused our chemical analysis on APs in an attempt to better understand the long term effects of APs from PW on the biology of fish. In this study, cod were exposed to several concentrations of real PW and 17 β -oestradiol (E₂), a natural oestrogen, at different developmental stages. Cod were exposed to PW either during the embryo and early larvae stage (up to 3 months of age) or during the early juvenile stage (from 3 – 6 months of age). Results showed that, in general, APs bioconcentrate in fish tissue in a dose and developmental stage dependent manner during PW exposure. However, juveniles appeared able to effectively metabolise the short chain APs. Importantly, PW exposure had no effect on embryo survival or hatching success. However, 1% PW clearly interfered with the development of normal larval pigmentation. After hatching most of the larvae exposed to 1% PW failed to begin feeding and died of starvation. This inability to feed may be linked to the increased incidence of jaw deformities seen in these larvae. In addition, cod exposed to 1% PW, had significantly higher levels of the biomarkers vitellogenin and CYP1A in plasma

43 and liver, respectively. No similar effects were seen in cod exposed to either 0.1% or 0.01%
44 PW.

45

46

47 Key words: Alkylphenols, Early lifestages, Embryo, Endocrine disruption, Cod, *Gadus morhua*,

48 Larvae, Produced water, Pigmentation, Sex differentiation.

1. Introduction

Produced water (PW) is a combination of formation water, condensation water, injection water and water used for desalting oil. Therefore, PW is a complex mixture of seawater, dispersed oil, PAHs and other dissolved hydrocarbons, organic acids, alkylphenols (APs), metals, and traces of production chemicals (Neff, 2002). As oilfields age, the amount of water injected into the reservoir to help maintain reservoir pressure increases. Since many oil fields in the North Sea are approaching the end of their productive lives the amount of water injected into the reservoirs has increased rapidly, resulting in a huge increase in the amount of PW discharged into the sea (Durell et al., 2006). From 1990 to 2006 the annual discharge of PW from the Norwegian sector increased from 10 mill m³ to 162 mill m³, and together with PW from the British, Danish and Dutch sectors, the total amount of PW released into the North Sea is estimated to be more than 500 mill m³ per year.

APs are natural components of crude oil (Ioppolo-Armanios et al., 1992), and as a result of their solubility in water high concentrations are still present in the aqueous phase after oil/water separation (Boitsov et al., 2007). Oil production platforms, therefore, release large amounts of APs into the seas via PW. APs have been shown to mimic the effects of the natural female sex hormone oestrogen, resulting in disruption of the endocrine and reproductive systems (Meier et al., 2007; Tollefsen and Nilsen, 2008). Previously, the main focus of research has been on the effects of long-chain APs (octyl-, and nonylphenols) which are degradation products of non-ionic surfactants known as AP-ethoxylates. More recently, it has been shown that APs with shorter chain lengths can also bind to the oestrogen receptor and produce mild oestrogenic effects (Tollefsen and Nilsen, 2008).

72

73 There is a huge amount of literature available describing the effects of APs on the endocrine
74 system of vertebrates, and on gonadal development in fish in particular (Arukwe and
75 Goksøyr, 1998; Servos, 1999). APs affect a number of reproductive parameters in fish,
76 including induction of plasma vitellogenin (Vtg) in male and juvenile fish (Jobling and
77 Sumpter, 1993; White et al., 1994), inhibition of spermatogenesis (Jobling and Sumpter,
78 1993; Gimeno et al., 1998; Miles-Richardson et al., 1999; Weber et al., 2002), and oogenesis
79 (Tanaka and Grizzle, 2002; Weber et al., 2003). In addition to binding to the oestrogen
80 receptor, APs have also been shown to affect the brain-pituitary-gonad axis via
81 induction/inhibition of the gonadotropins (Harris et al., 2001; Yadetie and Male, 2002) and
82 to have direct effects on steroidogenesis (Yokota et al., 2005; Arukwe, 2005).

83

84 In many fish species sexual differentiation appears to be a relatively plastic process that has
85 proved to be especially vulnerable to the endocrine-disrupting effects of APs (Gimeno et al.,
86 1996; Seki et al., 2003; Gronen et al., 1999; Kang et al., 2003; Knorr and Braunbeck, 2002).
87 Very little is known about the developmental processes that govern sexual differentiation in
88 fish and especially in marine fish like Atlantic cod (*Gadus morhua*). Like several other fish
89 species, cod likely exhibit specific “developmental windows” during larval development in
90 which they are especially vulnerable to the effects of endocrine-disrupting chemicals and
91 other environmental toxins (Gimeno et al., 1996; Devlin and Nagahama, 2002; van Aerle et
92 al., 2002). In salmonids like rainbow trout (*Oncorhynchus mykiss*), a sensitive developmental
93 window affecting sexual differentiation is known to occur early in the larval stage (Billard,
94 1992). However, in several marine fish the developmental window important for sexual

differentiation occurs later, at the juvenile stage after metamorphosis (Blazquez et al., 1998; Hendry et al., 2002; Chiasson et al., 2008)

Recently, Chiasson et al. (2008) have studied gonadal differentiation in Atlantic cod and haddock (*Melanogrammus aeglefinus*). They found that female cod appear before males, with ovarian cavities first observed at 102 days post hatch (dph) when the fish were 27 mm long (total length, TL). This is similar to what is seen in other marine species like Atlantic halibut (*Hippoglossus hippoglossus*) (Hendry et al., 2002). Male cod were first positively identified (by the appearance of testis containing primary spermatogonia) at 221 dph (TL = 94 mm).

Although a large amount of literature exists concerning the short-term toxic effects of exposure to high concentrations of APs and PW, only a few studies have examined more long-term effects. This study was specifically designed to examine the long-term effects on cod exposed to real PW during specific stages in their early development. Our chemical analysis focused mainly on the AP fraction of PW due to their known oestrogenic properties. The aim of this study was to investigate whether cod exposed to realistic concentrations of PW during early life suffer from decreased fitness. As measures of fitness, survival, growth, and frequency of malformations were studied. In addition, expression of the protein biomarkers vitellogenin (Vtg) and cytochrome P450 (CYP1A) were analysed by ELISA.

The fish described in this study were monitored until sexual maturation and the long term effects of PW on gonad development and reproductive success analysed. The results of that study will be published elsewhere.

2. Materials and methods

2.1. Rationale for exposure regimens

The aim of this study was to investigate the effects of realistic doses of PW on the sexual differentiation and fitness of cod exposed to PW during early life. PW, released from an oil-production platform into the sea is quickly diluted. Previous computer-modelling studies have concluded that PW is diluted approximately 1:30 at 10m, 1:100 at 100m, and 1:1000 at 1 km from the outlet pipe (Neff, 2002). Additional results from modelling and field measurements have shown that the concentration of dispersed oil present in the North Sea in an area of 50-100 km surrounding the largest oilfields is approximately 1-3 ppb, which roughly corresponds to a dilution factor of 1:10,000 (Rye et al., 1998). In the study described herein, fish were exposed to three different concentrations of PW that reflects the estimated concentrations found in the North Sea (Table 1). Cod were exposed to PW diluted 1:100 (1% v/v, high dose group, H-PW), 1:1000 (0.1% v/v, medium dose group, M-PW), and 1:10,000 (0.01% v/v, low dose group, L-PW). In addition to the three PW exposed groups, another group of fish were exposed to $10\mu\text{g l}^{-1}$ 17 β -oestradiol (E_2) to study the effects of high concentrations of oestrogenic compounds. A further group of fish were maintained in clean seawater throughout the experiment (unexposed fish, U).

2.2. Cod eggs

The eggs used in this study were obtained from wild cod caught in Tysfjorden in Lofoten, Norway. For spawning, one male and one female fish were placed in a spawning tank and the resultant eggs collected from a filter placed under the runoff outlet. To ensure a realistic

level of biological variation in this study, eggs collected from five separate pairs of spawning cod were mixed. The average egg diameter (D) from each spawning pair was measured, and the number of eggs calculated according to the following formula: N (number of eggs per mL) = $1222 \times D^{-2.71}$, (Kjesbu, 1989). Using this formula, 60,000 eggs (12,000 from each of the five pairs) were added to 100 L of water in each exposure tank.

2.3. Description of the PW exposure experiments

The experiments were conducted from March to December 2004. Fertilized eggs were collected over 2 days in March 2004 (25-26/03). Fifty percent hatching occurred on 13/04. Three separate exposure experiments were performed as described below (see also Figure 1).

2.3.1 Experiment 1: PW exposure of cod eggs and yolk-sac larvae in multi-well trays

Fertilized cod eggs (1-2 days old) from each of the 5 spawning pairs were transferred to the IMR laboratory in Bergen and incubated in 24-well plates (NUNC) (one egg per well). Five parallel exposure experiments were conducted: one for each individual family (spawning pair). For each family 16 plates containing 384 eggs in total were set up. The plates were divided into four groups and exposed to H-PW, M-PW, L-PW and normal seawater. Therefore, each treatment group consisted of 480 eggs divided into 20 plates (four plates from each of the 5 families per treatment group). The plates were incubated at 5°C, and the water was changed every second day. The eggs, and subsequent larvae, were examined microscopically every second day. Survival rate, hatching success, and frequency of lethal deformities were recorded until 3 days post-hatch (dph).

164

165 **2.3.2. Experiment 2A: Exposure of cod eggs and early larvae to PW**

166 Three parallel 100 L tanks, each containing 60,000 fertilized cod eggs (12,000 eggs from each
167 family; 1-2 days post fertilization), were subjected to one of the five different exposure
168 regimens described in section 2.1 for 23 days (in March/April 2004), until 3 dph. The larvae
169 were counted and 6500 from each tank were transferred to a fresh 100 L tank and the
170 exposure continued for 67 days (473 degree-days), through the whole start-feeding phase.
171 At the end of this 90 day period (in June 2004) the fish were approximately 2 cm in length
172 and 10 mg in weight (dry weight). The surviving fish from each exposure regimen were
173 pooled and 200 randomly selected early juveniles transferred into 500 L tanks containing
174 normal seawater. All fish in the H-PW group died during this 90 day exposure regimen so no
175 longer-term studies could be performed on this group. The fish (from the remaining four
176 treatment groups) were maintained in these 500 L tanks for approximately 7 months (until
177 December 2004).

178

179 **2.3.3. Experiment 2B: Exposure of cod to PW from 3 – 6 months of age**

180 For this experiment 100,000 cod eggs (20,000 eggs from each of the 5 spawning pairs) were
181 used. The eggs and larvae were maintained in an identical manner to those in Experiment 2A
182 except that they were incubated in clean seawater for 3 months until the early juvenile stage
183 (average length 3.1 ± 0.5 cm). After 3 months, 250 fish were randomly transferred into each
184 of ten 500 L tanks. Separate groups of fish were then exposed to one of the five treatments
185 as described in section 2.1 (2 parallel tanks per treatment). The exposure lasted for 78 days
186 from the early juvenile to juvenile phase (from July to September 2004), at which time the

fish were approximately 11 cm in length and around 13 g in weight. The fish were maintained in these tanks for approximately 3 months (until December 2004).

2.4. Tank setup and dilution of PW in experiments 2A and 2B

The fish in experiments 2A and 2B were maintained in 100 L and 500 L tanks during the PW exposure as described. Clean seawater was added to all treatment tanks from a shared header-tank by gravity flow. Water from all PW header-tanks was diluted 1:100 before being added to the exposure tanks by adjusting the flow rates of the feeder pumps. In experiment 2A the flow rate during the embryo phase was 25ml/min from the seawater header tank and 0.25ml/min from the PW or 17 β -oestradiol (E₂) header tanks. During the larval phase the initial flow rate started at 50ml/min clean seawater and 0.5ml/min diluted PW or E₂, and was gradually raised to 500ml/min seawater and 5ml/min diluted PW or E₂ as the fish increased in size. One PW header tank was set up for each exposure group (three PW header tanks for experiment 2A and three for experiment 2B). The header tank feeding the H-PW exposure tanks contained undiluted PW; the header tank feeding the M-PW exposure tanks contained 100 ml PW per L seawater; and the header tank feeding the L-PW exposure tanks contained 10ml PW per L seawater. Header tanks containing 1mg l⁻¹ E₂ were also set up to feed the E₂ exposure tank (E). Ten mg E₂ was first dissolved in one ml ethanol and 100 μ L of this stock solution was added per L of seawater. The nominal concentration of E₂ was therefore 10 μ g l⁻¹. The flow rate from all header tanks was monitored daily.

2.5. Collection and treatment of PW

The exposure regimen in this study was carried out with PW obtained from the Oseberg C oil production platform located in the North Sea off the west coast of Norway. The PW was collected in four 1000 L tanks and transferred to land by boat. The PW was then aliquoted into 25 L containers and frozen at -30°C until needed. Frozen PW was allowed to thaw over a period of 24 hours, bubbled with air for 5 minutes using an aquarium pump to remove accumulated hydrogen sulphide (H₂S), before dilution and addition to the exposure tanks.

2.6. Feeding

Newly hatched larvae were fed with natural zooplankton, collected by filtering seawater through a Hydrotech drum filter. The collected zooplankton consisted predominantly of copepods at various developmental stages. The zooplankton concentration was measured daily and kept at approximately 1000 zooplankton per L. For the first four weeks the larvae were fed with zooplankton retained between 80 and 250 µm mesh size. As the fish increased in size they were eventually fed with zooplankton retained between 350 µm and 1000 µm. Two types of cultivated plankton algae (*Isochrysis galbana* and *Rhodomonas sp.*) were also added to the tanks daily. Fish were subsequently weaned onto marine fish feed pellets (Dana Feed, Horsens, Denmark), and fed eight times every day from automatic feeders placed over the tanks. The fish were fed with successively larger pellets as they grew.

In experiment 2A, 40 eggs from each treatment group were incubated in a 1 L beaker. After hatching these larvae were not fed, but were monitored daily until all the larvae died of hunger (unfed group).

2.7. Temperature and light regimen

Experiment 1 was performed at a constant 5°C in a climate room. In experiments 2A and 2B, the average temperature in the tanks during the embryo phase was 4.9°C. The temperature was then gradually increased to 8.6°C throughout the larval phase and to 12°C during the early juvenile stage. The fish in the tanks were maintained under day length conditions for Bergen, Norway.

2.8. Chemical analysis of water and fish

The concentration of APs present in the diluted PW header tanks and in the exposure tanks was monitored regularly. Using these data the empirical dilution factor of APs in the exposure tanks relative to undiluted PW was calculated. The uptake of APs into fish tissue was measured at three different time points: yolk sack larvae from experiment 2A were analysed at 3 dph (after 22 days of exposure to PW, E₂, or clean seawater); early juveniles from experiment 2A were analysed at 71 dph (after 90 days of exposure); and late juveniles from experiment 2B were analysed directly following the 78 day exposure period detailed above. The concentrations of 52 APs were measured in the water and in the fish tissue according to previously published methods (Boitsov et al., 2004; Meier et al., 2005). Concentrations of PAHs present in the PW and in the exposure tanks was also measured according to previously published methods (Boitsov et al., 2004), except that PAH standards labelled with stable isotopes were used in addition to the internal AP standard.

2.9. Sampling

Fish were sampled multiple times during the course of experiments 2A and 2B as follows. In experiment 2A, larvae were sampled 7 times between March and June (3 dph to 67 dph) (see Figure 1). At each sampling, 15 larvae from each tank were randomly selected, euthanized, and their length and dry weight measured. Whether the larvae had begun feeding was assessed by microscopic examination of the stomach. The fish in experiment 2A were sampled 5 times (roughly once a month) between June and December (between 3 and 9 months post hatch). In experiment 2B fish were sampled 6 times between June and December. During these samplings, 50 fish were randomly selected, removed from the tank, individually weighed and returned to the tank. During the last sampling (in December) all the fish in each tank (approx. 200-250) were removed and individually weighed and measured. Finally, 125 fish from each exposure group were transferred to sea cages at the Austevoll research station as part of a longer term study (*manuscript in preparation*). The remaining fish in each group were slaughtered, and their sex was determined by morphological examination.

2.10. Sampling for biomarker analysis

At the end of the exposure period in experiment 2A, whole larvae were sampled for proteomic studies and the results are presented elsewhere (Bohne-Kjersem et al., 2010). At the end of the exposure period in experiment 2B, liver samples were taken for proteomic analysis (Bohne-Kjersem *et al.*, *in preparation*). Blood and liver samples from experiments 2B were also analysed for the presence of the biomarkers Vtg and CYP1A, respectively (Nilsen et al., 1998).

278

279 **2.11. ELISA of CYP1A and Vtg**

280 A quantitative ELISA kit (Biosense, Bergen, Norway) was used to measure the Vtg levels in
281 plasma of juvenile cod (Scott *et al.*, 2006a). A semi-quantitative ELISA using a monoclonal
282 anti-cod CYP1A antibody (clone NP7; Biosense, Bergen, Norway) diluted 1:1000 was used to
283 measure CYP1A levels in the livers of juvenile cod (Nilsen *et al.*, 1998).

284

285 **2.12. Statistical analyses**

286 The normality of each data sample and homogeneity of group variances were examined and
287 when necessary the data were log transformed prior to testing. Differences between groups
288 were analyzed by one-way ANOVA followed by a Dunnett's post test for normally distributed
289 data or Kruskal-Wallis non-parametric test followed by Dunn's post test when data failed
290 tests of normality. The statistical analyses were all performed using XLSTAT software
291 (Addinsoft, US).

3. Results

3.1. Chemical analysis of the PW

Continual analysis of the PW obtained from Osberg C throughout the exposure phase demonstrated that the AP profile and concentration remained constant over time (Figure 2). Our data also showed that freezing and thawing the PW had no effect on the AP profile or concentration (Figure 2). The average concentration of all APs of each chain length (from phenol to C9) was also measured (Table 2). The AP profile and relative concentrations in the PW obtained from Oseberg C for this study are similar to those measured previously in PW from the same platform (Boitsov et al., 2004; Boitsov et al., 2007). The AP data also showed how the AP concentration in PW falls as the solubility in water decreased (Table 2). The PW contained high concentrations of the most water soluble APs, phenol and cresol (C0, C1), but the relatively insoluble long-chain APs ($\geq C6$) were present at much lower concentrations (down to the ng l^{-1} range) (Table 2).

3.2. Chemical analysis of the water in the PW exposure tanks

A clear concentration gradient of APs between the different treatment groups (H-PW>M-PW>L-PW $\approx$ U) was observed. In addition, the dilution factor of the most volatile components, phenol and cresol, was markedly lower than the nominal concentration. This can most likely be explained by the fact that these compounds are extremely volatile and evaporate faster than they can be replenished despite the continual flow of PW into the exposure tanks. For the long-chain APs (C2-C5), the actual measured concentration was similar to the nominal concentration, at least in the H-PW and M-PW tanks. The dilution factor of the L-PW was so

great that the AP concentrations measured were similar to the background levels present in the normal seawater added to the tanks containing unexposed fish (U).

3.3. PAH measurements

Typically, for PW originating from Oseberg C, approximately 30% of the PAHs present consisted of naphthalene and methyl-naphthalene (Durell et al., 2006). The relative absence of the highly volatile light PAHs was most likely due to the “airing” of the PW to remove H₂S gas. The highly volatile nature of naphthalene also likely explains why the measured dilution factor in the exposure tanks was only roughly one third of the nominal concentration (Table 5).

3.4. Uptake of APs into the tissue of PW exposed fish

Analysis of the AP levels in cod tissue showed a dose related uptake. For yolk-sac larvae the AP concentration was measured in a pooled sample made up of approximately 2000 individual larvae (approx. wet weight 1 g) from all the parallel exposure tanks and sampled 3 dph. For early juveniles (sampled 78 dph) the AP analysis was performed on a pooled sample consisting of 10 individuals (approx. wet weight 1 g) from each of the 3 parallel exposure tanks. For late juveniles (sampled in September 2004) the AP analysis was performed on liver samples from 5 individual fish from each of the two parallel exposure tanks. Our analysis clearly showed that the fish in the H-PW group had taken up the highest levels of APs. Detectable levels of APs were also seen on the M-PW group, but the levels present of the L-PW group were not significantly different from the background levels found in the fish maintained in normal seawater (Figure 3A – C).

In addition, there were clear differences in the levels of APs found in the different developmental stages. The average bioconcentration factor (BCF) for each species of AP (C_2 - C_5) at each developmental stage (where $BCF = \frac{\text{AP concentration } (\mu\text{g kg}^{-1} \text{ wet weight}) \text{ in tissue}}{\text{AP concentration } (\mu\text{g l}^{-1}) \text{ in seawater}}$) was also calculated. There was a notably higher BCF in the yolk-sac larvae compared to the early and late juveniles for the short chain APs. The average BCF for C_2 phenols (9 different isomers) was 12 for the yolk-sac larvae, 2 for the early juveniles, and 0.6 for late juveniles. A similar pattern was also seen for the C_3 phenols (10 different isomers), but not for the C_4 (10 different isomers) or for the C_5 phenols (6 different isomers) (Table 6).

3.5. Effects of PW on cod embryos and yolk sac larvae

The effects of PW on cod eggs and embryos were studied both in multi-well dishes (Experiment 1) and in tanks (Experiments 2A). None of the PW concentrations had any apparent effect on survival or hatching success (an average of between 30-35% hatched in all treatment groups) of cod eggs in experiments 1 and experiment 2A (Figure 4 and data not shown). However, 100% of the embryos and newly hatched larvae exposed to H-PW (1% PW) in the Nunc trays (experiment 1) lacked pigmentation (Figure 5), and a similar lack of pigmentation was also observed in the larvae exposed to H-PW in the tanks (experiment 2A). This effect on pigment development appeared to be only transient, as fish sampled at later time points had apparently normal pigmentation.

Experiment 1 also revealed a large difference in egg quality between the 5 different spawning pairs (SP). Each individual egg was inspected microscopically a total of 5 times up to 3 dph and the percentage of visible deformities resulting in death were recorded. While one pair (SP5) demonstrated an average deformity rate of only 10%, one pair (SP3) had an average deformity rate of approximately 60%. However, there was no increase in the rate of embryo mortality as a result of the PW treatment (Figure 4).

3.6. Effects of PW on larvae and early juveniles

The larvae exposed to the highest concentration of PW (H-PW) showed clear effects: just after start-feeding the larvae were still noticeably less pigmented and the stomach was often empty or contained only a small amount of food (Figure 6). Interestingly, many larvae in the H-PW group appeared to have deformed jaws (Figures 5 and 6), and this may partly explain why these larvae are unable to feed efficiently.

At 19 dph large numbers of the larvae in the H-PW group died. This corresponds with the time of death observed in all the unfed groups. Therefore, exposure to 1% PW adversely affected the ability of cod larvae to begin feeding, and most larvae died of starvation. At the end of the larval exposure regimen described in Experiment 2A (in June 2004) the surviving fish in each of the three parallel tanks of the H-PW group were counted. In one tank no fish survived and in the other two tanks only 12 and 15 fish survived, respectively. In the unexposed control group (U) the number of surviving fish in each of the three parallel tanks was 693, 561, and 440 respectively. The percentage survival of the unexposed fish (U) was, therefore, slightly less than 10% (Figure 7). This enormous level of mortality and relatively

low percentage survival is a normal feature of development in batch spawners such as cod, that release millions of eggs every 3 to 4 days during the spawning period (Kristiansen et al., 1997). The percentage survival in the L-PW, M-PW, and E₂ treatment groups was slightly lower than the U group, but the differences observed were not significant (Figure 7).

At the start of experiment 2A (in April) the standard length of all fish was approximately 5 mm. At the end of the experiment (in June) the standard length ranged from 11-28 mm. Visually, the few surviving H-PW exposed fish were larger than the fish in the other groups at the end of experiment 2A (*data not shown*). However, due to the low number of survivors in the H-PW group, statistical significance could not be shown. This suggests that there has been a selective mortality and density dependent growth in the H-PW exposed group. No differences in growth were found in any of the other groups.

After the exposure ended (in June 2004) the surviving fish were transferred to clean seawater and growth and survival was monitored until December 2004. There were no significant differences in mortality throughout the experiment (which was less than 10% in all groups from June to December 2004). But the fish in the M-PW (19.2 ± 1.9 mm) and L-PW (19.4 ± 1.9 mm) groups were slightly smaller than the E₂ (20.0 ± 1.9 mm) and U (19.9 ± 2.0 mm) groups at the end of the experiment (in December 2004). However, we feel that this small difference in growth (which is statistically significant) has little if any biological relevance, and the statistical significance is partly due to the large number of observations ($n=200$ in each group).

3.7. Effects on cod juveniles exposed to PW between 3 – 6 months of age

The percentage survival during experiment 2B was between 86-91%, and there was no significant difference between any of the groups (data not shown). There was also no significant difference in the growth rate of the unexposed fish (U) and any of the PW exposed groups (Figure 8). E₂ on the other hand, had a huge effect on growth. At the end of the exposure regimen (in September 2004) the average weight of the E₂ exposed fish was only 43% of the average weight of the unexposed fish. This difference in weight was still evident in December after the fish had been kept in normal seawater for nine weeks (Figure 8).

3.8. Expression of Vtg and CYP1A

A significant induction of plasma Vtg levels in cod treated with H-PW and E₂ was observed (Figure 9A). In the unexposed fish (U) and the L-PW, and M-PW groups Vtg concentrations ranged from 8-14 ng ml⁻¹. In the H-PW group the Vtg concentration was 5 mg ml⁻¹, and in the E₂ treated group 150 mg ml⁻¹. Therefore, the Vtg concentration in the plasma of H-PW and E₂ treated fish is 0.5x10⁶ - 1x10⁶ times greater than the levels in the plasma of unexposed fish. However, although the Vtg induction in the H-PW group is clearly significant, the levels in plasma are still only 3% of the levels present in the plasma of E₂ treated fish.

CYP1A levels were significantly increased in the H-PW group probably due to the PAHs in the PW. Levels of CYP1A in cod liver were significantly down-regulated following E₂ exposure (Figure 9B).

4. Discussion

4.1. PW exposure and rationale behind the experimental design

The exposure experiments described here were performed using real produced water, collected immediately prior to discharge, from the Oseberg C oil production platform (StatoilHydro, Norway) located in the Norwegian sector of the North Sea. The advantage of using real produced water is that it more closely represents the complex mixture of chemical components (both known and unknown) that are discharged into the sea from oil platforms. The PW was transported to shore in 1000 L polyethylene (PE) containers and was then aliquoted into 25 L PE containers and frozen at -30°C. The PW was flushed with air for 5 minutes before freezing and after thawing in order to remove the highly toxic hydrogen sulphide gas (H₂S). A similar procedure has recently been used to study the effects of PW on adult cod (Sundt et al., 2009). However, the chemical makeup of PW is not stable and our treatment would be expected to alter the composition. Flushing with air will likely remove the most volatile components like benzene, toluene, ethylbenzene, and xylenes (BTEX). These compounds will also likely evaporate very quickly after the PW is released into the sea. Importantly, our analysis showed that the concentration of the AP fraction remains stable over time and throughout at least one freezing and thawing cycle (Figure 2). Furthermore, our analysis of the water chemistry during the actual exposure experiments showed the presence of a clear gradient in the AP concentration between the H-PW, M-PW and L-PW groups. Moreover, the measured AP concentrations were very close to the nominal concentrations specified in the experimental design (1%, 0.1%, and 0.01% PW, respectively) (Table 5).

450

451 **4.2. Uptake of APs into fish tissues**

452 The range of bioconcentration factors (BCF) observed in this study was similar to those that
453 have been reported previously (Servos, 1999). In addition, our results showed that
454 bioaccumulation was dependent on the developmental stage of the fish. Newly hatched yolk
455 sac larvae had a higher BCF of short chain APs (C₂ AP, C₃ AP) compared with early and late
456 juveniles. This may be partly explained by the fact that yolk sac larvae have a higher surface
457 area to weight ratio than later developmental stages (Petersen and Kristensen, 1998).
458 However, it is also likely that the detoxification enzyme system is less well developed in
459 newly hatched cod, as has been shown to be the case for the cytochrome P-450 system
460 (Goksøyr et al., 1988). The relative lack of short chain APs (C₂ AP and C₃ AP) in the liver tissue
461 of late juveniles supports the theory that the detoxification enzyme system is much more
462 developed at this later stage. Generally, fish are quite efficient at metabolizing APs, mainly
463 by phase II enzymes that conjugate intact APs to their corresponding glucuronides (Ferreira-
464 Leach and Hill, 2000).

465

466 **4.3. Effects of PW on cod embryos**

467 The concentrations of PW used in this study (up to 1% in the H-PW group) were not acutely
468 toxic to cod embryos. Although there was no increase in lethal malformations after PW
469 exposure, several sub-lethal effects were noted. Embryos exposed to PW failed to develop
470 pigmentation at the same time as embryos incubated in clean seawater. This effect on
471 pigmentation has been observed previously in marine embryos exposed to oil-related
472 hydrocarbons (Falk-Petersen et al., 1985; Paine et al., 1992). A similar effect is seen in

zebrafish (*Danio rerio*) embryos treated with 1-phenyl-2-thiourea (PTU). PTU is used routinely by zebrafish researchers to inhibit pigmentation (Karlsson et al., 2001). This compound inhibits melanogenesis by reducing the activity of the tyrosinase enzyme that converts tyrosine into melanin. Interestingly, it has also been shown that PTU is a weak activator of the aryl hydrocarbon receptor signaling pathway and induces weak transcription of CYP1A1 in zebrafish embryos (Wang et al., 2004). PTU has also been suggested to block production of thyroid hormone in developing zebrafish (Elsalini and Rohr, 2003). These effects have also been seen after exposure of fish embryos to oil hydrocarbons (Alkindi et al., 1996; Stephens et al., 1997; Billiard et al., 2008).

During vertebrate embryogenesis melanoblasts migrate from the neural crest and out into the developing tissue where they ultimately develop into mature melanocytes. This migration is normally a tightly controlled process, but the cellular signals that direct this migration are not well understood (Sulaimon and Kitchell, 2003). It has been suggested that abnormal development and migration of neural crest derived melanocytes may be a highly sensitive indicator of exposure to estrogenic contaminants in the environment (Bevan et al., 2003; Bevan et al., 2006). Clearly, more research on the effects of PW on the development of pigmentation and other developmental process is needed.

4.4. Effects of PW on cod larvae and early juveniles

Our findings strongly suggest that exposure to 1% PW prevents the larvae from beginning to feed on their own, leading to death by starvation. The results presented here are also in agreement with our own earlier studies that showed that yolk sac larvae exposed to 1.5%

PW for five days also failed to begin feeding on their own and died of starvation (*unpublished observations*). No apparent effects on survival and growth were seen in either the M-PW or L-PW exposed fish.

Larvae exposed to 1.5% PW show reduced swimming ability and at PW concentrations higher than 4% they go into narcosis and became unresponsive to stimuli (*unpublished observations*). This narcotic effect may explain why the larvae are not able to startfeed. Alternatively, the inability to startfeed could also be explained by the increased incidence of lower jaw deformities seen in larvae exposed to 1% PW. Lower jaw deformities are a well characterised effect of exposure to oil compounds (Tilseth et al., 1984; Pollino and Holdway, 2002; Carls et al., 1999; Heintz et al., 1999; Debruyn et al., 2007).

Our own unpublished observations, suggesting a narcotic effect, are in agreement with earlier work on the effects of PW exposure. In one such study, turbot (*Scophthalmus maximus*) larvae were exposed to concentrations of PW ranging from 0.001% to 1% for a 6 week period early in development (from 53 dph) (Stephens et al., 2000). The authors reported no increases in mortality in any of the exposure groups but found a reduction in the swimming activity of larvae exposed to 1% PW. Larvae exposed to 0.1% and 1% PW showed changes in the ultrastructure of the cell membranes of the gills which could result in a reduced ability to take up oxygen. In addition, numerous other signs of chronic stress were observed, including increased levels of cortisol and cytochrome P450, and increased activities of CYP1A and 7-ethoxyresorufin-O-deethylase (EROD) (Stephens et al., 2000).

Similarly, a reduction on growth was also observed in cod and herring (*Clupea harengus*) cultivated in a mesocosmos system and exposed to PW diluted 400-800 times (Gamble et al., 1987). A tenfold increase in the cytochrome P450 activity in herring larvae exposed to PW was also reported, indicating an increase in the metabolism of hydrocarbons. In addition, cod larvae exposed to the water soluble fraction (WSF) of crude oil displayed a decreased ability to begin feeding. This effect was observed at a total hydrocarbon concentration (THC) of 250 $\mu\text{g l}^{-1}$, a THC that corresponds to approximately 1% PW, assuming a typical THC of 20 mg l^{-1} (Tilseth et al., 1984).

Concentrations of PW above 1% prevent cod larvae from beginning to feed on their own. However, although the presence of APs in the water during the PW exposure and bioaccumulation of APs in the fish was clearly shown, PW contains many other compounds which likely also contribute to the toxicity. The total PAH concentration in the H-PW tanks was approximately 2 $\mu\text{g l}^{-1}$. From the literature the lowest observed effect concentrations (LOEC) of PAHs on fish larvae are reported to range from 1-23 $\mu\text{g l}^{-1}$ (reviewed in (Carls et al., 2008). In addition to APs and PAHs, PW also contains a large “hump” of unresolved complex mixture (UCM), which is composed of a large number of unknown compounds that likely also contribute to the toxicity (Neff et al., 2000; Rowland et al., 2001; Booth et al., 2007; Melbye et al., 2009).

4.5. Effects of PW on cod juveniles

Cod at later developmental stages (after metamorphosis) appear to be more robust and their survival and growth are not affected by exposure to PW (up to 1 %). Plasma and liver

tissue from cod juveniles was investigated for expression of two well known biomarkers, Vtg and CYP1A, respectively. These biomarker studies showed a significant induction of CYP1A in the livers of cod in the H-PW group (no effect was found in the M-PW or L-PW groups). Up-regulation of CYP1A is a marker for an increase in PAH metabolism (Billiard et al., 2002; Whyte et al., 2000). Vtg was also up-regulated in the fish exposed to H-PW. Vtg is a biomarker that is up-regulated after exposure to oestrogen and other oestrogenic chemicals (Rotchell and Ostrander, 2003). Although the Vtg up-regulation in the H-PW group was only 3% of that observed in the E₂ treated fish (Figure 9), it is still a clear indication that PW is a potent source of oestrogenic compounds.

Livers from the cod juveniles contain approximately 20 µg kg⁻¹ of C₂-C₅ APs (Figure 3C). Until recently, it was the release of long chain APs (octylphenol and nonylphenol) into the environment that was considered to be of most concern, as they were thought to be the most potent agonists of the oestrogen receptor (ER). However, Tollefsen and Nilsen (2008) have now shown that a large spectrum of AP isomers, including short chain APs, can act as ER agonists. Previous *in vitro* studies have also shown that PW contains oestrogenic compounds, and most studies have focussed on the well known oestrogenic effects of APs (Thomas et al., 2004a; Thomas et al., 2004b; Tollefsen et al., 2007). However, PW contains many unknown compounds that might have oestrogenic effects. For example, naphthenic acids present in PW can function as xeno-estrogens (Thomas et al., 2009). Many laboratories, including our own, are currently working towards a better understanding of the mechanisms behind the effects of PW in fish. For example, material from this study has been analysed using proteomics in order to search for new biomarkers, and changes in the larval

protein profile were observed even at the lowest PW concentration (0.001%) (Bohne-Kjersem et al., 2010). Several reports have also been published which have used microarray analysis to study the effects of PW at the mRNA level (Olsvik et al., 2007; Holth et al., 2008).

4.6. Effects of E₂ exposure on the early life stages of cod.

Severe effects were observed in the cod that had been exposed to E₂, but different effects were seen depending on the developmental stage of the fish at the time of exposure. In late juveniles, E₂ inhibited growth by more than 50% (Expt. 2B), but no effect on growth was seen in fish that had been exposed to E₂ during the embryo and larval stages (Expt. 2A). Previous studies with tilapia (*Oreochromis niloticus*) have shown similar growth-inhibiting effects of oestrogens (E₂ and 17 α -ethinylestradiol). Oestrogen exposure produced significant changes in the expression of insulin-like growth factor I (IGF-I) mRNA in the liver, and also of IGF-I and growth hormone (GH) mRNA in the brain (Shved et al., 2007; Davis et al., 2008; Shved et al., 2008).

The biomarker analysis of cod plasma (from Expt. 2B) showed an enormous induction of Vtg in the plasma of E₂ treated fish. These observations are in agreement with our own previously published work, which showed that Vtg levels were induced to extreme levels in adult male cod following E₂ treatment (Meier et al., 2007). E₂ also strongly down-regulated expression of CYP1A via a mechanism that likely involves crosstalk between the ER and the AhR (Safe et al., 2000; Navas and Segner, 2001).

4.7. Environmental implications of our results compared to the situation in the field

Field studies conducted with fish, mussels, and passive samplers in cages have shown that it is possible to detect an increased uptake of PAHs and APs up to several kilometres from the platform (Hylland et al., 2008; Durell et al., 2006; Johnsen and Røe, 1998; Tollefsen et al., 2005; Harman et al., 2009). Biomarker analyses have also shown that fish held in cages close to the PW discharge outlet show a small but detectable induction of Vtg and CYP1A (King et al., 2005; Scott et al., 2006a; Zhu et al., 2008). In addition, a size dependent increase in the levels of Vtg in plasma has been found in wild male cod from the North Sea (Scott et al., 2006b). Scott *et al* (2006b) suggest that oestrogenic exposure may originate from compounds that are biomagnified up through the food chain, but their results also showed that cod with increased Vtg levels were not only found in areas with high oil production activity.

It is important to point out that no definite proof that wild fish caught between North Sea oil fields are adversely affected by PW has been found (Grøsvik et al., 2007; Hylland et al., 2008). On the other hand, wild haddock caught in the Tampen region (an area with high oil production activity) showed increased levels of PAH metabolites in the bile and an increase in the occurrence of DNA adducts in the liver when compared with haddock caught in the Egersund Bank (an area with no oil activity). No similar adverse effects were found in cod or saithe (*Pollachius virens*) caught in the Tampen region (Hylland et al., 2006; Grøsvik et al., 2007).

Results presented here show that the cod yolk sac larvae stage were the most sensitive to the harmful effects of PW. Our data demonstrate that the lowest observable effect

concentration (LOEC) on yolk sac larvae is between 0.1% and 1% PW. But due to the dilution factor this concentration can only be expected to be found very close to oil platforms. After PW is discharged into the sea it is quickly diluted and the bioactive compounds will most likely be diluted to a concentration that does not produce any large scale harmful biological effects (Durell et al., 2006; Neff et al., 2006).

However, due to the sheer volume discharged into the North Sea (and discharges are forecast to continue rising until at least 2012-2014), PW is still considered to be a major source of environmental pollution. The concerns about PW have led the Norwegian government to enforce a strict “zero-discharge” policy for all oil exploration activities in the Norwegian Arctic areas.

622 **Acknowledgements**

623 This work was supported by the Norwegian Research Council, NFR project no. 141213/720
624 and NFR 159189/720. Torill Røe Utvik, Statoil Hydro is acknowledged for supplying the
625 project with PW from the Oseberg C field. The authors would also like to thank Dr. Anders
626 Thorsen for photographing the cod larvae and Frank Midtøy, Turid Solbakken, Kjersti Eline
627 Larsen, Vibeke Lokøy, Julie Skadal, and Anders Fuglevik for technical assistance.

628

Figure legends

Figure 1

Schematic representation of experiments 1, 2A and 2B. Experiment 1 was conducted in 24-well trays and ended on 16/04 (3 dph). Fish in experiment 2A were maintained in 100 L tanks during the PW exposure phase. At the end of the exposure, 200 randomly selected fish from the M-PW, L-PW, U and E₂ treatment groups were transferred to four 500 L tanks containing normal seawater, where they were maintained till December 2004. The fish in the H-PW group died during the exposure, so we were unable to study these fish further (*). In experiment 2B, the fish were maintained in 500 L tanks throughout the experiment. The fish were exposed to PW or E₂ for 78 days, after which they were maintained in normal seawater until December 2004. In the schematic diagram above filled boxes indicate exposure to PW or E₂, while open boxes indicate normal seawater. The time frame of the experiment is shown at the top, where the months of the year are indicated by capital letters. Fertilization (F) and three days post hatch (3 dph) are marked above the relevant dates. Dashed lines indicate the beginning and end of experiments and exposure periods (see Materials and Methods for a detailed description). The total number of fish used at each stage of the experiments is shown by the first line of numbers under the exposure regimen. The second line of text indicates the number of parallel tanks and their volume (or wells for Expt. 1) within the brackets, and the number of groups is indicated outside the brackets.

Figure 2

Stability of the AP concentrations ($\mu\text{g l}^{-1}$) in the undiluted PW over time. The results are given as the sum of all isomers at each molecular size.

Figure 3

Tissue concentration of APs ($\mu\text{g kg}^{-1}$ wet weight) in yolk sac larvae (A), early juveniles (B), and juveniles (C). The results are given as the sum of all isomers at each molecular size. For yolk sac larvae measurements were performed on only one sample ($n=1$) pooled from each of the three parallel tanks corresponding to approximately 2000 larvae (1 g of tissue, wet weight). For early juveniles, measurements were performed on one sample from each of the three parallel tanks ($n=3$). Each sample was made up of ten individual fish (1 g of tissue, wet weight). Results are shown as the average $\pm$ SD. For juveniles the measurements were performed on approximately 0.5 g (wet weight) of liver from 5 individual fish from each of 2 parallel tanks ($n=10$).

Figure 4

Percentage of lethal deformities seen throughout the entire embryo stage in experiment 1. Eggs from each of the 5 spawning pairs (SP1-5) were incubated individually in the wells of 24 well plates (4 parallel plates were analysed for each treatment) filled with the same PW concentrations used in experiments 2A and 2B.

Figure 5

Representative larvae (4 dph) from the U (A) and H-PW (B) groups following exposure to PW at the embryo stage. The average length of the larvae at 4 dph was 4.5 mm. The lack of

pigment and the jaw deformation can clearly be seen on the larvae from the H-PW group. Pictures were retouched in Adobe Photoshop CS4. The clone stamp brush was used to remove foreign particles and contrast was increased using the levels function. As a final retouch sharpening was performed using the unsharp mask filter.

Figure 6

Representative larvae (14 dph) from the U (A) and H-PW (B) groups. The average length of larvae at 14 dph was 5 mm. An arrow shows the stomach. Zooplankton can clearly be seen in the stomach of the larvae from the U group (A), but not in the stomach of the larvae from the H-PW group (B). The larvae from the H-PW group also has deformed jaws.

Figure 7

Survival and growth in Experiment 2A. Survival after 90 days of exposure during the egg and larval stage.

Figure 8

Growth in Experiment 2B. Growth of juveniles, measured as total length (cm), both during and after exposure (July-December).

Figure 9

A. Concentration of Vtg in serum as measured by ELISA. B. Levels of CYP1A in liver as measured by ELISA.

Reference List

- Alkindi, A.Y., Brown, J.A., Waring, C.P., and Collins, J.E., 1996. Endocrine, osmoregulatory, respiratory and haematological parameters in flounder exposed to the water soluble fraction of crude oil. *Journal of Fish Biology* 49, 1291-1305.
- Arukwe, A., 2005. Modulation of brain steroidogenesis by affecting transcriptional changes of steroidogenic acute regulatory (StAR) protein and cholesterol side chain cleavage (P450_{scc}) in juvenile Atlantic salmon (*Salmo salar*) is a novel aspect of nonylphenol toxicity. *Environmental Science & Technology* 39, 9791-9798.
- Arukwe, A. and Goksøyr, A., 1998. Xenobiotics, xenoestrogens and reproduction disturbances in fish. *SARSIA* 83, 225-241.
- Bevan, C.L., Porter, D.M., Prasad, A., Howard, M.J., and Henderson, L.P., 2003. Environmental estrogens alter early development in *Xenopus laevis*. *Environmental Health Perspectives* 111, 488-496.
- Bevan, C.L., Porter, D.M., Schumann, C.R., Bryleva, E.Y., Hendershot, T.J., Liu, H.B., Howard, M.J., and Henderson, L.P., 2006. The endocrine-disrupting compound, nonylphenol, inhibits neurotrophin-dependent neurite outgrowth. *Endocrinology* 147, 4192-4204.
- Billard, R., 1992. Reproduction in rainbow trout: Sex differentiation, dynamics of gametogenesis, biology and preservation of gametes. *Aquaculture* 100, 263-298.
- Billiard, S.M., Hahn, M.E., Franks, D.G., Peterson, R.E., Bols, N.C., and Hodson, P.V., 2002. Binding of polycyclic aromatic hydrocarbons (PAHs) to teleost aryl hydrocarbon receptors (AHRs). *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 133, 55-68.

- 718 Billiard, S.M., Meyer, J.N., Wassenberg, D.M., Hodson, P.V., and Di Giulio, R.T., 2008. Nonadditive
719 effects of PAHs on early vertebrate development: Mechanisms and implications for risk assessment.
720 Toxicological Sciences 105, 5-23.
- 721 Blazquez, M., Zanuy, S., Carrillo, M., and Piferrer, F., 1998. Structural and functional effects of early
722 exposure to estradiol-17 β and 17 α -ethynylestradiol on the gonads of the gonochoristic teleost
723 *Dicentrarchus labrax*. Fish Physiology and Biochemistry 18, 37-47.
- 724 Bohne-Kjersem, A., Bache, N., Meier, S., Nyhammer, G., Roepstorff, P., Sæle, O., Goksøyr, A., and
725 Grøsvik, B.E., 2010. Biomarker candidate discovery in Atlantic cod (*Gadus morhua*) continuously
726 exposed to North Sea produced water from egg to fry. Aquatic Toxicology 96, 280-289.
- 727 Boitsov, S., Meier, S., Klungsøyr, J., and Svardal, A., 2004. Gas chromatography-mass spectrometry
728 analysis of alkylphenols in produced water from offshore oil installations as pentafluorobenzoate
729 derivatives. Journal of Chromatography A 1059, 131-141.
- 730 Boitsov, S., Mjøs, S.A., and Meier, S., 2007. Identification of estrogen-like alkylphenols in produced
731 water from offshore oil installations. Marine Environmental Research 64, 651-665.
- 732 Booth, A.M., Sutton, P.A., Lewis, C.A., Lewis, A.C., Scarlett, A., Chau, W., Widdows, J., and Rowland,
733 S.J., 2007. Unresolved complex mixtures of aromatic hydrocarbons: Thousands of overlooked
734 persistent, bioaccumulative, and toxic contaminants in mussels. Environmental Science & Technology
735 41, 457-464.
- 736 Carls, M.G., Holland, L., Larsen, M., Collier, T.K., Scholz, N.L., and Incardona, J.P., 2008. Fish embryos
737 are damaged by dissolved PAHs, not oil particles. Aquatic Toxicology 88, 121-127.

- 738 Carls, M.G., Rice, S.D., and Hose, J.E., 1999. Sensitivity of fish embryos to weathered crude oil: Part I.
739 Low-level exposure during incubation causes malformations, genetic damage, and mortality in larval
740 Pacific herring (*Clupea pallasii*). Environmental Toxicology and Chemistry 18, 481-493.
- 741 Chiasson, M., Benfey, T.J., and Martin-Robichaud, D.J., 2008. Gonadal differentiation in Atlantic cod,
742 *Gadus Morhua* L., and haddock, *Melanogrammus Aeglefinus* (L.). Acta Ichthyologica et Piscatoria 38,
743 127-133.
- 744 Davis, L.K., Pierce, A.L., Hiramatsu, N., Sullivan, C.V., Hirano, T., and Grau, E.G., 2008. Gender-specific
745 expression of multiple estrogen receptors, growth hormone receptors, insulin-like growth factors
746 and vitellogenins, and effects of 17 β -estradiol in the male tilapia (*Oreochromis mossambicus*).
747 General and Comparative Endocrinology 156, 544-551.
- 748 Debruyne, A.M.H., Wernick, B.G., Stefura, C., McDonald, B.G., Rudolph, B.L., Patterson, L., and
749 Chapmant, P.M., 2007. *In situ* experimental assessment of lake whitefish development following a
750 freshwater oil spill. Environmental Science & Technology 41, 6983-6989.
- 751 Devlin, R.H. and Nagahama, Y., 2002. Sex determination and sex differentiation in fish: an overview
752 of genetic, physiological, and environmental influences. Aquaculture 208, 191-364.
- 753 Durell, G.S., Utvik, T.R., Johnsen, S., Frost, T., and Neff, J.M., 2006. Oil well produced water discharges
754 to the North Sea. Part I: Comparison of deployed mussels (*Mytilus edulis*), semi-permeable
755 membrane devices, and the DREAM model predictions to estimate the dispersion of polycyclic
756 aromatic hydrocarbons. Marine Environmental Research 62, 194-223.
- 757 Elsalini, O.A. and Rohr, K.B., 2003. Phenylthiourea disrupts thyroid function in developing zebrafish.
758 Development Genes And Evolution 212, 593-598.

- 759 Falk-Petersen, I.B., Kjørsvik, E., Lonning, S., Naley, A.M., and Sydnes, L.K., 1985. Toxic effects of
760 hydroxylated aromatic hydrocarbons on marine embryos. SARSIA.
- 761 Ferreira-Leach, A.M.R. and Hill, E.M., 2000. Bioconcentration and metabolism of 4-tert-octylphenol in
762 roach (*Rutilus rutilus*) fry. *Analisis* 28, 789-792.
- 763 Gamble, J.C., Davies, J.M., HAY, S.J., and Dow, F.K., 1987. Mesocosm experiments on the effects of
764 produced water discharges from offshore oil platforms in the northern North sea. SARSIA 72, 383-
765 386.
- 766 Gimeno, S., Gerritsen, A., Bowmer, T., and Komen, H., 1996. Feminization of male carp. *Nature* 384,
767 221-222.
- 768 Gimeno, S., Komen, H., Jobling, S., Sumpter, J., and Bowmer, T., 1998. Demasculinisation of sexually
769 mature male common carp, *Cyprinus carpio*, exposed to 4-tert-pentylphenol during spermatogenesis.
770 *Aquatic Toxicology* 43, 93-109.
- 771 Goksøyr, A., Serigstad, B., Solberg, T.S., and Stegeman, J.J., 1988. Response of cod (*Gadus morhua*)
772 larvae and juveniles to oil exposure detected with anti-cod cytochrome-P-450c IgG and anti-scup
773 cytochrome-P-450E MAb 1-12-3. *Marine Environmental Research* 24, 31-35.
- 774 Gronen, S., Denslow, N., Manning, S., Barnes, S., Barnes, D., and Brouwer, M., 1999. Serum
775 vitellogenin levels and reproductive impairment of male Japanese medaka (*Oryzias latipes*) exposed
776 to 4-tert-octylphenol. *Environmental Health Perspectives* 107, 385-390.
- 777 Grøsvik, B.E., Meier, S., Westrheim, K., Skarphedinsdottir, H., Liewenborg, B., Balk, L., and Klungsøyr,
778 J., 2007. Monitoring of the environmental condition in the water column 2005: Oil hydrocarbons in
779 fish from Norwegian waters. IMR report, Bergen, Norway.

- 780 Harman, C., Thomas, K.V., Tollefsen, K.E., Meier, S., Boyum, O., and Grung, M., 2009. Monitoring the
781 freely dissolved concentrations of polycyclic aromatic hydrocarbons (PAH) and alkylphenols (AP)
782 around a Norwegian oil platform by holistic passive sampling. *Marine Pollution Bulletin* 58, 1671-
783 1679.
- 784 Harris, C.A., Santos, E.M., Janbakhsh, A., Pottinger, T.G., Tyler, C.R., and Sumpter, J.P., 2001.
785 Nonylphenol affects gonadotropin levels in the pituitary gland and plasma of female rainbow trout.
786 *Environmental Science & Technology* 35, 2909-2916.
- 787 Heintz, R.A., Short, J.W., and Rice, S.D., 1999. Sensitivity of fish embryos to weathered crude oil: Part
788 II. Increased mortality of pink salmon (*Oncorhynchus gorbuscha*) embryos incubating downstream
789 from weathered Exxon Valdez crude oil. *Environmental Toxicology and Chemistry* 18, 494-503.
- 790 Hendry, C.I., Martin-Robichaud, D.J., and Benfey, T.J., 2002. Gonadal sex differentiation in Atlantic
791 halibut. *Journal of Fish Biology* 60, 1431-1442.
- 792 Holth, T.F., Nourizadeh-Lillabadi, R., Blaesbjerg, M., Grung, M., Holbech, H., Petersen, G.I., Alestrøm,
793 P., and Hylland, K., 2008. Differential gene expression and biomarkers in zebrafish (*Danio rerio*)
794 following exposure to produced water components. *Aquatic Toxicology* 90, 277-291.
- 795 Hylland, K., Beyer, J., Berntssen, M., Klungsøyr, J., Lang, T., and Balk, L., 2006. May organic pollutants
796 affect fish populations in the North Sea? *Journal of Toxicology and Environmental Health-Part A-
797 Current Issues* 69, 125-138.
- 798 Hylland, K., Tollefsen, K.E., Ruus, A., Jonsson, G., Sundt, R.C., Sanni, S., Røe Utvik, T.I., Johnsen, S.,
799 Nilssen, I., Pinturier, L., Balk, L., Barsiene, J., Marigomez, I., Feist, S.W., and Borseth, J.F., 2008. Water
800 column monitoring near oil installations in the North Sea 2001-2004. *Marine Pollution Bulletin* 56,
801 414-429.

- 802 Ioppolo-Armanios, M.I., Alexander, R., and Kagi, R.I., 1992. Identification and analysis of C₀-C₃ phenol
803 in Australian crude oils. *Organic Geochemistry* 18, 603-609.
- 804 Jobling, S. and Sumpter, J.P., 1993. Detergent components in sewage effluent are weakly oestrogenic
805 to fish: An *in vitro* study using rainbow trout (*Oncorhynchus mykiss*) hepatocytes. *Aquatic Toxicology*
806 27, 361-372.
- 807 Johnsen, S. and Røe, T.I., 1998. Dilution and bioavailability of produced water compounds in the
808 northern North Sea. A combined modeling and field study. SPE 46269. Society of Petroleum
809 Engineers, Richardson, TX 1-6.
- 810 Kang, I.J., Yokota, H., Oshima, Y., Tsuruda, Y., Hano, T., Maeda, M., Imada, N., Tadokoro, H., and
811 Honjo, T., 2003. Effects of 4-nonylphenol on reproduction of Japanese medaka, *Oryzias latipes*.
812 *Environmental Toxicology and Chemistry* 22, 2438-2445.
- 813 Karlsson, J., von Hofsten, J., and Olsson, P.E., 2001. Generating transparent zebrafish: A refined
814 method to improve detection of gene expression during embryonic development. *Marine*
815 *Biotechnology* 3, 522-527.
- 816 King, S.C., Johnson, J.E., Haasch, M.L., Ryan, D.A.J., Ahokas, J.T., and Burns, K.A., 2005. Summary
817 results from a pilot study conducted around an oil production platform on the Northwest Shelf of
818 Australia. *Marine Pollution Bulletin* 50, 1163-1172.
- 819 Kjesbu, O.S., 1989. The spawning activity of cod, *gadus-morhua* L. *Journal of Fish Biology* 34, 195-206.
- 820 Knorr, S. and Braunbeck, T., 2002. Decline in reproductive success, sex reversal, and developmental
821 alterations in Japanese medaka (*Oryzias latipes*) after continuous exposure to octylphenol.
822 *Ecotoxicology and Environmental Safety* 51, 187-196.

- 823 Kristiansen, T.S., Jorstad, K.E., Ottera, H., Paulsen, O.I., and Svasand, T., 1997. Estimates of larval
824 survival of cod by releases of genetically marked yolk-sac larvae. *Journal of Fish Biology* 51, 264-283.
- 825 Meier, S., Andersen, T.E., Norberg, B., Thorsen, A., Taranger, G.L., Kjesbu, O.S., Dahle, R., Morton,
826 H.C., Klungsøyr, J., and Svardal, A., 2007. Effects of alkylphenols on the reproductive system of
827 Atlantic cod (*Gadus morhua*). *Aquatic Toxicology* 81, 207-218.
- 828 Meier, S., Klungsøyr, J., Boitsov, S., Eide, T., and Svardal, A., 2005. Gas chromatography-mass
829 spectrometry analysis of alkylphenols in cod (*Gadus morhua*) tissues as pentafluorobenzoate
830 derivatives. *Journal of Chromatography A* 1062, 255-268.
- 831 Melbye, A.G., Brakstad, O.G., Hokstad, J.N., Gregersen, I.K., Hansen, B.H., Booth, A.M., Rowland, S.J.,
832 and Tollefsen, K.E., 2009. Chemical and toxicological characterization of an unresolved complex
833 mixture-rich biodegraded crude oil. *Environmental Toxicology and Chemistry* 28, 1815-1824.
- 834 Miles-Richardson, S.R., Pierens, S.L., Nichols, K.M., Kramer, V.J., Snyder, E.M., Snyder, S.A., Render,
835 J.A., Fitzgerald, S.D., and Giesy, J.P., 1999. Effects of waterborne exposure to 4-nonylphenol and
836 nonylphenol ethoxylate on secondary sex characteristics and gonads of fathead minnows
837 (*Pimephales promelas*). *Environmental Research* 80, S122-S137.
- 838 Navas, J.M. and Segner, H., 2001. Estrogen-mediated suppression of cytochrome P4501A (CYP1A)
839 expression in rainbow trout hepatocytes: role of estrogen receptor. *Chemico-Biological Interactions*
840 138, 285-298.
- 841 Neff, J.M., 2002. Bioaccumulation in marine organisms. Effect of contaminants from oil well
842 produced water. Elsevier Science Ltd, Oxford, UK.

- 843 Neff, J.M., Johnsen, S., Frost, T.K., Utvik, T.I.R., and Durell, G.S., 2006. Oil well produced water
 844 discharges to the North Sea. Part II: Comparison of deployed mussels (*Mytilus edulis*) and the DREAM
 845 model to predict ecological risk. *Marine Environmental Research* 62, 224-246.
- 846 Neff, J.M., Ostazeski, S., Gardiner, W., and Stejskal, I., 2000. Effects of weathering on the toxicity of
 847 three offshore Australian crude oils and a diesel fuel to marine animals. *Environmental Toxicology*
 848 *and Chemistry* 19, 1809-1821.
- 849 Nilsen, B.M., Berg, K., and Goksøyr, A., 1998. Induction of Cytochrome P450 1A (CYP1A) in Fish: A
 850 biomarker for Environmental Pollution. in: Phillips, I.R. and Shephard, E.A. (Eds.), *Methods in*
 851 *Molecular Biology*, Vol 107: Cytochrome P450 Protocols. Humana Press Inc, Totowa, NJ, pp. 423-438.
- 852 Olsvik, P.A., Lie, K.K., Stavrum, A.K., and Meier, S., 2007. Gene-expression profiling in gill and liver of
 853 zebrafish exposed to produced water. *International Journal of Environmental Analytical Chemistry*
 854 87, 195-210.
- 855 Paine, M.D., Leggett, W.C., McRuer, J.K., and Frank, K.T., 1992. Effects of hibernia crude-oil on capelin
 856 (*mallothus-villosus*) embryos and larvae. *Marine Environmental Research* 33, 159-187.
- 857 Petersen, G.I. and Kristensen, P., 1998. Bioaccumulation of lipophilic substances in fish early life
 858 stages. *Environmental Toxicology and Chemistry* 17, 1385-1395.
- 859 Pollino, C.A. and Holdway, D.A., 2002. Toxicity testing of crude oil and related compounds using early
 860 life stages of the crimson-spotted rainbowfish (*Melanotaenia fluviatilis*). *Ecotoxicology and*
 861 *Environmental Safety* 52, 180-189.
- 862 Rotchell, J.M. and Ostrander, G.K., 2003. Molecular markers of endocrine disruption in aquatic
 863 organisms. *Journal Of Toxicology And Environmental Health-Part B-Critical Reviews* 6, 453-495.

- 864 Rowland, S., Donkin, P., Smith, E., and Wraige, E., 2001. Aromatic hydrocarbon "humps" in the
865 marine environment: Unrecognized toxins? Environmental Science & Technology 35, 2640-2644.
- 866 Rye, H., Reed, M., and Ekrol, N., 1998. Sensitivity analysis and simulation of dispersed oil
867 concentrations in the North Sea with the PROVANN model. Environmental Modeling & Software 13,
868 421-429.
- 869 Safe, S., Wormke, M., and Samudio, I., 2000. Mechanisms of inhibitory aryl hydrocarbon receptor-
870 estrogen receptor crosstalk in human breast cancer cells. Journal Of Mammary Gland Biology And
871 Neoplasia 5, 295-306.
- 872 Scott, A.P., Kristiansen, S.I., Katsiadaki, I., Thain, J., Tollefsen, K.E., Goksøyr, A., and Barry, J., 2006a.
873 Assessment of Oestrogen Exposure in Cod (*Gadus morhua*) and Saithe (*Pollachius Virens*) in Relation
874 to their Proximity to an Oilfield. in: Hylland, K., Lang, T., and Vethaak, A.D. (Eds.), Biological Effects of
875 Contaminants in Marine Pelagic Ecosystems Society of Environmental Toxicology and Chemistry
876 (SETAC), pp. 329-339.
- 877 Scott, A.P., Katsiadaki, I., Witthames, P.R., Hylland, K., Davies, I.M., McIntosh, A.D., and Thain, J.,
878 2006b. Vitellogenin in the blood plasma of male cod (*Gadus morhua*): A sign of oestrogenic
879 endocrine disruption in the open sea? Marine Environmental Research 61, 149-170.
- 880 Seki, M., Yokota, H., Maeda, M., Tadokoro, H., and Kobayashi, K., 2003. Effects of 4-nonylphenol and
881 4-tert-octylphenol on sex differentiation and vitellogenin induction in medaka (*Oryzias latipes*).
882 Environmental Toxicology and Chemistry 22, 1507-1516.
- 883 Servos, M.R., 1999. Review of the aquatic toxicity, estrogenic responses and bioaccumulation of
884 alkylphenols and alkylphenol polyethoxylates. Water Quality Research Journal of Canada 34, 123-
885 177.

- 886 Shved, N., Berishvili, G., Baroiller, J.F., Segner, H., and Reinecke, M., 2008. Environmentally relevant
887 concentrations of 17 α -ethinylestradiol (EE2) interfere with the growth hormone (GH)/insulin-like
888 growth factor (IGF)-I system in developing bony fish. *Toxicological Sciences* 106, 93-102.
- 889 Shved, N., Berishvili, G., D'Cotta, H., Baroiller, J.F., Segner, H., Eppler, E., and Reinecke, M., 2007.
890 Ethinylestradiol differentially interferes with IGF-I in liver and extrahepatic sites during development
891 of male and female bony fish. *Journal of Endocrinology* 195, 513-523.
- 892 Stephens, S.M., Alkindi, A.Y., Waring, C.P., and Brown, J.A., 1997. Corticosteroid and thyroid
893 responses of larval and juvenile turbot exposed to the water-soluble fraction of crude oil. *Journal of*
894 *Fish Biology* 50, 953-964.
- 895 Stephens, S.M., Frankling, S.C., Stagg, R.M., and Brown, J.A., 2000. Sub-lethal effects of exposure of
896 juvenile turbot to oil produced water. *Marine Pollution Bulletin* 40, 928-937.
- 897 Sulaimon, S.S. and Kitchell, B.E., 2003. The biology of melanocytes. *Veterinary Dermatology* 14, 57-
898 65.
- 899 Sundt, R.C., Meier, S., Jonsson, G., Sanni, S., and Beyer, J., 2009. Development of a laboratory
900 exposure system using marine fish to carry out realistic effect studies with produced water
901 discharged from offshore oil production. *Marine pollution bulletin* 58, 1382-1388.
- 902 Tanaka, J.N. and Grizzle, J.M., 2002. Effects of nonylphenol on the gonadal differentiation of the
903 hermaphroditic fish, *Rivulus marmoratus*. *Aquatic Toxicology* 57, 117-125.
- 904 Thomas, K.V., Balaam, J., Hurst, M.R., and Thain, J.E., 2004a. Bio-analytical and chemical
905 characterisation of offshore produced water effluents for estrogen receptor (ER) agonists. *Journal of*
906 *Environmental Monitoring* 6, 593-598.

- 907 Thomas, K.V., Balaam, J., Hurst, M.R., and Thain, J.E., 2004b. Identification of *in vitro* estrogen and
908 androgen receptor agonists in North Sea offshore produced water discharges. Environmental
909 Toxicology and Chemistry 23, 1156-1163.
- 910 Thomas, K.V., Langford, K., Petersen, K., Smith, A.J., and Tollefsen, K.E., 2009. Effect-directed
911 identification of naphthenic acids as important *in vitro* xeno-estrogens and anti-androgens in North
912 Sea offshore produced water discharges. Environmental Science and Technology 43, 8066-8071.
- 913 Tilseth, S., Solberg, T.S., and Westrheim, K., 1984. Sublethal effects of the water-soluble fraction of
914 Ekofisk crude oil on early larval stages of cod (*Gadus morhua* L.). Marine Environmental Research 11,
915 1-16.
- 916 Tollefsen, K.E., Bøyum, O., Kringstad, A., Finne, E.F., Odden, E., and Grung, M., 2005. Water Column
917 Survey 2004. Water column monitoring of polar chemicals by use of POCIS passive samplers. NIVA
918 Report: 4940-2005 , Oslo, Norway.
- 919 Tollefsen, K.E., Harman, C., Smith, A., and Thomas, K.V., 2007. Estrogen receptor (ER) agonists and
920 androgen receptor (AR) antagonists in effluents from Norwegian North Sea oil production platforms.
921 Marine Pollution Bulletin 54, 277-283.
- 922 Tollefsen, K.E. and Nilsen, A.J., 2008. Binding of alkylphenols and alkylated non-phenolics to rainbow
923 trout (*Oncorhynchus mykiss*) hepatic estrogen receptors. Ecotoxicology and Environmental Safety 69,
924 163-172.
- 925 van Aerle, R., Pounds, N., Hutchinson, T.H., Maddix, S., and Tyler, C.R., 2002. Window of sensitivity
926 for the estrogenic effects of ethinylestradiol in early life-stages of fathead minnow, *Pimephales*
927 *promelas*. Ecotoxicology 11, 423-434.

- 928 Wang, W.D., Wang, Y., Wen, H.J., Buhler, D.R., and Hu, C.H., 2004. Phenylthiourea as a weak activator
929 of aryl hydrocarbon receptor inhibiting 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced CYP1A1
930 transcription in zebrafish embryo. *Biochemical Pharmacology* 68, 63-71.
- 931 Weber, L.P., Hill, R.L., Jr., and Janz, D.M., 2003. Developmental estrogenic exposure in zebrafish
932 (*Danio rerio*): II. Histological evaluation of gametogenesis and organ toxicity. *Aquatic Toxicology* 63,
933 431-446.
- 934 Weber, L.P., Kiparissis, Y., Hwang, G.S., Niimi, A.J., Janz, D.M., and Metcalfe, C.D., 2002. Increased
935 cellular apoptosis after chronic aqueous exposure to nonylphenol and quercetin in adult medaka
936 (*Oryzias latipes*). *Comparative Biochemistry and Physiology C-Pharmacology Toxicology &*
937 *Endocrinology* 131, 51-59.
- 938 White, R., Jobling, S., Hoare, S.A., Sumpter, J.P., and Parker, M.G., 1994. Environmentally persistent
939 alkylphenolic compounds are estrogenic. *Endocrinology* 135, 175-182.
- 940 Whyte, J.J., Jung, R.E., Schmitt, C.J., and Tillitt, D.E., 2000. Ethoxyresorufin-O-deethylase (EROD)
941 activity in fish as a biomarker of chemical exposure. *Critical Reviews in Toxicology* 30, 347-570.
- 942 Yadetie, F. and Male, R., 2002. Effects of 4-nonylphenol on gene expression of pituitary hormones in
943 juvenile Atlantic salmon (*Salmo salar*). *Aquatic Toxicology* 58, 113-129.
- 944 Yokota, H., Abe, T., Nakai, M., Murakami, H., Eto, C., and Yakabe, Y., 2005. Effects of 4-tert-
945 pentylphenol on the gene expression of P450 11 β -hydroxylase in the gonad of medaka (*Oryzias*
946 *latipes*). *Aquatic Toxicology* 71, 121-132.
- 947 Zhu, S.Q., King, S.C., and Haasch, M.L., 2008. Biomarker induction in tropical fish species on the
948 Northwest Shelf of Australia by produced formation water. *Marine Environmental Research* 65, 315-
949 324.

Groups	Dilution factor	Estimated distance from platform (m)
High (H-PW)	1:100 (1 %)	0-50
Medium (M-PW)	1:1000 (0.1 %)	50-1000
Low (L-PW)	1:10000 (0.01 %)	>2000
Oestrogen (E)	10 µg/l 17β-estradiol	-

Table 1

Theoretical dilution factors and estimated environmental relevance of the PW concentrations used in this study.

	Average $\pm$ SD	RSD (%)
Phenol	4696 $\pm$ 967	21
Σ C1	7542 $\pm$ 1334	18
Σ C2	638 $\pm$ 202	32
Σ C3	216 $\pm$ 58	27
Σ C4	53 $\pm$ 15	28
Σ C5	4,7 $\pm$ 2,5	53
Σ C6	0,10 $\pm$ 0,04	35
Σ C7	0,02 $\pm$ 0,01	70
Σ C8	0,02 $\pm$ 0,02	96
Σ C9	0,09 $\pm$ 0,05	61

Table 2

Average AP concentrations ($\mu\text{g/L}$) in the undiluted PW throughout the whole exposure period. The values shown are the average concentrations of 11 separate measurements ($n=11$) taken over a period of 154 days. The results are given as the sum of all isomers at each molecular size.

Experiment 2A

Concentration ($\mu\text{g l}^{-1}$)	U	L-PW	M-PW	H-PW
Phenol	$1,04 \pm 0,00$	$3,60 \pm 3,83$	$3,41 \pm 3,60$	$7,33 \pm 3,33$
Σ C1	$0,81 \pm 0,98$	$0,75 \pm 0,70$	$0,89 \pm 0,79$	$7,41 \pm 1,63$
Σ C2	$0,04 \pm 0,00$	$0,13 \pm 0,04$	$0,42 \pm 0,08$	$3,90 \pm 0,20$
Σ C3	$0,04 \pm 0,02$	$0,09 \pm 0,01$	$0,34 \pm 0,20$	$2,58 \pm 1,84$
Σ C4	$0,015 \pm 0,007$	$0,027 \pm 0,002$	$0,067 \pm 0,008$	$0,399 \pm 0,109$
Σ C5	$0,001 \pm 0,001$	$0,003 \pm 0,002$	$0,004 \pm 0,001$	$0,041 \pm 0,013$
$\Sigma \geq \text{C6}$	ND	ND	ND	ND

Experiment 2B

Concentration ($\mu\text{g l}^{-1}$)	U	L-PW	M-PW	H-PW
Phenol	$2,96 \pm 0,66$	$5,33 \pm 1,84$	$10,29 \pm 6,68$	$29,74 \pm 11,33$
Σ C1	$0,46 \pm 0,10$	$0,81 \pm 0,49$	$11,03 \pm 12,16$	$51,09 \pm 24,64$
Σ C2	$0,04 \pm 0,00$	$0,08 \pm 0,03$	$1,67 \pm 1,93$	$9,22 \pm 3,76$
Σ C3	$0,01 \pm 0,00$	$0,02 \pm 0,01$	$0,27 \pm 0,18$	$2,24 \pm 0,46$
Σ C4	$0,014 \pm 0,002$	$0,018 \pm 0,005$	$0,060 \pm 0,033$	$0,401 \pm 0,089$
Σ C5	$0,006 \pm 0,004$	$0,007 \pm 0,005$	$0,010 \pm 0,005$	$0,034 \pm 0,007$
$\Sigma \geq \text{C6}$	ND	ND	ND	ND

Table 3

Concentration of APs in each exposure group in experiment 2A and 2B. Individual measurements were made on two separate days in experiment 2A (n=2) and on four separate days in experiment 2B (n=4). The results are given as the sum of all isomers at each molecular size.

Experiment 2A				
Dilution factor (%)	U	L-PW	M-PW	H-PW
Phenol	0,02 ± 0,02	0,08 ± 0,07	0,07 ± 0,07	0,17 ± 0,03
Σ C1	0,01 ± 0,01	0,01 ± 0,01	0,01 ± 0,01	0,11 ± 0,05
Σ C2	0,01 ± 0,00	0,03 ± 0,01	0,11 ± 0,02	0,97 ± 0,08
Σ C3	0,02 ± 0,00	0,04 ± 0,01	0,14 ± 0,04	1,07 ± 0,43
Σ C4	0,03 ± 0,02	0,06 ± 0,01	0,14 ± 0,03	0,86 ± 0,33
Σ C5	0,02 ± 0,02	0,09 ± 0,08	0,12 ± 0,05	1,41 ± 0,65
Σ ≥C6	-	-	-	-
Experiment 2B				
Dilution factor (%)	U	L-PW	M-PW	H-PW
Phenol	0,06 ± 0,01	0,11 ± 0,05	0,20 ± 0,10	0,60 ± 0,24
Σ C1	0,01 ± 0,00	0,01 ± 0,00	0,13 ± 0,11	0,66 ± 0,27
Σ C2	0,00 ± 0,00	0,01 ± 0,00	0,19 ± 0,20	1,12 ± 0,32
Σ C3	0,01 ± 0,00	0,01 ± 0,01	0,16 ± 0,11	1,47 ± 0,66
Σ C4	0,03 ± 0,01	0,04 ± 0,02	0,12 ± 0,05	0,83 ± 0,31
Σ C5	0,13 ± 0,08	0,15 ± 0,10	0,21 ± 0,11	0,73 ± 0,17
Σ ≥C6	-	-	-	-

Table 4

Empirical dilution factors of the APs in each exposure group in experiments 2A and 2B. The results are given as the sum of all isomers at each molecular size.

	PW	H-PW	M-PW	L-PW	U
Naphthalene	242,9	0,791	0,173	0,046	0,020
C1-naphthalene	237,5	0,584	0,288	0,121	0,231
C2-naphthalene	97,5	0,303	0,072	ND	ND
Acenaphthylene	2,7	ND	ND	ND	ND
Acenaphthene	4,4	ND	ND	ND	ND
Fluorene	8,1	0,027	0,017	ND	ND
Anthracene	1,3	ND	ND	ND	ND
Phenanthrene	17,8	0,050	0,028	ND	0,032
C1-Phenanthrene	16,6	ND	ND	ND	ND
Pyrene	0,5	ND	ND	ND	ND
Benz(a)anthracene	0,8	ND	ND	ND	ND
Perylene	ND	ND	ND	ND	ND
Dibenz(a,h)anthracene	ND	ND	ND	ND	ND
Total PAHs	630,1	1,8	0,6	0,2	0,3

Table 5

Actual concentrations of PAHs ($\mu\text{g l}^{-1}$) measured in the undiluted PW and in the H-PW, M-PW, and L-PW exposure tanks. The background concentration of PAHs in the normal seawater used in this study was also measured (U). The results shown are from one single measurement (n=1) in August 2004.

	Yolk sac larvae	Early juveniles	Juveniles
C ₂ -phenol	12 ± 7	2 ± 2	0,6 ± 0,4
C ₃ -phenol	14 ± 5	10 ± 7	3 ± 3
C ₄ -phenol	31 ± 15	30 ± 23	30 ± 40
C ₅ -phenol	68 ± 15	74 ± 46	175 ± 169

Table 6

Average bioconcentration factor (BCF = concentration in wet weight tissue/concentration in seawater) at each molecular size for the H-PW group. For C₂, the concentration of 8 isomers were measured and the average is shown in the figure. Six isomers were measured for C₃, seven for C₄ and 2 for C₅. For yolk sac larvae and early juveniles the measurements were performed on pooled samples of whole individuals. One gram of tissue was used for analysis which corresponded to approximately 2000 yolk sac larvae and 10 early juveniles. For juveniles the measurements were performed on approximately 0.5g of liver from individual fish.

Figure 1