Feasibility of diagnosing influenza within 24 hours of symptom onset in children 1-3 years of age

S. Heinonen, H. Silvennoinen, P. Lehtinen, R. Vainionpää, T. Heikkinen

► To cite this version:

S. Heinonen, H. Silvennoinen, P. Lehtinen, R. Vainionpää, T. Heikkinen. Feasibility of diagnosing influenza within 24 hours of symptom onset in children 1-3 years of age. *European Journal of Clinical Microbiology and Infectious Diseases*, 2010, 30 (3), pp.387-392. 10.1007/s10096-010-1098-5 . hal-00636636

HAL Id: hal-00636636

<https://hal.science/hal-00636636>

Submitted on 28 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Feasibility of diagnosing influenza within 24 hours of symptom onset in children 1-3 years of age

Santtu Heinonen, MD,^{1*} Heli Silvennoinen, MD,¹ Pasi Lehtinen, MD, PhD¹

Raija Vainionpää, PhD,² Terho Heikkinen, MD, PhD¹

¹ Department of Pediatrics, Turku University Hospital,

Kiinamyllynkatu 4-8, FI-20520, Turku, Finland

² Department of Virology, University of Turku,

Kiinamyllynkatu 13, FI-20520, Turku, Finland

*Corresponding author.

Mailing address: Department of Pediatrics, Turku University Hospital, Kiinamyllynkatu 4-8,

FI-20520 Turku, Finland.

Phone: +358-2-313-0000.

Fax: +358-2-313-1460.

Email: shhein@utu.fi

Running title:

Early diagnosis of influenza in young children

ABSTRACT

Purpose

Diagnosing influenza at an early stage of illness is important for the initiation of effective antiviral treatment. However, especially in young children, influenza often commences with an abrupt onset of fever, with full-blown respiratory symptoms developing only later. We determined the feasibility of diagnosing influenza in young children already during the first signs of the illness.

Methods

During confirmed influenza activity, we obtained nasal swabs from children aged 1-3 years who presented as outpatients within 24 hours of the onset of fever ($\geq 38.0^{\circ}\text{C}$). The specimens were tested for influenza viruses with viral culture, antigen detection, PCR, and a rapid point-of-care test (Actim Influenza A&B, Medix Biochemica, Finland). In addition, follow-up specimens were obtained from a proportion of children 3-7 days later.

Results

Influenza virus was detected already within 24 hours of symptom onset in 56 of 61 (92%; 95% CI, 82-97%) children in whom influenza was eventually confirmed in the laboratory. A total of 158 rapid tests performed within 24 hours of symptom onset yielded a sensitivity of 90% (95% CI, 74-98%) for influenza A viruses but only 25% (95% CI, 3-61%) for influenza B viruses

($P < 0.001$), resulting in an overall sensitivity of 77% (95% CI, 61-89%) and specificity of 99% (95% CI, 95-100%) for any influenza viruses.

Conclusions

In most young children, influenza can be accurately diagnosed already within 24 hours of symptom onset. The rapid point-of-care test used was sensitive and specific for diagnosing influenza A, but its sensitivity for influenza B was limited.

KEYWORDS:

Influenza

Children

Diagnostic methods

Rapid test

INTRODUCTION

The burden of influenza is substantial in children [1, 2]. The attack rates are highest in this age group [3], and young children are hospitalized at rates similar to those for adults with high risk conditions [4]. Further, influenza predisposes children to bacterial complications such as acute otitis media [1]. Neuraminidase inhibitors oseltamivir and zanamivir can be used for treating influenza in children from 1 and 5 years of age, respectively [5]. These drugs reduce the duration and severity of symptoms when started within 48 hours of symptom onset [6, 7]. The mechanism of action of neuraminidase inhibitors and the results of a previous study in adults [8] suggest that the clinical benefits could be greater if the treatment were initiated at an earlier stage of illness. According to CDC recommendations, the antiviral treatment should be started as quickly as possible after the onset of symptoms [5].

Distinguishing influenza from other viral respiratory infections is difficult in young children because of the nonspecific signs and symptoms of influenza [9], children's inability to describe their subjective symptoms, and high frequency of other viral infections even during an influenza outbreak [2, 10, 11]. In a previous study among children, the overall sensitivity of the clinical diagnosis of influenza was only 38% and the positive predictive value 32% [12]. Therefore, virological methods are needed to optimize the use of antiviral drugs in children. For clinical decision-making, rapid point-of-care tests that provide almost immediate results are most useful. Previous studies have demonstrated that these tests are generally specific, but their sensitivity has varied greatly [13-19].

Infants and young children are frequently brought for clinical examination during an early stage of influenza illness. The clinician's dilemma is that influenza often commences with an abrupt onset of fever and ill appearance, with full-blown respiratory symptoms developing only later. Occasionally, febrile convulsion is the first sign of influenza in young children [20]. There are scarce data on whether influenza in children can be reliably diagnosed from upper respiratory specimens during the early stage of the illness, and limited data exist on the performance of influenza rapid tests in this clinical situation. We determined the feasibility of diagnosing influenza in outpatient children 1-3 years of age who presented within 24 hours of the onset of fever. In addition, we evaluated the diagnostic accuracy of influenza rapid testing performed in this setting.

MATERIALS AND METHODS

Study design

This study was conducted as part of a randomized placebo-controlled trial assessing the effectiveness of oseltamivir treatment started within 24 hours of the onset of symptoms in children (ClinicalTrials.gov; NCT00593502) during the influenza seasons of 2007-08 (predominance of A/H1N1 strains) and 2008-09 (predominance of A/H3N2 strains) at a single site in Turku, Finland [21]. Before each influenza season, children 1-3 years of age were recruited into follow-up cohorts (N=631 in 2007-08 and N=554 in 2008-09). No exclusion

criteria were employed for enrollment. The study was approved by the Ethics Committee of the Hospital District of Southwest Finland. Written informed consent was obtained from the parents of the participating children.

Study procedures

When local surveillance indicated that influenza was circulating in the area, a study clinic was opened. The parents were asked to bring their child to the study clinic within 24 hours if the child developed fever or any signs of respiratory infection. During each visit (n=764), the child was examined by a study physician, and nasal swabs were obtained for virological analyses. Rapid influenza tests (n=188) were routinely performed (according to the study protocol of the oseltamivir trial) during the first visit to all febrile children without respiratory symptoms, but also to any other children according to the decision by the attending physician who considered e.g. time constraints for performing the rapid tests during the busy hours of the influenza epidemic. A follow-up visit was scheduled on day 5-8 for all children who received the study medication. In addition, the parents were invited to bring their child to the study clinic whenever they deemed it necessary. Nasal swabs were obtained during all follow-up visits if the child was symptomatic. All visits were free of charge to the parents.

Virological methods

Three flocked swab specimens (Copan, Italy) were obtained from a depth of 2-3 cm in the nostrils in non-standardized order. The specimens were refrigerated at +6°C until analyzed in the virological laboratory next working day. The samples were subjected to viral culture in Madin-Darby canine kidney cells followed by immunoperoxidase staining with monoclonal antibodies [22] and antigen detection using time-resolved fluoroimmunoassay (TR-FIA) [23]. All samples that remained negative for influenza viruses with these methods were further tested with reverse-transcriptase polymerase-chain-reaction (RT-PCR) assays for influenza A and B viruses.

After collection of specimens for the virological laboratory, another nasal swab specimen for rapid testing was obtained from a proportion of children (as described earlier in “Study procedures”) using the polyester swabs provided with the kit (Actim Influenza A & B, Medix Biochemica, Finland). The manufacturer of these rapid tests was not involved in this study; the authors purchased the test kits at market price. The rapid test was performed at the point of care immediately after the sampling according to the manufacturer’s instructions. All samples were obtained and the rapid tests performed by one of four trained study nurses who also interpreted the results. The personnel performing the other virological analyses were unaware of the rapid test results.

Data analysis

Detection of influenza virus within 24 hours of symptom onset

In this analysis, we included all children in the placebo arm of the treatment trial in whom influenza was eventually confirmed by any of the conventional laboratory methods (viral isolation, TR-FIA, or RT-PCR) at any time within 8 days of the onset of symptoms. Children who had received oseltamivir were excluded from this analysis because the treatment might have decreased the viral load in the children and consequently led to increased numbers of negative test results during the follow-up visits. The sensitivity of detecting influenza virus within 24 hours of symptom onset was determined as the percentage of children (among all influenza-positive children) in whom influenza could be diagnosed in the nasal swabs collected during the first 24 hours of symptoms.

Performance of the influenza rapid test

The sensitivity, specificity, and predictive values of the rapid tests were assessed by comparing the results of the rapid tests obtained within 24 hours of symptom onset (i.e. before the start of any study medication) to those of the simultaneously obtained reference test samples. A reference test was considered positive if any of the methods (viral isolation, TR-FIA, or RT-PCR) yielded a positive result for influenza viruses.

Statistical analysis

Sensitivities, specificities, and predictive values and their binomial 95% confidence intervals were calculated for any influenza virus and for influenza A and B viruses separately. Fisher's exact test was used to compare proportions. Two-sided P values <0.05 were considered statistically significant. All analyses were performed using StatsDirect (version 2.7.7) software.

RESULTS

Detection of influenza virus within 24 hours of symptom onset

Of 409 children randomized to receive the study medication, influenza was eventually confirmed by any of the conventional laboratory methods (viral isolation, TR-FIA, or RT-PCR) in 98 children. Thirty-seven children who had received oseltamivir were excluded from this analysis. Of the remaining 61 placebo-treated children included in the analysis, 50 (82%) had influenza A and 11 (18%) influenza B. The diagnosis was based on viral culture in 54 (89%) children; antigen detection (TR-FIA) provided the diagnosis in 2 (3%) children with negative viral culture, and RT-PCR in 5 (8%) children in whom both viral culture and antigen detection were negative. In 57 (93%) children, nasal swabs were collected also during the follow-up visits. The mean age of the children was 2.5 (SD±0.8) years, and the median time from the onset of fever to the collection of the first nasal swab specimen was 6.5 hours (interquartile range, 3.4-13.3 h). Influenza virus was detected within 24 hours of the onset of fever in 56 of the 61 children, yielding a sensitivity of 92% (95% CI, 82-97%). The sensitivity was 94% (95% CI, 83-99%) for influenza A virus and 82% (95% CI, 48-98 %) for influenza B virus (P=0.2). There were no

statistically significant differences in the detection of the virus between children presenting <12 hours and 12-24 hours from the onset of symptoms ($P>0.9$); between vaccinated and unvaccinated children ($P=0.5$); or between children <2 years and 2-3 years of age ($P>0.9$).

Performance of the influenza rapid test

Among a total of 764 visits to the study clinic for new respiratory symptoms or fever, rapid tests were performed in 188 (25%) cases. The results of 30 rapid tests were excluded from this analysis because of the following reasons: absence of fever (18); duration of fever >24 h (3); absence of reference samples (4); rapid test expired (5). Among the remaining 158 children, influenza virus was detected with one or more of the reference methods in 39 (25%) children (Table 1). The rapid test was positive in 31 children (including one false positive), resulting in an overall sensitivity of 77% (95% CI, 61-89%) and specificity of 99% (95% CI, 95-100%) for any influenza virus. The sensitivity of the rapid test was 90% (95% CI, 74-98%) for influenza A virus and 25% (95% CI, 3-61%) for influenza B virus ($P<0.001$).

DISCUSSION

In children in whom influenza was eventually confirmed, 92% of nasal swab specimens that were obtained within 24 hours of the onset of fever were positive by at least one of the conventionally used laboratory methods. This suggests that children shed viruses in their nasal secretions in sufficient amounts to be detected already during the first day of illness. This

finding is clinically important because accurate diagnosis of influenza during the early stage of illness enables the rational use of antiviral drugs [6, 7] and proper cohorting of hospitalized patients to prevent nosocomial infections [24]. Rapid influenza diagnosis has also been associated with reduced use of antibiotics and shorter duration of hospitalization [25, 26].

The influenza rapid test that we used had an overall sensitivity of 77% and specificity of 99%. Our results are concordant with several previous studies in children, in which the sensitivities of different rapid tests have ranged from 63% to 85% and the specificities from 97% to 99% [13-15, 27]. Two previous studies assessing the performance of the same rapid test that we used found sensitivities of 65% [16] and 53% [17] in children. There are, however, reports of significantly lower sensitivities (ranging from 20% to 33%) for various rapid tests in populations consisting mainly of adults [28, 29]. Lower sensitivities (27-44 %) have been observed also in some studies with mainly pediatric populations [18, 19].

There are several potential explanations for the great variability between the sensitivities observed in different studies. Children shed influenza viruses in larger amounts and for longer periods than adults [30-32], which may account for the increased sensitivity observed in children [17, 32-34]. Also, the delay between the onset of symptoms and the collection of the specimen may affect the sensitivity of the rapid test. In experimental volunteer studies in adults, the viral load peaked on the second day after virus inoculation, and the shedding lasted for approximately 5 days [35]. In children, the viral shedding has been described to peak 1-3 days after the onset of illness and to last for 6-7 days [30, 31, 36]. In a Japanese pediatric study

in which most samples were obtained within 24 hours of the onset of fever, the overall sensitivity of an influenza rapid test was the same as in the present study (77%) [37]. In that study, the sensitivity of the rapid test was lower for specimens collected within 9 hours (64-69%) compared with specimens collected within 9-24 hours (83 -100%) of the onset of fever. A study conducted in Nicaraguan children found a higher sensitivity (74-75%) of influenza rapid tests for samples obtained 2-3 days after symptom onset, compared with samples obtained either 1 or 4 days after symptom onset (52-58%) [38]. In a recent study from Greece, the sensitivity of a rapid test was 76% when performed within 2 days of symptom onset and 65% when performed later during the course of the illness [27].

Presumably the most important factor affecting the yield of influenza virus is the quality of the sample, which, in turn, is dependent on the site and method of collection. A nasopharyngeal aspirate is often considered as the specimen of choice for everyday clinical purposes in children. The sensitivity of a nasal swab is, however, 91-92% when compared with a nasopharyngeal aspirate for the detection of influenza viruses in children, and the method is more pleasant for the patient and easy to perform in any setting [39, 40]. The sensitivity of nasal swabs may be further enhanced by using flocked swabs [41, 42] or collecting the swab from the nasopharynx [15].

We found the sensitivity of the rapid test to be high for influenza A virus but significantly lower for influenza B virus. Such significant differences were not seen in another study with Actim Influenza A&B rapid test, in which the sensitivities for influenza A and B viruses were 69% and

60%, respectively [16]. There are, however, previous reports of a lower sensitivity for influenza B virus with other rapid tests performed at different stages of illness. [13, 43, 44]. One potential explanation for the varying sensitivities of rapid tests for influenza A and B viruses was provided by a recent study from Hong Kong that demonstrated that viral shedding in naturally acquired influenza A infections peaked already at the time of symptom onset, whereas in influenza B infections the peak was reached only 3 days after the onset of symptoms [45]. Besides varying rates of viral replication, different assay methods and immunological differences might potentially interfere with the detection of influenza B viruses. In our study, however, the total number of children infected with influenza B viruses was too low for any firm conclusions to be made, and further studies in this area are warranted.

The main strength of the present study is that it was performed in real-life conditions. The patients were unselected young children who were examined at a primary care clinic during the acute phase of their illness. The rapid tests were performed immediately at the point of care by trained nurses, which reflects the actual situation in most clinical settings. In many previous studies, the rapid tests have been evaluated by using stored specimens in controlled laboratory conditions.

Our study has some limitations. We used flocked swabs for obtaining samples for the virology laboratory but a polyester swab included in the kit for the rapid test. This might have resulted in better-quality reference samples and thus underestimation of the sensitivity of the rapid test. However, using the exact swabs provided with the kit more likely represents the test's true

performance in the clinical practice. A total of three or four nasal swab specimens were obtained from the children during each visit, and consequently the representativeness of the samples may have varied. In addition, most of the follow-up samples were collected on study days 6-7. At that time, the viral load may have already been declining, and therefore we may have missed some children who actually had influenza. However, all samples that remained negative with viral culture and TR-FIA were further tested with RT-PCR and therefore we believe that potential variations in the quality of the samples or in the concentrations of viruses in the specimens are unlikely to have significantly affected our results. Finally, due to practical reasons, we obtained specimens for rapid testing only in approximately one-fourth of all cases. However, we have no reason to believe that this would have resulted in any major selection bias because the decisions to perform the rapid tests were made independently by several study physicians who worked at varying times, and there were no recommendations about which children should be preferably sampled for rapid testing.

In conclusion, our results indicate that influenza can be reliably diagnosed in children 1-3 years of age already during the first 24 hours of illness. Further, the influenza rapid test had a high specificity and sensitivity for diagnosing influenza A viruses, but its sensitivity for influenza B viruses was poor.

ACKNOWLEDGEMENTS

F. Hoffmann-La Roche Ltd. (Basel, Switzerland) provided financial support to the Hospital District of Southwest Finland (Turku University Hospital) for carrying out this investigator-initiated study. Dr Heinonen was supported by grants from Turku University Hospital Foundation and the Foundation for Paediatric Research. The sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation of the manuscript. The authors declare that they have no conflict of interest.

REFERENCES

- [1] Heikkinen T, Silvennoinen H, Peltola V, Ziegler T, Vainionpää R, Vuorinen T, Kainulainen L, Puhakka T, Jartti T, Toikka P, Lehtinen P, Routi T, Juven T (2004) Burden of influenza in children in the community. *J Infect Dis* 190(8):1369-1373
- [2] Poehling KA, Edwards KM, Weinberg GA, Szilagyi P, Staat MA, Iwane MK, Bridges CB, Grijalva CG, Zhu YW, Bernstein DI, Herrera G, Erdman D, Hall CB, Seither R, Griffin MR, for the New Vaccine Surveillance Network (2006) The underrecognized burden of influenza in young children. *N Engl J Med* 355(1):31-40
- [3] Monto AS, Sullivan KM (1993) Acute respiratory illness in the community. Frequency of illness and the agents involved. *Epidemiol Infect* 110 (1):145-160
- [4] Neuzil KM, Mellen BG, Wright PF, Mitchel EF, Griffin MR (2000) The effect of influenza on hospitalizations, outpatient visits, and courses of antibiotics in children. *N Engl J Med* 342(4):225-231
- [5] CDC (2009) Centers for Disease Control and Prevention: Updated interim recommendations for the use of antiviral medications in the treatment and prevention of influenza for the 2009-2010 season. <http://www.cdc.gov/H1N1flu/recommendations.htm>. Accessed 15 March 2010
- [6] Whitley RJ, Hayden FG, Reisinger KS, Young N, Dutkowski R, Ipe D, Mills RG, Ward P (2001) Oral oseltamivir treatment of influenza in children. *Pediatr Infect Dis J* 20(2):127-133
- [7] Hedrick JA, Barzilai A, Behre U, Henderson FW, Hammond J, Reilly L, Keene O (2000) Zanamivir for treatment of symptomatic influenza A and B infection in children five to twelve years of age: a randomized controlled trial. *Pediatr Infect Dis J* 19(5):410-417

- [8] Aoki FY, Macleod MD, Paggiaro P, Carewicz O, El Sawy A, Wat C, Griffiths M, Waalberg E, Ward P, on behalf of the IMPACT study group (2003) Early administration of oral oseltamivir increases the benefits of influenza treatment. *J Antimicrob Chemother* 51(1):123-129
- [9] Silvennoinen H, Peltola V, Lehtinen P, Vainionpää R, Heikkinen T (2009) Clinical presentation of Influenza in unselected children treated as outpatients. *Pediatr Infect Dis J* 28(5):372-375
- [10] Heikkinen T, Ziegler T, Peltola V, Lehtinen P, Toikka P, Lintu M, Jartti T, Juven T, Kataja J, Pulkkinen J, Kainulainen L, Puhakka T, Routi T (2003) Incidence of influenza in Finnish children. *Pediatr Infect Dis J* 22(10):S204-S206
- [11] Zambon MC, Stockton JD, Clewley JP, Fleming DM (2001) Contribution of influenza and respiratory syncytial virus to community cases of influenza-like illness: an observational study. *Lancet* 358(9291):1410-1416
- [12] Peltola V, Reunanen T, Ziegler T, Silvennoinen H, Heikkinen T (2005) Accuracy of clinical diagnosis of influenza in outpatient children. *Clin Infect Dis* 41(8):1198-1200
- [13] Cazacu AC, Greer J, Taherivand M, Demmler GJ (2003) Comparison of lateral-flow immunoassay and enzyme immunoassay with viral culture for rapid detection of influenza virus in nasal wash specimens from children. *J Clin Microbiol* 41(5):2132-2134
- [14] Grijalva CG, Poehling KA, Edwards KM, Weinberg GA, Staat MA, Iwane MK, Schaffner W, Griffin MR (2007) Accuracy and interpretation of rapid influenza tests in children. *Pediatrics* 119(1):e6-e11. doi: 10.1542/peds.2006-1694

- [15] Agoritsas K, Mack K, Bonsu BK, Goodman D, Salamon D, Marcon MJ (2006) Evaluation of the Quidel QuickVue test for detection of influenza A and B viruses in the pediatric emergency medicine setting by use of three specimen collection methods. *J Clin Microbiol* 44(7):2638-2641
- [16] Ghebremedhin B, Engelmann I, König W, König B (2009) Comparison of the performance of the rapid antigen detection *actim* Influenza A&B test and RT-PCR in different respiratory specimens. *J Med Microbiol* 58(3):365-370
- [17] Steininger C, Redlberger M, Graninger W, Kundi M, Popow-Kraupp T (2009) Near-patient assays for diagnosis of influenza virus infection in adult patients. *Clin Microbiol Infect* 15(3):267-273
- [18] Uyeki TM, Prasad R, Vukotich C, Stebbins S, Rinaldo CR, Ferng YH, Morse SS, Larson EL, Aiello AE, Davis B, Monto AS (2009) Low sensitivity of rapid diagnostic test for influenza. *Clin Infect Dis* 48(9):e89-e92
- [19] Cazacu AC, Chung SE, Greer J, Demmler GJ (2004) Comparison of the directigen flu A+B membrane enzyme immunoassay with viral culture for rapid detection of influenza A and B viruses in respiratory specimens. *J Clin Microbiol* 42(8):3707-3710
- [20] Chiu SS, Tse CYC, Lau YL, Peiris M (2001) Influenza A infection is an important cause of febrile seizures. *Pediatrics* 108(4):e63. doi:10.1542/peds.108.4.e63
- [21] Heinonen S, Silvennoinen H, Lehtinen P, Vainionpää R, Vahlberg T, Ziegler T, Ikonen N, Puhakka T, Heikkinen T (2010) Early oseltamivir treatment of influenza in children 1-3 years of age: a randomized controlled trial. *Clin Infect Dis* 51(8):887-894

- [22] Waris M, Ziegler T, Kivivirta M, Ruuskanen O (1990) Rapid detection of respiratory syncytial virus and influenza-A virus in cell-cultures by immunoperoxidase staining with monoclonal-antibodies. *J Clin Microbiol* 28(6):1159-1162
- [23] Nikkari S, Halonen P, Kharitononkov I, Kivivirta M, Khristova M, Waris M, Kendal A (1989) One-incubation time-resolved fluoroimmunoassay based on monoclonal antibodies in detection of influenza A and B viruses directly in clinical specimens. *J Virol Methods* 23(1):29-40
- [24] Serwint JR, Miller RM (1993) Why diagnose influenza infections in hospitalized pediatric-patients. *Pediatr Infect Dis J* 12(3):200-204
- [25] Noyola DE, Demmler GJ (2000) Effect of rapid diagnosis on management of influenza A infections. *Pediatr Infect Dis J* 19(4):303-307
- [26] Woo PCY, Chiu SS, Seto WH, Peiris M (1997) Cost-effectiveness of rapid diagnosis of viral respiratory tract infections in pediatric patients. *J Clin Microbiol* 35(6):1579-1581
- [27] Stripeli F, Sakkou Z, Papadopoulos N, Georgiou V, Gratsia P, Christodoulou I, Tsolia M (2010) Performance of rapid influenza testing in hospitalized children. *Eur J Clin Microbiol Infect Dis* 29(6):683-688
- [28] Rouleau I, Charest H, Douville-Fradet M, Skowronski DM, De Serres G (2009) Field performance of a rapid diagnostic test for influenza in an ambulatory setting. *J Clin Microbiol* 47(9):2699-2703
- [29] Stein J, Louie J, Flanders S, Maselli J, Hacker JK, Drew WL, Gonzales R (2005) Performance characteristics of clinical diagnosis, a clinical decision rule, and a rapid influenza test in the detection of influenza infection in a community sample of adults. *Ann Emerg Med* 46(5):412-419

- [30] Hall CB, Douglas RG, Geiman JM, Meagher MP (1979) Viral shedding patterns of children with influenza B infection. *J Infect Dis* 140(4):610-613
- [31] Frank AL, Taber LH, Wells CR, Wells JM, Glezen WP, Paredes A (1981) Patterns of shedding of myxoviruses and paramyxoviruses in children. *J Infect Dis* 144(5):433-441
- [32] Cheng CKY, Cowling BJ, Chan KH, Fang VJ, Seto WH, Yung R, Uyeki TM, Houck PM, Peiris JSM, Leung GM (2009) Factors affecting QuickVue Influenza A + B rapid test performance in the community setting. *Diagn Microbiol Infect Dis* 65(1):35-41
- [33] Steininger C, Kundi M, Aberle SW, Aberle JH, Popow-Kraupp T (2002) Effectiveness of reverse transcription-PCR, virus isolation, and enzyme-linked immunosorbent assay for diagnosis of influenza A virus infection in different age groups. *J Clin Microbiol* 40(6):2051-2056
- [34] Ruest A, Michaud S, Deslandes S, Frost EH (2003) Comparison of the directigen Flu A+B test, the QuickVue influenza test, and clinical case definition to viral culture and reverse transcription-PCR for rapid diagnosis of influenza virus infection. *J Clin Microbiol* 41(8):3487-3493
- [35] Carrat F, Vergu E, Ferguson NM, Lemaître M, Cauchemez S, Leach S, Valleron AJ (2008) Time lines of infection and disease in human influenza: A review of volunteer challenge studies. *Am J Epidemiol* 167(7):775-785
- [36] Sato M, Hosoya M, Kato K, Suzuki H (2005) Viral shedding in children with influenza virus infections treated with neuraminidase inhibitors. *Pediatr Infect Dis J* 24(10):931-932
- [37] Watanabe M, Nakagawa N, Ito M, Ihara T (2009) Sensitivity of rapid immunoassay for influenza A and B in the early phase of the disease. *Pediatr Int* 51(2):211-215

- [38] Gordon A, Videa E, Saborio S, Lopez R, Kuan G, Reingold A, Balmaseda A, Harris E (2009) Performance of an influenza rapid test in children in a primary healthcare setting in Nicaragua. *PLoS ONE* 4:e7907. doi:10.1371/journal.pone.0007907
- [39] Heikkinen T, Salmi AA, Ruuskanen O (2001) Comparative study of nasopharyngeal aspirate and nasal swab specimens for detection of influenza. *BMJ* 322(7279):138
- [40] Heikkinen T, Marttila J, Salmi AA, Ruuskanen O (2002) Nasal swab versus nasopharyngeal aspirate for isolation of respiratory viruses. *J Clin Microbiol* 40(11):4337-4339
- [41] Abu-Diab A, Azzeh M, Ghneim R, Zoughbi M, Turkuman S, Rishmawi N, Issa AER, Siriani I, Dauodi R, Kattan R, Hindiye MY (2008) Comparison between pernasal flocked swabs and nasopharyngeal aspirates for detection of common respiratory viruses in samples from children. *J Clin Microbiol* 46(7):2414-2417
- [42] Chan KH, Peiris JSM, Lim W, Nicholls JM, Chiu SS (2008) Comparison of nasopharyngeal flocked swabs and aspirates for rapid diagnosis of respiratory viruses in children. *J Clin Virol* 42(1):65-69
- [43] Hurt AC, Alexander R, Hibbert J, Deed N, Barr IG (2007) Performance of six influenza rapid tests in detecting human influenza in clinical specimens. *J Clin Virol* 39(2):132-135
- [44] Cruz AT, Cazacu AC, Greer JM, Demmler GJ (2008) Rapid assays for the diagnosis of influenza A and B viruses in patients evaluated at a large tertiary care children's hospital during two consecutive winter seasons. *J Clin Virol* 41(2):143-147
- [45] Lau LLH, Cowling BJ, Fang VJ, Chan KH, Lau EHY, Lipsitch M, Cheng CKY, Houck PM, Uyeki TM, Peiris JSM, Leung GM (2010) Viral shedding and clinical illness in naturally acquired influenza virus infections. *J Infect Dis* 201(10):1509-1516

TABLE 1. Sensitivity, specificity, and predictive values, according to influenza virus type, of 158 rapid influenza tests performed within 24 hours of the onset of symptoms.

Influenza virus type	Rapid test positive	Reference methods positive ^a	Sensitivity (95% CI)	Specificity (95% CI)	Positive predictive value (PPV)	Negative Predictive value (NPV)
Influenza A	29 ^b	31	90 % (74 – 98 %)	99 % (96 – 100 %)	97 % (82 – 100%)	98 % (93 – 100 %)
Influenza B	2	8	25 % (3 – 61 %)	100 % (98 – 100 %)	100% (16 – 100 %)	96 % (92 – 99 %)
Any influenza	31 ^b	39	77 % (61 – 89 %)	99 % (95 – 100 %)	97% (83 – 100 %)	93% (87 – 97 %)

^a Composite of viral culture, antigen detection and RT-PCR was used as the reference test. The reference test was considered positive if influenza was detected by any of these methods.

^b Including one false positive that remained negative with all reference methods.