Accepted Manuscript

Title: Expression, biological activities and mechanisms of action of A20 (TNFAIP3)

Authors: Lynn Verstrepen, Kelly Verhelst, Geert van Loo, Isabelle Carpentier, Steven C. Ley, Rudi Beyaert

PII: S0006-2952(10)00478-8
DOI: doi:10.1016/j.bcp.2010.06.044
Reference: BCP 10625

To appear in: *BCP*

Received date: 1-5-2010
Revised date: 22-6-2010
Accepted date: 25-6-2010

Please cite this article as: Verstrepen L, Verhelst K, van Loo G, Carpentier I, Ley SC, Beyaert R, Expression, biological activities and mechanisms of action of A20 (TNFAIP3), *Biochemical Pharmacology* (2010), doi:10.1016/j.bcp.2010.06.044

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Expression, biological activities and mechanisms of action of A20
(TNFAIP3)**

Lynn Verstrepen^{a,b}, Kelly Verhelst^{a,b}, Geert van Loo^{a,b}, Isabelle Carpentier^{a,b}, Steven C.

Ley^c and Rudi Beyaert^{a,b}

^aUnit of Molecular Signal Transduction in Inflammation, Department for Molecular
Biomedical Research, VIB, Technologiepark 927, 9000 Ghent, Belgium,

^bDepartment of Biomedical Molecular Biology, Ghent University, Technologiepark 927, 9000
Ghent, Belgium

&

^cDivision of Immune Cell Biology, National Institute for Medical Research, London, UK

Correspondence: Rudi.Beyaert@dmbr.vib-UGent.be; Technologiepark 927; 9000 Ghent;
Belgium. Tel.: ++32 (0)9 33 13770; fax: ++32 (0)9 33 13609

Abstract

A20 (also known as TNFAIP3) is a cytoplasmic protein that plays a key role in the negative regulation of inflammation and immunity. Polymorphisms in the *A20* gene locus have been identified as risk alleles for multiple human autoimmune diseases, and A20 has also been proposed to function as a tumor suppressor in several human B-cell lymphomas. A20 expression is strongly induced by multiple stimuli, including the proinflammatory cytokines TNF and IL-1, and microbial products that trigger pathogen recognition receptors, such as Toll-like receptors. A20 functions in a negative feedback loop, which mediates its inhibitory functions by downregulating key proinflammatory signaling pathways, including those controlling NF- κ B- and IRF3-dependent gene expression. Activation of these transcription factors is controlled by both K48- and K63- polyubiquitination of upstream signaling proteins, respectively triggering proteasome-mediated degradation or interaction with other signaling proteins. A20 turns off NF- κ B and IRF3 activation by modulating both types of ubiquitination. Induction of K48-polyubiquitination by A20 involves its C-terminal zinc-finger ubiquitin-binding domain, which may promote interaction with E3 ligases, such as Itch and RNF11 that are involved in mediating A20 inhibitory functions. A20 is thought to promote de-ubiquitination of K63-polyubiquitin chains either directly, due to its N-terminal deubiquitinase domain, or by disrupting the interaction between E3 and E2 enzymes that catalyze K63-polyubiquitination. A20 is subject to different mechanisms of regulation, including phosphorylation, proteolytic processing, and association with ubiquitin binding proteins. Here we review the expression and biological activities of A20, as well as the underlying molecular mechanisms.

Keywords: A20; NF- κ B; inflammation; autoimmunity; ubiquitination

1. Introduction

In vertebrates, the innate and adaptive immune systems protect against invading pathogens and tissue damage. Innate immunity forms the first line of defence and is triggered by recognition of pathogen-associated molecular patterns (PAMPs) by germline-encoded receptors on phagocytic cells, such as macrophages, neutrophils and dendritic cells. Recognition of PAMPs involves a variety of receptors, including Toll-like receptors (TLRs), Nod-like receptors (NLRs) and RIG-I-like receptors (RLRs), which recognize their ligands either at the cell surface or intracellularly [1]. Upon activation of these receptors, multiple signaling cascades are triggered, leading to the activation of nuclear factor- κ B (NF- κ B), interferon regulatory factor 3 (IRF3) and activator protein 1 (AP-1) transcription factors. Together, these transcription factors induce expression of genes which mediate the immediate effector functions of innate immune cells, such as iNOS and tissue destructive enzymes, together with inflammatory cytokines and chemokines, which are important in attracting T cells to sites of infection to initiate the adaptive immune response.

Activation of innate and adaptive immune responses is tightly regulated, as insufficient activation could result in defective clearance of pathogens, while excessive activation might lead to lethal systemic inflammation or autoimmunity. A20 is an important negative regulator of innate and adaptive immunity, both after acute infection and in immune homeostasis. The importance of A20 in limiting inflammation is underscored by the association of polymorphisms in the A20 genomic region with multiple human autoimmune and inflammatory diseases, including Crohn's disease, rheumatoid arthritis, systemic lupus erythematosus, psoriasis and type 1 diabetes (Figure 1). Furthermore, A20 also functions as a tumor suppressor in several B cell lymphomas. The role of A20 as a disease associated gene

has recently been reviewed in detail elsewhere [2, 3]. Here we summarize what is known about the expression, biological activities, and mechanism of action of A20.

2. Expression of A20

A20 was initially identified as a primary response gene following stimulation of human umbilical vein endothelial cells (HUVEC) with TNF, IL-1 or LPS [4]. Subsequently, A20 was also found to be transcriptionally upregulated in other cell types, by many other agonists (see Table 1). In contrast, human immature CD4⁺CD8⁺ double positive thymocytes and single positive thymocytes constitutively express significant levels of A20, but upon stimulation with TPA/ionomycin, A20 expression declines in the mature single-positive cells. Similarly, A20 is expressed in resting peripheral T-cells, and is down-regulated after activation [5].

Dysregulation of A20 expression in certain transformed cells suggests that A20 may play a role in the pathogenesis of cancer. A20 mRNA is upregulated in undifferentiated nasopharyngeal carcinoma and poorly differentiated head and neck squamous cell carcinomas (SCCs) of the skin, suggesting that A20 has a part in the pathogenesis of these epithelial malignancies. In contrast, no A20 mRNA is detected in normal samples of squamous epithelial tissues or in well-differentiated SCCs [6]. A20 expression in estrogen receptor (ER) and progesterone receptor (PR) negative breast tumour cells *in vivo* has been associated with poor prognosis. A20 mRNA levels are significantly higher in ER-negative breast cancer cell lines and tamoxifen resistant ER-positive tumours. Furthermore, estrogen stimulation of the MVLM cell line, an ER-positive human breast carcinoma, downregulates A20 mRNA expression [7]. A20 expression is also increased in inflammatory breast cancer, a frequently hormone receptor negative progressive form of cancer [8]. Together, these data demonstrate that a high expression of A20 is often linked with poor prognosis of epithelial malignancies.

On the other hand, a large number of B-cell lymphomas contain mutations in *A20* that result in loss of *A20* protein expression, indicative for a tumor suppressor role for *A20* [9], and suggesting that the role of *A20* in tumorigenesis might be cell type dependent.

In mice, constitutive *A20* mRNA expression can be detected in epithelial cells involved in early hair follicle development, but the highest levels of *A20* mRNA are seen in lymphoid tissues, like thymus, gut-associated lymphoid aggregates (containing B- and T-cells) and spleen [5]. Moreover, TNF injection in mice results in a substantial upregulation of *A20* mRNA in liver, kidney, spleen, thymus, lung, colon and lymph nodes [10].

The human *A20* gene is located on chromosome 6q23 [11], while the murine *A20* gene is mapped to chromosome 10 at 13 cM [5, 12]. Human *A20* mRNA is 4 kb long, containing an open reading frame of 2370 nucleotides, encoding a protein of 790 amino acids with a molecular weight of 90 kDa [13]. The 3'-untranslated region contains four copies of the sequence ATTTA, an mRNA destabilizing motif that is frequently found in the 3'UTR of early response inflammatory mediators. *A20* mRNA is superinduced by cycloheximide [4], suggesting a role for protein synthesis in the destabilization of *A20* mRNA. Reporter assays indicate that transcriptional activation of the *A20* gene is mediated by two NF- κ B binding sites in its promoter [14]. Consistently, *A20* expression is reduced in cells genetically deficient in NF- κ B activation [15]. Furthermore, *A20* negatively regulates its own expression as a consequence of its NF- κ B inhibitory activity [14], resulting in *A20* mRNA only being transiently induced following TNF stimulation [4]. The rapid induction of the *A20* gene upon TNF stimulation has been suggested to involve the constitutive association of the general transcription apparatus and co-activators, such as CBP and p300, with the *A20* promoter, mediated by the transcription factor Sp-1 [16]. While Sp-1 activity is necessary for the efficient transcription initiation, NF- κ B binding to the promoter enhances the re-initiation rate, resulting in multiple transcription rounds and augmenting expression levels. The basal

activity of the *A20* gene is repressed at the elongation step by DRB sensitivity-inducing factor (DSIF), whose activity is controlled by a negative upstream promoter element called ELIE (elongation-inhibitory element) [17]. Remarkably, following stimulation, inhibition of the *A20* promoter by DSIF persists, but it is now regulated by NF- κ B rather than ELIE. Recently, an E-box, overlapping with ELIE, in the *A20* promoter was demonstrated to bind upstream stimulatory factor (USF)1 [18]. USF1 mediates DSIF inhibition and recruitment to the *A20* promoter under basal conditions. Once NF- κ B is induced, the E-box is no longer involved in DSIF inhibition. Consistent with that, USF1 is displaced from the promoter by NF- κ B. (Figure 2)

3. *A20* structure

The amino acid sequence of *A20* is highly conserved (more than 90% identical when compared to human *A20*) in various mammalian species, and clear orthologs are present in Zebrafish and *Xenopus* (50% identical to human *A20*, but up to 90% in the N-terminal *A20* domain). In contrast, no such clear orthologs can be found in *Drosophila* and *C. elegans* [19]. Homology comparisons have revealed the presence of an ovarian tumor (OTU) domain in the N-terminal part of *A20* (Figure 3) [20], suggesting that *A20* functions as a deubiquitinase, which has been confirmed biochemically [21]. The crystal structure of the OTU-domain shows ten β -strands and ten α -helices, which are arranged to form an expansive surface on which ubiquitin binding, interaction with substrates and catalysis can take place [22, 23]. The catalytic site is composed of a catalytic triad (Cys103, His256, Asp70), with a highly conserved surface patch, forming the ubiquitin-binding site and a separate region for binding target proteins. Cezanne and TRABID are two other OTU-domain containing deubiquitinases

that show considerable similarity with A20 within their OTU-domain, and have therefore been proposed as A20-like proteins [24, 25].

The C-terminal domain of A20 contains seven zinc finger (ZnF) structures, of which 6 have the sequence CX₄CX₁₁CX₂C and one is similar to the sequence CX₂CX₁₁CX₂C (Figure 3) [13]. The human and mouse A20 ZnFs are completely identical, except for a single substitution of methionine for valine in ZnF4 [5]. It has been proposed that ZnF4 confers intrinsic ubiquitin ligase activity to A20 (see 4.3). The C-terminal ZnF containing domain is also required for homo-oligomerization of A20, although it is not clear whether this is important for A20 function [13]. The A20-related protein Cezanne only contains a single A20-like zinc finger in its C-terminus, which is most similar to ZnF4 and ZnF7 of A20 [25]. Also the A20-related protein TRABID contains three ZnFs at its N-terminus, but these are unrelated to those of A20. Finally, A20-like ZnFs have also been described in other proteins such as ZNF216 and Rabex-5 [26, 27].

Between ZnF3 and ZnF4 of A20, there is a perfect 14-3-3 binding sequence (RSKSDP), which is conserved between human and mouse (Figure 3) [28, 29]. Consistently, A20 specifically interacts with the η and ζ isoforms of 14-3-3, as mutations in the consensus 14-3-3 binding site abolish these interactions [28]. Interaction of A20 with 14-3-3 alters the localization of A20 from a punctuate cytoplasmic staining to a more diffuse cytoplasmic pattern, which is also associated with decreased amounts of A20 in the insoluble cell fraction [29]. Mutations in the 14-3-3 binding domain of A20 lead to proteolysis of A20 in MCF-7 cells [30], suggesting that the binding to 14-3-3 protects A20 from degradation. Although A20 is mostly localized in the cytosol, a fraction of A20 can associate with a lysosome-interacting compartment in a manner that requires its carboxy terminal ZnFs [31]. This lysosome-associated A20 has been shown to target the A20-associated signaling molecule TRAF2 to the lysosomes for degradation [32].

4. Biological activities of A20

4.1. Cytoprotective effects of A20

The first function described for A20 was its cytoprotective effect upon TNF stimulation of cells, based on the effect of A20 overexpression [33]. This was later confirmed genetically as $A20^{-/-}$ murine embryonic fibroblasts and thymocytes were found to be more sensitive to TNF-induced cell death than wild type (WT) cells [10]. However, the anti-apoptotic function of A20 is not a general feature, as A20 only protects some cell types from specific death inducing agents. For example, overexpression of A20 protects MCF7S1 and WEHI-S cells from TNF-mediated apoptosis, but not from killing by anti-Fas, activated monocytes, serum starvation or H_2O_2 [34]. In contrast, A20 protects endothelial cells against apoptosis induced by TNF and Fas triggering [35], whereas in A549, HepG2 and HeLa cells A20 is not able to block TNF-induced apoptosis [36, 37]. Induction of A20 after viral infection has also been suggested as a mechanism to protect infected cells from p53-induced apoptosis [38], and A20 downregulation may be important in activation-induced cell death (AICD) in T-cells [39]. Table 2 summarizes the effects of A20 on cell survival in different cell types. It should be mentioned that our knowledge of the cytoprotective effects of A20 and the underlying mechanisms (see below) is mostly based on studies using A20 overexpression in cell lines, implicating the need to confirm their physiological relevance using A20-deficient primary cells.

The underlying mechanism by which A20 inhibits apoptosis is still largely unclear. Overexpression experiments indicate that the C-terminal ZnF containing domain of A20 is necessary and sufficient for its anti-apoptotic activity following TNF stimulation, but the 14-

3-3 binding site is dispensable [30, 36]. In Jurkat T-cells, ectopic expression of A20 inhibits TNF-induced apoptosis by preventing recruitment of RIP1 and TRADD to the TNF-RI complex, which mediate the recruitment of FADD and the initiator procaspase 8, leading to its autoactivation by proteolytic cleavage [15]. In endothelial cells, however, A20 overexpression was shown to restrict not only TNF- but also Fas-induced apoptosis, associated with the inhibition of proteolytic cleavage of apical caspases 8 and 2, executioner caspases 3 and 6, Bid cleavage, and release of cytochrome c, thus preserving mitochondrion integrity [35]. Inhibition of both TNF-RI- and Fas-mediated apoptosis suggests that A20 interferes at a common step in their downstream signaling pathways. The two death receptors differ only in their use of proximal signaling molecules. TNF-RI requires the adapters TRADD and RIP1 to transduce signals to FADD, whereas Fas interacts directly with FADD and activates caspase 8. Inhibition of both of these death pathways in endothelial cells supports an effect of A20 at the level of FADD/caspase 8. Recently, A20 was shown to reverse death receptor-induced polyubiquitination of caspase 8 [40]. As the latter is necessary for caspase 8 aggregation, leading to its full activation and processing, it is tempting to speculate that A20 mediates its anti-apoptotic activities by deubiquitinating caspase 8.

A20 not only inhibits TNF-induced apoptosis, but also TNF-induced necrosis as shown by overexpression of A20 in the murine fibrosarcoma L929 cell line [41], which is associated with reduced activation of phospholipases and reduced induction of reactive oxygen species. Similarly, A20 overexpression protects endothelial cells from complement-mediated, but not heat-induced necrosis [35]. Recently, A20 was shown to inhibit both the caspase 8 dependent and independent cell death pathway in macrophages stimulated with TNF [42]. Overexpression of A20 in these cells protected against the loss of lysosomal integrity and cathepsin B release (caspase 8 independent and typical for necrosis) and DNA

fragmentation (RIP1 and caspase 8 dependent and typical for apoptosis). The mechanism for the anti-necrotic action of A20 is still unknown.

In contrast to its cytoprotective effects, A20 has also been reported to sensitize vascular smooth muscle cells to cytokine- and Fas-mediated apoptosis [35]. In addition, cells expressing high levels of A20 were shown to be more sensitive to oxidative stress-mediated cell death than cells with low A20 levels [43]. The reason for these paradoxical effects is still unclear, but might reflect the delicate balance between a direct anti-apoptotic effect of A20 and an indirect apoptosis-sensitizing effect due to its NF- κ B inhibitory function.

4.2. Negative regulation of inflammation by A20

Early experiments in cell lines suggested an anti-inflammatory role for A20, since overexpressed A20 was found to substantially inhibit NF- κ B activation in response to several different stimuli (see Table 2). Consistently, inhibition of NF- κ B activation by A20 expression decreases expression of several NF- κ B target genes, including E-selectin, ICAM-1, IL-8, I κ B α , tissue factor, VCAM-1, IL-6 and GM-CSF [44-50]. A physiological function of A20 in the negative regulation of inflammation was confirmed by the phenotype of A20 knockout mice [10]. A20^{-/-} mice die shortly after birth by severe inflammation and tissue damage in multiple organs, including liver, kidneys, intestine, joints and bone marrow. A20^{-/-} RAG1^{-/-} mice, which are devoid of T- and B-cells, display the same phenotype, indicating that A20 regulates innate immune homeostasis independent of adaptive immune cells [51]. In accordance with A20 overexpression experiments, analysis of MEFs from A20^{-/-} mice demonstrates a role for A20 in regulating NF- κ B activation [10]. TNF stimulation of A20-deficient MEFs results in prolonged IKK activation compared to WT cells. This causes sustained phosphorylation and degradation of I κ B α , and consequent increased NF- κ B activity

in A20-deficient cells. The finding that A20/TNF-RI and A20/TNF double knockout mice have a similar phenotype as the A20 single knockout mice, suggest a critical role for A20 in the regulation of TNF-independent pro-inflammatory signals and the maintenance of immune homeostasis [51]. LPS activation of IKK is prolonged in macrophages generated from A20/TNF double knockout mice, indicating that A20 negatively regulates TLR4 activation of NF- κ B, independently of TNF. Indeed, recent experiments have suggested that the inflammatory phenotype of *A20*^{-/-} mice is due to excessive signaling from MyD88-dependent TLRs (such as TLR3, TLR4 and TLR9); unlike, *A20*^{-/-} mice, *A20*^{-/-}*MyD88*^{-/-} mice do not die prematurely or exhibit severe cachexia and have reduced inflammation in their spleens [52]. Using broad spectrum antibiotics treatment it was further shown that commensal intestinal bacteria drive these constitutive TLR signals in the absence of A20. It should be noted that *A20*^{-/-} macrophages also express more TNF, IL-6 and NO after triggering of TLR3, which is MyD88-independent, suggesting that A20 also interferes with the TRIF-dependent TLR signaling axis [51].

Clear inhibitory effects of A20 have also been observed on IRF3 activation. For example, A20 overexpression strongly inhibits RIG-I mediated activation of IRF3 [53]. A20 overexpression has also been found to inhibit ISRE- and IFN- β promoter-dependent luciferase activity upon poly(I:C) stimulation of TLR3 [54], as well as Newcastle disease virus (NDV)-induced IRF3 activation in HEK293 cells [55]. Consistent with these data, A20 knock down in HEK293T, HeLa and THP-1 cells enhances ISRE-dependent transcription after poly(I:C) stimulation, infection with NDV [55] or Sendai virus [53]. Co-transfection data suggest that A20 inhibits TLR3-mediated IRF3 activation upstream or at the level of TBK1/IKK ϵ kinases, possibly involving direct interaction of A20 with TBK1, IKK ϵ and TRIF. However, analyses of *A20*^{-/-} mice suggest that the effects of A20 on expression of IRF3-regulated genes may be indirect. Irradiated mice reconstituted with hematopoietic stem cells derived from *A20*^{-/-}

1 *MyD88*^{-/-} mice have higher levels of IFN- β and MCP-1, two TRIF-dependent pro-
 2 inflammatory cytokines, after LPS injection [52]. However, in BMDMs generated from these
 3 mice, the kinetics of IRF3 phosphorylation after LPS stimulation are comparable to those in
 4 wild-type cells, while I κ B α phosphorylation is sustained due to prolonged IKK activation.
 5 These results suggest that A20 may restrict TRIF-dependent signaling to IFN- β by regulating
 6 NF- κ B activation and not IRF3. A20 may control common signaling proteins involved in
 7 MyD88- and TRIF-dependent signaling to NF- κ B, but not implicated in IRF3 activation.
 8 Possible candidates are TRAF6 and RIP1 (see 4.3).
 9

10 A20 overexpression has additionally been reported to inhibit AP-1 activation
 11 following TNF, IL-1 and LPS stimulation [34, 56]. However, AP-1 inhibition by A20 has not
 12 always been detected [48, 57] and may be cell type- or stimulus-specific. The AP-1 inhibitory
 13 effect can be explained by the inhibition of JNK activation by A20 [45], although this has not
 14 been detected in all cells [47]. Results obtained from A20-deficient MEF cells support a role
 15 for A20 in the negative regulation of TNF-induced JNK activation [10].
 16

17 4.3. Mechanisms involved in the NF- κ B inhibitory effect of A20

18 As discussed in 4.2, analysis of A20-deficient cells has indicated that A20 negatively
 19 regulates TNF and TLR4 activation of IKK. Consistent with these data, A20 has been shown
 20 to inhibit NF- κ B activation by co-expressed signaling intermediates which function upstream
 21 of the IKK complex, such as TAK1, TRAF2/6, TRADD and RIP1 [37, 48, 57, 58], but does
 22 not affect NF- κ B activation by overexpressed RelA [37, 48]. However, contradictory data
 23 have been generated on the effect of A20 on nuclear translocation and DNA-binding of NF-
 24 κ B. For example, overexpression of A20 in HEK293T and L929sA cells inhibits TNF-
 25 induced NF- κ B dependent luciferase expression without affecting nuclear translocation and
 26

DNA-binding of NF- κ B, suggesting that A20 specifically interferes with an NF- κ B transactivation signal [48, 57]. It will clearly be important to obtain complementary evidence for A20 regulation of NF- κ B transactivation using A20-deficient cells.

Co-expression experiments have demonstrated that several signaling intermediates upstream of the IKK complex can bind to A20 (see Table 3). For example, A20 interacts with TRAF1, TRAF2 and TRAF6, which was shown to be mediated by its N-terminal domain in the case of TRAF1 and TRAF2 [58, 59]. A20 has also been shown to interact with IKK1 and NEMO via its C-terminal ZnF domain [60, 61], but not with IKK2 [57]. In accordance with these data, endogenous A20 is recruited to the TNF-RI signaling complex after TNF stimulation [61]. The ability of A20 to bind to proximal components of the TNF-RI signaling pathway to NF- κ B is consistent with data indicating that A20 regulates IKK activation after TNF stimulation.

Wertz et al. proposed that A20 contains two ubiquitin-editing domains, an N-terminal deubiquitinating (DUB) domain of the OTU family and a C-terminal ubiquitin ligase domain [62]. The N-terminal domain was shown to deubiquitinate K63-polyubiquitinated RIP1, an essential mediator of the proximal TNF-RI signaling complex. The C-terminal ZnF containing domain functions as an ubiquitin ligase that catalyzes the addition of K48-polyubiquitin chains to RIP1, thereby targeting RIP1 for proteasomal degradation [62, 63] (Figure 4). A20 mutated at the OTU-domain catalytic cysteine residue is defective in adding K48-polyubiquitin chains to RIP1, suggesting that RIP1 K63 deubiquitination is a prerequisite for A20-mediated RIP1 K48-polyubiquitination and subsequent degradation. The catalytic cysteine is also necessary for deubiquitination of K63-polyubiquitinated TRAF6 by A20 (Figure 4) and inhibition of NF- κ B activation after LPS stimulation [51]. Furthermore, stimulation of NOD2 with MDP leads to K63-polyubiquitination of RIP2, which can be inhibited by A20 depending on its OTU domain [64]. A20 has also been reported to remove

K63-polyubiquitin chains from NEMO in TNF-stimulated cells. However, the role of NEMO ubiquitination in NF- κ B activation is unclear [65, 66]. As discussed earlier, *in vitro* the N-terminal domain of A20 has been shown to hydrolyse both K48- and K63-polyubiquitin [21], and hydrolysis of K48-polyubiquitin chains was later shown to be much more efficient than K63-polyubiquitin chains [22, 23]. This suggests that other factors such as the polyubiquitinated protein itself or possibly other A20-binding proteins might determine the DUB specificity of A20 for K63-polyubiquitinated substrates in intact cells.

Reconstitution of $A20^{-/-}$ MEF cells with the OTU catalytic cysteine mutant of A20 is much less effective in down-regulating TNF-induced NF- κ B activity than wild type A20 [62], suggesting that DUB activity is essential for A20's inhibitory function. However, several studies showed that the OTU domain is not important for the NF- κ B or IRF3 inhibitory activity of overexpressed A20. For example, overexpression of the C-terminal domain or the OTU catalytic cysteine mutant of A20 still inhibits TNF-, IL-1- and Pam3Cys-induced NF- κ B activation, as well as poly-(I:C)- and NDV-induced IRF3 activation, as potently as the wild type molecule [21, 30, 48, 55, 58]. The physiological relevance of such overexpression experiments remain to be demonstrated.

ZnF4 is essential for retrovirally expressed A20 to trigger RIP1 proteolysis in $A20^{-/-}$ MEFs and it has been suggested that ZnF4 confers ubiquitin ligase activity to A20 [62]. However, it is unclear whether A20 has intrinsic E3 ligase activity or if this activity is mediated by an associated protein such as Itch and RNF11, which are two A20-binding E3 ubiquitin ligases that have been reported to be essential for A20's NF- κ B inhibitory function [67, 68]. It is also evident that ZnF4 is not always required for A20 to inhibit NF- κ B. For example, the inhibitory effect of overexpressed A20 on RIG-I- and virus-induced NF- κ B and IRF3 activation was found to involve ZnF7 and not ZnF4 [53]. Likewise, ZnF7 contributes to the inhibition of TNF-induced NF- κ B activation by overexpressed A20 [37], as well as to the

localization of A20 to lysosome-associated compartments [31]. Furthermore, oligomerization of ZnF7 as such inhibits NF- κ B activation [37]. Klinkenberg et al. demonstrated that there is redundancy between the ZnFs of A20 for inhibition of TNF-induced NF- κ B activation, with 4 ZnFs being sufficient for inhibition [60]. However, it should be noted that either ZnF4 or ZnF7 were still present in inhibitory A20 mutants.

Recently, A20 was found to use also alternative mechanisms to interfere with NF- κ B activation [69]. More specifically, upon LPS or IL-1 stimulation, binding of the E3 ligase TRAF6 to A20 prevents TRAF6 from interacting with the E2 conjugating enzyme Ubc13, which normally triggers TRAF6 autoubiquitination and activation. Similarly, A20 disrupts the interaction between the E2 enzyme Ubc5 and the E3 ligases cIAP1 and TRAF2 upon TNF stimulation, as well as Ubc5/TRAF6 binding in response to IL-1. Moreover, A20 also interacts with Ubc5 and Ubc13 and triggers their K48-polyubiquitination and proteasomal degradation at later time points. Remarkably, reconstitution of A20^{-/-} MEFs with A20 carrying mutations in respectively its OTU domain or ZnF4 indicated that both are somehow involved in controlling E2-E3 enzyme interactions and E2 degradation. Clearly, knockin mice with a mutation in the OTU active site or specific ZnFs will be very useful to reveal the DUB-dependent and -independent functions of A20 under physiological conditions.

4.4. Mechanisms regulating A20 activity

As mentioned before, an important mechanism regulating A20 activity is the regulation of its expression. Most cells express no detectable or only limited amounts of A20 under basal conditions and A20 expression is rapidly induced upon NF- κ B activation. However, T-cells and B-cells constitutively express considerable high levels of A20, which inhibits ubiquitination of the NF- κ B signaling protein MALT1 [70]. In this case, antigen

receptor stimulation triggers a rapid decrease and subsequent reappearance of A20, suggesting that A20 removal could be a crucial step in eliminating the negative regulatory activity of A20. Proteasomal degradation of A20, as well as its cleavage by the paracaspase MALT1, contribute to the decrease in A20, resulting in enhanced IL-2 production and NF- κ B activation [70, 71].

A20 activity can also be regulated by phosphorylation. TNF as well as LPS stimulation leads to the phosphorylation of human A20 at Ser381 by IKK2, increasing the ability of A20 to inhibit NF- κ B signaling [72]. A20 phosphorylation might thus allow the downregulation of NF- κ B activity to be tailored more precisely to the needs of the cell, depending on the strength and duration of the inflammatory signal. The mechanism by which phosphorylation modulates A20 function is not known.

The activity of A20 is also regulated by its binding to specific proteins that have essential functions in limiting cytokine- and LPS-induced NF- κ B activation and inflammation. The A20-binding protein TAX1BP1 interacts with RIP1 and TRAF6 and recognizes K63-polyubiquitin chains through a C-terminal ZnF domain. Therefore, TAX1BP1 most likely functions as an 'ubiquitin receptor' protein that connects A20 with ubiquitinated substrates [73, 74]. Similarly, A20 also requires TAX1BP1 to terminate IRF3 signaling, where TAX1BP1 has been proposed to function as an adaptor molecule between A20 and TBK1/IKK ϵ , preventing the interaction of the E3 ubiquitin ligase TRAF3 with its cognate substrates [75]. Consistent with its role in mediating A20 function, *TAX1BP1*^{-/-} mice die from inflammatory cardiac valvulitis and are hypersensitive to the toxic effects of low doses of TNF and IL-1 [73]. In contrast, Shembade et al. reported that *TAX1BP1*^{-/-} mice are embryonic lethal [74]. It should be mentioned that in the latter study, *TAX1BP1*^{-/-} mice were generated from a gene trap clone, which might explain the different phenotype. TAX1BP1 and A20 also form a complex with the E3 ubiquitin ligases Itch and RNF11, which are

required for A20 to inactivate NF- κ B signaling [67, 68]. Interestingly, the human T cell leukemia virus type I (HTLV-I) Tax oncoprotein disrupts formation of the A20-TAX1BP1-Itch ternary complex, which may allow constitutive NF- κ B activation in HTLV-I infected cells [67]. Finally, A20 also interacts with ABIN-1, -2, and -3. Because overexpression of ABINs mimics the NF- κ B inhibitory function of A20, they have been proposed to cooperate with A20 in the regulation of NF- κ B signaling [76]. ABINs also function as ubiquitin-binding proteins via a conserved UBAN domain and ABIN-1 has been shown to interact with polyubiquitinated NEMO [77]. Since small interfering RNA targeting of ABIN-1 abrogates A20-dependent deubiquitination of NEMO and RNA interference of A20 impairs the ability of ABIN-1 to inhibit NF- κ B activation [65], ABIN-1 has been suggested to physically link A20 to NEMO, hereby facilitating A20-mediated deubiquitination of NEMO, resulting in inhibition of NF- κ B. However, ABIN-1 or ABIN-2 knockout mice did not confirm a role for ABIN-1 and -2 in the regulation of NF- κ B signaling [78, 79], possibly due to redundancy of different ABINs. In contrast, ABIN-1 knockout mice confirmed the previously described anti-apoptotic function of ABIN-1 [78, 80]. Although the anti-apoptotic effect of ABIN-1 makes it tempting to speculate that ABIN-1 mediates the anti-apoptotic effect of A20, this was also ruled out [78].

4.5. Other functions of A20 (see table 2)

Experiments testing the effects of A20 overexpression or deficiency suggest that A20 has functions in addition to the negative regulation of pro-inflammatory gene expression. For example, *A20*^{-/-} mice show thickening of epidermal and dermal layers without apparent inflammation, suggesting a role for A20 in skin differentiation [10]. Consistent with this idea, A20 is found in the basal layer of the forming skin at 15 dpc, and in epithelial cells at

epidermal invaginations at dpc 17, suggesting a role for A20 in early hair follicle development [5]. As *IκBα*^{-/-} mice have a similar skin phenotype [81] to *A20*^{-/-} mice, A20 may affect skin differentiation via NF-κB regulation. However, IKK1, which interacts with A20 [57], controls skin development independently of NF-κB activation, by stimulating keratinocyte differentiation [82, 83]. Therefore, it is possible that A20 affects this NF-κB-independent function of IKK1 to modulate skin differentiation.

A20 expression has also been shown to exert anti-proliferative effects on cells. Adenoviral transfection of A20 inhibits vascular smooth muscle cell (VSMC) proliferation induced by TNF by arresting cells in the G₀/G₁ phase [84]. In VSMCs transfected with A20, expression of p53 and the p53 target genes p21^{waf1} and p27^{kip} is increased [85], which would be expected to contribute to the anti-proliferative effect of A20. The effect of A20-deficiency on cell proliferation has not yet been tested, which is essential to confirm that A20 regulates cell proliferation under physiological conditions. Recent data suggest that A20 is involved in neovascularisation. A20 knockdown in HUVEC cells reduces both tubule area and length in *in vitro* angiogenesis assays [86]. Furthermore, A20 expression is largely restricted to endothelial cells of new vasculature in breast cancers [86]. It is unclear whether either of these novel functions of A20 is due to elevated NF-κB activity.

5. Conclusions and future perspectives

Immune signaling pathways are highly complex and tight control is needed in order to maintain homeostasis. The NF-κB pathway serves as a paradigm and has drastically increased our understanding of signal propagation and termination mechanisms. Part of the complexity comes from the many posttranslational modifications which occur. Of these, polyubiquitination has emerged as an exciting area of regulation for NF-κB signaling, and it

1 can be expected that other signaling pathways use similar regulatory mechanisms. Mounting
2 evidence indicates that A20 interferes with NF- κ B, IRF3 and cell death signaling to limit
3 inflammation and immunity by modulating the ubiquitination status of specific signaling
4 proteins via multiple mechanisms. What determines whether A20 removes K48- or K63-
5 polyubiquitin chains, as well as the underlying mechanism of its ubiquitin ligase activity
6 remain intriguing questions. Posttranslational modifications of A20, including
7 phosphorylation and proteolytic processing, appear to be critical for its function and activity.
8 It will be of interest to analyze whether A20 can also undergo other posttranslational
9 modifications and if there is cross-talk between the different A20 modifications.
10
11
12
13
14
15
16
17
18
19
20
21

22 A20 has been associated with several diseases. Importantly, polymorphisms in the
23 A20 genomic region (in most cases located outside the A20 protein encoding sequence) are
24 associated with autoimmunity [3]. Moreover, A20 is inactivated in a substantial number of
25 lymphomas by deletion, promoter methylation, frame shift mutations and/or non-sense
26 mutations that result in truncations and point mutations in A20 [2, 87-89]. Because both A20
27 alleles are often affected in these lymphomas, A20 was proposed as a new tumor suppressor.
28 It will be important to study the effect of the corresponding mutations on the expression and
29 biological activity of A20, using several cellular and biochemical assays. As many mutations
30 in the A20 locus are located outside the mRNA encoding region, also epigenetic regulation of
31 A20 expression is likely to be an important issue. Finally, what is the relationship of A20 with
32 other NF- κ B regulating DUBs such as CYLD, which shares several substrates with A20? This
33 question, as well as improved understanding of the physiological functions of A20, might be
34 addressed by studying the cell type and tissue specific functions of A20, using a combination
35 of mouse genetics, primary cells, cell lines, and patient samples. A detailed understanding of
36 the molecular and cellular function of A20 is likely to facilitate the development of novel
37 therapeutics for inflammatory diseases and cancer.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

ACKNOWLEDGEMENTS

L. Verstrepén and G. van Loo are postdoctoral researchers with the “Fund for Scientific Research of Flanders” (FWO) and K. Verhelst is a PhD fellow with the “Institute for the promotion of Innovation by Science and Technology” (IWT). G. van Loo also holds an FWO Odysseus Grant. Work in the authors’ lab is further supported by research grants from the “Interuniversity Attraction Poles program” (IAP6/18), the FWO, the “Belgian Foundation against Cancer”, the “Strategic Basis Research program” of the IWT, the “Centrum voor Gezwelziekten” and the “Concerted Research Actions” (GOA grant 01G06B6) and ‘Group-ID MRP’ initiative of the Ghent University. S. Ley is funded by the UK Medical Research Council.

REFERENCES

- [1] Barton GM. A calculated response: control of inflammation by the innate immune system. *J Clin Invest* 2008;118:413-20.
- [2] Hymowitz SG, Wertz IE. A20: from ubiquitin editing to tumour suppression. *Nat Rev Cancer* 2010;10:332-41.
- [3] Vereecke L, Beyaert R, van Loo G. The ubiquitin-editing enzyme A20 (TNFAIP3) is a central regulator of immunopathology. *Trends Immunol* 2009;30:383-91.
- [4] Dixit VM, Green S, Sarma V, Holzman LB, Wolf FW, O'Rourke K, et al. Tumor necrosis factor- α induction of novel gene products in human endothelial cells including a macrophage-specific chemotaxin. *J Biol Chem* 1990;265:2973-8.
- [5] Tewari M, Wolf FW, Seldin MF, O'Shea KS, Dixit VM, Turka LA. Lymphoid expression and regulation of A20, an inhibitor of programmed cell death. *J Immunol* 1995;154:1699-706.
- [6] Codd JD, Salisbury JR, Packham G, Nicholson LJ. A20 RNA expression is associated with undifferentiated nasopharyngeal carcinoma and poorly differentiated head and neck squamous cell carcinoma. *J Pathol* 1999;187:549-55.
- [7] Vendrell JA, Ghayad S, Ben-Larbi S, Dumontet C, Mechti N, Cohen PA. A20/TNFAIP3, a new estrogen-regulated gene that confers tamoxifen resistance in breast cancer cells. *Oncogene* 2007;26:4656-67.
- [8] Lerebours F, Vacher S, Andrieu C, Espie M, Marty M, Lidereau R, et al. NF-kappa B genes have a major role in inflammatory breast cancer. *BMC Cancer* 2008;8:41.
- [9] Malynn BA, Ma A. A20 takes on tumors: tumor suppression by an ubiquitin-editing enzyme. *J Exp Med* 2009;206:977-80.
- [10] Lee EG, Boone DL, Chai S, Libby SL, Chien M, Lodolce JP, et al. Failure to regulate TNF-induced NF-kappaB and cell death responses in A20-deficient mice. *Science* 2000;289:2350-4.
- [11] Boonyasrisawat W, Eberle D, Bacci S, Zhang YY, Nolan D, Gervino EV, et al. Tag polymorphisms at the A20 (TNFAIP3) locus are associated with lower gene expression and increased risk of coronary artery disease in type 2 diabetes. *Diabetes* 2007;56:499-505.
- [12] Idel S, Dansky HM, Breslow JL. A20, a regulator of NFkappaB, maps to an atherosclerosis locus and differs between parental sensitive C57BL/6J and resistant FVB/N strains. *Proc Natl Acad Sci U S A* 2003;100:14235-40.
- [13] Opipari AW, Jr., Boguski MS, Dixit VM. The A20 cDNA induced by tumor necrosis factor α encodes a novel type of zinc finger protein. *J Biol Chem* 1990;265:14705-8.
- [14] Krikos A, Laherty CD, Dixit VM. Transcriptional activation of the tumor necrosis factor α -inducible zinc finger protein, A20, is mediated by kappa B elements. *J Biol Chem* 1992;267:17971-6.
- [15] He KL, Ting AT. A20 inhibits tumor necrosis factor (TNF) α -induced apoptosis by disrupting recruitment of TRADD and RIP to the TNF receptor 1 complex in Jurkat T cells. *Mol Cell Biol* 2002;22:6034-45.
- [16] Ainbinder E, Revach M, Wolstein O, Moshonov S, Diamant N, Dikstein R. Mechanism of rapid transcriptional induction of tumor necrosis factor α -responsive genes by NF-kappaB. *Mol Cell Biol* 2002;22:6354-62.
- [17] Ainbinder E, Amir-Zilberstein L, Yamaguchi Y, Handa H, Dikstein R. Elongation inhibition by DRB sensitivity-inducing factor is regulated by the A20 promoter via a novel negative element and NF-kappaB. *Mol Cell Biol* 2004;24:2444-54.

- [18] Amir-Zilberstein L, Dikstein R. Interplay between E-box and NF-kappaB in regulation of A20 gene by DRB sensitivity-inducing factor (DSIF). *J Biol Chem* 2008;283:1317-23.
- [19] Saitoh T, Yamaoka S. A20, UCSD-Nature Molecule Pages 2010, doi:10.1038/mp.a000159.01
- [20] Makarova KS, Aravind L, Koonin EV. A novel superfamily of predicted cysteine proteases from eukaryotes, viruses and *Chlamydia pneumoniae*. *Trends Biochem Sci* 2000;25:50-2.
- [21] Evans PC, Ovaa H, Hamon M, Kilshaw PJ, Hamm S, Bauer S, et al. Zinc-finger protein A20, a regulator of inflammation and cell survival, has de-ubiquitinating activity. *Biochem J* 2004;378:727-34.
- [22] Komander D, Barford D. Structure of the A20 OTU domain and mechanistic insights into deubiquitination. *Biochem J* 2008;409:77-85.
- [23] Lin SC, Chung JY, Lamothe B, Rajashankar K, Lu M, Lo YC, et al. Molecular basis for the unique deubiquitinating activity of the NF-kappaB inhibitor A20. *J Mol Biol* 2008;376:526-40.
- [24] Evans PC, Smith TS, Lai MJ, Williams MG, Burke DF, Heyninck K, et al. A novel type of deubiquitinating enzyme. *J Biol Chem* 2003;278:23180-6.
- [25] Evans PC, Taylor ER, Coadwell J, Heyninck K, Beyaert R, Kilshaw PJ. Isolation and characterization of two novel A20-like proteins. *Biochem J* 2001;357:617-23.
- [26] Huang J, Teng L, Li L, Liu T, Chen D, Xu LG, et al. ZNF216 Is an A20-like and IkappaB kinase gamma-interacting inhibitor of NFkappaB activation. *J Biol Chem* 2004;279:16847-53.
- [27] Lee S, Tsai YC, Mattera R, Smith WJ, Kostelansky MS, Weissman AM, et al. Structural basis for ubiquitin recognition and autoubiquitination by Rabex-5. *Nat Struct Mol Biol* 2006;13:264-71.
- [28] De Valck D, Heyninck K, Van Crielinge W, Vandenabeele P, Fiers W, Beyaert R. A20 inhibits NF-kappaB activation independently of binding to 14-3-3 proteins. *Biochem Biophys Res Commun* 1997;238:590-4.
- [29] Vincenz C, Dixit VM. 14-3-3 proteins associate with A20 in an isoform-specific manner and function both as chaperone and adapter molecules. *J Biol Chem* 1996;271:20029-34.
- [30] Lademann U, Kallunki T, Jaattela M. A20 zinc finger protein inhibits TNF-induced apoptosis and stress response early in the signaling cascades and independently of binding to TRAF2 or 14-3-3 proteins. *Cell Death Differ* 2001;8:265-72.
- [31] Li L, Hailey DW, Soetandyo N, Li W, Lippincott-Schwartz J, Shu HB, et al. Localization of A20 to a lysosome-associated compartment and its role in NFkappaB signaling. *Biochim Biophys Acta* 2008;1783:1140-9.
- [32] Li Y, Begovich AB. Unraveling the genetics of complex diseases: susceptibility genes for rheumatoid arthritis and psoriasis. *Semin Immunol* 2009;21:318-27.
- [33] Pipari AW, Jr., Hu HM, Yabkowitz R, Dixit VM. The A20 zinc finger protein protects cells from tumor necrosis factor cytotoxicity. *J Biol Chem* 1992;267:12424-7.
- [34] Jaattela M, Mouritzen H, Elling F, Bastholm L. A20 zinc finger protein inhibits TNF and IL-1 signaling. *J Immunol* 1996;156:1166-73.
- [35] Daniel S, Arvelo MB, Patel VI, Longo CR, Shrikhande G, Shukri T, et al. A20 protects endothelial cells from TNF-, Fas-, and NK-mediated cell death by inhibiting caspase 8 activation. *Blood* 2004;104:2376-84.
- [36] De Valck D, Jin DY, Heyninck K, Van de Craen M, Contreras R, Fiers W, et al. The zinc finger protein A20 interacts with a novel anti-apoptotic protein which is cleaved by specific caspases. *Oncogene* 1999;18:4182-90.

- [37] Natoli G, Costanzo A, Guido F, Moretti F, Bernardo A, Burgio VL, et al. Nuclear factor κ B-independent cytoprotective pathways originating at tumor necrosis factor receptor-associated factor 2. *J Biol Chem* 1998;273:31262-72.
- [38] Fries KL, Miller WE, Raab-Traub N. Epstein-Barr virus latent membrane protein 1 blocks p53-mediated apoptosis through the induction of the A20 gene. *J Virol* 1996;70:8653-9.
- [39] Malewicz M, Zeller N, Yilmaz ZB, Weih F. NF κ B controls the balance between Fas and tumor necrosis factor cell death pathways during T cell receptor-induced apoptosis via the expression of its target gene A20. *J Biol Chem* 2003;278:32825-33.
- [40] Jin Z, Li Y, Pitti R, Lawrence D, Pham VC, Lill JR, et al. Cullin3-based polyubiquitination and p62-dependent aggregation of caspase-8 mediate extrinsic apoptosis signaling. *Cell* 2009;137:721-35.
- [41] Heyninck K, Denecker G, De Valck D, Fiers W, Beyaert R. Inhibition of tumor necrosis factor-induced necrotic cell death by the zinc finger protein A20. *Anticancer Res* 1999;19:2863-8.
- [42] Tran TM, Temkin V, Shi B, Pagliari L, Daniel S, Ferran C, et al. TNF α -induced macrophage death via caspase-dependent and independent pathways. *Apoptosis* 2009;14:320-32.
- [43] Storz P, Doppler H, Ferran C, Grey ST, Toker A. Functional dichotomy of A20 in apoptotic and necrotic cell death. *Biochem J* 2005;387:47-55.
- [44] Cooper JT, Stroka DM, Brostjan C, Palmetshofer A, Bach FH, Ferran C. A20 blocks endothelial cell activation through a NF- κ B-dependent mechanism. *J Biol Chem* 1996;271:18068-73.
- [45] Eliopoulos AG, Stack M, Dawson CW, Kaye KM, Hodgkin L, Sihota S, et al. Epstein-Barr virus-encoded LMP1 and CD40 mediate IL-6 production in epithelial cells via an NF- κ B pathway involving TNF receptor-associated factors. *Oncogene* 1997;14:2899-916.
- [46] Ferran C, Stroka DM, Badrichani AZ, Cooper JT, Bach FH. Adenovirus-mediated gene transfer of A20 renders endothelial cells resistant to activation: a means of evaluating the role of endothelial cell activation in xenograft rejection. *Transplant Proc* 1997;29:879-80.
- [47] Ferran C, Stroka DM, Badrichani AZ, Cooper JT, Wrighton CJ, Soares M, et al. A20 inhibits NF- κ B activation in endothelial cells without sensitizing to tumor necrosis factor-mediated apoptosis. *Blood* 1998;91:2249-58.
- [48] Heyninck K, De Valck D, Vanden Berghe W, Van Crielinge W, Contreras R, Fiers W, et al. The zinc finger protein A20 inhibits TNF-induced NF- κ B-dependent gene expression by interfering with an RIP- or TRAF2-mediated transactivation signal and directly binds to a novel NF- κ B-inhibiting protein ABIN. *J Cell Biol* 1999;145:1471-82.
- [49] Longo CR, Arvelo MB, Patel VI, Daniel S, Mahiou J, Grey ST, et al. A20 protects from CD40-CD40 ligand-mediated endothelial cell activation and apoptosis. *Circulation* 2003;108:1113-8.
- [50] Zhu CH, Ying DJ, Mi JH, Zhang W, Dong SW, Sun JS, et al. The zinc finger protein A20 protects endothelial cells from burns serum injury. *Burns* 2004;30:127-33.
- [51] Boone DL, Turer EE, Lee EG, Ahmad RC, Wheeler MT, Tsui C, et al. The ubiquitin-modifying enzyme A20 is required for termination of Toll-like receptor responses. *Nat Immunol* 2004;5:1052-60.
- [52] Turer EE, Tavares RM, Mortier E, Hitotsumatsu O, Advincula R, Lee B, et al. Homeostatic MyD88-dependent signals cause lethal inflammation in the absence of A20. *J Exp Med* 2008;205:451-64.

- [53] Lin R, Yang L, Nakhaei P, Sun Q, Sharif-Askari E, Julkunen I, et al. Negative regulation of the retinoic acid-inducible gene I-induced antiviral state by the ubiquitin-editing protein A20. *J Biol Chem* 2006;281:2095-103.
- [54] Wang YY, Li L, Han KJ, Zhai Z, Shu HB. A20 is a potent inhibitor of TLR3- and Sendai virus-induced activation of NF-kappaB and ISRE and IFN-beta promoter. *FEBS Lett* 2004;576:86-90.
- [55] Saitoh T, Yamamoto M, Miyagishi M, Taira K, Nakanishi M, Fujita T, et al. A20 is a negative regulator of IFN regulatory factor 3 signaling. *J Immunol* 2005;174:1507-12.
- [56] O'Reilly SM, Moynagh PN. Regulation of Toll-like receptor 4 signalling by A20 zinc finger protein. *Biochem Biophys Res Commun* 2003;303:586-93.
- [57] Zetoune FS, Murthy AR, Shao Z, Hlaing T, Zeidler MG, Li Y, et al. A20 inhibits NF-kappa B activation downstream of multiple Map3 kinases and interacts with the I kappa B signalosome. *Cytokine* 2001;15:282-98.
- [58] Song HY, Rothe M, Goeddel DV. The tumor necrosis factor-inducible zinc finger protein A20 interacts with TRAF1/TRAF2 and inhibits NF-kappaB activation. *Proc Natl Acad Sci U S A* 1996;93:6721-5.
- [59] Heyninck K, Beyaert R. The cytokine-inducible zinc finger protein A20 inhibits IL-1-induced NF-kappaB activation at the level of TRAF6. *FEBS Lett* 1999;442:147-50.
- [60] Klinkenberg M, Van Huffel S, Heyninck K, Beyaert R. Functional redundancy of the zinc fingers of A20 for inhibition of NF-kappaB activation and protein-protein interactions. *FEBS Lett* 2001;498:93-7.
- [61] Zhang SQ, Kovalenko A, Cantarella G, Wallach D. Recruitment of the IKK signalosome to the p55 TNF receptor: RIP and A20 bind to NEMO (IKKgamma) upon receptor stimulation. *Immunity* 2000;12:301-11.
- [62] Wertz IE, O'Rourke KM, Zhou H, Eby M, Aravind L, Seshagiri S, et al. De-ubiquitination and ubiquitin ligase domains of A20 downregulate NF-kappaB signalling. *Nature* 2004;430:694-9.
- [63] Newton K, Matsumoto ML, Wertz IE, Kirkpatrick DS, Lill JR, Tan J, et al. Ubiquitin chain editing revealed by polyubiquitin linkage-specific antibodies. *Cell* 2008;134:668-78.
- [64] Hitotsumatsu O, Ahmad RC, Tavares R, Wang M, Philpott D, Turer EE, et al. The ubiquitin-editing enzyme A20 restricts nucleotide-binding oligomerization domain containing 2-triggered signals. *Immunity* 2008;28:381-90.
- [65] Mauro C, Pacifico F, Lavorgna A, Mellone S, Iannetti A, Acquaviva R, et al. ABIN-1 binds to NEMO/IKKgamma and co-operates with A20 in inhibiting NF-kappaB. *J Biol Chem* 2006;281:18482-8.
- [66] Stilo R, Varricchio E, Liguoro D, Leonardi A, Vito P. A20 is a negative regulator of BCL10- and CARMA3-mediated activation of NF-kappaB. *J Cell Sci* 2008;121:1165-71.
- [67] Shembade N, Harhaj NS, Parvatiyar K, Copeland NG, Jenkins NA, Matesic LE, et al. The E3 ligase Itch negatively regulates inflammatory signaling pathways by controlling the function of the ubiquitin-editing enzyme A20. *Nat Immunol* 2008;9:254-62.
- [68] Shembade N, Parvatiyar K, Harhaj NS, Harhaj EW. The ubiquitin-editing enzyme A20 requires RNF11 to downregulate NF-kappaB signalling. *EMBO J* 2009;28:513-22.
- [69] Shembade N, Ma A, Harhaj EW. Inhibition of NF-kappaB signaling by A20 through disruption of ubiquitin enzyme complexes. *Science* 2010;327:1135-9.

- [70] Duwel M, Welteke V, Oeckinghaus A, Baens M, Kloo B, Ferch U, et al. A20 negatively regulates T cell receptor signaling to NF-kappaB by cleaving Malt1 ubiquitin chains. *J Immunol* 2009;182:7718-28.
- [71] Coornaert B, Baens M, Heyninck K, Bekaert T, Haegman M, Staal J, et al. T cell antigen receptor stimulation induces MALT1 paracaspase-mediated cleavage of the NF-kappaB inhibitor A20. *Nat Immunol* 2008;9:263-71.
- [72] Hutti JE, Turk BE, Asara JM, Ma A, Cantley LC, Abbott DW. IkappaB kinase beta phosphorylates the K63 deubiquitinase A20 to cause feedback inhibition of the NF-kappaB pathway. *Mol Cell Biol* 2007;27:7451-61.
- [73] Iha H, Peloponese JM, Verstrepen L, Zapart G, Ikeda F, Smith CD, et al. Inflammatory cardiac valvulitis in TAX1BP1-deficient mice through selective NF-kappaB activation. *EMBO J* 2008;27:629-41.
- [74] Shembade N, Harhaj NS, Liebl DJ, Harhaj EW. Essential role for TAX1BP1 in the termination of TNF-alpha-, IL-1- and LPS-mediated NF-kappaB and JNK signaling. *EMBO J* 2007;26:3910-22.
- [75] Parvatiyar K, Barber GN, Harhaj EW. TAX1BP1 and A20 inhibit antiviral signaling by targeting TBK1-IKKi kinases. *J Biol Chem* 2010;285:14999-5009.
- [76] Verstrepen L, Carpentier I, Verhelst K, Beyaert R. ABINs: A20 binding inhibitors of NF-kappa B and apoptosis signaling. *Biochem Pharmacol* 2009;78:105-14.
- [77] Wagner S, Carpentier I, Rogov V, Kreike M, Ikeda F, Lohr F, et al. Ubiquitin binding mediates the NF-kappaB inhibitory potential of ABIN proteins. *Oncogene* 2008;27:3739-45.
- [78] Oshima S, Turer EE, Callahan JA, Chai S, Advincula R, Barrera J, et al. ABIN-1 is a ubiquitin sensor that restricts cell death and sustains embryonic development. *Nature* 2009;457:906-9.
- [79] Papoutsopoulou S, Symons A, Tharmalingham T, Belich MP, Kaiser F, Kioussis D, et al. ABIN-2 is required for optimal activation of Erk MAP kinase in innate immune responses. *Nat Immunol* 2006;7:606-15.
- [80] Wullaert A, Wielockx B, Van Huffel S, Bogaert V, De Geest B, Papeleu P, et al. Adenoviral gene transfer of ABIN-1 protects mice from TNF/galactosamine-induced acute liver failure and lethality. *Hepatology* 2005;42:381-9.
- [81] Beg AA, Sha WC, Bronson RT, Baltimore D. Constitutive NF-kappa B activation, enhanced granulopoiesis, and neonatal lethality in I kappa B alpha-deficient mice. *Genes Dev* 1995;9:2736-46.
- [82] Hu Y, Baud V, Delhase M, Zhang P, Deerinck T, Ellisman M, et al. Abnormal morphogenesis but intact IKK activation in mice lacking the IKKalpha subunit of IkappaB kinase. *Science* 1999;284:316-20.
- [83] Hu Y, Baud V, Oga T, Kim KI, Yoshida K, Karin M. IKKalpha controls formation of the epidermis independently of NF-kappaB. *Nature* 2001;410:710-4.
- [84] Wang AB, Li HL, Zhang R, She ZG, Chen HZ, Huang Y, et al. A20 attenuates vascular smooth muscle cell proliferation and migration through blocking PI3k/Akt signaling in vitro and in vivo. *J Biomed Sci* 2007;14:357-71.
- [85] Patel VI, Daniel S, Longo CR, Shrikhande GV, Scali ST, Czismadia E, et al. A20, a modulator of smooth muscle cell proliferation and apoptosis, prevents and induces regression of neointimal hyperplasia. *FASEB J* 2006;20:1418-30.
- [86] Chng HW, Camplejohn RS, Stone MG, Hart IR, Nicholson LJ. A new role for the anti-apoptotic gene A20 in angiogenesis. *Exp Cell Res* 2006;312:2897-907.
- [87] Chanudet E, Huang Y, Ichimura K, Dong G, Hamoudi RA, Radford J, et al. A20 is targeted by promoter methylation, deletion and inactivating mutation in MALT lymphoma. *Leukemia* 2010;24:483-7.

- [88] Kato M, Sanada M, Kato I, Sato Y, Takita J, Takeuchi K, et al. Frequent inactivation of A20 in B-cell lymphomas. *Nature* 2009;459:712-6.
- [89] Schmitz R, Hansmann ML, Bohle V, Martin-Subero JI, Hartmann S, Mechttersheimer G, et al. TNFAIP3 (A20) is a tumor suppressor gene in Hodgkin lymphoma and primary mediastinal B cell lymphoma. *J Exp Med* 2009;206:981-9.
- [90] Wu WS, Xu ZX, Chang KS. The promyelocytic leukemia protein represses A20-mediated transcription. *J Biol Chem* 2002;277:31734-9.
- [91] Mustapha S, Kirshner A, De Moissac D, Kirshenbaum LA. A direct requirement of nuclear factor-kappa B for suppression of apoptosis in ventricular myocytes. *Am J Physiol Heart Circ Physiol* 2000;279:H939-45.
- [92] Ladner KJ, Caligiuri MA, Guttridge DC. Tumor necrosis factor-regulated biphasic activation of NF-kappa B is required for cytokine-induced loss of skeletal muscle gene products. *J Biol Chem* 2003;278:2294-303.
- [93] Qin YJ, Zhang ZL, Yu LY, He JW, Hou YN, Liu TJ, et al. A20 overexpression under control of mouse osteocalcin promoter in MC3T3-E1 cells inhibited tumor necrosis factor-alpha-induced apoptosis. *Acta Pharmacol Sin* 2006;27:1231-7.
- [94] Arvelo MB, Cooper JT, Longo C, Daniel S, Grey ST, Mahiou J, et al. A20 protects mice from D-galactosamine/lipopolysaccharide acute toxic lethal hepatitis. *Hepatology* 2002;35:535-43.
- [95] Liuwantara D, Elliot M, Smith MW, Yam AO, Walters SN, Marino E, et al. Nuclear factor-kappaB regulates beta-cell death: a critical role for A20 in beta-cell protection. *Diabetes* 2006;55:2491-501.
- [96] Gon Y, Asai Y, Hashimoto S, Mizumura K, Jibiki I, Machino T, et al. A20 inhibits toll-like receptor 2- and 4-mediated interleukin-8 synthesis in airway epithelial cells. *Am J Respir Cell Mol Biol* 2004;31:330-6.
- [97] Song XT, Evel-Kabler K, Shen L, Rollins L, Huang XF, Chen SY. A20 is an antigen presentation attenuator, and its inhibition overcomes regulatory T cell-mediated suppression. *Nat Med* 2008;14:258-65.
- [98] Oshima N, Ishihara S, Rumi MA, Aziz MM, Mishima Y, Kadota C, et al. A20 is an early responding negative regulator of Toll-like receptor 5 signalling in intestinal epithelial cells during inflammation. *Clin Exp Immunol* 2010;159:185-98.
- [99] Wang J, Ouyang Y, Guner Y, Ford HR, Grishin AV. Ubiquitin-editing enzyme A20 promotes tolerance to lipopolysaccharide in enterocytes. *J Immunol* 2009;183:1384-92.
- [100] Sarma V, Lin Z, Clark L, Rust BM, Tewari M, Noelle RJ, et al. Activation of the B-cell surface receptor CD40 induces A20, a novel zinc finger protein that inhibits apoptosis. *J Biol Chem* 1995;270:12343-6.
- [101] Hess S, Gottfried E, Smola H, Grunwald U, Schuchmann M, Engelmann H. CD40 induces resistance to TNF-mediated apoptosis in a fibroblast cell line. *Eur J Immunol* 1998;28:3594-604.
- [102] Laherty CD, Perkins ND, Dixit VM. Human T cell leukemia virus type I Tax and phorbol 12-myristate 13-acetate induce expression of the A20 zinc finger protein by distinct mechanisms involving nuclear factor kappa B. *J Biol Chem* 1993;268:5032-9.
- [103] Sporn SA, Eierman DF, Johnson CE, Morris J, Martin G, Ladner M, et al. Monocyte adherence results in selective induction of novel genes sharing homology with mediators of inflammation and tissue repair. *J Immunol* 1990;144:4434-41.
- [104] Kumar-Sinha C, Varambally S, Sreekumar A, Chinnaiyan AM. Molecular cross-talk between the TRAIL and interferon signaling pathways. *J Biol Chem* 2002;277:575-85.

- [105] Romero DG, Plonczynski M, Vergara GR, Gomez-Sanchez EP, Gomez-Sanchez CE. Angiotensin II early regulated genes in H295R human adrenocortical cells. *Physiol Genomics* 2004;19:106-16.
- [106] Golovko O, Nazarova N, Tuohimaa P. A20 gene expression is regulated by TNF, Vitamin D and androgen in prostate cancer cells. *J Steroid Biochem Mol Biol* 2005;94:197-202.
- [107] Riachy R, Vandewalle B, Kerr Conte J, Moerman E, Sacchetti P, Lukowiak B, et al. 1,25-dihydroxyvitamin D3 protects RINm5F and human islet cells against cytokine-induced apoptosis: implication of the antiapoptotic protein A20. *Endocrinology* 2002;143:4809-19.
- [108] Zhang A, Zhang M, Wang Y, Xie H, Zheng S. Calcitriol prolongs recipient survival by inducing expression of zinc-finger protein A20 and inhibiting its downstream gene following rat orthotopic liver transplantation. *Immunopharmacol Immunotoxicol* 2006;28:591-600.
- [109] Prasad AS, Bao B, Beck FW, Kucuk O, Sarkar FH. Antioxidant effect of zinc in humans. *Free Radic Biol Med* 2004;37:1182-90.
- [110] Hayakawa K, Hiramatsu N, Okamura M, Yamazaki H, Nakajima S, Yao J, et al. Acquisition of anergy to proinflammatory cytokines in nonimmune cells through endoplasmic reticulum stress response: a mechanism for subsidence of inflammation. *J Immunol* 2009;182:1182-91.
- [111] Laherty CD, Hu HM, Opipari AW, Wang F, Dixit VM. The Epstein-Barr virus LMP1 gene product induces A20 zinc finger protein expression by activating nuclear factor kappa B. *J Biol Chem* 1992;267:24157-60.
- [112] Miller WE, Earp HS, Raab-Traub N. The Epstein-Barr virus latent membrane protein 1 induces expression of the epidermal growth factor receptor. *J Virol* 1995;69:4390-8.
- [113] Yurochko AD, Huang ES. Human cytomegalovirus binding to human monocytes induces immunoregulatory gene expression. *J Immunol* 1999;162:4806-16.
- [114] Onose A, Hashimoto S, Hayashi S, Maruoka S, Kumasawa F, Mizumura K, et al. An inhibitory effect of A20 on NF-kappaB activation in airway epithelium upon influenza virus infection. *Eur J Pharmacol* 2006;541:198-204.
- [115] Yokota S, Okabayashi T, Yokosawa N, Fujii N. Measles virus P protein suppresses Toll-like receptor signal through up-regulation of ubiquitin-modifying enzyme A20. *FASEB J* 2008;22:74-83.
- [116] Nguyen H, Sankaran S, Dandekar S. Hepatitis C virus core protein induces expression of genes regulating immune evasion and anti-apoptosis in hepatocytes. *Virology* 2006;354:58-68.
- [117] Maeda S, Otsuka M, Hirata Y, Mitsuno Y, Yoshida H, Shiratori Y, et al. cDNA microarray analysis of Helicobacter pylori-mediated alteration of gene expression in gastric cancer cells. *Biochem Biophys Res Commun* 2001;284:443-9.
- [118] Mysorekar IU, Mulvey MA, Hultgren SJ, Gordon JI. Molecular regulation of urothelial renewal and host defenses during infection with uropathogenic Escherichia coli. *J Biol Chem* 2002;277:7412-9.
- [119] Yurochko AD, Liu DY, Eierman D, Haskill S. Integrins as a primary signal transduction molecule regulating monocyte immediate-early gene induction. *Proc Natl Acad Sci U S A* 1992;89:9034-8.
- [120] Hjelmeland AB, Wu Q, Wickman S, Eyler C, Heddleston J, Shi Q, et al. Targeting A20 decreases glioma stem cell survival and tumor growth. *PLoS Biol* 2010;8:e1000319.

- [121] Honma K, Tsuzuki S, Nakagawa M, Tagawa H, Nakamura S, Morishima Y, et al. TNFAIP3/A20 functions as a novel tumor suppressor gene in several subtypes of non-Hodgkin lymphomas. *Blood* 2009.
- [122] Miao HS, Yu LY, Hui GZ, Guo LH. Antiapoptotic effect both in vivo and in vitro of A20 gene when transfected into rat hippocampal neurons. *Acta Pharmacol Sin* 2005;26:33-8.
- [123] Sass G, Shembade ND, Haimeri F, Lamoureux N, Hashemolhosseini S, Tannapfel A, et al. TNF pretreatment interferes with mitochondrial apoptosis in the mouse liver by A20-mediated down-regulation of Bax. *J Immunol* 2007;179:7042-9.
- [124] Honma K, Tsuzuki S, Nakagawa M, Tagawa H, Nakamura S, Morishima Y, et al. TNFAIP3/A20 functions as a novel tumor suppressor gene in several subtypes of non-Hodgkin lymphomas. *Blood* 2009;114:2467-75.
- [125] Ramsey HE, Da Silva CG, Longo CR, Csizmadia E, Studer P, Patel VI, et al. A20 protects mice from lethal liver ischemia/reperfusion injury by increasing peroxisome proliferator-activated receptor- α expression. *Liver Transpl* 2009;15:1613-21.
- [126] Daniel S, Patel VI, Shrikhande GV, Scali ST, Ramsey HE, Csizmadia E, et al. The universal NF-kappaB inhibitor a20 protects from transplant vasculopathy by differentially affecting apoptosis in endothelial and smooth muscle cells. *Transplant Proc* 2006;38:3225-7.
- [127] Eliopoulos AG, Blake SM, Floettmann JE, Rowe M, Young LS. Epstein-Barr virus-encoded latent membrane protein 1 activates the JNK pathway through its extreme C terminus via a mechanism involving TRADD and TRAF2. *J Virol* 1999;73:1023-35.
- [128] De Valck D, Heyninck K, Van Crielinge W, Contreras R, Beyaert R, Fiers W. A20, an inhibitor of cell death, self-associates by its zinc finger domain. *FEBS Lett* 1996;384:61-4.
- [129] Hu X, Yee E, Harlan JM, Wong F, Karsan A. Lipopolysaccharide induces the antiapoptotic molecules, A1 and A20, in microvascular endothelial cells. *Blood* 1998;92:2759-65.
- [130] Heyninck K, Kreike MM, Beyaert R. Structure-function analysis of the A20-binding inhibitor of NF-kappa B activation, ABIN-1. *FEBS Lett* 2003;536:135-40.
- [131] Van Huffel S, Delaei F, Heyninck K, De Valck D, Beyaert R. Identification of a novel A20-binding inhibitor of nuclear factor-kappa B activation termed ABIN-2. *J Biol Chem* 2001;276:30216-23.
- [132] Wullaert A, Verstrepen L, Van Huffel S, Adib-Conquy M, Cornelis S, Kreike M, et al. LIND/ABIN-3 is a novel lipopolysaccharide-inducible inhibitor of NF-kappaB activation. *J Biol Chem* 2007;282:81-90.
- [133] Bohgaki M, Tsukiyama T, Nakajima A, Maruyama S, Watanabe M, Koike T, et al. Involvement of Ymer in suppression of NF-kappaB activation by regulated interaction with lysine-63-linked polyubiquitin chain. *Biochim Biophys Acta* 2008;1783:826-37.

Figure legends

Figure 1: A20 is a susceptibility gene for multiple diseases

Figure 2: Structure and activation of the A20 promoter

A. Structure of A20 promoter. A20 contains upstream of the core promoter two binding sites for NF- κ B, six Sp-1 sites, and an E-box overlapping the ELIE site. B. Under basal conditions, the general transcription apparatus (composed of RNA polymerase II, TFIID, TAFs, TBP and others), together with the co-activators CBP and p300, is constitutively associated with the core promoter. Together with Sp-1, this allows a single round of transcription initiation. In addition, A20 mRNA elongation is inhibited by DSIF, whose activity is controlled by the E-box binding protein USF1. C. Upon stimulation, NF- κ B is activated and binds to the κ B sites in the A20 promoter, leading to re-initiation and multiple rounds of transcription. In addition, NF- κ B displaces USF1 from the promoter and takes over the control of DSIF activity, permitting mRNA elongation by RNA polymerase II.

Figure 3: Domain structure of A20

A20 consists of an N-terminal OTU domain and 7 C-terminal zinc fingers (z), mediating respectively K63 deubiquitination and K48-polyubiquitination. Regions of phosphorylation, proteolytic cleavage and binding of other proteins are indicated.

Figure 4: Inhibition of TNF and LPS-induced NF- κ B signaling by A20

NF- κ B signaling involves the interaction of multiple proteins, which can be subdivided in adaptors (orange), kinases (green), and E3 ubiquitin ligases (blue). The latter can mediate three different types of polyubiquitination: K63 linked, K48 linked, linear (depicted in different colors). Binding of TNF or LPS to respectively TNF-RI or TLR4 induces the recruitment of different adaptor proteins to the receptor. These form a platform for the binding of E3 ubiquitin ligases and kinases. The E3 ubiquitin ligases together with E2 ligases (Ubc5 and Ubc13) then mediate the K63 polyubiquitination of themselves and the kinases RIP1 and IRAK1. This forms a signal for the recruitment of specific ubiquitin binding adaptor proteins that function to recruit the kinases TAK1 and IKK, resulting in TAK-1 mediated IKK phosphorylation and activation. Similarly, the ubiquitin ligase complex LUBAC is recruited and is responsible for the modification of the adaptor protein NEMO with a linear

polyubiquitin chain of which the function is still unclear. Activated IKK then phosphorylates the NF- κ B inhibitor protein I κ B α , leading to its K48-polyubiquitination by the SCF/ β TRCP ubiquitin ligase complex. This type of modification is recognized by the proteasome and results in I κ B α degradation, thus allowing NF- κ B (depicted here as a p50/p65 dimer) to translocate to the nucleus. One of the many NF- κ B responsive genes encodes the NF- κ B inhibitor protein A20, which provides a negative feedback mechanism. A20 is a deubiquitinase that inhibits NF- κ B signaling by multiple mechanisms: (1) ubiquitin-editing of RIP1 (K63-deubiquitination followed by its K48-polyubiquitination, resulting in RIP1 degradation); (2) K63-deubiquitination of TRAF6; (3,4) disruption of the interaction of the E3 ubiquitin ligases cIAP and TRAF6 with the E2 enzymes Ubc5 and Ubc13; (5,6) K48-polyubiquitination of the E2 enzymes Ubc5 and Ubc13, leading to their proteasomal degradation. A20 binds to specific ubiquitin binding adaptor proteins (ABINs, TAX1BP1), which recruit A20 to specific ubiquitinated substrates or mediate the binding of other proteins such as the E3 ubiquitin ligases RNF11 and ITCH, which are essential for A20 functioning.

Table 1: Expression pattern of A20

Stimulus	Cell type	Ref.
TNF	Osteosarcoma cell line U2OS	[90]
	Vascular smooth muscle cells (VSMC)	[85]
	Jurkat T-cells	[15]
	Human microvascular endothelial cells (HMEC-1)	[82]
	Ventricular myocytes	[91]
	Myotubes	[92]
	Osteoblastic MC3T3-E1 cell line	[93]
	Primary human hepatocytes, hepatocellular carcinoma cell line HepG2	[94]
	MCF-7 breast carcinoma cell line	[33]
	β cells (primary islets, Min6)	[95]
IL-1	β cells (primary islets, Min6)	[95]
LPS	Bone marrow-derived macrophages (BMDMs)	[51]
	Primary human airway epithelial cells	[96]
	HMEC-1 cells	[82]
	Bone marrow-derived dendritic cells (BMDCs)	[97]
	Freshly isolated human blood monocytes	[98]
	Enterocytes (IEC-6, IEC-18, RIE-1, SW480, HCT-15)	[98, 99]
CpG DNA	Enterocytes (IEC-18)	[99]
Flagellin	Enterocytes (HCT-15, HT-29, mouse primary cells, Colon-26)	[98]
CD40	B-cells	[100]
	Fibroblast cell line SV80	[101]
TPA	Jurkat T-cells, U937 promonocytic cells	[102]
	Freshly isolated human blood monocytes	[103]
TRAIL	MCF-7	[104]
Angiotensin II	Human adrenocortical cells	[105]
Peptidoglycan	Primary human airway epithelial cells	[96]
H₂O₂	HeLa cells	[43]
Serum from patients with burn injuries	Human umbilical vein endothelial cells (HUVEC)	[50]

TNF-RII	Murine CT6 cells	[58]
Androgen	LNCaP prostate cancer cells	[106]
CD3	Do-11.10 T cell hybridomas, CD4 ⁺ and CD8 ⁺ T-cells	[39]
Vitamin D₃	Increase of expression: * rat insulin-producing β -cells (RINm5F) * human islet cells * liver allografts Decrease of expression: * LNCaP cells	[107, 108] [106]
Zinc depletion	Diminished LPS- and TPA-induced expression in promyelocytic leukemia HL-60 cells	[109]
ER stress	Mesangial cells, NRK-52E cells	[110]
Epstein-Barr virus latent membrane protein LMP1	B-cells Human epithelial cells	[111] [112]
Human T-cell leukemia virus type I oncoprotein Tax	Jurkat T cells	[102]
Human cytomegalovirus	Human peripheral blood monocytes	[113]
Influenza virus	Human bronchial epithelial cells, lung homogenates from mice	[114]
Sendai virus	U937	[54]
Measles virus P protein	Monocytes (THP1, U937)	[115]
Hepatitis C virus core protein	HepG2	[116]
Helicobacter pylori	Human gastric cancer cells	[117]
Uropathogenic Escherichia coli	Mice bladder tissues	[118]
VLA-4 stimulation	Monocytes	[119]

Table 2: Biological activities of A20

Function	Stimulus	Cells	Evidence	Mechanisms	Ref
Anti-apoptotic	TNF	Thymocytes, MCF7S1, WEHI-S, endothelial cells, β -cells, Jurkat, rat primary hippocampal neurons, NIH3T3, the osteoblastic MC3T3-E1, SV80, primary hepatocytes, NIH3T3, primary glioma stem cells	knockout, overexpression, RNAi	* prevention of recruitment of RIP1 and TRADD to TNF-RI * inhibition of p53 * inhibition of Bax expression	[10, 15, 33, 36, 38, 94, 96, 101, 120-123]
	Fas	Endothelial cells	overexpression		[35]
	TRAIL	Lung carcinoma (H460)	overexpression	*Caspase 8 deubiquitination	[40]
	Natural killer cells	Endothelial cells	overexpression		[35]
	Serum deprivation	B-cells	overexpression		[100]
	Serum from patients with burn injuries	Human umbilical vein endothelial cells (HUVEC)	overexpression		[50]
	LPS	Human microvascular endothelial cells (HMEC-1)	overexpression		[82]
	CD40	Human coronary artery endothelial cells (HCAEC)	overexpression		[49]
	AICD	Do-11.10 T cell hybridomas	overexpression		[39]
	etoposide	Lymphoblastoid cell lines	RNAi		[124]
	H ₂ O ₂	Lymphoblastoid cell lines	RNAi		[124]
Anti-necrosis	TNF	L929	overexpression	* inhibition of phospholipases; production of reactive oxygen species	[41, 43]
	Complement	Endothelial cells	overexpression		[35]
	Hypoxia-	Mouse hepatocytes (NMuLi)	overexpression	* increase in PPAR α expression	[125]

	reoxygenation				
	H ₂ O ₂	Mouse hepatocytes (NMuLi)	overexpression	* increase in PPAR α expression	[125]
Pro-apoptotic	Fas-and cytokine mediated	Vascular smooth muscle cells (VSMC)	overexpression	*NF- κ B inhibition	[43, 126]
NF-κB inhibition	TNF	MCF7S1, bovine aortic endothelial cells (BAEC), HEK293T, MEF, HeLa	overexpression, knockout, RNAi	* ubiquitin editing of RIP1, leading to its degradation * deubiquitination of NEMO * inhibition of TRAF/cIAP1 E3 ligase activity by antagonizing the interaction with E2 enzymes * induction of proteasomal degradation of E2 enzymes * degradation of TRAF2 via the lysosomes	[10, 34, 44, 58, 62, 65, 96] [69] [32]
	IL-1 LPS	MCF7S1, HEK293T, human airway epithelial cells, HEK293, BAEC, macrophages	overexpression, knockout	* deubiquitination of TRAF6 * inhibition of TRAF6 E3 ligase activity by antagonizing the interaction with E2 enzymes * induction of proteasomal degradation of E2 enzymes	[34, 44, 51, 56, 58, 96] [69]
	H ₂ O ₂	HeLa, endothelial cells	overexpression, RNAi		[43, 47]
	TPA	BAEC, HEK293T	overexpression		[44] [71]
	anti-CD3	Jurkat	RNAi	*MALT1 deubiquitination	[70]
	API-MALT1	HEK293T	overexpression		[71]
	RIG-I	HEK293	overexpression		[53]
	Helicobacter pylori	Human gastric cancer cells	overexpression		[117]

	Influenza virus	Human bronchial epithelial cells	overexpression	[114]
	Peptidoglycan	Human airway epithelial cells	overexpression	[96]
	Serum from patients with burn wounds	HUVEC	overexpression	[50]
	CD40	HEK293T	overexpression	[58]
	TNF-RII	HEK293T	overexpression	[58]
	NOD1/NOD2	HEK293T	overexpression	* deubiquitination of RIP2 [64]
	LMP1	Human keratinocytes, HEK293	overexpression	[45, 127]
AP-1 inhibition	TNF	MCF7S1	overexpression	[34]
	IL-1	MCF7S1	overexpression	[34]
	LMP1	HEK293	overexpression	[127]
	TLR-4	HEK293	overexpression	[56]
IRF3 inhibition	RIG-I	HEK293	overexpression	[53]
	TRIF	HEK293	overexpression	* reduction of TRIF expression [53]
	Sendai virus	HEK293	overexpression, RNAi	[53, 54]
	Poly(I:C)	HEK293/TLR3, HeLa, THP-1	overexpression, RNAi	[54, 55]
	Newcastle disease virus	HEK293/TLR-3	overexpression, RNAi	[55]
	IKK ϵ /TBK1 VSV	HEK293	overexpression	*Inhibition of IKK ϵ /TBK1 ubiquitination by disrupting their interaction with TRAF3 E3 ligase [75]
Skin differentiation			knockout mice	* impaired NF- κ B activation [5, 10, 57] * interaction with IKK1
Cell cycle arrest		VSMC	overexpression	* arrest in G ₀ /G ₁ by inhibition of PI3K/Akt/GSK3 β signaling [84, 85] * upregulation of p53, p21 ^{waf1}

				and p27 ^{k_p}
Mitogenic in the liver				[94]
Neovascularisation	HUVEC	RNAi		[86]

Table 3: A20-interacting proteins

Interacting protein	Evidence	Interacting domain in A20	Interacting domain in partner protein	Ref
A20	Y2H	C-terminal ZnF-domain (AA 373-790, human)		[128]
14-3-3	Y2H, in vitro binding assay, COIP (overexpression, semi-endogenous)	RSKSDP sequence (human and murine) between ZnF3 and ZnF4, with the second serine being phosphorylated		[28, 29]
NEMO	Y2H, COIP (overexpression, semi-endogenous upon overexpression of TNF-RI)	C-terminal ZnF-domain (AA 387-790, human) (AA 367-468, murine)	AA 95-218 (human)	[60, 61]
TAX1BP1	Y2H, COIP (overexpression)	C-terminal ZnF-domain (AA 468-699, murine) (AA 373-790, human)		[60, 129]
TRIF	COIP (overexpression)			[54]
TBK1	COIP (overexpression)	N- and C-terminus (AA 1-378 and AA 379-790, human)		[55]
IKK ϵ	COIP (overexpression)	N- and C-terminus (AA 1-378 and AA 379-790, human)		[55]
IKK1	COIP (overexpression)	C-terminal ZnF-domain (AA 387-790, human)		[57]
TRAF1	Y2H, COIP (overexpression, endogenous)	N-terminal domain (AA 1-386, human)	TRAF-N domain	[58]
TRAF2	Y2H, COIP (overexpression)	N-terminal domain (AA 1-386, human)	TRAF-N domain (AA 264-358)	[58]

TRAF6	COIP (overexpression)			[59]
ABIN-1	Y2H, COIP (overexpression)	C-terminal ZnF-domain (AA 369-775, murine)	AA 420-647, containing AHD1	[48, 130]
ABIN-2	Y2H, COIP (overexpression)	C-terminal ZnF-domain (AA 369-775, murine)	AA 251-430, containing AHD1	[131]
ABIN-3	COIP (overexpression)	C-terminal ZnF-domain	AHD1	[132]
MALT1	COIP (endogenous)		C-terminal part, containing the caspase- like domain	[71]
Ymer	Y2H, COIP (overexpression, endogenous)	N-terminal domain (AA 91-305) containing the OTU-domain	AA 65-134 containing the ubiquitin- binding region	[133]
RNF11	COIP (overexpression, endogenous upon TNF stimulation)		RING-H2 and PPXY motif	[68]
Caspase 8	COIP (overexpression)			[40]

Figure 1

Figure 3

Lys63 deubiquitination

Lys48 ubiquitination

Figure 4