



Lectures on F-theory compactifications and model building

Timo Weigand

► **To cite this version:**

Timo Weigand. Lectures on F-theory compactifications and model building. Classical and Quantum Gravity, IOP Publishing, 2010, 27 (21), pp.214004. 10.1088/0264-9381/27/21/214004. hal-00638670

HAL Id: hal-00638670

<https://hal.archives-ouvertes.fr/hal-00638670>

Submitted on 7 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lectures on F-theory compactifications and model building

Timo Weigand¹

¹ *ITP Heidelberg, Philosophenweg 19,
69120 Heidelberg, Germany*

`tweigand@thphys.uni-heidelberg.de`

Abstract

These lecture notes are devoted to formal and phenomenological aspects of F-theory. We begin with a pedagogical introduction to the general concepts of F-theory, covering classic topics such as the connection to Type IIB orientifolds, the geometry of elliptic fibrations and the emergence of gauge groups, matter and Yukawa couplings. As a suitable framework for the construction of compact F-theory vacua we describe a special class of Weierstrass models called Tate models, whose local properties are captured by the spectral cover construction. Armed with this technology we proceed with a survey of F-theory GUT models, aiming at an overview of basic conceptual and phenomenological aspects, in particular in connection with GUT breaking via hypercharge flux.

Contents

1	Introduction	2
2	A first encounter with F-theory	4
2.1	The need for a non-perturbative formulation of Type IIB with 7-branes	4
2.2	F/M-theory duality and Calabi-Yau 4-folds	10
2.3	The geometry of elliptic fibrations	11
2.4	Sen’s orientifold limit	15
2.5	Gauge symmetry from degenerations	16
3	Technology for F-theory compactifications	18
3.1	Tate models	18
3.2	Fluxes and 3-brane tadpoles	21
3.3	Matter curves and Yukawa points	22
3.4	F-theory-heterotic duality	26
3.5	The spectral cover construction for F-theory models	30
4	Phenomenological applications to GUT model building	34
4.1	SU(5) GUT models and the principle of decoupling	35
4.2	Options for GUT breaking	37
4.3	Some constraints from hypercharge flux	38
4.4	Proton decay	40
4.5	Further developments	41

1 Introduction

String phenomenology is the branch of string theory that takes the theory seriously as a consistent, fundamental theory of gravitational and particle physics interactions in four dimensions. Its primary goal is to understand the solutions of this theory and their implications for four-dimensional particle phenomenology and cosmology. This includes the investigation of both the mathematical structure of the space of string vacua and of their concrete phenomenological properties.

In the perturbative region of the string landscape, two corners exhibit particularly attractive four-dimensional solutions: The heterotic string heavily exploits the availability of exceptional gauge symmetry, into which the observed gauge group of the Standard Model can be embedded in elegant ways realising the idea of a grand unified theory (GUT). Many different techniques have been developed to study these solutions in various regions of the moduli space, ranging from heterotic orbifolds and free-fermionic constructions to smooth Calabi-Yau compactifications with vector bundles, see e.g. the review [1] and references therein. Since both gravity and gauge dynamics descend from the closed string sector, all aspects of four-dimensional physics are sensitive to the global structure of the compactification space.

Perturbative Type II orientifolds with D-branes, on the other hand, are based on the classical gauge groups $U(N)$, $SO(2N)$ and $Sp(2N)$. This type of constructions can directly accomodate the Standard Model gauge group, while GUT physics is not immediate. The huge literature on Type IIA orientifolds with intersecting D6-branes and on Type IIB orientifolds with D7/D3-branes (reviewed e.g. in [2–4]) exploits furthermore the localisation of the gauge degrees of freedom along the D-branes, which are therefore of a different physical origin than

the gravity sector in the bulk. This opens up the possibility of exploring a certain subclass of phenomenological questions in the context of local models, while other aspects of the associated four-dimensional physics remain sensitive to the full compactification details.

The strict separation between the phenomenological properties of the heterotic and brane constructions ceases to exist at finite values of the string coupling g_s . In fact, in all but a minute class of brane constructions g_s is dynamical and varies over the compactification space in such a way as to leave the perturbative regime somewhere. The correct way to describe Type IIB compactifications with 7-branes in this generic situation is called F-theory [5]. F-theory is a fascinating subject by itself because it geometrises the backreaction of the 7-branes on the ambient space and is therefore, in some sense, *the* way to think about 7-branes. It incorporates certain strong coupling effects with breath-taking elegance. The rich and by now classic literature on the more formal aspects of F-theory reflects the amount to which these non-perturbative phenomena have mesmerized string theorists.

More recently it has been exploited that the four-dimensional solutions of F-theory are also interesting from a phenomenological viewpoint [6–9]. This is because at strong coupling new degrees of freedom - (p,q) strings - become light and realise exceptional gauge symmetries even in a theory based on branes. The perturbative dichotomy between localisation of gauge degrees of freedom and exceptional gauge dynamics is therefore resolved. This bears exciting prospects for GUT model building and has been the subject of fruitful and intensive investigations in recent times. Most importantly, F-theory compactifications also inherit the favourable properties of M-theory and of Type IIB orientifolds with respect to moduli stabilisation: The combination of 3-brane instanton effects and background fluxes allows in principle for the stabilisation of Kähler and complex structure moduli within the framework of warped Calabi-Yau compactifications. This is motivation enough to take F-theory seriously as a promising corner for string phenomenology.

When dealing with F-theory compactifications one must be aware that to date there exists no description of F-theory as a fundamental theory. In this respect F-theory has a status very different from M-theory, which can - at least conjecturally - be conceived as the theory reducing in its long-wavelength limit to eleven-dimensional supergravity coupled to membranes. Early ideas [5, 10, 11] to define an analogous twelve-dimensional theory whose fundamental objects are 3-branes have not lead to a consistent picture. Rather F-theory should be thought of as a genuinely non-perturbative description of a class of string vacua which, in certain limits, is accessible by string dualities from three different corners of the M-theory star. These are

- F-theory as backreacted Type IIB orientifolds with D7-branes and varying dilaton,
- F-theory as dual to $E_8 \times E_8$ heterotic theory,
- F-theory as dual to M-theory on a vanishing T^2 .

Of these three, the F/M-theory duality probably captures the dynamics in the most general way. While particularly fruitful for concrete applications, the first two dualities can sometimes be misleading because they describe only certain aspects of the dynamics of a typical F-theory compactification.

These lectures intend to provide a pedagogical introduction to some of the technology and the phenomenological applications of F-theory model building. A complimentary set of lecture notes covering general aspects of F-theory is [12], while the specifics of F-theory GUT model building are surveyed in the review [13].

In section 2.1 we begin with a definition of F-theory from the perspective of Type IIB orientifolds with 7-branes. The F/M-theory duality is briefly sketched in section 2.2. Given

the geometric nature of F-theory, the bread and butter of the business is to understand the geometry of elliptic fibrations, which we approach in an elementary manner in section 2.3. The connection to the Type II orientifold picture is made via Sen’s orientifold limit in section 2.4. A closer investigation of the appearance of gauge degrees of freedom from the singularities of the elliptic fibration concludes our first encounter with F-theory in section 2.5.

In section 3 we introduce some of the more advanced technology of F-theory compactifications. The basic algorithm to read off the gauge groups from a given model is presented in section 3.1. The corresponding Tate models represent a convenient framework for F-theory compactifications. Besides the pure geometry, gauge flux is an essential ingredient in F-theory models (see section 3.2). Note that it is this aspect that is currently the least understood. An account of charged matter and Yukawa couplings follows in section 3.3. In section 3.4 we recall basic aspects of F-theory/heterotic duality; an important ingredient is the construction of vector bundles on elliptically fibered Calabi-Yau 3-folds via the spectral cover construction. Section 3.5 concludes our tour through the model building rules of F-theory with a review of the recent application of the spectral cover construction in general F-theory models with no heterotic dual.

In the final part of these lecture notes we apply what we have learned in sections 2 and 3 to the construction of $SU(5)$ GUT vacua. This is by now a vast and dynamical field, and rather than aiming at completeness we focus on some fundamental aspects. After briefly describing the decoupling idea underlying local models in section 4.1, we investigate various options for GUT symmetry breaking in section 4.2. GUT breaking by hypercharge flux is critically assessed in section 4.3. The *experimentum crucis* in GUT model building is proton decay, and we explain some of the challenges within the F-theory context in section 4.4. An incomplete list of further phenomenological topics can be found in 4.5.

2 A first encounter with F-theory

2.1 The need for a non-perturbative formulation of Type IIB with 7-branes

Backreaction from 7-branes

In this section we approach F-theory as the strong coupling limit of Type IIB orientifolds with O7/O3-planes and D7/D3-branes. For background on Type II orientifolds we refer to existing reviews such as [2–4]. The usual philosophy in the description of perturbative Type II orientifolds with D-branes is to neglect the backreaction of the branes and the orientifold planes on the background geometry in the spirit of a probe approximation. This is justified as long as asymptotically away from the brane the backreaction becomes negligible. In this case, one can consider a large volume limit in which knowledge of the detailed form of the solution is not required at least to understand the main properties of the string vacuum.

To see when this approximation is justified, we consider a p -brane in ten dimensions. It represents a source term in the normal $n = 9 - p$ spatial directions. At a heuristic level, this leads to a Poisson equation for the background fields sourced by the brane. Schematically denoting these sourced fields as Φ , one can write this as

$$\Delta\Phi(r) \simeq \delta(r) \implies \Phi(r) \simeq \frac{1}{r^{n-2}} \quad n > 2. \quad (1)$$

More precisely, within Type II supergravity the BPS solution for a stack of N p -branes along directions $\mu = 0, 1, \dots, p$, $p < 7$, takes the form [14]

$$ds^2 = H_p^{-1/2} \eta_{\mu\nu} dx^\mu dx^\nu + H_p^{1/2} \sum_i dx^i dx^i, \quad (2)$$

$$e^{2\phi} = e^{2\phi_0} H_p^{\frac{3-p}{2}}, \quad C_{p+1} = \frac{H_p^{-1} - 1}{e^{\phi_0}} dx^0 \wedge \dots \wedge dx^p, \quad (3)$$

$$H_p = 1 + \left(\frac{r_p}{r}\right)^{(7-p)}, \quad r_p^{(7-p)} = \text{const } e^{\phi_0} N. \quad (4)$$

Here ϕ denotes the ten-dimensional dilaton and the Ramond-Ramond $(p+1)$ -form potential C_{p+1} couples electrically to the p -brane. The backreaction is governed by the harmonic function H_p , which asymptotes to unity away from the brane. In particular, e^{ϕ_0} denotes the asymptotic value of e^ϕ and can be taken as the value of the string coupling g_s relevant in the large volume limit.

However, the above logic goes through only if the codimension of the brane $n > 2$. The critical case $n = 2$ corresponds precisely to D7-branes in Type IIB theory. A D7-brane along, say, dimensions $0, \dots, 7$ looks like a charged point particle localised in the two normal directions $8, 9$ - a cosmic string [15]. Solutions to the two-dimensional Poincaré equation scale *logarithmically* with the distance to the source. Such a logarithmic profile is in sharp contrast with the favourable asymptotics for lower dimensional branes encountered above.

Let us see how this heuristic argument applies in more detail. Recall, e.g. from [4], that the string frame Type IIB effective action in the democratic formulation is given by

$$\begin{aligned} S_{IIB}^{(S)} = & \frac{2\pi}{\ell_s^8} \left(\int d^{10}x e^{-2\phi} (\sqrt{-g} R + 4\partial_M \phi \partial^M \phi) - \frac{1}{2} e^{-2\phi} \int H_3 \wedge *H_3 \right. \\ & \left. - \frac{1}{4} \sum_{p=0}^4 \int F_{2p+1} \wedge *F_{2p+1} - \frac{1}{2} \int C_4 \wedge H_3 \wedge F_3 \right). \end{aligned} \quad (5)$$

Here $\ell_s = 2\pi\sqrt{\alpha'}$ and the field strengths are defined as

$$\begin{aligned} H_3 &= dB_2, \quad F_1 = dC_0, \quad F_3 = dC_2 - C_0 dB_2, \\ F_5 &= dC_4 - \frac{1}{2} C_2 \wedge dB_2 + \frac{1}{2} B_2 \wedge dC_2, \end{aligned} \quad (6)$$

supplemented by the duality relations $F_9 = *F_1$, $F_7 = -*F_3$, $F_5 = *F_5$ at the level of equations of motion.

The RR-field sourced electrically by a D7-brane is C_8 and is dual to the axion C_0 that combines with the string coupling $g_s = e^\phi$ into the complex axio-dilaton

$$\tau = C_0 + \frac{i}{g_s}. \quad (7)$$

The D7-brane action is the sum of the two terms

$$S_{\text{DBI}} = -\frac{2\pi}{\ell_s^8} \int d^7\zeta e^{-\phi} \sqrt{\det(g + 2\pi\alpha'\mathcal{F}/2\pi)}, \quad (8)$$

$$S_{\text{CS}} = -\frac{2\pi}{\ell_s^8} \int \text{tr} \exp(2\pi\alpha'\mathcal{F}) \sum_p C_{2p} \sqrt{\frac{\hat{A}(T)}{\hat{A}(N)}}, \quad (9)$$

where $2\pi\alpha'\mathcal{F} = 2\pi\alpha'F + B_2$ in terms of the Yang-Mills field strength F and the last factor denotes curvature contribution in terms of the A-roof-genus (see e.g. [16]) of the tangent and normal bundle to the D7-brane. Define the complex coordinate $z = x^8 + ix^9$ for the dimensions perpendicular to the 7-brane. Taking into account constraints from supersymmetry, it turns out that the axio-dilaton must be a holomorphic function in z . Therefore the Poincaré equation for C_8 in presence of a 7-brane at $z = z_0$ takes the form

$$d * F_9 = \delta^{(2)}(z - z_0). \quad (10)$$

The integrated form of (10) is

$$1 = \int_{\mathcal{C}} d * F_9 = \oint_{S^1} F_1 = \oint_{S^1} dC_0. \quad (11)$$

A simple solution can be found that is valid close to the brane at $z = z_0$,

$$\tau(z) = \tau_0 + \frac{1}{2\pi i} \ln(z - z_0) + \dots, \quad (12)$$

where we have omitted possible regular terms in z . Note in particular that $g_s \rightarrow 0$ at the position of the brane $z = z_0$.¹

Far away from the brane, the simple solution of τ must be modified. Nonetheless we see already that the background exhibits a logarithmic branch cut in the complex plane normal to the D7-brane. A careful analysis of the Einstein equations reveals [15] that asymptotically away from the brane, spacetime becomes locally flat, but suffers a deficit angle. Contrary to the backreaction of lower-dimensional branes, this effect does not asymptote away and the probe philosophy is, strictly speaking, not applicable in this region.

Despite the non-trivial deficit angle at infinity, one can identify a region where the backreaction of the brane on the geometry is negligible. Following the lucid discussion in [17] one can, ignoring the higher terms, rewrite the solution (12) as

$$\tau(z) = \frac{1}{2\pi i} \ln \frac{z - z_0}{\lambda} \quad \Longrightarrow \quad e^{-\phi} = -\frac{1}{2\pi} \ln \left| \frac{z - z_0}{\lambda} \right| \quad (13)$$

in the vicinity of the brane, where λ is related to τ_0 . At the point $z - z_0 = \lambda$ one encounters $g_s = e^\phi \rightarrow \infty$. The presence of this special point breaks the naively expected rotational invariance around the source, a clear sign of backreaction. However, for $|z - z_0| \ll |\lambda|$ the geometry is approximately flat. It is this region that is the analogue of the asymptotic large distance limit where the backreaction of p -branes with $p < 7$ discussed above is negligible. The limit of weak coupling is the one where this region (and distances therein) is large enough to trust effective supergravity, i.e. where $\lambda \rightarrow \infty$.² I.e. in this limit the inevitable increase of g_s away from the brane happens at larger and larger distances, and one remains at weak coupling as long as one focuses on suitable radius around the brane.

Generically, however, the fact that g_s develops a strongly varying non-trivial profile obscures an interpretation of the background in terms of perturbation theory. As just discussed,

¹This does not mean that the gauge theory on the 7-brane is trivial: The relevant frame is the Einstein frame, where the 7-brane gauge coupling is independent of g_s and given to leading order by the volume of the internal 4-cycle wrapped by the brane (for the case of compactification to 4 dimensions).

²In compactifications, while the profile of τ is set by the brane configurations, the integration constant τ_0 or overall scale λ remains a modulus of the low-energy effective theory.

in generic situations τ inevitably takes values of order 1 and larger in some regions of the compactification manifold. Besides, the expression for τ must receive strong stringy corrections in the vicinity of an orientifold-plane, which carries -4 units of C_8 -charge. A naive solution $\tau(z) \simeq -\frac{4}{2\pi i} \ln(z - z_0)$ which would be suggested by the above arguments is unphysical as the dilaton would become negative close to the O-plane.

An exception to the inevitable variation of τ in the presence of 7-branes and O-planes is the very special situation where all 7-brane charges cancel *locally* because a suitable number of D7-branes and O7-planes lie on top of each other. In this case, g_s is constant on the entire compactification manifold and in absence of stabilising effects such as fluxes it can freely be chosen to lie in the perturbative window $g_s \ll 1$ everywhere. In fact, one should reserve the term 'perturbative' for this non-generic configuration. This establishes that a proper treatment of backgrounds with varying axio-dilaton is not an option, but rather required for a satisfactory definition of generic configurations with 7-branes.

Monodromies and $SL(2, \mathbb{Z})$ invariance

A peculiar feature of the 7-brane backreaction is the appearance of monodromies. The monodromy associated with the simple solution (12) implies that as one encircles the position of the D7-brane in the z -plane, the axio-dilaton transforms as

$$\tau \rightarrow \tau + 1. \quad (14)$$

At first sight this might come as a shock as it seems to make a consistent interpretation of the background solution impossible. The *deus ex machina* approaching to our rescue is the fact that Type IIB is invariant under $SL(2, \mathbb{Z})$ transformations, of which (14) is the simplest example. As is most readily verified after transforming (5) into Einstein frame (see e.g. [4]), the classical Type IIB action enjoys the $SL(2, \mathbb{R})$ invariance

$$\tau \rightarrow \frac{a\tau + b}{c\tau + d}, \quad \begin{pmatrix} C_2 \\ B_2 \end{pmatrix} \rightarrow \begin{pmatrix} aC_2 + bB_2 \\ cC_2 + dB_2 \end{pmatrix} = M \begin{pmatrix} C_2 \\ B_2 \end{pmatrix}, \quad \det M = 1. \quad (15)$$

Note that C_4 is invariant under these transformations. This classical symmetry is broken at the non-perturbative level to $SL(2, \mathbb{Z})$. The reason is that $D(-1)$ instanton effects involve a factor $e^{2\pi i \tau}$. Invariance of such quantum effects under transformations of the type

$$M = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} : C_0 \rightarrow C_0 + b$$

restricts $b \in \mathbb{Z}$, thus reducing $SL(2, \mathbb{R})$ to $SL(2, \mathbb{Z})$.

Therefore as one encircles a D7-brane, the full background transforms by the $SL(2, \mathbb{Z})$ action $T = \begin{pmatrix} 1 & \\ 0 & 1 \end{pmatrix}$, and the monodromy action is merely a symmetry of the theory.

$[p, q]$ -branes

Once we have accepted that a consistent interpretation of D7-branes forces us to take the $SL(2, \mathbb{Z})$ symmetry at face value, we are led to more exotic objects than D7-branes which have no interpretation in terms of perturbation theory. The new objects we must include are $[p, q]$ -branes and corresponding $\binom{p}{q}$ strings [18]. To see this, recall from the Polyakov worldsheet action that the fundamental superstring is charged electrically under the NS-NS B_2 -field. Given the mixing of B_2 with the RR 2-form C_2 under general $SL(2, \mathbb{Z})$ transformations as in

(15) there must also exist an analogous object charged electrically under C_2 . This, of course, is nothing other than the D1-string, and its corresponding coupling to C_2 is the Chern-Simons action. It is therefore appropriate to combine the F1- and the D1-string into an $SL(2, \mathbb{Z})$ doublet: An F1-string is represented as the vector $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and a D1-string is a $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ object. A general $\begin{pmatrix} p \\ q \end{pmatrix}$ string carries p units of electric B_2 -charge and q units of electric C_2 -charge. Such objects exist as supersymmetric bound states for p, q co-prime [19]. Note that in perturbative Type IIB theory only $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ strings are present as the fundamental objects, while the D1-string enters as a solitonic, non-perturbative object.

In perturbative Type IIB theory, a D7-brane is by definition a hypersurface on which fundamental strings can end. This motivates the definition, in strongly-coupled Type IIB theory, of a $[p, q]$ -7-brane as the hypersurface on which $\begin{pmatrix} p \\ q \end{pmatrix}$ strings can end. Two $[p, q]$ -7-branes of different p, q are called mutually non-local.

A general $[p, q]$ -brane induces an $SL(2, \mathbb{Z})$ monodromy $M_{p,q}$ on the background fields as one encircles the location of the brane. This monodromy generalises the perturbative action (14) induced by $T = M_{1,0}$ and can be shown to take the form

$$M_{p,q} = g_{p,q} M_{1,0} g_{p,q}^{-1} = \begin{pmatrix} 1 - pq & p^2 \\ -q^2 & 1 + pq \end{pmatrix}. \quad (16)$$

Here $g_{p,q} = \begin{pmatrix} p & r \\ q & s \end{pmatrix}$ is the $SL(2, \mathbb{Z})$ matrix that transforms a $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ string into a $\begin{pmatrix} p \\ q \end{pmatrix}$ string.³ The only eigenvector of $M_{p,q}$ is a $\begin{pmatrix} p \\ q \end{pmatrix}$ -string itself, any other type of string gets transformed as transported around the location of the $[p, q]$ -7-brane.

Every $[p, q]$ -brane as such can be mapped into a $[1, 0]$ -brane, i.e. a conventional D7-brane, by an $SL(2, \mathbb{Z})$ transformation. Locally around each single 7-brane the geometry therefore is indistinguishable from the one backreacted by a D7-brane. However, in the presence of mutually non-local $[p, q]$ -branes in the above sense, new phenomena arise because the various branes cannot be *simultaneously* transformed into a D7-brane. And, most importantly, a consistent compactification necessarily includes 7-branes of different type.

The simplest example where this becomes apparent starts from a perturbative Type IIB orientifold on T^2 modded out by $\Omega(-1)^{F_L}\sigma$. The geometric orientifold action σ transforms the complex coordinate of the torus as $z \rightarrow -z$. Its four fixed points are the location of O7-branes. Local tadpole cancellation in perturbative models (in the sense introduced above) requires 4 D7-branes (plus their image branes) on top of each O7-plane, resulting in a famous $SO(8)^4$ gauge group. The $[p, q]$ -brane interpretation of this configuration was given by Sen [20], who showed that eight-dimensional Type II compactifications require a set of three different $[p, q]$ -branes with $[p, q]$ -labels

$$A : [1, 0], \quad B : [3, -1], \quad C : [1, -1]. \quad (17)$$

In this notation A-type branes correspond to perturbative D7-branes. In fact the branes B and C are chosen such that the combined monodromy $M_{BC} = M_{3,-1} M_{1,-1}$ acts on a $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ string by orientation reversal,

$$M_{3,-1} M_{1,-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = - \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (18)$$

as in the context of an orientifold theory. This identifies the BC-system as a perturbative O7-plane.

³Note that r and s are not uniquely determined. This ambiguity drops out in all physically relevant quantities.

In generic configurations, however, objects with general $SL(2, \mathbb{Z})$ monodromies will play a role, simply because the action of $M_{1,0}$ and M_{BC} does not generate the full $SL(2, \mathbb{Z})$. This raises a puzzle: What are the consistency conditions for theories with mutually non-local $[p, q]$ -branes? Is any configuration of coincident branes like the BC system allowed? Given the dyonic nature of $[p, q]$ -branes, perturbative methods are bound to fail. For compactifications to eight dimensions one might try to keep track of the $SL(2, \mathbb{Z})$ monodromies by hand and search for intrinsically consistent configurations, but already in six dimensions this becomes intractable.

Even more spectacularly, there exists a miraculous reformulation of the problem that allows us to almost blindly read off the consistent configurations of 7-branes. This formulation is the much sought-after F-theory.

Towards a geometric description

The crucial insight [5] that underlies the formulation of such a non-perturbative theory is the identification of the $SL(2, \mathbb{Z})$ symmetry of ten-dimensional Type IIB supergravity with the geometric $SL(2, \mathbb{Z})$ action on the complex structure of a two-torus T^2 . Inspired by the $SL(2, \mathbb{Z})$ transformation of τ one interprets the axio-dilaton as the complex structure of a fictitious elliptic curve.




Figure 1: $SL(2, \mathbb{Z})$ action $M_{1,0}$ on complex structure τ of a T^2 .

The variation of τ in presence of a set of D7-branes is therefore modelled as the variation of the complex structure of an elliptic curve transverse to the location of the 7-branes. Such a structure defines an elliptic fibration, and the non-triviality of the fibration is a measure for how strongly the axio-dilaton varies as a consequence of the backreaction of the branes.

The F-theory conjecture states that the physics of Type IIB orientifold compactifications with 7-branes on the complex n -fold B_n is encoded in the geometry of an $(n+1)$ -fold Y_{n+1} which is elliptically fibered over B_n . The elliptic fiber itself is not part of the physical spacetime but merely a book-keeping device that accounts for the variation of τ . In particular, at the location of 7-branes the axio-dilaton τ diverges as seen for the solution 12. If the complex structure of an elliptic curves diverges, this indicates the shrinking of a one-cycle and thus the degeneration of the elliptic curve. Thus the degeneration locus of the elliptic curve describes the presence of 7-branes.

In fact, duality with M-theory yields additional restrictions on the relevant elliptic fibrations. First, for $\mathcal{N} = 1$ supersymmetry to be conserved, the space Y_{n+1} has to be Calabi-Yau. Further, only the limit of vanishing elliptic fiber is to be considered. Both these facts will become apparent in the context of F/M-theory duality reviewed in the next section.

To return to the simple example alluded to before, understanding compactifications of Type IIB orientifolds to eight dimensions involves F-theory on a Calabi-Yau 2-fold Y_2 . In this case there is only one such Ricci flat 2-fold, the famous $K3$. The physically most interesting situation of course corresponds to F-theory on an elliptically fibered 4-fold Y_4 , which is supposed to capture Type IIB orientifolds compactified to four dimensions on the 3-fold B_3 .

2.2 F/M-theory duality and Calabi-Yau 4-folds

So far we have motivated F-theory as a clever way to think about Type IIB orientifolds with 7-branes. To determine the details of the effective action and of the physical degrees of freedom though, it is useful to approach F-theory via duality with eleven-dimensional M-theory, taken as its long-wavelength limit of eleven-dimensional supergravity coupled to M2/M5-branes. This duality also provides a more direct way to uncover the appearance of the elliptic curve in F-theory.

The starting point is therefore compactification of eleven-dimensional supergravity on $\mathbb{R}^{1,9} \times T^2$. Let τ be the complex structure of the torus $T^2 = S_A^1 \times S_B^1$. E.g. for the special case of a rectangular torus we have $\tau = i \frac{R_A}{R_B}$ in terms of the radii R_A and R_B of S_A^1 and S_B^1 . The M/F-theory duality consists in taking the limit of vanishing torus volume in the following two-step procedure [21]:

- The circle S_A^1 with radius R_A is identified as the M-theory circle in the reduction from M-theory to Type IIA theory in ten dimensions so that $g_{IIA} \simeq \frac{R_A}{\ell_s}$. The limit $R_A \rightarrow 0$ therefore corresponds to taking the perturbative IIA limit of M-theory, and we recover weakly coupled Type IIA theory on $\mathbb{R}^{1,9} \times S_B^1$.
- Now perform a T-duality along the remaining circle S_B^1 with radius R_B . This yields Type IIB theory on $\mathbb{R}^{1,9} \times \tilde{S}_B^1$ with dual radius $\tilde{R}_B = \frac{\ell_s^2}{R_B}$. The limit $R_B \rightarrow 0$ results in ten-dimensional Type IIB theory.

Application of the T-duality transformation identifies the IIB coupling as $g_{IIB} \simeq g_{IIA} \frac{\ell_s}{R_B} \simeq \frac{R_A}{R_B} \simeq \text{Im}(\tau)$. The last step applies to rectangular tori but has simple generalisations. A more precise analysis as worked out in detail e.g. in [12] allows one to trace back the Type IIB RR axion C_0 to the type IIA RR one-form C_1 . The latter in turn derives from the real part of the complex structure of the M-theory torus.

What we have sketched is a description of Type IIB theory in ten dimensions in terms of M-theory on $\mathbb{R}^{1,9} \times T^2$, where the IIB axio-dilaton $\tau = C_0 + \frac{i}{g_s}$ is identified with the complex structure of the M-theory T^2 . F-theory on T^2 can therefore be defined as the ten-dimensional IIB theory which is dual, in the above sense, to M-theory on $\mathbb{R}^{1,9} \times T^2$. Note that for the duality to work the volume of the M-theory T^2 has to vanish. This is the physical reason why only the complex structure τ , but not the volume of the elliptic curve appears as a physical field in Type IIB/F-theory.

The above logic extends adiabatically to the more general case of a non-trivial elliptic fibration rather than a direct product $\mathbb{R}^{1,9} \times T^2$. The physically most interesting situation is of course that of M-theory compactification on $\mathbb{R}^{1,2} \times Y_4$, where $Y_4 : T^2 \rightarrow B_3$ is a complex 4-fold elliptically fibered over a 3-complex dimensional base B_3 . If the 4-fold Y_4 is Calabi-Yau this yields a three-dimensional effective theory with four supercharges [22–24]. In the limit of vanishing fiber volume the fourth dimension grows large and this setup is dual to the four-dimensional effective theory obtained by Type IIB compactification on B_3 , or, by definition,

F-theory on Y_4 . The four supercharges lead to an $\mathcal{N} = 1$ effective theory in four dimensions. Thus the Calabi-Yau property of Y_4 is an obvious requirement from the M-theory perspective.

Kaluza-Klein reduction of the M-theory 3-form allows one to recover also the higher Type IIB RR-forms. We denote by α and β the periodic coordinates along the M-theory torus $T^2 = S_A^1 \times S_B^2$ and focus on a generic non-singular fiber for the time being. Reduction of C_3 yields

$$C_3 = \tilde{C}_3 + B_2 \wedge d\alpha + C_2 \wedge d\beta + B_1 \wedge d\alpha \wedge d\beta. \quad (19)$$

After T-duality along β and decompactification to four dimensions, \tilde{C}_3 furnishes the degrees of freedom of the RR 4-form $C_4 = \tilde{C}_3 \wedge d\beta$, while B_1 becomes part of the four-dimensional metric $g_{i\beta}$. B_2 and C_2 , on the other hand, translate into the NS-NS and R-R 2-forms in Type IIB. Their transformation properties as a doublet under the $SL(2, \mathbb{Z})$ group derive immediately from the geometric $SL(2, \mathbb{Z})$ transformation of the A and B cycle of the M-theory torus.

It is beyond the scope of these lectures to enter a detailed discussion of the F-theory effective action. An in-depth derivation of the four-dimensional F-theory effective action via M/F-theory duality can be found in [25] (see also [12]), which we recommend for more details. An important ingredient in Kaluza-Klein reduction, e.g. to determine the precise massless supergravity spectrum, are the topological properties of Calabi-Yau 4-folds. Many aspects of the geometry and topology of (elliptic) Calabi-Yau 4-folds can be found e.g. in [26]. Suffice it here to recall for completeness that the Hodge diagram of Calabi-Yau 4-folds is characterised by three independent Hodge numbers $h^{1,1}$, $h^{2,1}$, $h^{3,1}$, to which $h^{2,2}$ is related via

$$h^{2,2} = 2(22 + 2h^{1,1} + 2h^{3,1} - h^{2,1}). \quad (20)$$

An important quantity for model building, which will appear prominently in the sequel, is the Euler characteristic

$$\chi(Y_4) = \int_{Y_4} c_4(T_{Y_4}) = 6(8 + h^{1,1} + h^{3,1} - h^{2,1}). \quad (21)$$

2.3 The geometry of elliptic fibrations

The above considerations show that understanding the non-perturbative F-theoretic region of the string landscape requires familiarity with the concept of elliptic fibrations, to which we now turn in some detail. The problem splits into two parts: We first need to understand elliptic curves as such, and then find a way to describe their fibration over the base B_n such as to form an elliptically fibered Calabi-Yau manifold Y_{n+1} ,

$$\pi : Y_{n+1} \rightarrow B_n. \quad (22)$$

Elliptic curves as $\mathbb{P}_{2,3,1}[6]$

There are different ways to describe an elliptic curve, the simplest being as a hypersurface or, more generally, as a complete intersection of some weighted projective space. As an example consider the weighted projective space $\mathbb{P}_{2,3,1}$ spanned by the homogeneous coordinates $(x, y, z) \simeq (\lambda^2 x, \lambda^3 y, \lambda z)$, where (x, y, z) are complex coordinates and $\lambda \in \mathbb{C}^*$. Due to the scaling relation this space is 2-complex dimensional. An elliptic curve is a flat 1-complex dimensional space, i.e. a Calabi-Yau 1-fold. It can thus be described by the hypersurface cut out by the vanishing locus of a homogenous polynomial in (x, y, z) of degree 6 under

rescaling by λ . Here we used the general fact that a hypersurface in weighted projective space is Ricci-flat whenever the degree of its defining polynomial equals the sum of degrees of the homogenous coordinates. A degree-six polynomial in $\mathbb{P}_{2,3,1}$ defines the space $\mathbb{P}_{2,3,1}[6]$. One can show that after suitable coordinate redefinitions such a polynomial can always be brought into the so-called Weierstrass form

$$P_W = y^2 - x^3 - fxz^4 - gz^6 = 0, \quad (23)$$

where $f, g \in \mathbb{C}$ specify the shape of the elliptic curve as detailed below. Note at this stage that alternative representations of elliptic curves along these lines are the hypersurfaces $\mathbb{P}_{1,1,2}[4]$ or $\mathbb{P}_{1,1,1}[3]$.

As argued at the end of section 2.1 the physics of the 7-branes is encoded in the degenerations of the elliptic fiber. A general mathematical fact is that a hypersurface described by the equation $P_W = 0$ becomes singular whenever

$$P_W = 0 \quad \text{and} \quad dP_W = 0, \quad (24)$$

where the last condition indicates a degeneration of the tangent space. It is easy to work out these conditions for the Weierstrass curve (23). First use the scaling relations to go to inhomogeneous coordinates by setting $z = 1$. Then the above conditions are equivalent to

$$\begin{aligned} y &= 0, \\ -x^3 - fx - g &= (x - a_1)(x - a_2)(x - a_3) = 0, \\ (x - a_1)(x - a_2) + (x - a_2)(x - a_3) + (x - a_1)(x - a_3) &= 0, \end{aligned} \quad (25)$$

where a_1, a_2, a_3 are the three complex roots of the cubic polynomial $-x^3 - fx - g$. Clearly the above requires that two or more of these roots coincide. As always, the structure of the roots of this cubic polynomial is encoded in the discriminant Δ , which vanishes if and only if at least two roots coincide. For the Weierstrass model the discriminant of the cubic $-x^3 - fx - g$ takes the simple form

$$\Delta = 27g^2 + 4f^3, \quad (26)$$

which shows that the structure of the elliptic curve indeed depends on the parameters f and g .

In fact, the dependence of the complex structure τ on f and g can be made more precise. Since τ undergoes $SL(2, \mathbb{Z})$ monodromies it is more convenient to work with the $SL(2, \mathbb{Z})$ invariant Jacobi j -function, which yields an isomorphism from the fundamental domain of $SL(2, \mathbb{Z})$ to the Riemann sphere. Instead of introducing the formal definition of $j(z)$ in terms of theta-functions it suffices here to note the expansion

$$j(z) = \exp(-2\pi iz) + 744 + (\exp(2\pi iz)) + \dots \quad (27)$$

A classical mathematical result is that the complex structure τ of a Weierstrass elliptic curve is determined by f, g via the relation

$$j(\tau) = \frac{4(24f)^3}{\Delta}. \quad (28)$$

From elliptic curves to elliptic fibrations

Armed with this representation of a single elliptic curve, we can proceed to elliptic fibrations. Suppose we have some n -complex dimensional manifold B_n , covered by local coordinates u_i . Then a fibration of the Weierstrass curve (23) over B_n is obtained by promoting the constants f and g in (23) to suitable polynomials in the coordinates u_i of B_n , $f = f(u_i)$, $g = g(u_i)$. This defines the corresponding Weierstrass model, also known as an E_8 -fibration. Note that in principle one can also consider fibrations of the elliptic curves $\mathbb{P}_{1,1,2}[4]$ or $\mathbb{P}_{1,1,1}[3]$ (called E_7 - and, respectively, E_6 -fibrations) and their generalisations. The crucial point, however, is the following mathematical theorem (see e.g. [27]): Every elliptic fibration with a section can be represented by a Weierstrass model defined in terms of the equation (23) with varying f and g . All other representations such as the E_7 - and E_6 -fibrations are birationally equivalent (equivalent up to a flop transition).

Via the relation (28) the complex structure τ is now dependent on the base coordinates u_i . In particular, the elliptic fiber degenerates on a codimension-one sublocus on B_n determined by the vanishing of the likewise u_i -dependent discriminant $\Delta = 27g^2 + 4f^3$. In view of what we said before, this vanishing locus must be interpreted as a divisor wrapped by a stack of 7-branes. Indeed, suppose Δ vanishes to order N along the divisor S on B_n defined by $w = 0$. Then eqns. (27) and (28) imply that in the directions normal to this divisor

$$\tau \simeq \frac{N}{2\pi i} \ln(w) \quad (29)$$

up to regular terms, which is exactly the behaviour expected for N coincident 7-branes. The connection between this geometric description of 7-branes as the degeneration locus of the elliptic fibration and the concept of $[p, q]$ -7-branes comes about as follows: A single $[p, q]$ -7-brane is the locus on the base B_n along which a (p, q) -cycle on the elliptic fiber degenerates. This can be shown by matching the geometric $SL(2, \mathbb{Z})$ monodromies with the action (16) of $M_{p,q}$. Locally around each 7-brane we are free to declare the degenerate 1-cycle in the fiber to correspond to the $(1, 0)$ fiber by a choice of basis for $H_2(T^2, \mathbb{Z})$. This corresponds to the assertion that locally each brane looks like a D7-brane by an $SL(2, \mathbb{Z})$ transformation. In IIB theory, a stack of N coincident 7-branes in generic position with respect to the O7-plane gives rise to gauge group $U(N)$. In the F-theory picture the gauge group along a divisor is described by the details of the degeneration of the elliptic fiber as we will see below.

First, however, we need to discuss the Calabi-Yau property in greater detail. The first Chern class of an elliptic fibration Y_{n+1} is related to the first Chern class of the base space B_n and the degeneration locus of the fibration as [28]

$$c_1(T_{Y_{n+1}}) \simeq \pi^* \left(c_1(T_{B_n}) - \sum_i \frac{\delta_i}{12} [\Gamma_i] \right), \quad \delta_i = \mathcal{O}(\Delta)|_{\Gamma_i}. \quad (30)$$

Here

$$\pi : Y_{n+1} \rightarrow B_n \quad (31)$$

denotes the projection from the fibration to the base, and its pullback π^* maps $H^2(B_n, \mathbb{Z}) \rightarrow H^2(Y_{n+1}, \mathbb{Z})$. Furthermore, the discriminant is supposed to vanish to order δ_i along the divisors Γ_i on the base, with dual 2-form $[\Gamma_i]$. Strictly speaking this formula is correct only for elliptic $K3$ -manifolds, but the extra complications due to higher co-dimension degenerations on the base do not affect the present argument [29]. We see that the Calabi-Yau property of Y_{n+1}

forces B_n to acquire positive curvature such as to compensate for the degenerations of the fiber. In particular, B_n itself is not Calabi-Yau. Since the discriminant locus describes the positions of the 7-branes, we find that the total amount of 7-branes leading to a consistent compactification is thus constrained by the curvature of the base space via

$$\sum_i \delta_i [\Gamma_i] = 12 c_1(T_{B_n}). \quad (32)$$

This equation is reminiscent of the 7-brane tadpole cancellation condition in perturbative Type IIB theory

$$\sum_i N_i [\Gamma_i] = 4[O_7]. \quad (33)$$

This demonstrates the very different nature of string model building in perturbative Type IIB orientifolds and F-theory in that the 7-brane consistency conditions are *automatically* incorporated in a well-defined geometry.

From a mathematical perspective, the relation (32) constrains the degree of the polynomials f and g that appear in the Weierstrass model as follows: The class of the left-hand side is just the class of the full discriminant Δ , and (32) tells us that $[\Delta] = 12c_1(T_{B_n})$. In view of $\Delta = 27g^2 + 4f^3$ this determines the class of f, g . In general f and g are not globally defined functions of the base space, but rather sections of a line bundle \mathcal{L} . In terms of the canonical bundle K_{B_n} , which has first Chern class $c_1(K_{B_n}) = -c_1(T_{B_n})$, it follows that f and g are sections of $K_{B_n}^{-4}$ and $K_{B_n}^{-6}$, respectively. Homogeneity of the defining Weierstrass polynomial then forces also x and y to transform as sections of the base. In all, we find

$$x \in H^0(B_n, K_{B_n}^{-2}), \quad y \in H^0(B_n, K_{B_n}^{-3}), \quad z \in H^0(B_n, \mathcal{O}), \quad (34)$$

$$f \in H^0(B_n, K_{B_n}^{-4}), \quad g \in H^0(B_n, K_{B_n}^{-6}). \quad (35)$$

Example: F-theory on $K3$

As a simple example consider $K3$ on the locus of its moduli space where it arises as an elliptic fibration over $B_1 = \mathbb{P}^1$. In terms of the normalised volume form t of \mathbb{P}^1 , the first Chern class of $K_{\mathbb{P}^1}$ is simply $c_1(K_{\mathbb{P}^1}) = -2t$. This follows from the Hirzebruch-Riemann-Roch theorem, whereby

$$\chi = 2 - 2g = \int_{\mathbb{P}^1} c_1(T_{\mathbb{P}^1}). \quad (36)$$

The standard notation is of course $K_{\mathbb{P}^1} = \mathcal{O}(-2)$. More generally a section of the line bundle $\mathcal{O}(-n)$ is a homogenous polynomial of degree n in the homogeneous coordinates $[u_0, u_1]$ of \mathbb{P}^1 . Together with the original scaling relation of the Weierstrass model the elliptically fibered K_3 is spanned by the coordinates

$$(u_0, u_1; x, y, z) \simeq (u_0, u_1; \lambda^2 x, \lambda^3 y, \lambda z) \simeq (\mu u_0, \mu u_1; \mu^4 x, \mu^6 y, z). \quad (37)$$

Famously, the discriminant Δ is thus a polynomial of degree 24 on the base, with 24 zeroes known as the position of the 24 7-branes of compactification of IIB theory on \mathbb{P}^1 . Not all of these 24 branes describe perturbative D7-branes. As discussed already around eqn. (18), in the strict perturbative limit the tadpoles are cancelled locally by grouping the 24 branes into four stacks of six branes. Each stack corresponds to 4 D7-branes, i.e. A-type branes, on top

of a system of BC branes. The latter furnish the F-theoretic description of the IIB O7-plane. This can be derived via a monodromy analysis [20].

At a more advanced level, the F-theory solution teaches us several exciting lessons: In [20] Sen showed how the O7-brane splits as one removes one or more A-branes from the A^4BC system. Recall from the discussion after (12) that in such a situation where tadpoles are no longer cancelled locally the naive supergravity solution must break down close to the O7-plane. In F-theory, the geometry adjusts itself via this non-perturbative brane split so as to render the configuration self-consistent. This not only reveals that the O7-plane is a dynamical object per se; it also furnishes a highly non-trivial check that F-theory correctly captures the non-perturbative degrees of the theory. Indeed it is quite remarkable that a comparatively straightforward analysis of the geometry of $K3$ alone dictates all consistent configurations $[p, q]$ -brane configurations. Along these lines, the authors of [30] found more general solutions with constant, but non-perturbatively large axio-dilaton for which the 24 branes group e.g. into three bunches of the type A^5BC^2 and more. This shows that the number of branes of a given $[p, q]$ -type on a given compactification space changes from configuration to configuration due to the appearance of monodromies as we start moving the branes around.

2.4 Sen's orientifold limit

After this first encounter with the geometric description of F-theory, let us go back and analyse more closely how to recover the weakly coupled Type IIB orientifold picture. This is accomplished by a procedure due to Sen [31]. Recall from the discussion after eqn. (12) that in the IIB limit one considers the 7-branes as probe objects and takes the axio-dilaton as non-varying and perturbatively small. We already noted that strictly speaking this can only be realised if all 7-brane tadpoles are cancelled locally by placing a suitable amount of 7-branes on top of the orientifold planes. More generally, there can however exist a limit in which one can *approximately* take τ small and non-varying even though not all tadpoles are cancelled locally. As argued after eqn. (13) this corresponds to the limit $\lambda \rightarrow \infty$. Since the profile of τ is encoded in the Jacobi function (28) a constant τ can be achieved as long as $j(\tau)$ is non-varying over the base B_3 of the F-theory 4-fold. To arrange for this one Sen makes the general ansatz for the sections f and g

$$f = -3h^2 + \epsilon\eta, \quad g = -2h^3 + \epsilon h\eta - \frac{\epsilon^2}{12}\chi, \quad (38)$$

where ϵ is an arbitrary constant. Sen's orientifold limit corresponds to the limit $\epsilon \rightarrow 0$ with h , η and χ generically non-vanishing. In this case one finds that indeed the string coupling becomes arbitrarily weak everywhere except at the location $h = 0$, where $g_s \rightarrow \infty$. It can thus be considered as a perturbative parameter of the theory. The discriminant locus factorises to leading order in ϵ as

$$\Delta_\epsilon = -9\epsilon^2 h^2 (\eta^2 - h\chi) + \mathcal{O}(\epsilon^3) \quad (39)$$

so that the fibration degenerates at $h = 0$ and at $\eta^2 - h\chi$. The latter may factorise further depending on the particular form of the polynomials η and χ . By closer inspection of the monodromies around these loci it can be shown that $h = 0$ corresponds to the location of an O7-plane, while the latter locus is that of an ordinary D7-brane. The inclusion of higher order terms in ϵ , on the other hand, destroys this factorisation of Δ into orientifold and brane piece. This process can be thought of as recombination of the O7-plane and the D7-brane.

Since in perturbation theory there are no light strings stretching between the O7-plane and a D7-brane, this recombination is truly non-perturbative. The involved recombination modes are expected to be $\binom{p}{q}$ strings.

The Type IIB theory is defined on the Calabi-Yau 3-fold X which is a double cover of the base B branched over $h = 0$. Concretely, suppose the base B_3 is embedded into some projective space with projective coordinates u_i . Then one can define X as the complete intersection in the projective space spanned by the coordinates u_i and ξ which is defined by the original hypersurface polynomial and the additional constraint

$$\xi^2 - h(u_i) = 0. \quad (40)$$

Conversely, given a Type IIB orientifold on X together with a \mathbb{Z}_2 action σ , one can construct the F-theory uplift by reversing Sen's limit above. This has been worked out and applied in the more recent literature [32–34].

Note that the form of the 7-brane locus $\eta^2 - \xi^2 \chi = 0$ (where we replace h in $\eta^2 - h\chi = 0$ by ξ^2 according to (40)) shows that for generic polynomials η, χ the defining equation for a single D7-brane configuration with a well-defined F-theory uplift is non-generic [17,35]. Brane-image brane configurations require a specialisation of these polynomials, e.g. by factorisation of χ . The relation of such non-generic brane splits to elliptic fibrations of E_7 - and E_6 -type has been investigated in [36,37].

Finally we would like to point out that the limit $\epsilon \rightarrow 0$ with all other polynomials generic is not the only way to make τ non-varying along B_3 . There exist in addition the two possibilities of taking either $f \equiv 0$ or $g \equiv 0$ [30]. In these cases, τ is constant and fixed at non-perturbatively large values, in agreement with the appearance of exceptional gauge groups. These configurations correspond to generalisations of the \mathbb{Z}_2 orientifold action of the strongly coupled Type IIB string to \mathbb{Z}_n actions.

2.5 Gauge symmetry from degenerations

From a physics perspective the most essential data of an elliptic fibration are the locus and the type of fiber degenerations because these allow us to deduce the nature of the 7-branes wrapping the corresponding divisor. The different ways how the complex structure of the elliptic fiber can degenerate have been classified by Kodaira [28] for the case of a Weierstrass model of $K3$. The relevant criterion to be imposed here is that the singularities can be resolved without destroying the Calabi-Yau property. These results have subsequently been generalised to higher Calabi-Yau spaces (see [38]⁴ and references therein for a physics discussion).

Consider a Weierstrass model Y_{n+1} . One distinguishes degenerations that merely render the fibration singular without inducing an actual singularity in Y_{n+1} versus loci where the Calabi-Yau $(n+1)$ -fold becomes singular itself. The simplest kind of degeneration of the first type corresponds to a so-called I_1 singularity, which arises in the fiber over just a single 7-brane. For a $[p, q]$ -7-brane the (p, q) 1-cycle of the elliptic fiber pinches off and the elliptic fiber forms a sausage-type \mathbb{P}^1 whose north and south pole touch each other. Following Kodaira, I_1 singularities arise in a Weierstrass model whenever Δ develops a simple zero while f and g are non-zero. In this case two of the roots a_i in the cubic (25) coincide. From eqs. (27), (28) the simple zero in Δ leads to the logarithmic profile (12) of τ characteristic of a single 7-brane.

⁴ Note that this reference is devoted to an analysis of elliptic Calabi-Yau 3-folds. For higher dimensional manifolds extra effects due to a more complicated monodromy structure can occur, and as of this writing no complete classification exists in the literature. Also, the results of ref. [38] apply to situations where the discriminant locus itself is non-singular as a divisor of Y_3 .

When Δ vanishes to higher order, generically also Y_{n+1} develops an actual singularity as a manifold. For the simplest case of $Y_2 = K_3$ the possible singularities that can occur are precisely the ones listed in the famous ADE classification [28]. For higher dimensional Y_{n+1} with $n > 2$ also the action of monodromies along the brane have to be taken into account. This can create more general singularities including non-simply laced examples of B and C type and such exceptional cases as G_2 or F_4 .

Let us focus for simplicity on Calabi-Yau 4-folds Y_4 . One intuitive way to understand the connection between a singularity over the divisor $S \subset B_3$ and a Lie algebra G is as follows: The singularity in Y_4 can be resolved by standard methods of algebraic geometry. For our purposes it suffices to consider cases that allow for a so-called split simultaneous resolution. As described e.g. in [39] this means that there exists a new, non-singular Calabi-Yau \bar{Y}_4 defined by replacing the singular fiber over S by a tree of \mathbb{P}^1 s which we call in the sequel Γ_i , $i = 1, \dots, \text{rk}(G)$. Note that in this process $h^{1,1}$ increases by $\text{rk}(G)$. These \mathbb{P}^1 s have the property that they intersect one another like the simple roots of the Lie algebra G in the following sense: The resolution 2-cycles Γ_i , are fibered over the 7-brane $S \subset B_3$. Denote the resulting divisors of \bar{Y}_4 by

$$\hat{D}_i : \Gamma_i \rightarrow S. \quad (41)$$

By construction these are \mathbb{P}^1 -fibrations. Consider furthermore the linear combination $\hat{D}_0 = \hat{S} - \sum_i a_i D_i$, where \hat{S} is the elliptic fibration over S and a_i denote the Dynkin labels for the Lie algebra G associated with the singularity over S . Then \hat{D}_0, \hat{D}_i encode the extended Dynkin diagram of G in the sense that their Poincaré dual 2-forms $[\hat{D}_i] \in H^2(\bar{Y}_4)$ satisfy

$$\int_{\bar{Y}_4} [\hat{D}_i] \wedge [\hat{D}_j] \wedge \tilde{\omega} = -C_{ij} \int_S \tilde{\omega}, \quad i, j, = 0, 1, \dots, \text{rk}(G) \quad (42)$$

for $\tilde{\omega} \in H^2(B_3)$ and C_{ij} the Cartan matrix of G . Conversely, the singular F-theory limit can be understood as the limit in which the Γ_i shrink to zero volume. The singularity is the result of a collision of $\text{rk}(G)$ zero size \mathbb{P}^1 s.

The appearance of the corresponding gauge groups along the 7-brane wrapping the singular locus can be understood by F/M-theory duality. As sketched in (19), Kaluza-Klein reduction of C_3 along non-singular fibers results in the Type IIB closed string 2-forms B_2 and C_2 . As we have just seen, in the fiber above the discriminant locus extra two-cycles appear along which C_3 can be reduced. For simplicity consider an A_{n-1} singularity associated with gauge group $SU(n)$. As argued above, the resolution \mathbb{P}^1 s $\Gamma_i, i = 1, \dots, (n-1)$ correspond to the nodes of the Dynkin diagram of A_{n-1} . Massless vector states arise from two sources:

- The M-theory 3-form C_3 can be reduced along Γ_i , leading to 1-forms $A_i = \int_{\Gamma_i} C_3$ along the 7-brane. These are interpreted as the gauge potentials for the abelian group factors in the Cartan subalgebra of $SU(n)$.
- The M-theory M2-brane can wrap chains of 2-cycles $S_{ij} = \Gamma_i \cup \Gamma_{i+1} \cup \dots \cup \Gamma_j$ for any $i \leq j$. Taking into account the two possible orientations this gives rise to $n^2 - n$ states along the brane. In the singular limit of vanishing volume of the Γ_i these states become massless and lead to the W -bosons of $SU(n)$.

In all this yields the $n^2 - 1$ generators of $SU(n)$. This logic can be generalised to other simple groups.

In addition there can be extra abelian gauge factors which do not arise as the Cartan generators of a non-abelian gauge group and which are strictly speaking not localised along individual divisors. An unambiguous way to determine the total rank of the gauge group of a singular Weierstrass model is to note that all $U(1)$ generators derive from reduction of C_3 as $C_3 = A_i \wedge \omega_i + \dots$, where the two-forms ω_i have one leg along the T^2 fiber and one leg along the base [29, 40]. Therefore

$$r = h^{1,1}(\overline{Y}_{n+1}) - h^{1,1}(B_n) - 1 \quad (43)$$

gives the total rank of the F-theory gauge group including possible abelian gauge factors.⁵ The physics of these extra $U(1)$ s in compactifications to four dimensions has been discussed in [37].

Let us stress that for model purposes it is essential to have control over the resolution \overline{Y}_4 of the singular fibration Y_4 , e.g. to determine the complete rank of the gauge group or in the context of 3-brane tadpole cancellation, see section 3.2. In fact, for the class of Calabi-Yau 4-folds constructed as hypersurfaces or complete intersections of toric spaces powerful tools have been developed to perform this resolution in terms of the divisors \hat{D}_i explicitly [41, 42]. These have recently been exploited in the context of F-theory model building in [37, 43–45].

To conclude this section, we point out that the appearance of exceptional and more general gauge groups has a beautiful interpretation in terms of $\binom{p}{q}$ -strings [46, 47]. This can be understood already for F-theory on $K3$, where the possible gauge groups are exhausted by the ADE classification. By definition, only $\binom{p}{q}$ -strings can end on $[p, q]$ -7-branes. Due to the appearance of monodromies, however, $\binom{p}{q}$ -strings stretched between two branes along different paths can lead to inequivalent states. A mild version of this phenomenon was encountered already in the perturbative limit. A $\binom{1}{0}$ -string encircling a BC system gets orientation reversed, see eqn. (18). The resulting unoriented strings correspond to strings between a stack of branes and their orientifold images. More generally, one can classify allowed paths that lead to BPS-strings and determine their interactions via consistent splitting and joining. Suffice it here to highlight from [46] that the various $\binom{p}{q}$ -strings occurring for the $[p, q]$ -7-brane system $A^n BC^2$ indeed span the adjoint of E_{5+n} , $n = 1, 2, 3$, in agreement with the monodromies for the generalised orientifolds of [30]. The underlying reason for this new richness is that $\binom{p}{q}$ -strings of different types can form string junctions with more than just two endpoints. This overcomes the limitation to two-index representations as carried by the Chan-Paton factors of perturbative $\binom{1}{0}$ -strings. The latter allow only for the construction of $U(N)$, $Sp(2N)$ and $SO(N)$ groups.

3 Technology for F-theory compactifications

3.1 Tate models

Given the importance of the singularity structure for an analysis of the F-theory landscape, we find it useful to present the methods to detect the singularities in slightly greater detail. There exists an algorithm due to Tate that allows one to read off the singularity structure of an elliptic fibration, say $Y_4 : T^2 \rightarrow B_3$. Tate's formalism consists in a local coordinate redefinition that brings the Weierstrass constraint $P_W = 0$ in (23) into the Tate form

$$P_W = x^3 - y^2 + x y z a_1 + x^2 z^2 a_2 + y z^3 a_3 + x z^4 a_4 + z^6 a_6 = 0. \quad (44)$$

⁵In addition, there are $h^{2,1}(B_n)$ bulk $U(1)$ fields, corresponding to reduction of the RR 4-form C_4 along 3-cycles on B_n .

For a general Weierstrass model this can be achieved *locally* via the so-called Tate algorithm described in detail in [38]. In the sequel we will only be working with the inhomogeneous Tate form by setting $z = 1$. The $a_n(u_i)$ depend on the complex coordinates u_i of the base B . They encode the discriminant locus of the elliptic fibration. To recover the Weierstrass model one first introduces the combinations

$$\beta_2 = a_1^2 + 4a_2, \quad \beta_4 = a_1a_3 + 2a_4, \quad \beta_6 = a_3^2 + 4a_6. \quad (45)$$

One can check by completing the square and the cube in (44) that the Weierstrass sections f and g are related to these via

$$f = -\frac{1}{48}(\beta_2^2 - 24\beta_4), \quad g = -\frac{1}{864}(-\beta_2^3 + 36\beta_2\beta_4 - 216\beta_6). \quad (46)$$

The discriminant $\Delta = 27g^2 + 4f^3$ can then be expressed as

$$\Delta = -\frac{1}{4}\beta_2^2(\beta_2\beta_6 - \beta_4^2) - 8\beta_4^3 - 27\beta_6^2 + 9\beta_2\beta_4\beta_6. \quad (47)$$

In general, the discriminant Δ will factorize with each factor describing the location of a 7-brane on a divisor S_i in B_3 . The precise group is encoded by the vanishing degree of the a_i and Δ . This has been classified in Table 2 of ref. [38], which we are reproducing (in the form given in [33]) for convenience of the reader in table 1.⁶

For example, an $SU(5)$ gauge group along the divisor

$$S : \quad w = 0 \quad (48)$$

corresponds to

$$a_1 = \mathfrak{b}_5, \quad a_2 = \mathfrak{b}_4w, \quad a_3 = \mathfrak{b}_3w^2, \quad a_4 = \mathfrak{b}_2w^3, \quad a_6 = \mathfrak{b}_0w^5, \quad (49)$$

where the \mathfrak{b}_i generically depend on all coordinates u_i of the base B but do not contain an overall factor of w . It is straightforward to evaluate the discriminant Δ of the elliptic fibration in terms of the new sections \mathfrak{b}_i as

$$\Delta = -w^5 (\mathfrak{b}_5^4 P + w\mathfrak{b}_5^2(8\mathfrak{b}_4 P + \mathfrak{b}_5 R) + w^2(16\mathfrak{b}_3^2\mathfrak{b}_4^2 + \mathfrak{b}_5 Q) + \mathcal{O}(w^3)) \quad (50)$$

with

$$P = \mathfrak{b}_3^2\mathfrak{b}_4 - \mathfrak{b}_2\mathfrak{b}_3\mathfrak{b}_5 + \mathfrak{b}_0\mathfrak{b}_5^2, \quad R = 4\mathfrak{b}_0\mathfrak{b}_4\mathfrak{b}_5 - \mathfrak{b}_3^3 - \mathfrak{b}_2^2\mathfrak{b}_5. \quad (51)$$

Generically the expression in brackets in (50), denoted as S_1 , does not factorize further and thus constitutes the single-component locus of an I_1 singularity. Cohomologically, one thus finds that the class $[\Delta]$ splits as $[\Delta] = 5[S] + [S_1]$.

As stressed in section 2.3, the most general elliptic fibration with a section can always be written as a Weierstrass model. A special subclass of such models, however, can be brought into the Tate form not just locally around the discriminant locus, but globally. I.e. for those models the defining constraint is (44) in terms of globally defined sections $a_i \in H^0(B, K_B^{-i})$. Clearly each such global Tate model defines also a Weierstrass model in that each set of a_i defines the Weierstrass sections f and g as in (46). The converse, however, is not true globally as the transformation from f and g to a_i involves branch cuts.

⁶Note that this table was derived for the case of elliptic Calabi-Yau 3-folds, see footnote 4.

sing. type	discr. deg(Δ)	gauge enhancement		coefficient vanishing degrees				
		type	group	a_1	a_2	a_3	a_4	a_6
I_0	0	—	—	0	0	0	0	0
I_1	1	—	—	0	0	1	1	1
I_2	2	A_1	$SU(2)$	0	0	1	1	2
I_3^{ns}	3		[unconv.]	0	0	2	2	3
I_3^{s}	3		[unconv.]	0	1	1	2	3
I_{2k}^{ns}	$2k$	C_{2k}	$SP(2k)$	0	0	k	k	$2k$
I_{2k}^{s}	$2k$	A_{2k-1}	$SU(2k)$	0	1	k	k	$2k$
I_{2k+1}^{ns}	$2k+1$		[unconv.]	0	0	$k+1$	$k+1$	$2k+1$
I_{2k+1}^{s}	$2k+1$	A_{2k}	$SU(2k+1)$	0	1	k	$k+1$	$2k+1$
II	2	—	—	1	1	1	1	1
III	3	A_1	$SU(2)$	1	1	1	1	2
IV^{ns}	4		[unconv.]	1	1	1	2	2
IV^{s}	4	A_2	$SU(3)$	1	1	1	2	3
$I_0^{*\text{ns}}$	6	G_2	G_2	1	1	2	2	3
$I_0^{*\text{ss}}$	6	B_3	$SO(7)$	1	1	2	2	4
$I_0^{*\text{s}}$	6	D_4	$SO(8)$	1	1	2	2	4
$I_1^{*\text{ns}}$	7	B_4	$SO(9)$	1	1	2	3	4
$I_1^{*\text{s}}$	7	D_5	$SO(10)$	1	1	2	3	5
$I_2^{*\text{ns}}$	8	B_5	$SO(11)$	1	1	3	3	5
$I_2^{*\text{s}}$	8	D_6	$SO(12)$	1	1	3	3	5
$I_{2k-3}^{*\text{ns}}$	$2k+3$	B_{2k}	$SO(4k+1)$	1	1	k	$k+1$	$2k$
$I_{2k-3}^{*\text{s}}$	$2k+3$	D_{2k+1}	$SO(4k+2)$	1	1	k	$k+1$	$2k+1$
$I_{2k-2}^{*\text{ns}}$	$2k+4$	B_{2k+1}	$SO(4k+3)$	1	1	$k+1$	$k+1$	$2k+1$
$I_{2k-2}^{*\text{s}}$	$2k+4$	D_{2k+2}	$SO(4k+4)$	1	1	$k+1$	$k+1$	$2k+1$
$IV^{*\text{ns}}$	8	F_4	F_4	1	2	2	3	4
$IV^{*\text{s}}$	8	E_6	E_6	1	2	2	3	5
III*	9	E_7	E_7	1	2	3	3	5
II*	10	E_8	E_8	1	2	3	4	5
non-min	12	—	—	1	2	3	4	6

Table 1: Refined Kodaira classification resulting from Tate’s algorithm. In order to distinguish the “semi-split” case $I_{2k}^{*\text{ss}}$ from the “split” case I_{2k}^{s} one has to work out a further factorization condition, see §3.1 of [38].

Models which are globally of the Tate form are very convenient because one can immediately read off the gauge group from table 1 without going through the algorithm of [38] to bring the Weierstrass polynomial locally into the form (44). It is important to keep in mind, though, that they do not give rise to the most general singularity structure. In particular, there is a sense in which such models are based on an underlying gauge group E_8 . Consider the situation that the gauge group G realised along the divisor S is contained within E_8 . Then in the global Tate model the dynamics of the gauge sector can be understood via breaking an

original E_8 symmetry down to the commutant of a complimentary gauge group H such that $E_8 \rightarrow G \times H$. This can be seen by organising the sections of the Tate model as a sum of two terms, the first of which corresponds to singularity enhancement E_8 , while the second encodes an H -bundle responsible for the breaking down to G . For more technical information on this point we refer to [37] and references therein.

3.2 Fluxes and 3-brane tadpoles

While the entire information of 7-brane charge is contained in the geometric data of the F-theory Calabi-Yau 4-fold, the induced 5- and 3-brane charge is extra data encoded in the background flux. In the Type IIB orientifold picture one distinguishes closed string fluxes - most notably the three-form flux $G_3 = F_3 - \tau H_3$ - and brane fluxes F . The latter are the background value of the Yang-Mills field strength on the 7-branes. In F-theory, both RR 3-form fluxes and brane fluxes enjoy a uniform description in terms of the M-theory four-form flux G_4 . In principle, brane fluxes arise by the reduction

$$G_4 = \sum_i F^{(i)} \wedge \omega_i + \dots, \quad (52)$$

where ω_i denote those normalisable harmonic 2-forms of Y_{n+1} that are neither elements of $H^{1,1}(B_n)$ nor represent the fiber class itself. For example, this expression straightforwardly describes abelian gauge flux associated with a Cartan $U(1)$ generator contained in the gauge group G along S . In this case $\omega_i \equiv \omega_i^G$ are the 2-forms Poincaré dual to the resolution divisors \hat{D}_i introduced in (41). Such flux breaks the gauge group G to the commutant of the Cartan generator, a mechanism that will be heavily exploited in the context of GUT symmetry breaking in section 4.2.

More general fluxes, however, are extremely hard to describe in detail because they encode the information of a non-abelian gauge bundle. This is best understood in the context of the global Tate model introduced before. The flux we have in mind is associated with the orthogonal gauge group $H \subset E_8$ and therefore does not affect the gauge group G . An auxiliary tool to study such fluxes is given by the spectral cover construction, which is the subject of section 3.5. Note that gauge flux is essential to achieve a chiral matter spectrum, see the discussion in the next section.

The total amount of F-theory fluxes is constrained by the 3-brane tadpole cancellation condition, which is dual to M2-brane charge cancellation in M-theory. Here we simply quote the result [48]

$$\frac{\chi(\bar{Y}_4)}{24} = N_{M2} + \frac{1}{2} \int_{\bar{Y}_4} G_4 \wedge G_4, \quad (53)$$

where N_{M2} is the number of M2-branes filling the three non-compact dimensions. By F/M-duality $N_{M2} = N_{D3}$, the number of spacetime-filling D3-branes.

The object $\chi(\bar{Y}_4)$ is the Euler characteristic, introduced in eqn. (21), of the resolved Calabi-Yau 4-fold. It encodes all curvature induced 3-form charge of the 7-branes. Indeed in models with a weakly coupled Type IIB description one can match $\chi(\bar{Y}_4)$ with the curvature-dependent terms in the Chern-Simons action of the O7-plane and D7-branes [35]. Special care has to be taken in the computation of $\chi(\bar{Y}_4)$: As long as the 4-fold is non-singular, $\chi(Y_4) = \int c_4(Y)$, and for a Weierstrass model over B_3 a simple formula exists [26, 48]

$$\chi(Y_4) = 12 \int_{B_3} c_1(B_3)c_2(B_3) + 360 \int_{B_3} c_1^3(B_3). \quad (54)$$

For the phenomenologically relevant case of F-theory compactification with non-abelian gauge groups, however, Y_4 is singular and the computation of its Euler characteristic requires the resolution of Y_4 to \bar{Y}_4 . For general singular Calabi-Yau 4-folds the computation of $\chi/24$ can become quite involved [49, 50]. However, as stressed above, for the class of 4-folds that can be described with the help of toric methods a well-defined algorithm exists that allows one to deduce $\chi/24$, see [43–45] for examples of this type in the context of global F-theory GUT models. Furthermore for global Tate models with non-abelian singularities solely along a single divisor S , a simple closed formula for $\chi/24$ has been conjectured [43], see the discussion around (100).

Note that the value of χ for a singular 4-fold Y_4 is usually significantly lower than the expression (54) for the corresponding smooth Weierstrass model. This is of direct relevance for the construction of four-dimensional F-theory vacua because moduli stabilisation and the engineering of a realistic particle spectrum require the inclusion of four-form flux. In consistent models the flux is not allowed to overshoot $\chi/24$ as this would necessitate the inclusion of anti 3-branes, which would lead to instabilities.⁷ Typically, F-theory vacua with high rank non-abelian gauge groups are therefore much more constrained than vacua with lower rank.

3.3 Matter curves and Yukawa points

In previous sections we have described the localisation of the non-abelian gauge degrees of freedom in F-theory along divisors D_a of the base space B_n of the elliptic Calabi-Yau. Each such divisor carries a G_a gauge theory with four supercharges, corresponding to $\mathcal{N} = 1$ Super-Yang-Mills in four dimensions. Besides the vector multiplet in the adjoint representation of G_a there arises extra massless charged matter.

In Type IIB orientifolds on a Calabi-Yau 3-fold X , chiral multiplets in the adjoint representation of G_a correspond to D7-brane moduli [51]. These include brane deformation moduli and Wilson line moduli, counted respectively by $H_-^{(0,2)}(D_a) \simeq H_-^{(0)}(S, K_S)$ and $H_-^{(0,1)}(S)$. The subscripts indicate that only the odd eigenspaces of the cohomology groups under the orientifold action σ are relevant. In F-theory there is no clear separation between open and closed string moduli. In fact the brane position moduli become part of the complex structure moduli $H^{(3,1)}(Y_4)$. For a precise account of these fields and the Wilson line moduli in the effective action derived by dimensional reduction of the dual M-theory we refer to [25].

Apart from these non-chiral matter fields, there exist two types of potentially chiral charged matter: so-called bulk states that propagate along the whole divisor and matter at the intersection locus of two divisors D_a and D_b . The appearance of these chiral multiplets parallels the situation in Type IIB orientifolds, which is therefore worthwhile recalling in this context. For this we will follow mostly the discussion in [52].

Bulk matter

Consider two parallel stacks of Type IIB D7-branes along the same divisor S , each carrying internal gauge flux $2\pi\alpha'F_a$ and $2\pi\alpha'F_b$, respectively. This flux is the curvature of two vector bundles V_a and V_b with Chern character

$$\mathrm{tr} \left[e^{2\pi\alpha'F_i} \right] = \mathrm{ch}(V_i). \quad (55)$$

⁷One might argue that the anti 3-brane should be stabilised in a warped region as in KKLT and thus do no further harm, but this assumes a sufficient amount of 3-form flux allowed by the 3-brane tadpole to create a warped throat in the first place.

In the sequel we restrict ourselves to line bundles L_a, L_b characterised entirely by their first class $c_1(L_i) = \text{tr} \frac{1}{2\pi} \ell_s^2 F_i$. For simplicity we assume that S is at general position with respect to the O7-plane so that the gauge group is $U(N_a) \times U(N_b)$. The bulk matter describes chiral multiplets in the bifundamental representation propagating along the whole divisor S . Such matter is in general counted by so-called extension groups

$$\text{Ext}^n(\iota_* L_a, \iota_* L_b), \quad n = 0, \dots, 3, \quad (56)$$

where $\iota : S \rightarrow X$ denotes the inclusion of the divisor S into the Calabi-Yau 3-fold X . This was derived in detail from an analysis of B-model vertex operators in [53]. The value $n = 1$ refers to anti-chiral multiplets transforming as (\bar{N}_a, N_b) , while $n = 2$ corresponds to chiral multiplets in the same representation. For consistency, the states counted by the groups corresponding to $n = 0$ and $n = 3$ must be absent. These states do not describe matter fields but rather refer to gauge multiplets. One can show that the sheaf extension groups eqn. (56) translate into certain cohomology groups for the line bundles on the divisor S , concretely

$$\begin{aligned} \text{Ext}^0(\iota_* L_a, \iota_* L_b) &= H^0(S, L_a \otimes L_b^\vee), \\ \text{Ext}^1(\iota_* L_a, \iota_* L_b) &= H^1(S, L_a \otimes L_b^\vee) + H^2(S, L_a^\vee \otimes L_b), \\ \text{Ext}^2(\iota_* L_a, \iota_* L_b) &= H^2(S, L_a \otimes L_b^\vee) + H^1(S, L_a^\vee \otimes L_b), \\ \text{Ext}^3(\iota_* L_a, \iota_* L_b) &= H^0(S, L_a^\vee \otimes L_b). \end{aligned} \quad (57)$$

The net chirality follows as [52]

$$I_{ab}^{bulk} = \sum_{n=0}^3 (-1)^n \dim \text{Ext}^n(\iota_* L_a, \iota_* L_b) = - \int_X [S] \wedge [S] \wedge (c_1(L_a) - c_1(L_b)). \quad (58)$$

As a variant of this setup one can also consider a single brane stack with gauge group G along S and consider gauge flux in terms of a non-trivial embedding of a vector bundle with structure group $G_2 \subset G$. This breaks G to the commutant G_1 of G_2 . In this case chiral multiplets charged under G_2 arise, which originate from the adjoint representation of G . More precisely, the group theoretic decomposition

$$G \rightarrow G_1 \times G_2 \quad (59)$$

$$\text{ad}_G \rightarrow (\text{ad}_{G_1}, 1) \oplus (1, \text{ad}_{G_2}) \oplus \sum (R_x, U_x) \quad (60)$$

leads to bulk matter in suitable representations R_x under the visible gauge group G_1 . The individual states are counted by appropriate cohomology groups with values in the bundle representation U_x . For instance, a non-trivial gauge line bundle L associated with a Cartan $U(1)$ of G breaks $G \rightarrow \tilde{G} \times U(1)$, and if the representations of \tilde{G} carry $U(1)$ charge q , the relevant bundle is L^q . This generalises the appearance of $L_a \otimes L_b^\vee$ in (57) for bifundamental states.

Even though derived originally in the perturbative Type IIB context, these expressions continue to hold for general 7-branes in F-theory. This can be derived e.g. with the help of the eight-dimensional twisted field theory introduced in [7].

Localised matter

From the intersecting brane perspective it is natural to expect that additional massless matter appears at the intersection of two 7-branes. What happens in F-theory is that as two singular

loci collide transversely, the singularity type of the elliptic fibration enhances over intersection common locus [54]. This intersection occurs in complex codimension one along the base. In F-theory compactifications on Calabi-Yau 4-folds, two divisors D_a and D_b intersect along a complex curve $C_{ab} = D_a \cap D_b$. The appearance of extra matter can be understood in terms of the tree of zero-size \mathbb{P}^1 s over each singular locus. Along the intersection locus the two sets of \mathbb{P}^1 s unite to form the (affine) Dynkin diagram of a new gauge group G_{ab} . Its rank is the sum of the gauge groups G_a and G_b along the two divisors. Note that group theoretically there may be several types of enhancements possible, each leading to different types of matter states. By abuse of notation one calls the group G_{ab} the enhanced gauge group along the intersection locus even though in actuality there exists no $\mathcal{N} = 1$ SYM theory with a corresponding vector multiplet. However, from the reduction of the M-theory 3-form and from wrapped M2-branes along the various \mathbb{P}^1 s in the fiber one does find as many massless states along C_{ab} as is necessary to form the adjoint of G_{ab} . These states include the states propagating along D_a and D_b but also contain extra matter localised at C_{ab} . Under the group theoretic decomposition

$$G_{ab} \rightarrow G_a \times G_b \tag{61}$$

$$\text{ad}_{G_{ab}} \rightarrow (\text{ad}_{G_a}, 1) \oplus (1, \text{ad}_{G_b}) \oplus \sum (R_x, U_x) \tag{62}$$

the localised states transform in the representation (R_x, U_x) . These states descend from M2-branes wrapping chains of \mathbb{P}^1 s associated both with the Dynkin diagram of G_a and G_b , in analogy to the chains S_{ij} defined in the paragraph after (42). Away from the intersection locus C_{ab} , such chains are not of zero size and the extra matter becomes massive.

Note that for enhancements to non-classical groups G_{ab} the irreps (R_x, U_x) need not correspond to two-index representations, as would be the case for perturbative Type IIB models. The larger set of possibilities is again due to the multiple endpoints of multi-pronged $\binom{p}{q}$ strings, which furnish a genuine strong coupling effect. While the Chan-Paton factors of a fundamental $\binom{1}{0}$ string with two endpoints always give rise to two-index representations, a multi-pronged string can accommodate more general charges.

An analysis of the twisted defect SYM theory along the matter curves [7] confirms that this massless matter is counted by the same expressions familiar from perturbative Type IIB models [53],

$$H^i(C_{ab}, F_x \otimes K_{C_{ab}}^{\frac{1}{2}}), \quad i = 0, 1. \tag{63}$$

Here $i = 0$ and $i = 1$ respectively count chiral and anti-chiral multiplets in representation (R_x, U_x) , and F_x is schematic for suitable combinations of gauge flux restricted onto C_{ab} . The chiral index follows via Hirzebruch-Riemann-Roch as

$$\chi = \dim H^0(C_{ab}, F_x \otimes K_{C_{ab}}^{\frac{1}{2}}) - \dim H^1(C_{ab}, F_x \otimes K_{C_{ab}}^{\frac{1}{2}}) = \int_{C_{ab}} F_x. \tag{64}$$

We will be more specific about the concrete chirality formulae in the context of the spectral cover construction of gauge flux in section 3.5.

The type of singularity enhancement can be read off from the discriminant locus with the help of Tate's algorithm. As a non-trivial example we consider again the $SU(5)$ gauge theory along the divisor S introduced above. In this case the localised matter states arise at the intersection of the $SU(5)$ brane S with the I_1 -locus S_1 . Since the latter carries no non-abelian gauge group, the singularity gets enhanced by rank one. Two types of rank-one enhancements are possible for $SU(5)$ [6, 7]:

- Enhancement $A_4 \rightarrow D_5$ corresponding to $SU(5) \rightarrow SO(10)$. The extra matter transforms in the **10** representation of $SU(5)$ as can be determined from the branching rule

$$\mathbf{45} \rightarrow (\mathbf{24})_0 + (\mathbf{1})_0 + \mathbf{10}_2 + \overline{\mathbf{10}}_{-2}. \quad (65)$$

The subscripts denote the (formal) $U(1)$ charges under the decomposition $SO(10) \rightarrow SU(5) \times U(1)$.

From Tate's algorithm this D_5 enhancement occurs along the curve

$$P_{10}: \quad w = 0 \quad \cap \quad \mathfrak{b}_5 = 0, \quad (66)$$

along which the discriminant (50) scales like w^7 , as required for a D_5 singularity.

- Enhancement $A_4 \rightarrow A_5$ corresponding to $SU(5) \rightarrow SU(6)$. The $SU(6)$ locus hosts matter in the **5** arising via

$$\mathbf{35} \rightarrow (\mathbf{24})_0 + (\mathbf{1})_0 + \mathbf{5}_1 + \overline{\mathbf{5}}_{-1}. \quad (67)$$

This enhancement occurs whenever

$$P_5: \quad w = 0 \quad \cap \quad P = \mathfrak{b}_3^2 \mathfrak{b}_4 - \mathfrak{b}_2 \mathfrak{b}_3 \mathfrak{b}_5 + \mathfrak{b}_0 \mathfrak{b}_5^2 = 0, \quad (68)$$

consistent with the scaling $\Delta \propto w^6$ in (50).

In addition there can appear localised GUT singlets **1** on matter curves away from (but possibly intersecting) the $SU(5)$ brane S . These localised states appear at self-intersections of the I_1 -part S_1 of the discriminant, along which the gauge group enhances to A_1 [37].

Yukawa points

In compactifications on Calabi-Yau 4-folds, two or more matter curves can meet in points. Here the singularity type of the fiber enhances even further by the same mechanism that leads to the enhancement along matter curves. Consider for example the intersection of three matter curves C_{ab}, C_{bc}, C_{ac} in one point. Group theoretically the representations of the matter states along the three curves combine to form the adjoint ad_{abc} of an enhanced gauge group G_{abc} . Even though there exists no actual gauge theory associated with G_{abc} , the cubic interaction term for the adjoint of this (hypothetical) enhanced gauge group leads to Yukawa interactions for the massless matter [6,7]. This can be argued by a closer analysis of the local gauge theory description of the geometry responsible for the enhancements. The resulting cubic interaction in the product theory $G_a \times G_b \times G_c$ can be determined by decomposing the triple product ad_{abc}^3 into gauge invariant triple products for the irreducible representations of G_a, G_b and G_c . This picture is in perfect agreement with expectations from weakly-coupled Type IIB theory, where Yukawa couplings are known to be localised at the intersection of matter curves.

To illustrate this picture in the context of our $SU(5)$ gauge theory, we note that there exist three possible rank-two enhancements of A_4 to E_6, D_6 and A_7 with the following Yukawa structure [6, 7]:

- The **10 10 5** Yukawa is localized at a point of E_6 enhancement $\mathfrak{b}_5 = 0 = \mathfrak{b}_4$.
- The **10 $\overline{\mathbf{5}}$ $\overline{\mathbf{5}}$** is localized at a D_6 point $\mathfrak{b}_5 = 0 = \mathfrak{b}_3$.

	sing. type	discr. deg(Δ)	gauge enh.		coeff. vanish. deg					object equation
			type	group	a_1	a_2	a_3	a_4	a_6	
GUT:	I_5^s	5	A_4	$SU(5)$	0	1	2	3	5	$S : w = 0$
matter:	I_6^s	6	A_5	$SU(6)$	0	1	3	3	6	$P_5 : P = 0$
	I_1^{*s}	7	D_5	$SO(10)$	1	1	2	3	5	$P_{10} : b_5 = 0$
Yukawa:	I_2^{*s}	8	D_6	$SO(12)^*$	1	1	3	3	5	$\mathfrak{b}_3 = \mathfrak{b}_5 = 0$
	IV^{*s}	8	E_6	E_6	1	2	2	3	5	$\mathfrak{b}_4 = \mathfrak{b}_5 = 0$
extra:	I_7^s	7	A_6	$SU(7)$	0	1	3	4	7	$P = R = 0,$ $(\mathfrak{b}_4, \mathfrak{b}_5) \neq (0, 0)$

Table 2: Relevant gauge enhancements in a generic $SU(5)$ GUT geometry (borrowed from [43]).

Indeed $\mathfrak{b}_4 = 0 = \mathfrak{b}_5$ and $\mathfrak{b}_5 = 0 = \mathfrak{b}_3$ correspond to a single and double zero of P_5 in agreement with the order of $\overline{\mathbf{5}}$ representations appearing in the coupling.

- At $P = 0 = R$ but $(\mathfrak{b}_4, \mathfrak{b}_5) \neq (0, 0)$ the singularity type enhances to A_6 . This realizes the coupling $\mathbf{5} \overline{\mathbf{5}} \mathbf{1}$. The state $\mathbf{1}$ represents a (possibly localised) GUT singlet.

We summarise the various codimension singularities of generic $SU(5)$ models and their physical relevance in table 2.

3.4 F-theory-heterotic duality

Generalities on the duality

Thus far we have approached F-theory via Type IIB orientifolds and via duality with M-theory. In this section we highlight some of the most basic aspects of the duality with the heterotic string. This duality is a particularly fruitful source of inspiration for many applications to concrete model building.

The basic assertion is that F-theory on an elliptic $K3 : T^2 \rightarrow \mathbb{P}^1$ is dual to the heterotic string on T^2 [5]. This can be argued by comparing the moduli space of both theories. In particular, the heterotic string coupling $\exp(2\phi)$ is dual to the volume of the base \mathbb{P}^1 of the F-theory $K3$.

This basic duality can be extended "adiabatically" by fibering an elliptic $K3$ -surface and, respectively, an elliptic curve over a common n -complex dimensional base B_n in such a way as to produce Calabi-Yau manifolds on either side of the duality. Thus F-theory on a $K3$ -fibered Calabi-Yau $(n+2)$ -fold $Y_{n+2} : K3 \rightarrow B_2$ is dual to the heterotic string on the elliptic Calabi-Yau $(n+1)$ -fold $Z_{n+1} : T^2 \rightarrow B_n$. Since Y_{n+2} exhibits a double fibration structure as a $K3$ -fibration over B_2 and, as always, as an elliptic fibration $Y_{n+2} : T^2 \rightarrow B_{n+1}$, the base B_{n+1} is by itself \mathbb{P}^1 -fibered, $B_{n+1} : \mathbb{P}^1 \rightarrow B_n$. Therefore only a very special subclass of F-theory compactifications possesses a straightforward heterotic dual in terms of an elliptic fibration Z_{n+1} .

On the heterotic side, a perturbative $SO(32)$ or $E_8 \times E_8$ gauge group descends from the respective ten-dimensional theory by compactification on a smooth Calabi-Yau. If present, singularities on the heterotic compactification space lead to extra gauge group factors not related to this perturbative gauge group which can be of potentially huge rank and which are due to massless non-perturbative states. In the sequel we will focus on $E_8 \times E_8$ heterotic

models with only the perturbative group, i.e. on models for which the elliptic fibration Z_{n+1} is smooth. Heterotic models on smooth Calabi-Yau spaces are defined in terms of a holomorphic vector bundle $V_1 \oplus V_2$ with structure group $H_1 \times H_2$ embedded into $E_8 \times E_8$. This breaks the gauge group down to the commutant $G_1 \times G_2$. The vector bundle data must therefore map into the singularity structure of the elliptic fibration Y_{n+2} on the dual F-theory side.

Heterotic/F-theory duality is particularly powerful in compactifications to six dimensions, where it relates the heterotic string on a $K3$ which is elliptically fibered over \mathbb{P}^1 to F-theory on a $K3$ fibration over that same \mathbb{P}^1 . This implies that the base B_2 of the F-theory elliptic fibration is itself a \mathbb{P}^1 -fibration over \mathbb{P}^1 . Such a fibration is called a rationally ruled or Hirzebruch surface \mathbb{F}_k , where $k = 0, 1, \dots$ determines the structure of the fibration. The geometry of elliptic fibrations over \mathbb{F}_k and applications to F-theory/heterotic duality in six dimensions have been discussed in detail in the pioneering [29] (more general elliptic 3-folds are analysed in [40]). The value k of the F-theory base \mathbb{F}_k determines the second Chern class or instanton number of the heterotic bundle $V_1 \times V_2$ embedded into $E_8 \times E_8$ as $(12 - k, 12 + k)$, $k = 0, 1, \dots, 12$. The heterotic dilaton is now related to the volume of the fiber and of the base \mathbb{P}^1 of the F-theory B_2 as

$$\exp(2\phi) = \frac{\text{vol}_f}{\text{vol}_b}. \quad (69)$$

For compactifications to four dimensions, the set of possible F-theory $K3$ -fibrations is constrained by the fact that only a small number of complex base spaces B_2 allow for elliptic Calabi-Yau spaces $Z_3 : T^2 \rightarrow B_2$ as required on the heterotic side. In fact, for smooth heterotic compactifications with $\mathcal{N} = 1$ supersymmetry Z_3 must have $SU(3)$ holonomy and B_2 can only be a (blow-up in r points of) \mathbb{P}_2 , a (blow up of) \mathbb{F}_k or the Enriques surface $K3/\mathbb{Z}_2$ [55] (see [40] for more explanations). The base B_3 of the dual F-theory elliptic 4-fold is then a \mathbb{P}^1 -fibration over these complex surfaces. Such a fibration is characterised by a line bundle \mathcal{T} over B_2 with first Chern class $c_1(\mathcal{T}) = t$ as follows: Consider the rank 2 bundle $\mathcal{O} \oplus \mathcal{T}$ over B_2 . Its fiber consists of two copies of \mathbb{C} , the fibers of the trivial line bundle \mathcal{O} and of \mathcal{T} . In the same manner as one forms a \mathbb{P}^1 by projectivising two complex coordinates $(0, 0) \neq (z_1, z_2) \simeq \lambda(z_1, z_2)$, a \mathbb{P}^1 -fibration can be obtained as the projectivisation $\mathbb{P}(\mathcal{O} \oplus \mathcal{T}) = B_3$. I.e. one projectivises each fiber $\mathbb{C} \oplus \mathbb{C}$ and then fibers over B_2 . Let $r = c_1(\mathcal{O}(1))$, where by abuse of notation $\mathcal{O}(1)$ is the line bundle over B_3 that reduces, along each \mathbb{P}^1 fiber, to $\mathcal{O}_{\mathbb{P}^1}(1)$. Then one can show that [56]

$$r(r + t) = 0, \quad c_1(B_3) = c_1(B_2) + 2r + t. \quad (70)$$

The upshot is that the class t generalises the integer k characterising the Hirzebruch surface \mathbb{F}_k for F-theory in 6 dimensions. In order to understand the generalisation of the relation between k and the instanton number $(12 - k, 12 + k)$ of the heterotic bundle, we need to familiarize ourselves with the construction of vector bundles on elliptic 3-folds Z_3 . Thanks to the work of Donagi [57] and Friedman, Morgan and Witten [56] a large class of holomorphic vector bundles on elliptic fibrations are known in terms of the spectral cover construction. In the sequel we give a brief summary of the basic idea and quantities characterising a spectral cover bundle.

The heterotic spectral cover construction

The construction of a rank n spectral cover bundle involves two concepts: that of a spectral surface $\mathcal{C}^{(n)}$ and of a spectral line bundle \mathcal{N} , both describable in terms of cohomological data

on B_2 . Under F-theory/heterotic duality this data maps to the geometry of the singular Y_4 and to gauge flux G_4 .

The basic idea of the spectral cover construction is to first construct a stable $(S)U(n)$ bundle on the elliptic fibre over each point of the base, which is then extended over the whole manifold Z_3 by gluing the data together suitably. Recall that in general, an $(S)U(n)$ bundle defines a rank n complex vector bundle. Its restriction to the elliptic fiber E_b over $b \in B_2$ can be shown to be isomorphic to the direct sum of n complex line bundles

$$V|_{E_b} = \mathcal{N}_1 \oplus \dots \oplus \mathcal{N}_n, \quad (71)$$

each of which has to be of zero degree. If $G = SU(n)$ as opposed to $U(n)$, $V|_{E_b}$ must in addition be of trivial determinant, i.e. $\bigotimes_{i=1}^n \mathcal{N}_i = \mathcal{O}_{E_b}$. The zero degree condition on \mathcal{N}_i implies that there exists for each \mathcal{N}_i a meromorphic section with precisely one zero at some Q_i and a pole at p , the zero of the elliptic curve. I.e. $\mathcal{N}_i = \mathcal{O}_{E_b}(Q_i - p)$. Consequently, stable $(S)U(n)$ bundles on an elliptic curve are in one-to-one correspondence with the unordered n -tuple of points Q_i , and the reduction of $U(N)$ to $SU(n)$ is encoded in the additional requirement that $\sum_i (Q_i - p) = 0$ in the group law of the elliptic curve.

Having understood the restriction of a rank n bundle V to each elliptic fibre, [56] proceed to construct the whole of V . In intuitive terms, the above implies that over an elliptically fibered manifold a $U(n)$ vector bundle determines a set of n points, varying over the base. More precisely, the bundle V over Z_3 with the property

$$V|_{E_b} = \bigoplus_{i=1}^n \mathcal{O}(Q_i - p) \quad (72)$$

uniquely defines an n -fold (ramified) cover $\mathcal{C}^{(n)}$ of B_2 , the spectral cover. It is defined by a projection

$$\pi_n : \mathcal{C}^{(n)} \rightarrow B_2 \quad \text{such that} \quad \mathcal{C}^{(n)} \cap E_b = \bigcup_i Q_i. \quad (73)$$

$\mathcal{C}^{(n)}$ is conveniently described, as a hypersurface in Z_3 , by its Poincaré dual two-form. The Weierstrass model Z_3 possesses a section σ which identifies the base B_2 as an element of $H_4(B_2, \mathbb{Z})$. This section has the important property

$$\sigma \cdot \sigma = -\sigma c_1(S). \quad (74)$$

The class of $\mathcal{C}^{(n)}$ can then be written as

$$[\mathcal{C}^{(n)}] = n\sigma + \pi^*(\eta) \in H^2(Z_3, \mathbb{Z}) \quad (75)$$

for η some effective class in $H^2(B_2, \mathbb{Z})$. In particular the first piece $n\sigma$ shows that the spectral surface is an n -fold cover of the base B_2 .

Several distinct bundles over Z_3 may well give rise to the same spectral cover $\mathcal{C}^{(n)}$ since the latter only encodes the information about the restriction of V to the fibre E_b . To recover V from the spectral data we need to specify in addition how it varies over the base, i.e. $V|_{B_2}$. As discussed in [56] this is uniquely accomplished by the so-called spectral line bundle \mathcal{N} on $\mathcal{C}^{(n)}$ with the property

$$\pi_{n*} \mathcal{N} = \mathcal{V}|_{B_2}. \quad (76)$$

For the first Chern class $c_1(\mathcal{N}) \in H^2(\mathcal{C}^{(n)}; \mathbb{Z})$ of the spectral bundles one can make the general decomposition ansatz [56, 58]

$$c_1(\mathcal{N}) = \frac{r}{2} + \gamma. \quad (77)$$

Here we have abbreviated

$$r = -c_1(\mathcal{C}^{(n)}) + \pi_n^* c_1(S), \quad \gamma = \frac{1}{n} \pi_n^* c_1(V) + \gamma_u, \quad (78)$$

where γ_u is chosen such that it satisfies $\pi_{n*} \gamma_u = 0$. This yields

$$\gamma_u = \lambda (n\sigma - \pi_n^* \eta + n\pi_n^* c_1(S)), \quad (79)$$

for a number $\lambda \in \mathbb{Q}$. Let us further parametrize $c_1(V)$ by some element $\zeta \in H^2(S; \mathbb{Z})$ [58],

$$\zeta = c_1(V). \quad (80)$$

The parameter $\lambda \in \mathbb{Q}$ is subject to certain constraints to be discussed shortly. Note that this bundle exists for generic complex structure since it only involves σ and the pullback of classes from B_2 .

The parameter $\lambda \in \mathbb{Q}$ has to be chosen such that $c_1(\mathcal{N})$ defines an integer class in $H^2(\mathcal{C}^{(n)}; \mathbb{Z})$. On the non-Calabi-Yau space X the adjunction formula leads to

$$-c_1(\mathcal{C}^{(n)}) = (n-2)\sigma + \pi_n^*(\eta - 2c_1(B_2)). \quad (81)$$

Putting everything together, we have

$$c_1(\mathcal{N}) = -\sigma + n\left(\frac{1}{2} + \lambda\right)\sigma + \left(\frac{1}{2} - \lambda\right)\pi_n^*\eta \quad (82)$$

$$+ \left(-\frac{1}{2} + n\lambda\right)\pi_n^* c_1(B_2) + \frac{1}{n}\pi_n^* \zeta. \quad (83)$$

E.g. for an $SU(5)$ bundle, integrality of $c_1(\mathcal{N})$ therefore puts the value of $\lambda \in \mathbb{Q}$ subject to the constraints

$$5\left(\frac{1}{2} + \lambda\right) \in \mathbb{Z}, \quad \left(\frac{1}{2} - \lambda\right)\eta + \left(5\lambda - \frac{1}{2}\right)c_1(S) \in H^2(S; \mathbb{Z}). \quad (84)$$

From these data one can compute the higher Chern classes. For an $SU(n)$ bundle these are [56, 59]

$$\begin{aligned} \int_{B_2} c_2(V) &= \int_{B_2} \eta\sigma - \frac{1}{24}\chi_{SU(n)} - \frac{1}{2} \int_{B_2} \pi_{n*}(\gamma^2), \\ \int_{Z_3} c_3(V) &= \lambda \eta(\eta - nc_1(B_2)), \end{aligned} \quad (85)$$

where $\chi_{SU(n)} = \int_{B_2} c_1^2(B_2)(n^3 - n) + 3n\eta(\eta - nc_1(B_2))$.

To summarize, a $U(n)$ spectral cover bundle is characterised by the following topological data:

- the class of the spectral surface $[\mathcal{C}^{(n)}] = n\sigma + \pi^*\eta$, $\eta \in H^2(B_2, \mathbb{Z})$;
- the first Chern class of the spectral line bundle $c_1(\mathcal{N})$ as in (77).

Under F-theory-heterotic duality the bundle data map partly into the singular geometry of Y_4 and partly into gauge flux G_4 . The gauge groups G_1 and G_2 are localised on the divisors given by the base B_2 located at north and south pole of the \mathbb{P}^1 of the fiber of $B_3 : \mathbb{P}^1 \rightarrow B_2$.

- The classes η_i for the two bundles embedded into $E_8^{(1)}$ and $E_8^{(2)}$ correspond on the F-theory side to [56]

$$\eta_1 = 6c_1(S) - t, \quad \eta_2 = 6c_1(S) + t, \quad (86)$$

where $t = c_1(\mathcal{T})$, see the discussion around (70).

- The quantity γ_u appearing in (79) maps into G_4 flux; in particular [60]

$$\int_{Y_4} G_4 \wedge G_4 = - \int_{B_2} \pi_{n*}(\gamma_1^2 + \gamma_2^2). \quad (87)$$

In [56], also methods for the construction of gauge bundles with more general structure group including E_6, E_7, E_8 are developed. If one embeds a bundle with structure group $H_1 \times E_8$ into $E_8 \times E_8$, the visible heterotic gauge group is $G = G_1$. On the F-theory side this maps into a single gauge group G_1 along the base of the \mathbb{P}^1 -fibration B_3 . Note that this is precisely the structure encountered for global Tate models.

3.5 The spectral cover construction for F-theory models

Let us come back to general F-theory models on an elliptically fibration Y_4 . As we have reviewed, the global structure of the Weierstrass model Y_4 is specified by the sections $f \in H^0(B_3, K_{B_3}^{-4})$ and $g \in H^0(B_3, K_{B_3}^{-6})$. Locally this can be brought into the form of eqn. (44) of a Tate model. A special class of fibrations is even globally of the Tate form and thus based on an underlying E_8 gauge symmetry, broken to gauge group G along a single divisor S .

If one is interested not in the full details of the global 4-fold geometry, but merely in the physics on the divisor S , one can restrict the Tate model to the neighbourhood of $S \subset B_3$. This restriction of the Tate constraint to the neighbourhood of S likewise goes under the name spectral cover construction, whose application we just described in the context of heterotic model building. By heterotic/F-theory duality it is clear that spectral covers have a natural appearance also for F-theory compactifications with a heterotic dual. More recently, however, it has been appreciated [61, 62] that spectral covers are useful to describe the geometry and gauge flux of F-theory compactifications even without (simple) heterotic duals - at least in and to some amount even beyond a local picture.

The general philosophy

Before discussing the technicalities, let us try and gain an intuitive understanding of the appearance of the spectral cover construction. For general F-theory models, the essence of the spectral cover idea is to zoom into the local neighbourhood of the divisor $S : w = 0$ within B_3 by discarding all terms of higher power in the normal coordinate w that appear in the sections \mathfrak{b}_n , defined in (49) for the case of $SU(5)$. The restrictions of \mathfrak{b}_n to the divisor S ,

$$b_n = \mathfrak{b}_n|_{w=0}, \quad (88)$$

are therefore sections entirely on S . In this local picture the brane S is described as the base of the bundle $K_S \rightarrow S$, given by $s = 0$. The neighbourhood of S is then modelled by a spectral

surface viewed as a divisor of the total space of K_S . In the sequel we will concentrate on the Tate model for an $SU(5)$ GUT symmetry along S with associated spectral surface

$$\mathcal{C}^{(5)} : b_0 s^5 + b_2 s^3 + b_3 s^2 + b_4 s + b_5 = 0. \quad (89)$$

One can think of $\mathcal{C}^{(5)}$ as encoding the information about the discriminant locus in the local vicinity of S . In particular, as we will see the intersections of $\mathcal{C}^{(5)}$ and S determine the **10**-matter curves (66) on S . It is also clear from the relation (88), though, that all the information in \mathfrak{b}_n contained in terms higher in w is lost in the spectral cover approach.

The spectral cover approach to F-theory model building serves in particular as an auxiliary construction to construct the gauge flux required for chirality of the model. Given the local nature of the spectral covers, it is reasonable to suspect that this yields a correct description of gauge flux near the brane S and in particular along the matter curves on S . This is good news as it is the restriction of the fluxes onto these curves which governs the chirality of a model, but more work is needed to fully understand the continuation of the fluxes in a global construction.

Recall from section 3.4 that in models with a heterotic dual, the elliptic 4-fold Y_4 also has the structure of a $K3$ fibration $K3 \rightarrow B_2$ over a complex surface B_2 ; i.e. the base space B_3 of the elliptic fibration Y is itself globally \mathbb{P}^1 fibered over B_2 . If in addition on the heterotic side the second E_8 factor is broken by an E_8 bundle, then the only non-abelian gauge group is localised along B_2 , which in the $SU(5)$ example we would identify with the GUT divisor S . The GUT divisor is therefore the base of a *globally* defined fibration in models with heterotic dual. In general F-theory models, we have seen that this is not the case. However, from the discussion of the split resolution in section 2.5 we know that one can *locally* view S as the basis of an ALE fibration which describes the singularity structure along S [7]. The ALE fiber contains a distinguished set of two-cycles $\Gamma_{E_8}^i$ whose intersection form is related to the Cartan matrix of E_8 . For generic non-zero size of these two-cycles the E_8 symmetry is broken. If the divisor S exhibits enhanced gauge symmetry G this is because some of the $\Gamma_{E_8}^i$, called Γ_G^i in the sequel, in the fiber shrink to zero size. The intersection matrix of the two-cycles Γ_H^i with non-zero size is related to the Cartan matrix of the commutant $H \subset E_8$ of G .

This picture is of course very reminiscent of the breaking of E_8 to G in the heterotic string by means of a gauge bundle with structure group H . Furthermore, as seen at the end of section 3.4 in F-theory, some of the degrees of freedom of the heterotic vector bundle are encoded purely geometrically, while others map to gauge flux. The geometric part is interpreted in the local field theory of [7] as encoding the vacuum expectation value of the Higgs field $\varphi \in H^{(0)}(S, K_S)$ associated with the normal fluctuations of the 7-brane. The spectral cover now is designed to describe the size of the non-zero two-cycles responsible for the breaking of E_8 to G along S .

Finally we stress once again that for the special case of global Tate models the underlying E_8 structure is exact. It is clear then that the spectral cover construction, which geometrises the breaking of E_8 to G via the Higgs bundle of structure group H , has a chance to capture more than just the very local neighbourhood of the divisor S .

Technical details of spectral covers for an $SU(5)$ model

In what follows we restrict ourselves to the spectral cover description of a $G = SU(5)_{GUT}$ singularity along a divisor $S \subset B_3$. The complement of $SU(5)_{GUT}$ in E_8 is denoted by $H = SU(5)_\perp$. For more background and for details on more general configurations we refer to [61, 62].

The starting point is to construct an auxiliary non-Calabi-Yau 3-fold \mathcal{W} as a fibration over S which encodes the singular geometry of S in B_3 . This space \mathcal{W} serves as a compactification of the total space of the bundle $K_S \rightarrow S$ introduced before. We will therefore think of S either as a divisor on B_3 or as the base of a fictitious 3-fold \mathcal{W} . The definition of \mathcal{W} is as the projectivized bundle over the GUT divisor S

$$\mathcal{W} = \mathbb{P}(\mathcal{O}_S \oplus K_S), \quad p_{\mathcal{W}} : \mathcal{W} \rightarrow S, \quad (90)$$

where $p_{\mathcal{W}}\mathcal{W}$ is the projection to the base of the bundle. The base S is viewed as the vanishing locus of the section σ in \mathcal{W} . This section satisfies the important relation

$$\sigma \cdot \sigma = -\sigma c_1(S). \quad (91)$$

This should ring a bell and remind us of the relation (91) encountered for the heterotic spectral cover construction on the physical elliptic Calabi-Yau space Z_3 . Indeed from now on the construction proceeds in the same fashion as described in section 3.4, replacing the base B_2 of the elliptic Calabi-Yau fibration Z_3 by the GUT divisor S , viewed as the base of the \mathbb{P}^1 -fibration \mathcal{W} . Unlike Z_3 , \mathcal{W} is not Calabi-Yau and has first Chern class

$$c_1(\mathcal{W}) = 2\sigma + 2c_1(S). \quad (92)$$

The spectral cover is constructed as a 5-fold cover of S within \mathcal{W} ,

$$\pi_5 : \mathcal{C}^{(5)} \rightarrow S. \quad (93)$$

While in the heterotic context the intersection points of $\mathcal{C}^{(5)}$ with the elliptic fiber encode the information about the vector bundle V restricted to the fiber, see eqn. (73), here the 5 intersection points with the \mathbb{P}^1 -fiber denote the 5 eigenvalues of the $SU(5)_H$ -valued Higgs field φ along S [61, 62], which in turn specifies the local neighbourhood of S inside B_3 [7, 62].

Given the implicit underlying E_8 structure of the ALE fibration, the massless matter representations of $G = SU(5)$ can be understood as the irreducible representations R_x in the decomposition $\mathbf{248} \rightarrow \sum \text{ad}_i + \sum_x (R_x, U_x)$,

$$\mathbf{248} \mapsto (\mathbf{24}, 1) + (1, \mathbf{24}) + [(\mathbf{10}, \mathbf{5}) + (\overline{\mathbf{5}}, \mathbf{10}) + h.c.]. \quad (94)$$

The matter curve P_{10} is the locus $b_5 = 0$ on S . It is associated with the spectral cover in the fundamental representation of $H = SU(5)_{\perp}$ because the $\mathbf{10}$ appears as $(\mathbf{10}, \mathbf{5})$ in (94). Let us also define the object \mathcal{P}_{10} viewed as a curve in \mathcal{W} [62],

$$\mathcal{P}_{10} = \mathcal{C}^{(5)} \cap \sigma \subset \mathcal{W}. \quad (95)$$

Then the matter curve on S is related to \mathcal{P}_{10} as

$$[P_{10}] = [\mathcal{P}_{10}]|_{\sigma} = (5\sigma + \pi_5^* \eta)|_{\sigma} = \eta - 5c_1(S) \quad (96)$$

with the help of (91), i.e. the restriction of \mathcal{P}_{10} is cohomologically equivalent to the matter curve P_{10} .

The matter curve for the $\overline{\mathbf{5}}$ on X is more complicated and was analyzed in detail in the context of the heterotic string in [63–65], to which we refer for more details.

Gauge flux in $SU(5)$ GUT models

So far all that the spectral cover approach has done for us is to rewrite the geometric data in a seemingly more complicated manner. Its actual power, however, becomes apparent once one includes gauge flux into the compactification. In the context of $SU(5)$ GUT theories one distinguishes between the gauge flux along the GUT divisor S itself and the flux along the matter branes which constitute the I_1 locus of the discriminant. The first type of gauge flux takes values in $G = SU(5)_{GUT}$ and will thus further break the $SU(5)$ GUT symmetry. As for flux on the matter branes, the spectral cover approach makes use of the fact that $\mathcal{C}^{(5)}$ describes the geometry in the vicinity of S and in particular the *local* geometry of the I_1 component. In the machinery of the spectral cover, gauge flux on I_1 is therefore given in terms of the so-called spectral line bundle \mathcal{N} along $\mathcal{C}^{(5)}$ defined by its first Chern class

$$c_1(\mathcal{N}) \in H^{(1,1)}(\mathcal{C}^{(5)}; \mathbb{Z}). \quad (97)$$

Since $\mathcal{C}^{(5)}$ is a five-fold cover of S in \mathcal{W} one can push this line bundle forward to S via π_{5*} . This defines a rank-5 vector bundle

$$V = \pi_{5*}\mathcal{N} \quad (98)$$

on S . Following the general logic of the ALE fibration over S the structure group of this bundle V is identified with the commutant $SU(5)_\perp \subset E_8$ of the GUT $SU(5)$ along S and therefore indeed associated with the flux on I_1 .

Therefore the construction of gauge flux on the matter branes is reduced to the problem of constructing an $SU(5)_H$ bundle V on the basis of the \mathbb{P}^1 -fibration \mathcal{W} . At this point we can again refer to the discussion of the spectral line bundle and its associated spectral cover bundle in the heterotic context, which proceeds, *mutatis mutandis*, in the same manner. The spectral cover approach thus provides a concrete dictionary to describe a certain subclass of gauge fluxes in terms of a few parameters. Moreover, it allows for the computation of the chiral index for the various matter representations localised along the matter curves in terms of these input parameters and is therefore of considerable practical use for model building purposes. It would lead too far to derive these formulae for the chirality in full detail here. We simply state without proof that the chiral index of states in the **10** representation localised on the curve P_{10} on S is given by [62],

$$\chi_{\mathbf{10}} = \sigma \cdot \mathcal{C}^{(5)} \cdot \gamma_u = -\lambda \int_S \underbrace{\eta(\eta - 5c_1(S))}_{P_{10}}, \quad (99)$$

with γ_u as in eqn. (79) for $SU(5)_H$ bundles. Consistently, this is also the number of chiral families in the $\bar{\mathbf{5}}$ representation.

Let us conclude this general presentation of the spectral cover construction by taking up our initial concerns about its global validity. We have stressed several times that as of this writing it yet remains to find a general description of gauge flux in terms of global G_4 flux. Nonetheless it is remarkable that the spectral cover construction does capture some of the global aspects of the geometry correctly. More precisely this is the case for the special subclass of elliptic fibrations that can globally be described as a Tate model with non-abelian gauge group $G \subset E_8$ solely along a divisor S . If we take the spectral cover construction at face value for a second one can conjecture a simple closed expression for the Euler characteristic of the resolution \bar{Y}_G of the singular 4-fold given by [43]

$$\chi(\bar{Y}_G) = \chi^*(Y) + \chi_H - \chi_{E_8}. \quad (100)$$

Here $\chi^*(Y) = 12 \int_B c_1(B_3) c_2(B_3) + 360 \int_B c_1^3(B_3)$ denotes the expression valid for a smooth elliptic fibration over base B_3 , and the remaining two terms are listed in table 3. They involve data solely on S and reflect the underlying E_8 structure of the spectral cover construction, in which we think of first enhancing the singularity over S to E_8 and then breaking it via the H -bundle down to G . This formula can be derived for models with a heterotic dual by F-theory-heterotic duality. The point is now that for global Tate models the underlying E_8 structure is correct globally. Indeed for global Tate models in which an explicit resolution of the singularities is available, the result of (100) can be compared to the value of $\chi(\overline{Y}_G)$ computed explicitly via resolution. This has been performed in [43, 44] for a number of cases using the machinery of toric geometry, finding perfect agreement. For such geometries it is not unreasonable that also the gauge flux constructed via the spectral covers has a global extension. It is this expectation that underlies the global F-theory models as existent in the literature as of this writing, but a more complete understanding of gauge flux without relying on the spectral cover construction is desirable. For example [66] proposes a global extension of spectral cover fluxes in terms of a so-called spectral divisor designed so as to reproduce the expressions for the chirality of states along the GUT branes, i.e. the local aspects associated with non-abelian gauge symmetry. It remains to be seen how to capture also genuinely global aspects of the flux such as the integral $\int_{\overline{Y}_4} G_4 \wedge G_4$ appearing or the chiral index of singlets under the non-abelian group G .

$G = E_8/H$	H	χ_H
$E_{9-n}, n \leq 5$	$SU(n)$	$\int_S c_1^2(S)(n^3 - n) + 3n\eta(\eta - nc_1(S))$
$SU(3)$	E_6	$72 \int_S (\eta^2 - 7\eta c_1(S) + 13c_1^2(S))$
$SU(2)$	E_7	$18 \int_S (8\eta^2 - 64\eta c_1(S) + 133c_1^2(S))$
-	E_8	$120 \int_S (3\eta^2 - 27\eta c_1(S) + 62c_1^2(S))$

Table 3: Redefined Euler characteristic for E_n -type gauge groups. Here η is given by $\eta = 6c_1(S) + c_1(N_S)$.

4 Phenomenological applications to GUT model building

As stressed in the introduction, F-theory combines two characteristic aspects of Type II orientifolds with D-branes on the one hand and of heterotic string vacua on the other which are of general phenomenological interest and of significant use in the context of realistic GUT model building, in particular

- the appearance of exceptional gauge groups as in $E_8 \times E_8$ heterotic string constructions and
- the localisation of gauge degrees of freedom, matter states and Yukawa interactions as in perturbative D-brane models.

Beginning with [6–9] there have been intense recent investigations exploring the prospects of F-theory compactifications for GUT phenomenology. While in principle applicability of

F-theory is by no means restricted to GUT models, the relevance of exceptional gauge groups in the context of unification singles out this class of constructions as the one with the most distinctive F-theoretic features as compared with intersecting brane models in perturbative orientifolds. Promising starting points for the construction of GUT models are the gauge groups $SO(10)$ and $SU(5)$. Most efforts in the F-theory literature have focused on the latter, mainly due to complications with GUT breaking via internal fluxes.

Given the localisation of gauge degrees of freedom in F-theory, a considerable number of phenomenological questions can be discussed already at the level of local models. Among these are the structure of the GUT matter curves and the details of GUT matter Yukawa couplings. Several other issues, by contrast, can only be addressed in a satisfactory manner in the context of a globally defined compactification. This is true in particular for all aspects of $U(1)$ symmetries - including $SU(5)$ GUT breaking via hypercharge flux and abelian selection rules - and the physics of GUT singlets (e.g. certain aspects of neutrino physics), which are localised away from the GUT divisor.

It is beyond the scope of these lectures to survey all exciting aspects of F-theory GUT model building that have emerged recently; rather we will outline some of the general philosophy behind the construction of $SU(5)$ GUT models. Armed with this background the interested reader can easily delve into more advanced topics.

4.1 $SU(5)$ GUT models and the principle of decoupling

$SU(5)$ is the mother of all GUT groups. In Georgi-Glashow $SU(5)$ models [67], the embedding of the MSSM gauge group $SU(3) \times SU(2) \times U(1)_Y$ rests on the identification of the $U(1)_Y$ generator with the Cartan generator $\text{diag}(2, 2, 2, -3, -3)$ within $SU(5)$. The MSSM matter is organised into $SU(5)$ multiplets as

$$\begin{aligned} \mathbf{10} &\leftrightarrow (Q_L, u_R^c, e_R^c), & \bar{\mathbf{5}}_{\mathbf{m}} &\leftrightarrow (d_R^c, L), & \mathbf{1} &\leftrightarrow \nu_R^c, \\ \mathbf{5}_{\mathbf{H}} &\leftrightarrow (T_u, H_u), & \bar{\mathbf{5}}_{\mathbf{H}} &\leftrightarrow (T_d, H_d). \end{aligned} \quad (101)$$

The triplets T_u, T_d , which are not present in the MSSM, must receive high-scale masses via doublet-triplet splitting. An alternative $SU(5)$ GUT scenario called flipped $SU(5)$ [68, 69] starts from $SU(5) \times U(1)_X$ and the MSSM matter is related to the identifications (101) by "flipping" $e_R^c \leftrightarrow \nu_R^c$ and $d_R^c \leftrightarrow u_R^c$. Since $U(1)_Y$ arises as a combination of $U(1)_X$ and the Cartan generator $\text{diag}(2, 2, 2, -3, -3)$ of $SU(5)$, flipped $SU(5)$ is a unified model only if $SU(5) \times U(1)_X$ is itself embedded into a higher group such as $SO(10)$. For definiteness we focus for now on Georgi-Glashow $SU(5)$ models.

The geometric origin [6–9] of the $SU(5)$ GUT gauge group, the matter representations and the Yukawa couplings in terms of codimension one, two and three singularities of the elliptic fibration has been provided in sections 3.1 and 3.3, see table 2. Matter charged under $SU(5)$ localises on the curves $P_{\mathbf{10}}$ and $P_{\mathbf{5}}$ on the GUT divisor S , while the role of ν_R^c can be played by any GUT singlets with a suitable $\mathbf{5}_{\mathbf{H}} \bar{\mathbf{5}}_{\mathbf{m}} \mathbf{1}$ coupling. Note that for generic $SU(5)$ geometries, the matter curve for the $\mathbf{5}$ representation is a single connected object; in this situation all three generations of $\bar{\mathbf{5}}_{\mathbf{m}}$ and the vector-like pair $\mathbf{5}_{\mathbf{H}} + \bar{\mathbf{5}}_{\mathbf{H}}$ are localised on the same curve. We will see later that this is unacceptable for phenomenological reasons and the model must be further refined.

The MSSM Yukawa couplings follow from the $SU(5)$ GUT interactions by decomposition of the $SU(5)$ representations as

$$\mathbf{10} \mathbf{10} \mathbf{5}_{\mathbf{H}} \longrightarrow Q_L u_R^c H_u, \quad \mathbf{10} \bar{\mathbf{5}}_{\mathbf{m}} \bar{\mathbf{5}}_{\mathbf{H}} \longrightarrow L e_R^c H_d + Q_L d_R^c H_d. \quad (102)$$

The $\mathbf{10\ 10\ 5_H}$ Yukawa coupling localises at a point of E_6 singularity, which is a strong coupling phenomenon. It is for the sake of this coupling that exceptional symmetry is essential. In perturbative Type II orientifolds this interaction is forbidden by global $U(1)$ selection rules and can only be generated by D-brane instantons [70]. The natural presence of this crucial coupling, from which the up-quark masses descend, is the prime motivation to pursue F-theory as the framework for GUT models with branes; in perturbative models, the volume of D-brane instanton generating $\mathbf{10\ 10\ 5_H}$ Yukawa coupling would have to be tuned so as to prevent too large suppression. On the other hand, in flipped $SU(5)$ models, where $\mathbf{10\ 10\ 5_H}$ gives rise to down quark masses, such a non-perturbative suppression can be a welcome rationale to argue for the hierarchy between the top and bottom quark mass.

An important question concerns the nature of the GUT brane S , which must be a Kähler surface embedded into the base space B as a holomorphic divisor. In full generality no waterproof restrictions on S can be given other than it had better support one of the GUT breaking mechanisms which will be discussed in section 4.2. A reasonable, though not strictly necessary organising principle, however, is to require the existence of a well-defined decoupling limit for gravity [8]. This paradigm is inspired by the separation of the four-dimensional GUT scale $M_{\text{GUT}} = 10^{16}$ GeV and the Planck scale $M_{\text{Pl.}} = 10^{19}$ GeV as well as by UV completeness of GUT models.

The four-dimensional Planck scale arises from dimensional reduction of the Einstein-Hilbert term in Einstein frame

$$S = M_*^8 \int_{\mathbb{R}^{1,3} \times B} \sqrt{-g} R \quad \Rightarrow \quad M_{\text{Pl.}}^2 = M_*^8 \text{Vol}(B). \quad (103)$$

Here M_* is the M-theory fundamental length scale inherited via M/F -theory duality and can be viewed, in the IIB limit, as the value of ℓ_s^{-1} in the Einstein frame. On the other hand, the GUT scale, determined by the breaking scale of $SU(5)$ down to the Standard Model gauge group, is parametrically given by the volume of the GUT divisor S ,

$$M_{\text{GUT}}^4 \simeq \text{Vol}^{-1}(S). \quad (104)$$

For example in the context of GUT breaking via hypercharge flux as discussed in section 4.2 this approximate relation arises because the volume of S sets the flux induced mass of the XY gauge bosons within $SU(5)$; after all these states propagate along the whole divisor S . The observed hierarchy of about 10^{-3} between M_{GUT} and $M_{\text{Pl.}}$ translates into a small hierarchy between the typical radii of the GUT brane and the six-dimensional F-theory base of approximately

$$\underbrace{\ell_s}_{0.2x} < \underbrace{R_S}_{2.2x} < \underbrace{R_B}_{5.6x}, \quad x = 10^{-16} \text{ GeV}^{-1}. \quad (105)$$

The relation to ℓ_s has been chosen in such a way that in addition

$$\alpha_{\text{GUT}}^{-1} = M_*^4 \text{Vol}(S) \simeq 24. \quad (106)$$

This equality in turn follows parametrically by reduction of the eight-dimensional Yang-Mills action $S_{YM} = M_*^4 \int_{\mathbb{R}^{1,3} \times S} F^2$.

Note that for phenomenologically viable models it is enough to stabilise the moduli in agreement with the crude estimate of (105). *A fortiori*, it is often postulated [8,9,71] that the GUT brane S allow for a limit

$$\text{Vol}(B) \rightarrow \infty, \quad \text{Vol}(S) \text{ finite} \quad (107)$$

such as to decouple gravity completely - at least in principle. Alternatively to this physical decoupling limit one can consider the mathematical decoupling limit [71]

$$\text{Vol}(S) \rightarrow 0, \quad \text{Vol}(B) \text{ finite}, \quad (108)$$

even though the two are not completely equivalent [44]. This mathematical decoupling limit can be taken if the surface S is Fano, which amounts to requiring that $\int_C K_S^{-1} > 0$ for every holomorphic curve C in S . The list of such Fano surfaces is very restrictive and consists of the del Pezzo surfaces $\mathbb{P}^1 \times \mathbb{P}^1$, \mathbb{P}^2 and dP_r , $r = 1, \dots, 8$. The latter are defined as \mathbb{P}^2 with r points in generic position blown up to \mathbb{P}^1 . Since these surfaces are used extensively in the F-theory GUT literature, we briefly collect some of their basic topological properties: The second homology $H_2(dP_r, \mathbb{Z})$ is spanned by the elements l, E_1, \dots, E_r , where E_i denote the i -th blow-up \mathbb{P}^1 and l the hyperplane class inherited from \mathbb{P}^2 . The non-vanishing intersection numbers are $l \cdot l = 1$, $E_i \cdot E_j = -\delta_{ij}$. The anti-canonical bundle is given, by slight abuse of notation, by $c_1(K_{dP_r}^{-1}) = c_1(dP_r) = 3l - \sum_{i=1}^r E_i$ with $c_1(dP_r)^2 = 9 - r$ and $c_2(dP_r) = 3 + r$.

Generic Fano surfaces are shrinkable in the sense of (108). At closer inspection, however, the existence of suitable matter curves and possible global embeddings in $SU(5)$ models poses certain restrictions on the shrinkability of the GUT divisor and forbids shrinkability to a point in generic situations. Rather the GUT brane can only shrink to a curve or to a point in such a way that another divisor shrinks simultaneously. For more information on these restrictions we refer to [44, 71].

4.2 Options for GUT breaking

Let us now discuss possible ways to break $SU(5)$ to the observed $SU(3) \times SU(2) \times U(1)_Y$ MSSM gauge group. As in all constructions with D-branes there exist three options to accomplish this:

- **Via a GUT Higgs** (scalar field) in the adjoint representation **24** of $SU(5)$. This effectively realises the GUT breaking mechanism of conventional field-theoretic GUT models. The string theoretic origin of the Higgs would be a brane deformation modulus counted by $H^{(0)}(S, K_S)$. An obvious challenge associated with this approach is the generation of a suitable potential for the GUT Higgs field which leads to dynamical symmetry breaking. In principle 7-brane deformation moduli are stabilised by background fluxes, but no concrete setup has been described so far that incorporates dynamical GUT breaking.
- **Via Wilson line moduli.** These correspond to elements of $H^{(0,1)}(S)$ and likewise transform in the adjoint of $SU(5)$. The same remarks concerning the generation of a potential for dynamical symmetry breaking apply with the added complication that Wilson line moduli are not even sensitive to background fluxes. Alternatively one can consider GUT breaking by discrete Wilson lines. These are available whenever the brane S has a discrete, but non-trivial first homotopy group $\pi_1(S)$. The VEV of the discrete Wilson line is now topological and part of the defining data of the compactification. Note that this is exactly the same strategy as pursued in Calabi-Yau compactifications of the $E_8 \times E_8$ heterotic string with $SU(N)$ gauge bundles. In some sense this is the cleanest approach to GUT symmetry breaking. The implementation of F-theory models along GUT divisors with non-trivial $\pi_1(S)$ has not been achieved in the literature so far, but seems a promising avenue for future research.

- **Via hypercharge flux.** This corresponds to turning on non-trivial gauge flux F_Y associated with the hypercharge generator $T_Y = \text{diag}(2, 2, 2, -3, -3) \subset SU(5)$. As for Wilson lines the flux is part of the defining topological data of the compactification and circumvents the quest for the dynamical generation of a symmetry breaking scalar potential. This approach does not rest on any strong coupling effect and is equally possible for perturbative Type IIB models.

Note that the last two GUT breaking mechanisms are not available in conventional, four-dimensional field theoretic GUTs. Studying their possible consequences for the more detailed phenomenology of $SU(5)$ GUTs is therefore particularly interesting.

4.3 Some constraints from hypercharge flux

For definiteness we focus on the last GUT breaking mechanism. This is also the only possibility for del Pezzo surfaces, which have no geometric deformations and no Wilson line moduli. The use of hypercharge flux was suggested in the context of F-theory model building in [8, 9] and first realised in compact models with 7-branes within perturbative Type IIB orientifolds in [52]. In IIB language one simply embeds a non-trivial line bundle L_Y along the GUT brane into $SU(5)$ by identifying its generator with the above hypercharge generator T_Y . In F/M-theory language this corresponds to G_4 flux of the form $G_4 = F_Y \wedge \omega_Y$. Here ω_Y is the 2-form dual to the \mathbb{P}^1 in the fiber over S that corresponds to the node in the $SU(5)$ Dynkin diagram associated with the Cartan generator T_Y .

In presence of $U(1)_Y$ gauge flux the GUT matter decomposes into representations of $SU(3) \times SU(2) \times U(1)_Y$ as

$$\begin{aligned}
\mathbf{24} &\mapsto (\mathbf{8}, \mathbf{1})_{0_Y} + (\mathbf{1}, \mathbf{3})_{0_Y} + (\mathbf{1}, \mathbf{1})_{0_Y} + (\mathbf{3}, \mathbf{2})_{5_Y} + (\bar{\mathbf{3}}, \mathbf{2})_{-5_Y}, \\
\bar{\mathbf{5}} &\mapsto (\bar{\mathbf{3}}, \mathbf{1})_{2_Y} + (\mathbf{1}, \mathbf{2})_{-3_Y}, \\
\mathbf{10} &\mapsto (\mathbf{3}, \mathbf{2})_{1_Y} + (\bar{\mathbf{3}}, \mathbf{1})_{-4_Y} + (\mathbf{1}, \mathbf{1})_{6_Y}, \\
\mathbf{5}_H &\mapsto (\mathbf{3}, \mathbf{1})_{-2_Y} + (\mathbf{1}, \mathbf{2})_{3_Y}, \quad \bar{\mathbf{5}}_H \mapsto (\bar{\mathbf{3}}, \mathbf{1})_{2_Y} + (\mathbf{1}, \mathbf{2})_{-3_Y}.
\end{aligned} \tag{109}$$

The cohomology classes counting the MSSM matter contain factors of L_Y^q with q the $U(1)_Y$ charge of the state. In order to guarantee the same number of states within each $SU(5)$ family, the net $U(1)_Y$ flux through each matter curve must therefore vanish [8, 9]. As we will discuss in section 4.4, absence of dimension 4 proton decay operators requires that the $\mathbf{5}$ matter curve $P_{\mathbf{5}}$ split into a $\bar{\mathbf{5}}_{\mathbf{m}}$ matter curve and a Higgs curve. Then what we need is that

$$c_1(L_Y) \cdot P_{\mathbf{10}} = 0 = c_1(L_Y) \cdot P_{\bar{\mathbf{5}}_{\mathbf{m}}}. \tag{110}$$

On the other hand, if in addition the Higgs curve splits into two curves P_{H_u} and P_{H_d} , non-zero hypercharge flux through these Higgs curves allows for an elegant solution to the doublet-triplet splitting problem if it is chosen such as to project out the massless $(\mathbf{3}, \mathbf{1})_{2_Y}$ within $\bar{\mathbf{5}}_{\mathbf{H}}$ [8].

Three challenges have to be met for constructions with hypercharge flux breaking, two of which can be solved successfully, while one of them remains as an open question of intense current debate.

Massless $U(1)_Y$

$U(1)_Y$ must remain massless after GUT symmetry breaking. In the language of IIB 7-branes with gauge flux, it is well-known that the Chern-Simons coupling $\int 8 \text{Tr} F \wedge \text{Tr} F \wedge \text{Tr} F$ leads to a Stückelberg

mass for $U(1)_Y$ by dimensional reduction of the RR four-form C_4 along 2-cycles on S ,

$$S_{\text{Stueckelberg}} \simeq \int_{\mathbb{R}^{1,3}} F_Y^{4D} \wedge c_2^{(i)} \text{tr} T_Y^2 \int_S c_1(L_Y) \wedge \iota^* \omega_i. \quad (111)$$

Here ω_i denotes a basis of $H^2(B, \mathbb{Z})$ and we have decomposed $C_4 = c_2^{(i)} \wedge \omega_i$ with $c_2^{(i)}$ denoting two-forms in four dimensions. A mass term for $U(1)_Y$ can only be avoided if the gauge flux F_Y is switched on exclusively along 2-cycles in S which are homologically trivial as two-cycles in the ambient geometry [51, 72]; in this case $c_1(L_Y)$ is orthogonal to $\iota^* H^2(B, \mathbb{Z})$. Cycles of this type are said to lie in the relative cohomology of S with respect to B . Note that to ensure this topological constraint one needs full control of the global compactification geometry. This is because the question of triviality of a 2-cycle on S can only be answered by studying an implicit embedding of S into a compact geometry.

Absence of massless exotics

The decomposition of the $SU(5)$ GUT matter displayed in (109) contains the fields $(\mathbf{3}, \mathbf{2})_{5_Y} + (\overline{\mathbf{3}}, \mathbf{2})_{-5_Y}$. From the point of view of the MSSM these are exotic matter states which must be absent at the massless level for phenomenological viability of the model. Since they descend from the adjoint of $SU(5)$ they correspond to modes propagating along the entire GUT divisor S and are therefore "bulk" states. From the discussion of bulk matter states in section 3.3 we recall that these states are counted by the cohomology groups of the hypercharge flux L_Y^q along S , where q is the $U(1)_Y$ charge. Absence of exotic states $(\mathbf{3}, \mathbf{2})_{5_Y} + (\overline{\mathbf{3}}, \mathbf{2})_{-5_Y}$ therefore requires that $H^i(S, L_Y^{\pm 5}) = 0$.

Vanishing cohomology for such a high power of line bundles is difficult to engineer. A way out is to admit a suitably fractional line bundle \mathcal{L}_Y instead of the integer quantised bundle L_Y [8]. A clean way to define this is by a twisting procedure that works both in perturbative Type IIB [52] and in the E_8 based F-theory models [33] discussed in these lectures. For such embeddings, the potential exotics are eventually counted by $H^i(S, \mathcal{L}_Y^{\pm 1})$, which therefore has to vanish. Here \mathcal{L}_Y denotes the hypercharge bundle in the twisted embedding. This constraint poses certain restrictions on the type of hypercharge flux switched on along the GUT brane. e.g. for a del Pezzo surface it can be achieved if and only if $c_1(\mathcal{L}_Y) = E_i - E_j$ for $i \neq j$ [8].

As a final remark we note that it is this absence of bulk exotics that cannot be achieved for an analogous breaking of the GUT group $SO(10)$ by internal fluxes [8]. This is the technical reason why $SO(10)$ GUT models have received less attention in the F-theory literature. Alternatively, $SO(10)$ can be broken to flipped $SU(5) \times U(1)_X$ via suitable fluxes, and subsequently the standard field theoretic GUT Higgs mechanism [69] can be invoked to break $SU(5) \times U(1)_X$ to the MSSM gauge group. For recent realisations of this scenario see [45, 73].

Gauge coupling unification

The most subtle and controversial challenge of the hypercharge GUT breaking scenario arises in the context of gauge coupling unification [9, 74]. The problem is that hypercharge flux along the GUT divisor seems to spoil the tree-level equality of the MSSM gauge couplings at the GUT scale. To see this one must obtain the gauge kinetic function f of the four-dimensional GUT gauge theory, defined as

$$S_{YM}^{(4D)} = \frac{1}{2} \text{Re}(f) \int_{\mathbb{R}^{1,3}} \text{tr} F \wedge \star F + \frac{1}{2} \text{Im}(f) \int_{\mathbb{R}^{1,3}} \text{tr} F \wedge F, \quad (112)$$

from the 7-brane effective action by dimensional reduction. This task is most straightforwardly accomplished again in Type IIB 7-brane language. The kinetic Yang-Mills and the topological Chern-Simons term follow from reduction of S_{DBI} and S_{CS} in equ. (8), respectively. The flux induced corrections to the leading order gauge kinetic function

$$f_S = \frac{1}{g_s} \frac{\text{Vol}_S}{\ell_s^4} + i \int_S C_4 \quad (113)$$

can be deduced from the contribution to $\text{Im}(f)$ encoded in the Cherns-Simons term proportional to $\int C_0 \wedge \text{tr} F^4$. To this end one takes into account holomorphicity of f together with the fact that C_0 and g_s combine into the holomorphic field $\tau = C_0 + \frac{i}{g_s}$. What is important is that the contribution from hypercharge flux differs for the three gauge couplings α_s , α_w , α_Y of the MSSM gauge groups and distorts the gauge coupling relations at the Kaluza-Klein scale. The exact relation depends on the precise group theoretic embedding, see [74] and [9] for two different types of embedding. In addition, threshold corrections from Kaluza-Klein states have to be taken into account [9]. Final agreement on the interpretation of the physical consequences for unification has not yet been achieved in the literature; preliminary results suggest that indeed new physics such as extra thresholds below the GUT scale is required to reconcile this splitting of the gauge couplings with one-loop GUT unification. The concrete nature of these effects is model dependent. The minimal such threshold could be played by the unavoidable massive Higgs triplets [74], while more radical approaches consider incomplete multiplets of massive exotic matter [75, 76] (see also [77]).

4.4 Proton decay

A classic topic in GUT model building is proton stability. In particular, avoiding dimension 4 and 5 proton decay operators is key to the phenomenological viability of $SU(5)$ models [78, 79]. We therefore face the question if string compactifications - here those of F-theory - add new ingredients to achieve these requirements. On the one hand, we can seek for stringy realisations of known field theoretic mechanisms such as favourable symmetries. In a more ambitious vein, new types of selection rules might become available in string theory that have no obvious four-dimensional counterpart.

Let us begin with dimension 4 proton decay. Dangerous MSSM operators of the type $u_R^c d_R^c d_R^c$, $LL e_R^c$, $QL d_R^c$ descend from the coupling $\mathbf{10} \bar{\mathbf{5}}_m \bar{\mathbf{5}}_m$, which must therefore be prevented. The same holds for its unwanted cousin $\mathbf{10} \bar{\mathbf{5}}_H \bar{\mathbf{5}}_H$. It was noted already in [7] that a necessary condition for absence of $\mathbf{10} \bar{\mathbf{5}}_m \bar{\mathbf{5}}_m$ while allowing at the same time for the Yukawa couplings $\mathbf{10} \bar{\mathbf{5}}_m \bar{\mathbf{5}}_H$ and $\mathbf{10} \mathbf{10} \bar{\mathbf{5}}_H$ is that the $\mathbf{5}$ matter curve splits into (at least) two curves P_m for $\bar{\mathbf{5}}_m$ and P_H for $\mathbf{5}_H + \bar{\mathbf{5}}_H$. From the perspective of the Weierstrass model, the splitting of matter curves corresponds to a non-generic situation that requires the restriction of some of the complex structure moduli, more on that in a second. If this splitting were sufficient to prevent dimension 4 proton decay, it would furnish an example of a geometric selection rule.

As found in [61], however, typically P_m and P_H intersect at points without further singularity enhancements. If this happens, the wavefunction of $\bar{\mathbf{5}}_m$ and $\mathbf{5}_H + \bar{\mathbf{5}}_H$ obey the same boundary conditions and dangerous couplings of the type $\mathbf{10} \bar{\mathbf{5}}_m \bar{\mathbf{5}}_m$ are unavoidable. To be on the safe side, explicit field theoretic selection rules have to be implemented. The minimal such selection rule - R-parity - could descend from a geometric discrete \mathbb{Z}_2 symmetry that acts appropriately on the massless modes [61], but no concrete realisations of this idea have been constructed as of this writing. More radically, a (massive) $U(1)$ selection symmetry can forbid

unwanted Yukawa couplings [61]. The maybe simplest example of such a $U(1)$ symmetry is $U(1)_X$ with charge assignments [75]

$$\mathbf{10}_1, \quad (\bar{\mathbf{5}}_m)_{-3}, \quad (\mathbf{5}_H)_{-2} + (\bar{\mathbf{5}}_H)_2, \quad (114)$$

but other examples such as a Peccei-Quinn type $U(1)_{PQ}$ [76] with different charge assignments for H_u and H_d or models with several $U(1)$ s [80] have also been considered.

At the level of model building the implementation of such abelian symmetries requires a further specification of the complex structure moduli of the Weierstrass model. For global Tate models this leads to so-called $U(1)$ restricted Tate models [37]. In particular the engineering of, say, a $U(1)_X$ symmetry automatically leads to a split of the $\bar{\mathbf{5}}_m$ and the Higgs curve. Note that as remarked before all questions associated with $U(1)$ symmetries defy a local treatment and require knowledge of the full compactification data of the 4-fold. This is intuitively clear because a VEV of a $U(1)$ charged GUT singlet localised away from the GUT brane S can higgs the abelian symmetry. Previously, the implementation of $U(1)$ symmetries had been studied via so-called split spectral covers [75]. Split spectral covers can be regarded as the restriction of $U(1)$ restricted Tate models to the neighbourhood of the GUT brane. As such they are not sufficient to guarantee the presence of $U(1)$ symmetries [37, 81].

Dimension 5 proton can in principle be prevented via a missing partner mechanism [7] if also $\mathbf{5}_H$ and $\bar{\mathbf{5}}_H$ localise on separate curves. One possibility to achieve this is in the context of $U(1)_{PQ}$ extended models [76]. Depending on details of the model, exotics in incomplete GUT multiplets can result as a side-effect, which in turn affects gauge coupling unification as outlined in the previous section.

4.5 Further developments

Our presentation of the phenomenological properties of F-theory GUTs has only covered the crudest aspects, and many more advanced phenomenological topics have been studied. For a review devoted specifically to the phenomenology of F-theory constructions and a more complete list of references we recommend [13]. Topics worth highlighting include these:

- The local nature of brane models offers the possibility of studying the structure of Yukawa couplings without referring to the details of the global geometry. Investigations of the Yukawa and flavour structure of $SU(5)$ GUT models in this context include [61, 82–91].
- Possible connections with neutrino physics are the subject of [92, 93].
- On a more formal level, instanton effects in F-theory have been reconsidered in the recent literature [94–97] with special attention to the generation of phenomenologically viable matter couplings known from the weakly coupled Type II limit (see e.g. [98] for a review and references).

Another focus in the recent literature is the realisation of these and other model building ideas in concrete compact examples. The motivation behind this is, as was stressed already, that certain questions of phenomenological relevance cannot be disentangled from the global geometric structure. F-theory GUT vacua have been constructed in [43–45, 75, 76, 99]. In particular, methods of toric geometry, as applied in [43–45], allow one to explicitly construct fully-fledged singular Calabi-Yau 4-folds and their explicit resolution in a way that keeps full control of the singularities of the Tate model. As far as the construction of gauge flux is concerned, the models [43–45, 75, 76, 99] rely on the spectral cover approach outlined in section

3.5. Modulo the caveats alluded to there, 3-generation GUT models that satisfy the known global consistency conditions such as 3-brane tadpole cancellation and D-term supersymmetry have been achieved.

We hope we could convince the reader that F-theory is a fascinating subject that has a lot to offer both to phenomenologically-minded string theorists and to those with more formal interests. Many open questions in both directions are awaiting a satisfactory solution in the future, maybe by some participants of this winter school.

Acknowledgements

I would like to thank the organisers of the CERN Winter School on Supergravity, Strings and Gauge Theory 2010 for the invitation to lecture at this nice meeting, and the participants for their interest and their valuable questions and remarks. I am indebted to my collaborators over past couple of years who have been sharing my interest in F-theory model building: R. Blumenhagen, V. Braun, B. Jurke, T. Grimm and S. Krause. Special thanks to M. Kerstan and S. Krause for useful comments on a draft of these lecture notes. Finally I am grateful to the Max-Planck-Institute in Munich, the KITP Santa Barbara and the Ecole Normale, Paris, for hospitality during parts of this work. This project was supported in part by the SFB-Transregio 33 “The Dark Universe” by the DFG and the National Science Foundation under Grant No. PHY05-51164.

References

- [1] H. P. Nilles, S. Ramos-Sanchez, M. Ratz, and P. K. S. Vaudrevange, “From strings to the MSSM,” *Eur. Phys. J.* **C59** (2009) 249–267, 0806.3905.
- [2] D. Lust, “Intersecting brane worlds: A path to the standard model?,” *Class. Quant. Grav.* **21** (2004) S1399–1424, hep-th/0401156.
- [3] R. Blumenhagen, M. Cvetič, P. Langacker, and G. Shiu, “Toward realistic intersecting D-brane models,” *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 71–139, hep-th/0502005.
- [4] R. Blumenhagen, B. Kors, D. Lust, and S. Stieberger, “Four-dimensional String Compactifications with D-Branes, Orientifolds and Fluxes,” *Phys. Rept.* **445** (2007) 1–193, hep-th/0610327.
- [5] C. Vafa, “Evidence for F-Theory,” *Nucl. Phys.* **B469** (1996) 403–418, hep-th/9602022.
- [6] R. Donagi and M. Wijnholt, “Model Building with F-Theory,” 0802.2969.
- [7] C. Beasley, J. J. Heckman, and C. Vafa, “GUTs and Exceptional Branes in F-theory - I,” *JHEP* **01** (2009) 058, 0802.3391.
- [8] C. Beasley, J. J. Heckman, and C. Vafa, “GUTs and Exceptional Branes in F-theory - II: Experimental Predictions,” *JHEP* **01** (2009) 059, 0806.0102.
- [9] R. Donagi and M. Wijnholt, “Breaking GUT Groups in F-Theory,” 0808.2223.
- [10] A. A. Tseytlin, “Self-duality of Born-Infeld action and Dirichlet 3-brane of type IIB superstring theory,” *Nucl. Phys.* **B469** (1996) 51–67, hep-th/9602064.
- [11] C. M. Hull, “String Dynamics at Strong Coupling,” *Nucl. Phys.* **B468** (1996) 113–154, hep-th/9512181.
- [12] F. Denef, “Les Houches Lectures on Constructing String Vacua,” 0803.1194.
- [13] J. J. Heckman, “Particle Physics Implications of F-theory,” 1001.0577.
- [14] G. T. Horowitz and A. Strominger, “Black strings and P-branes,” *Nucl. Phys.* **B 360** (1991).
- [15] B. R. Greene, A. D. Shapere, C. Vafa, and S.-T. Yau, “Stringy Cosmic Strings and Noncompact Calabi-Yau Manifolds,” *Nucl. Phys.* **B337** (1990) 1.
- [16] M. Nakahara, “Geometry, topology and physics,”. Boca Raton, USA: Taylor and Francis (2003) 573 p.
- [17] A. P. Braun, A. Hebecker, and H. Triendl, “D7-Brane Motion from M-Theory Cycles and Obstructions in the Weak Coupling Limit,” *Nucl. Phys.* **B800** (2008) 298–329, 0801.2163.
- [18] J. H. Schwarz, “An $SL(2,Z)$ multiplet of type IIB superstrings,” *Phys. Lett.* **B360** (1995) 13–18, hep-th/9508143.

- [19] E. Witten, “Bound states of strings and p-branes,” *Nucl. Phys.* **B460** (1996) 335–350, [hep-th/9510135](#).
- [20] A. Sen, “F-theory and Orientifolds,” *Nucl. Phys.* **B475** (1996) 562–578, [hep-th/9605150](#).
- [21] E. Witten, “Non-Perturbative Superpotentials In String Theory,” *Nucl. Phys.* **B474** (1996) 343–360, [hep-th/9604030](#).
- [22] K. Becker and M. Becker, “M-Theory on Eight-Manifolds,” *Nucl. Phys.* **B477** (1996) 155–167, [hep-th/9605053](#).
- [23] K. Dasgupta, G. Rajesh, and S. Sethi, “M theory, orientifolds and G-flux,” *JHEP* **08** (1999) 023, [hep-th/9908088](#).
- [24] M. Haack and J. Louis, “M-theory compactified on Calabi-Yau fourfolds with background flux,” *Phys. Lett.* **B507** (2001) 296–304, [hep-th/0103068](#).
- [25] T. W. Grimm, “The N=1 effective action of F-theory compactifications,” [1008.4133](#).
- [26] A. Klemm, B. Lian, S. S. Roan, and S.-T. Yau, “Calabi-Yau fourfolds for M- and F-theory compactifications,” *Nucl. Phys.* **B518** (1998) 515–574, [hep-th/9701023](#).
- [27] P. Deligne, “Courbes elliptiques ,” *Lecture Notes in Math., Springer, Berlin* **476** (1975) 53–75.
- [28] K. Kodaira, “On compact analytic surfaces,” *Annals of Math.* **77** (1963) 563.
- [29] D. R. Morrison and C. Vafa, “Compactifications of F-Theory on Calabi-Yau Threefolds – I,” *Nucl. Phys.* **B473** (1996) 74–92, [hep-th/9602114](#).
- [30] K. Dasgupta and S. Mukhi, “F-theory at constant coupling,” *Phys. Lett.* **B385** (1996) 125–131, [hep-th/9606044](#).
- [31] A. Sen, “Orientifold limit of F-theory vacua,” *Phys. Rev.* **D55** (1997) 7345–7349, [hep-th/9702165](#).
- [32] A. Collinucci, “New F-theory lifts,” [0812.0175](#).
- [33] R. Blumenhagen, T. W. Grimm, B. Jurke, and T. Weigand, “F-theory uplifts and GUTs,” [0906.0013](#).
- [34] A. Collinucci, “New F-theory lifts II: Permutation orientifolds and enhanced singularities,” [0906.0003](#).
- [35] A. Collinucci, F. Denef, and M. Esole, “D-brane Deconstructions in IIB Orientifolds,” *JHEP* **02** (2009) 005, [0805.1573](#).
- [36] P. Aluffi and M. Esole, “New Orientifold Weak Coupling Limits in F-theory,” *JHEP* **02** (2010) 020, [0908.1572](#).
- [37] T. W. Grimm and T. Weigand, “On Abelian Gauge Symmetries and Proton Decay in Global F- theory GUTs,” [1006.0226](#).

- [38] M. Bershadsky *et al.*, “Geometric singularities and enhanced gauge symmetries,” *Nucl. Phys.* **B481** (1996) 215–252, [hep-th/9605200](#).
- [39] S. H. Katz and C. Vafa, “Geometric engineering of $N = 1$ quantum field theories,” *Nucl. Phys.* **B497** (1997) 196–204, [hep-th/9611090](#).
- [40] D. R. Morrison and C. Vafa, “Compactifications of F-Theory on Calabi–Yau Threefolds – II,” *Nucl. Phys.* **B476** (1996) 437–469, [hep-th/9603161](#).
- [41] P. Candelas and A. Font, “Duality between the webs of heterotic and type II vacua,” *Nucl. Phys.* **B511** (1998) 295–325, [hep-th/9603170](#).
- [42] P. Candelas, E. Perevalov, and G. Rajesh, “Toric geometry and enhanced gauge symmetry of F- theory/heterotic vacua,” *Nucl. Phys.* **B507** (1997) 445–474, [hep-th/9704097](#).
- [43] R. Blumenhagen, T. W. Grimm, B. Jurke, and T. Weigand, “Global F-theory GUTs,” [0908.1784](#).
- [44] T. W. Grimm, S. Krause, and T. Weigand, “F-Theory GUT Vacua on Compact Calabi-Yau Fourfolds,” *JHEP* **07** (2010) 037, [0912.3524](#).
- [45] C.-M. Chen, J. Knapp, M. Kreuzer, and C. Mayrhofer, “Global $SO(10)$ F-theory GUTs,” [1005.5735](#).
- [46] M. R. Gaberdiel and B. Zwiebach, “Exceptional groups from open strings,” *Nucl. Phys.* **B518** (1998) 151–172, [hep-th/9709013](#).
- [47] O. DeWolfe and B. Zwiebach, “String junctions for arbitrary Lie algebra representations,” *Nucl. Phys.* **B541** (1999) 509–565, [hep-th/9804210](#).
- [48] S. Sethi, C. Vafa, and E. Witten, “Constraints on low-dimensional string compactifications,” *Nucl. Phys.* **B480** (1996) 213–224, [hep-th/9606122](#).
- [49] B. Andreas and G. Curio, “On discrete twist and four-flux in $N = 1$ heterotic/F-theory compactifications,” *Adv. Theor. Math. Phys.* **3** (1999) 1325–1413, [hep-th/9908193](#).
- [50] B. Andreas and G. Curio, “From Local to Global in F-Theory Model Building,” [0902.4143](#).
- [51] H. Jockers and J. Louis, “The effective action of D7-branes in $N = 1$ Calabi-Yau orientifolds,” *Nucl. Phys.* **B705** (2005) 167–211, [hep-th/0409098](#).
- [52] R. Blumenhagen, V. Braun, T. W. Grimm, and T. Weigand, “GUTs in Type IIB Orientifold Compactifications,” *Nucl. Phys.* **B815** (2009) 1–94, [0811.2936](#).
- [53] S. H. Katz and E. Sharpe, “D-branes, open string vertex operators, and Ext groups,” *Adv. Theor. Math. Phys.* **6** (2003) 979–1030, [hep-th/0208104](#).
- [54] S. H. Katz and C. Vafa, “Matter from geometry,” *Nucl. Phys.* **B497** (1997) 146–154, [hep-th/9606086](#).
- [55] A. Grassi *Internat. J. Math.* **4** (1993) 203.

- [56] R. Friedman, J. Morgan, and E. Witten, “Vector bundles and F theory,” *Commun. Math. Phys.* **187** (1997) 679–743, [hep-th/9701162](#).
- [57] R. Donagi, “Principal bundles on elliptic fibrations,” *Asian J. Math.* **1** (1997) 214–223, [alg-geom/9702002](#).
- [58] B. Andreas and D. Hernandez Ruiperez, “U(n) vector bundles on Calabi-Yau threefolds for string theory compactifications,” *Adv. Theor. Math. Phys.* **9** (2005) 253–284, [hep-th/0410170](#).
- [59] G. Curio, “Chiral matter and transitions in heterotic string models,” *Phys. Lett.* **B435** (1998) 39–48, [hep-th/9803224](#).
- [60] G. Curio and R. Y. Donagi, “Moduli in N = 1 heterotic/F-theory duality,” *Nucl. Phys.* **B518** (1998) 603–631, [hep-th/9801057](#).
- [61] H. Hayashi, T. Kawano, R. Tatar, and T. Watari, “Codimension-3 Singularities and Yukawa Couplings in F- theory,” *Nucl. Phys.* **B823** (2009) 47–115, [0901.4941](#).
- [62] R. Donagi and M. Wijnholt, “Higgs Bundles and UV Completion in F-Theory,” [0904.1218](#).
- [63] R. Donagi, Y.-H. He, B. A. Ovrut, and R. Reinbacher, “The particle spectrum of heterotic compactifications,” *JHEP* **12** (2004) 054, [hep-th/0405014](#).
- [64] R. Blumenhagen, S. Moster, R. Reinbacher, and T. Weigand, “Massless spectra of three generation U(N) heterotic string vacua,” *JHEP* **05** (2007) 041, [hep-th/0612039](#).
- [65] H. Hayashi, R. Tatar, Y. Toda, T. Watari, and M. Yamazaki, “New Aspects of Heterotic/F-Theory Duality,” *Nucl. Phys.* **B806** (2009) 224–299, [0805.1057](#).
- [66] J. Marsano, N. Saulina, and S. Schafer-Nameki, “A Note on G-Fluxes for F-theory Model Building,” [1006.0483](#).
- [67] H. Georgi and S. L. Glashow, “Unity of All Elementary Particle Forces,” *Phys. Rev. Lett.* **32** (1974) 438–441.
- [68] S. M. Barr, “A new symmetry breaking pattern for SO(10) and proton decay,” *Phys. Lett.* **B112** (1982) 219.
- [69] I. Antoniadis, J. R. Ellis, J. S. Hagelin, and D. V. Nanopoulos, “Supersymmetric Flipped SU(5) Revitalized,” *Phys. Lett.* **B194** (1987) 231.
- [70] R. Blumenhagen, M. Cvetič, D. Lust, R. Richter, and T. Weigand, “Non-perturbative Yukawa Couplings from String Instantons,” *Phys. Rev. Lett.* **100** (2008) 061602, [0707.1871](#).
- [71] C. Cordova, “Decoupling Gravity in F-Theory,” [0910.2955](#).
- [72] M. Buican, D. Malyshev, D. R. Morrison, H. Verlinde, and M. Wijnholt, “D-branes at singularities, compactification, and hypercharge,” *JHEP* **01** (2007) 107, [hep-th/0610007](#).

- [73] C.-M. Chen and Y.-C. Chung, “Flipped SU(5) GUTs from E_8 Singularity in F-theory,” 1005.5728.
- [74] R. Blumenhagen, “Gauge Coupling Unification in F-Theory Grand Unified Theories,” *Phys. Rev. Lett.* **102** (2009) 071601, 0812.0248.
- [75] J. Marsano, N. Saulina, and S. Schafer-Nameki, “Monodromies, Fluxes, and Compact Three-Generation F-theory GUTs,” 0906.4672.
- [76] J. Marsano, N. Saulina, and S. Schafer-Nameki, “Compact F-theory GUTs with $U(1)_{PQ}$,” 0912.0272.
- [77] G. K. Leontaris and N. D. Tracas, “Gauge coupling flux thresholds, exotic matter and the unification scale in F-SU(5) GUT,” *Eur. Phys. J.* **C67** (2010) 489–498, 0912.1557.
- [78] H. Murayama and A. Pierce, “Not even decoupling can save minimal supersymmetric SU(5),” *Phys. Rev.* **D65** (2002) 055009, hep-ph/0108104.
- [79] S. Raby, “Proton decay,” hep-ph/0211024.
- [80] E. Dudas and E. Palti, “On hypercharge flux and exotics in F-theory GUTs,” 1007.1297.
- [81] H. Hayashi, T. Kawano, Y. Tsuchiya, and T. Watari, “More on Dimension-4 Proton Decay Problem in F-theory – Spectral Surface, Discriminant Locus and Monodromy,” 1004.3870.
- [82] A. Font and L. E. Ibanez, “Yukawa Structure from U(1) Fluxes in F-theory Grand Unification,” *JHEP* **02** (2009) 016, 0811.2157.
- [83] J. J. Heckman and C. Vafa, “Flavor Hierarchy From F-theory,” 0811.2417.
- [84] J. J. Heckman, A. Tavanfar, and C. Vafa, “The Point of E_8 in F-theory GUTs,” 0906.0581.
- [85] A. Font and L. E. Ibanez, “Matter wave functions and Yukawa couplings in F-theory Grand Unification,” 0907.4895.
- [86] S. Cecotti, M. C. N. Cheng, J. J. Heckman, and C. Vafa, “Yukawa Couplings in F-theory and Non-Commutative Geometry,” 0910.0477.
- [87] J. P. Conlon and E. Palti, “Aspects of Flavour and Supersymmetry in F-theory GUTs,” *JHEP* **01** (2010) 029, 0910.2413.
- [88] H. Hayashi, T. Kawano, Y. Tsuchiya, and T. Watari, “Flavor Structure in F-theory Compactifications,” 0910.2762.
- [89] F. Marchesano and L. Martucci, “Non-perturbative effects on seven-brane Yukawa couplings,” *Phys. Rev. Lett.* **104** (2010) 231601, 0910.5496.
- [90] E. Dudas and E. Palti, “Froggatt-Nielsen models from E_8 in F-theory GUTs,” 0912.0853.

- [91] S. F. King, G. K. Leontaris, and G. G. Ross, “Family symmetries in F-theory GUTs,” *Nucl. Phys.* **B838** (2010) 119–135, 1005.1025.
- [92] V. Bouchard, J. J. Heckman, J. Seo, and C. Vafa, “F-theory and Neutrinos: Kaluza-Klein Dilution of Flavor Hierarchy,” *JHEP* **01** (2010) 061, 0904.1419.
- [93] R. Tatar, Y. Tsuchiya, and T. Watari, “Right-handed Neutrinos in F-theory Compactifications,” 0905.2289.
- [94] R. Blumenhagen, A. Collinucci, and B. Jurke, “On Instanton Effects in F-theory,” 1002.1894.
- [95] M. Cvetič, I. Garcia-Etxebarria, and R. Richter, “Branes and instantons at angles and the F-theory lift of O(1) instantons,” *AIP Conf. Proc.* **1200** (2010) 246–260, 0911.0012.
- [96] M. Cvetič, I. Garcia-Etxebarria, and J. Halverson, “Global F-theory Models: Instantons and Gauge Dynamics,” 1003.5337.
- [97] R. Donagi and M. Wijnholt, “MSW Instantons,” 1005.5391.
- [98] R. Blumenhagen, M. Cvetič, S. Kachru, and T. Weigand, “D-Brane Instantons in Type II Orientifolds,” *Ann. Rev. Nucl. Part. Sci.* **59** (2009) 269–296, 0902.3251.
- [99] J. Marsano, N. Saulina, and S. Schafer-Nameki, “F-theory Compactifications for Supersymmetric GUTs,” 0904.3932.