The axonal guidance factor Netrin-1 as a potential modulator of swine follicular function

G. Basini, C. Cortimiglia, L. Baioni, S. Bussolati, S. Grolli, R. Ramoni, F. Grasselli

► To cite this version:

G. Basini, C. Cortimiglia, L. Baioni, S. Bussolati, S. Grolli, et al.. The axonal guidance factor Netrin-1 as a potential modulator of swine follicular function. *Molecular and Cellular Endocrinology*, 2010, 10.1016/j.mce.2010.08.001 . hal-00639771

HAL Id: hal-00639771

<https://hal.science/hal-00639771>

Submitted on 10 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: The axonal guidance factor Netrin-1 as a potential modulator of swine follicular function

Authors: G. Basini, C. Cortimiglia, L. Baioni, S. Bussolati, S. Grolli, R. Ramoni, F. Grasselli

PII: S0303-7207(10)00415-6
DOI: doi:10.1016/j.mce.2010.08.001
Reference: MCE 7613

To appear in: *Molecular and Cellular Endocrinology*

Received date: 31-3-2010
Revised date: 2-8-2010
Accepted date: 2-8-2010

Please cite this article as: Basini, G., Cortimiglia, C., Baioni, L., Bussolati, S., Grolli, S., Ramoni, R., Grasselli, F., The axonal guidance factor Netrin-1 as a potential modulator of swine follicular function, *Molecular and Cellular Endocrinology* (2010), doi:10.1016/j.mce.2010.08.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The axonal guidance factor Netrin-1 as a potential
modulator of swine follicular function

G. Basini¹, C. Cortimiglia¹, L. Baioni¹, S. Bussolati¹, S. Grolli², R.
Ramoni², F. Grasselli¹

¹Sezione di Fisiologia Veterinaria, ²Sezione di Biochimica Veterinaria, Dipartimento di Produzioni
Animali, Biotecnologie Veterinarie, Qualità e Sicurezza degli Alimenti, Via del Taglio 10,
Università degli Studi di Parma, 43126 Parma, Italy

ABSTRACT

This study was aimed to improve knowledge about swine ovarian follicular function, paying attention to angiogenesis, since new vessel growth is a fundamental event in ovarian function. In particular, we investigated a potential involvement of netrin-1, a protein known as a guidance axon factor.

Firstly, we studied the expression and immunolocalization of netrin-1 in swine ovarian follicle and its effect on cultured swine granulosa cell viability and steroidogenesis. Furthermore, aortic endothelial cells were employed to verify a possible netrin-1 effect on angiogenesis.

Our data demonstrate the expression and the presence of netrin-1 in swine follicular fluid; in addition, it was shown that netrin-1 inhibits granulosa cell viability and estradiol 17 β levels while it stimulates progesterone production. Netrin-1 also inhibits aortic endothelial cell growth in the angiogenesis bioassay.

1 This effect appears to be mediated by inhibiting Vascular Endothelial Growth
2 Factor and stimulating Nitric Oxide.
3 Therefore, we hypothesize that netrin-1 could be important for follicular
4 function in the swine.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1 1. Introduction

2 Netrins are a family of highly conserved secreted proteins structurally related to
 3 laminins. They were named after the Sanskrit word "netr", which means "one
 4 who guides", since they possess guidance cues in axon pathfinding. Recent
 5 studies report the involvement of netrins in other developmental process,
 6 including the formation of vessels (Cirulli and Yebra 2007). Both neuronal and
 7 vascular development require guidance to establish a precise branching pattern
 8 and it has been shown that several molecules implicated in axonal navigation
 9 have a similar role in regulating vessel sprouting (Larrivée et al., 2009). In
 10 particular, netrin-1 has been related to the fine tuning of the angiogenic
 11 process and published data demonstrate pro- and anti-angiogenic effects (Park
 12 et al., 2004; Larrivée et al., 2007; Bouvrée K et al., 2008; Castets M et al.,
 13 2009). Maeda et al. (2008) recently demonstrated that netrin-1 is expressed in
 14 swine granulosa cells suggesting that the protein could be involved in the
 15 modulation of ovarian follicle function. In particular, netrin-1 could play a role in
 16 the regulation of the marked cyclical angiogenesis which takes place in the
 17 developing follicle and represents an invaluable physiological model for the
 18 investigation of the molecular events responsible for new vessel growth.

19 On these bases, the present study was undertaken to explore the hypothesis
 20 that netrin-1 could act as a physiological modulator of swine follicular function.

21 The expression and the immunolocalization of netrin-1 in swine ovarian follicle
 22 were explored and the effect of different protein concentrations on the main
 23 parameters of swine granulosa cell function were examined. Moreover, an

angiogenesis bioassay set up in our laboratory (Basini et al., 2008a) was employed to evaluate the effect of netrin-1 on new vessel growth.

2. Materials and Methods

All reagents used in this study were obtained from Sigma (St. Louis, MO, USA) unless otherwise specified.

2.1 Collection of ovaries

Swine ovaries were collected at a local slaughterhouse from Large White cross-bred gilts, placed into cold PBS (4°C) supplemented with penicillin (500 IU/ml), streptomycin (500 µg/ml) and amphotericin B (3.75 µg/ml), maintained in a freezer bag and transported to the laboratory within 1 h.

2.2 Netrin-1 in swine ovarian follicle

2.2.1 Immunohistochemistry

Ovaries were placed in 10% buffered formalin and paraffin embedded. Following routine processing, 6 µm-thick sections were placed on polylysine-treated slides and stained with a polyclonal antiserum raised against a peptide corresponding to 501-604 amino acid sequence of human netrin-1 (Santa Cruz Biotechnology, Santa Cruz, California). Briefly, sections were blocked in 10% normal swine serum for 20 min (200 µl) and then the primary antibody was

applied (250 µl of a solution diluted 1:100 in PBS) and incubated overnight at 4 °C. After rinsing in PBS, the slides were incubated for 15 minutes with 200 µl of a biotinylated secondary antibody. After three 5-min washes in PBS, 200 µl of a solution of streptavidin-peroxidase complex (LSAB®+SYSTEM-HRP, Dako, Glostrup, Denmark) was applied for 15 min. Positive reactions were revealed by aminoethylcarbazole (AEC) (Dako, Glostrup, Denmark). Sections were washed three times for 5 min with PBS and counterstained with Gill's ematoxylin for 2 min. Control sections were subjected to the same procedure, except that diluted swine serum replaced the primary antibody.

2.2.2 Western Blotting of follicular fluid protein content

Follicular fluid was collected with a 26-gauge from follicles >5mm (Basini et al., 2008b), centrifuged ($300 \times g$; 10 min) to separate rapidly the fluid from the cell fraction and refrigerated at -80 °C until assayed. As a positive control, a sample of soluble protein extract from porcine spinal marrow was also prepared. To this aim few grams of tissue were collected from a freshly slaughtered pig and homogenized by Ultra-Turrax (one stroke of 30 seconds at maximum speed) in 20 mM Tris/HCl pH 7.8 (1 ml of buffer/gram of tissue). The suspension was centrifuged (18,000Xg) for 5 minutes and the supernatant was stored -80 °C.

The protein pattern of follicular fluid and spinal marrow samples were resolved by 10% sodium dodecyl-sulphate polyacrylamide gel electrophoresis (SDS-PAGE) according to Laemmli (1970). After electrophoresis, the protein bands were transferred onto a ProBlott membrane (Applied Biosystem, Carlsbad,

1 California) (Towbin et al., 1979) that was blocked with 5% skimmed dry milk
 2 dissolved in 20 mm Tris / HCl buffer, pH 7.8 containing 150 mm NaCl, and
 3 0.01% (w / v) Tween 20 (TTBS). After washing with TTBS the membrane was
 4 incubated overnight with an anti-netrin-1 polyclonal antibody (Santa Cruz
 5 Biotechnology, Santa Cruz, California) raised in rabbit diluted (1:50) in TTBS
 6 containing 1% (w / v) skimmed dry milk. After three washes in TTBS the
 7 membrane was incubated 45 min. with a solution of a biotinilated anti-rabbit
 8 IgG antibody (LSAB®+SYSTEM-HRP, Dako, Glostrup, Denmark) diluted 1:5 in
 9 TTBS. A solution of complex of streptavidin-peroxidase was then applied for 15
 10 min. and positive reactions were revealed by AEC. A replicate SDS-PAGE gel
 11 was stained with Comassie blue.

12

13 *2.2.3 RNA extraction and RT-PCR*

14 Follicles >5 mm were dissected from the ovaries according to Basini et al.
 15 (2010). Each follicle was cut open, granulosa cells were separated by means of
 16 using a siliconized Pasteur pipette and centrifuged at $300 \times g$ for 10 min. Theca
 17 tissue was peeled from each follicle using a forceps, deep frozen in liquid
 18 nitrogen and powdered in a mortar.

19 Total RNA was extracted using Nucleospin ® RNA II (Macherey-Nagel GmbH,
 20 Duren, Germany) according to the manufacturer's instructions. Total RNA was
 21 quantified by absorbance at 260 nm (Gen Quant Pro, Amersham Biosciences,
 22 Freiburg, Germany). Total RNA (2 µg) was reverse transcribed with Ready-to-
 23 Go You Prime First-Strands Beads (Amersham Biosciences, Freiburg, Germany).

1 Netrin-1 cDNA was PCR amplified in parallel with β actin cDNA as an internal
 2 positive control. For Netrin-1, the sense primer (5'-GAG ACC TTC AAC ACG
 3 CCG-3') and antisense primer (5'-GGA AGG TGG ACA GCG AGG-3') (MWG
 4 Biotec, Ebersberg, Germany) were derived from Gene Bank sequence of porcine
 5 Netrin-1 (accession number DQ368597). Swine actin was amplified using the
 6 primers pACTIN sense (5'- GAG ACC TTC AAC ACG CCG-3') and pACTIN
 7 antisense (5'-GGA AGG TGG ACA GCG AGG-3') (MWG Biotec, Ebersberg,
 8 Germany). An aliquot (5 μ l) of the cDNA template was amplified by PCR using
 9 0.5 μ l (500 mU) Taq polymerase (Fermentas, Hannover, MD, USA) in 50 μ l PCR
 10 buffer containing 10 mM dNTP mix (Fermentas), 25 μ M for Netrin-1 primers
 11 and 5 μ M for Actin primers . Amplification was carried out using the thermal
 12 cycler PTC-100 Peltier (MJ Research, San Francisco, CA, USA). After an initial
 13 denaturation step for 5 min at 95°C, target cDNA was amplified for 3 cycles
 14 with a denaturation step at 95 °C for 30 sec, annealing at 60 °C for 30 sec (-
 15 1°C in the second and third cycle), elongation at 72 °C for 30 sec, and for 35
 16 cycles with a denaturation step at 95 °C for 30 sec, annealing at 57 °C for 30
 17 sec, elongation at 72 °C for 30 sec. Reactions were terminated with a final
 18 elongation at 72 °C for 10 min. The PCR products were separated on 1.5%
 19 agarose gel stained with ethidium bromide (0.5 μ g/ml) and visualized under UV
 20 light. Digital images were captured by Power Shot A610 photo camera (Canon,
 21 Tokyo, Japan). Quantification of band intensity was performed with NIH Image
 22 software (Scion Corporation, MA, USA, <http://rsb.info.nih.gov/nih-image/>).

23

2.3 Netrin-1 effects on swine granulosa cell functions

2.3.1 Granulosa cell culture

Granulosa cells were aseptically harvested by aspiration of follicles with a 26-gauge needle, released in medium containing heparin (50 IU/ml), centrifuged for pelleting and then treated with 0.9% prewarmed ammonium chloride at 37°C for 1 min to remove red blood cells. Cell number and viability were estimated using a haemocytometer under a phase contrast microscope after vital staining with trypan blue (0.4%) of an aliquot of the cell suspension. Cells were seeded in DMEM/Ham's F12 supplemented with sodium bicarbonate (2.2 mg/ml), bovine serum albumin (BSA 0.1%), penicillin (100 IU/ml), streptomycin (100 µg/ml), amphotericin B (2.5 µg/ml), selenium (5 ng/ml) and transferrin (5 µg/ml). Once seeded, cells were incubated for 48 h at 37 °C under humidified atmosphere (5% CO₂) in the presence or absence of Netrin-1 at the concentration of 5, 30 and 100 ng/ml. The protein was produced as reported by Serafini et al., 1994. Briefly, cDNAs encoding chick Netrin-1 tagged with a c-myc epitope at its C-terminus and fused to the constant (Fc) region of the human IgG1, were subcloned into the expression vector pCEP4 (Invitrogen, Carlsbad, California) and used to transfect 293-EBNA cells (Invitrogen, Carlsbad, California). Protein was finally purified from conditioned media by heparin affinity chromatography to 85-90% homogeneity, as assessed by silver staining (Keini-Masu et al., 1996).

1 *2.3.2 Granulosa cell viability*

2 2×10^5 cells were seeded in 96-well plates in 200 μ l CM. Cell proliferation was
 3 assayed using a bioluminescent assay (ATP-lites; Packard Bioscience,
 4 Groningen, Netherlands) which measured intracellular ATP levels. ATP, being
 5 present in all metabolically active cells, is a viability marker whose
 6 concentration declines very rapidly when the cells undergo necrosis or
 7 apoptosis. The ATP lite-M assay system is based on the production of light
 8 caused by the reaction of ATP with added luciferase and D-luciferin. The
 9 emitted light is proportional to the ATP concentration. Briefly, 50 μ l of
 10 mammalian cell lysis solution were added to 100 μ l of cell suspension and the
 11 plate was shaken for 5 minutes in an orbital shaker at 700 rpm in order to lyse
 12 the cells and stabilize ATP. Then 50 μ l of substrate solution were added to the
 13 wells and the microplate was shaken for 5 minutes in an orbital shaker at 700
 14 rpm. The plate was placed in the dark for 10 minutes and the luminescence
 15 was measured in a luminometer (Victor, Packard Bioscience, Groningen,
 16 Netherlands).

17

18 *2.3.3 Granulosa cell steroid production.*

19 10^4 cells/well were seeded in 96-well plates in 200 μ l of CM supplemented with
 20 androstenedione (28 ng/ml). Culture media were then collected, frozen and
 21 stored at -20°C until progesterone (P4) and estradiol 17β (E2) were
 22 determined by a validated Radio Immuno Assays (Grasselli et al., 1993). P4
 23 assay sensitivity and ED_{50} were 0.24 and 1 nmol/l, respectively; E2 assay

sensitivity and ED_{50} were 0.05 and 0.2 nmol/l. The intra- and inter-assay coefficients of variation were less than 12% for both assays.

2.4 Netrin-1 effects on swine aortic endothelial cell functions

2.4.1 Aortic endothelial cell culture

An immortalized porcine aortic endothelial cell line (AOC) (Carrillo et al. 2002) was generously provided by José Yelamos (Hospital Universitario Virgen de la Arrixaca, El Palmar, 30120 Murcia, Spain). Cells were cultured in M199 medium supplemented with sodium bicarbonate (2.2 mg/ml), FCS 20%, penicillin (100 IU/ml), streptomycin (100 µg/ml), amphotericin B (2.5 µg/ml). In all experiments, AOC at 19th passage were used.

2.4.2 Angiogenesis bioassay

The microcarrier-based fibrin gel angiogenesis assay was performed as described by Basini et al. (2008) with some modifications. Briefly, 12.5 mg gelatin-coated cytodex-3 microcarriers in 1 ml PBS were incubated for 3 h to hydrate. After two washings in PBS and one in CM, the microcarriers were put in flasks containing 5 ml CM; AOC (5×10^5) were added and cultured for 24 h in order to let the endothelial cells coat the microcarriers. For the fibrin gel preparation, 40 µl microcarriers covered by AOC were pipetted into 6 well plates containing a solution of fibrinogen (1mg/ml PBS, pH 7.6), added with 1250 IU thrombin (250 µl). Fibrin gels were allowed to polymerize for 30 min

at 37°C, and then were equilibrated for 60 min with 2 ml M199. After a change of the medium, AOC were incubated in the presence or absence of 5, 30 or 100 ng/ml of Netrin-1. Plates were incubated at 37°C under humidified atmosphere (5% CO₂). AOC were cultured for 96 h, renewing totally the treatment after 48h as described above.

2.4.3 Quantification of AOC growth on fibrin gel matrix

Endothelial cell proliferation in the fibrin gel matrix was evaluated by means of the public domain NIH Program Scion Image Beta 4.02. Ten pictures were taken for each gel at 48 and 96 h; images were converted into gray scale, resized to 50% (Paintbrush Software, MS Office) and saved as Bitmap 24bit format compatible with Scion. The modified images were then imported into the program and measurements were made drawing the perimeter of the area occupied by AOC expressed as number of pixel. In order to validate the measurement of the area covered by AOC in fibrin gels as a reliable method to evaluate cell proliferation, fibrin gels were stained by the nuclear dye bis-benzimide (Hoechst 33258, 20 µg/ml in PBS for 60 min) and examined by the fluorescence microscope. This procedure was performed 20 times; for each experiment the number of nuclei was counted under fluorescence and pictures of the area covered by AOC were taken in order to measure the surface covered in the fibrin gel. A strong correlation was observed between the area covered by AOC and the number of nuclei found in the same area ($r = 0.96$).

2.4.5 Aortic endothelial cell VEGF production

2 2×10^5 cells were seeded in 96-well plates in 200 μ l M199, in the presence or
3 absence of 5, 30 or 100 ng/ml of Netrin-1 and incubated for 48 h. VEGF content
4 in culture media was quantified by an ELISA (Quantikine, R&D System,
5 Minneapolis, MI, USA); this assay, developed for human VEGF detection, has
6 been validated for pig VEGF (Barboni et al. 2000). The assay sensitivity was
7 8.74 pg/ml, the inter- and intra-assay CVs were always less than 7%. Victor
8 Reader set to read at a wavelength of 450 nm emission was used to quantify
9 the reaction product.

2.4.6 Aortic endothelial cell NO production

12 2×10^5 cells were seeded in 96-well plates in 200 μ l M199, in the presence or
13 absence of 5, 30 or 100 ng/ml of Netrin-1 and incubated for 48 h. Nitric oxide
14 (NO) was assessed by measuring nitrite levels in culture media by microplate
15 method based on the formation of a chromophore after reaction with Greiss
16 reagent, which was prepared fresh daily by mixing equal volumes of stock A
17 (1% sulfanilamide, 5% phosphoric acid) and stock B (0.1% N-[naphthyl]
18 ethylenediamine dhydrochloride). After incubation with Greiss reagent the
19 absorbance was determined with the Victor Reader using a 540 nm against 620
20 nm filter. A calibration curve ranging from 25 to 0.39 μ M was prepared by
21 diluting sodium nitrite in culture medium.

2.5 Statistical analysis.

Data are presented as means $\pm$ SEM. Statistical analysis was performed by means of ANOVA using Statgraphics package (STSC Inc., Rockville, MD, USA). When significant differences were found, means were compared by Scheffè's F test. P values < 0.05 were considered to be statistically significant.

3. Results

3.1 Netrin-1 in swine ovarian follicle

3.1.1 Immunohistochemistry

Immunostaining of Netrin-1 in swine ovarian follicle wall was undetectable. On the contrary, immunoreaction was localized in the ovarian vessels (Figure 1 b).

3.1.2 Follicular fluid Western Blotting

The immunoblotting showed the presence of Netrin-1 in swine follicular fluid (Figure 2). The band corresponding to Netrin-1, in fact, was resolved at the expected molecular weight of about 70 kDa both in follicular fluid and in a positive control consisting of a soluble protein extract from porcine spinal marrow.

3.1.3 Netrin-1 gene expression

Netrin-1 expression was evidenced both in granulosa and theca layer from swine follicles (Figure 3).

3.2 Netrin-1 effects on swine granulosa cell functions

3.2.1 Granulosa cell proliferation

Granulosa cell viability, evaluated as ATP content, was slightly but significantly ($p < 0.05$) inhibited by the treatment with 30 and 100 ng/ml of Netrin-1 for 48 h. The lowest concentration of Netrin-1 tested, 5 ng/ml, was ineffective (Figure 4).

3.2.2 Granulosa cell steroid production.

Granulosa cell steroid production, evaluated by RIA assays, was affected by Netrin-1 treatment for 48 h. E2 production was significantly inhibited ($p < 0.01$) by 30 and 100 ng/ml of Netrin-1 (Figure 5); the same concentrations, on the contrary, stimulated ($p < 0.001$) P4 production (Figure 6). The lowest concentration tested, 5 ng/ml, was ineffective on both steroid secretions (Figure 5 and 6).

3.3 Netrin-1 effects on swine aortic endothelial cell functions

3.3.1 Angiogenesis bioassay

Netrin-1 significantly inhibited ($p < 0.05$) new vessel growth both after 48 and 96 h of incubation. All the concentrations tested (5, 30 and 100 ng/ml) exerted the same effect. (Figure 7 and 8).

3.3.2 Aortic endothelial cell VEGF production

1 VEGF secretion by AOC was significantly inhibited ($p < 0.01$) by the 48 h
 2 treatment with all the examined concentrations of Netrin-1 (5, 30 and 100
 3 ng/ml) (Figure 9).

4 5 *3.3.3 Aortic endothelial cell NO production*

6 All Netrin-1 concentrations (5, 30 and 100 ng/ml) significantly stimulated
 7 ($p < 0.01$) NO generation after 48 h of incubation (Figure 10).

8 9 4. Discussion

10 Ovarian physiology is connoted by follicular growth, a fundamental event which
 11 is strictly dependent on cell proliferation, steroid production and new vessel
 12 growth. The molecular machinery which regulates ovarian angiogenesis, though
 13 extensively studied, is far to be completely clarified. In a recent review, Larrivée
 14 et al. (2009) observe that anatomical and structural similarities between blood
 15 vessels and nerves have been highlighted by several studies. Among the
 16 molecules with attractive and repulsive properties which have been found to
 17 modulate the proper guidance of both nerves and blood vessels, netrin-1
 18 displays both pro- and antiangiogenic effects (Park et al., 2004; Castets et al
 19 2009). Interestingly, netrin-1 has been found to be expressed in swine
 20 granulosa cells from 3-5 mm ovarian follicles, thus suggesting its potential
 21 function in follicular function regulation (Maeda et al., 2008).

22 In the present study netrin-1 was evidenced in ovarian vessels but was
 23 undetectable in follicular wall. In addition, we document its expression both in

1 granulosa and theca layer from swine ovarian follicles >5 mm as well as in
2 swine follicular fluid. These data lead us to hypothesize that netrin-1 could play
3 a regulatory role in the function of the ovarian follicle, a structure which
4 represents a typical example of events of proliferation and differentiation.
5 Similarly, Strizzi et al. (2008) demonstrated that mammary gland, a structure
6 subjected to different stages of development and differentiation, expresses
7 netrin-1, which appears to be involved in the fine-tuning of these complex
8 events. In particular, our data show that the protein inhibits granulosa cell
9 viability and E2 synthesis, while it increases P4 production, thus suggesting its
10 involvement in the induction of granulosa cell terminal differentiation. The
11 relationship between growth and differentiation has been a matter of great
12 interest in cellular biology. In granulosa cells, the differentiative function and
13 growth-related process appear inversely controlled (Basini et al., 1998): netrin-
14 1 could play a pivotal role in the paracrine/autocrine regulation of these events.
15 Since follicular growth is strictly dependent on adequate vascular development,
16 this structure has been suggested as an invaluable physiological model of
17 angiogenesis, a process which represents an important aspect of growth and
18 metastasis of solid tumors (Grimm et al 2009). The similarities between nerve
19 and blood vessel guidance have attracted considerable interest towards the
20 study of a potential involvement of neurotropic factors in angiogenesis control
21 (Strizzi et al., 2009).
22 Among these, netrin-1 deserves particular attention even if its exact role in
23 angiogenesis is still doubtful since numerous studies report both pro- or

1 antiangiogenic activities (Bradford et al., 2009). The results obtained in the
2 fibrin bioassay demonstrate an inhibition of angiogenesis by netrin-1. This is in
3 agreement with Lu et al. (2004), but contrary to Park et al. (2004) and Wilson
4 et al. (2006) who suggested that netrin-1 stimulates new vessel growth.
5 Therefore, it appears that netrin-1 function, as already demonstrated in the
6 nervous system (Masuda et al., 2009), could be dependent on receptor type as
7 well as on the intracellular signal transduction pathways activated by each
8 receptor. Further studies will be needed in order to investigate this fundamental
9 topic in our experimental model. Our data also suggest, for the first time, that
10 this effect most likely appears to be mediated by an inhibition of VEGF
11 production by endothelial cells. Moreover, our data are in agreement with
12 Nguyen et al. (2006) demonstrate that netrin-1 appears to increase NO output
13 by endothelial cells. It should be noted that the role of NO in angiogenesis
14 regulation is far from being clarified. Several studies suggest a stimulatory
15 effect (Kimura et al., 2000; Dulak et al., 2000; Sandau et al., 2000) while
16 others point out inhibitory effects (Powell et al., 2000; Santini et al., 2009). The
17 reasons for these discrepancies are still unknown, even if NO levels could play
18 an important role (Kimura et al. 2003).

19 Taken together our data demonstrate that netrin-1 is physiologically present in
20 swine ovary and it is likely involved in modulating follicular function. In
21 particular, data obtained in granulosa cells let us to hypothesize that netrin-1
22 could be important for granulosa cell terminal differentiation; moreover, the
23 observations in endothelial cells seem to suggest its involvement in the control

of new vessel growth. Further studies are strongly advisable to address netrin-1 function since the molecular machinery responsible for angiogenesis inhibition is far to be clarified.

Acknowledgements

We would like to thank Professor Hinck, Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, California and Professor Yelamos (Department of Biochemistry, Molecular Biology and Immunology, Facultad de Medicina, Universidad de Murcia, Spain) for supplying AOC. This research was supported by a FIL and MIUR PRIN grants.

References

- Barboni, B., Turriani, M., Galeati, G., Spinaci, M., Bacci, M.L., Forni, M., Mattioli, M., 2000. Vascular endothelial growth factor production in growing pig antral follicles. *Biol. Reprod.* 63, 858-864.
- Basini, G., Baratta, M., Bussolati, S., Tamanini, C., 1998. Interleukin-1 beta fragment (163-171) modulates bovine granulosa cell proliferation in vitro: dependence on size of follicle. *J. Reprod. Immunol.* 37, 139-153.
- Basini, G., Bussolati, S., Santini, S.E., Bianchi, F., Careri, M., Mangia, A., Musci, M., Grasselli, F., 2008a. Hydroxyestrogens inhibit angiogenesis in swine ovarian follicles. *J. Endocrinol.* 199, 127-135.

- 1 Basini, G., Bussolati S., Santini S.E., Grasselli F., 2008b. Reactive oxygen
2 species and anti-oxidant defences in swine follicular fluids. *Reprod. Fertil. Dev.*
3 20, 269-274.
- 4 Basini, G., Baioni, L., Bussolati, S., Grolli, S., Kramer, LH, Wagner, G.F.,
5 Grasselli F., 2010. Expression and localization of stanniocalcin 1 in swine ovary.
6 *Gen. Comp. Endocrinol.* 166, 404-408.
- 7 Bouvrée, K., Larrivée, B., Lv, X., Yuan, L., DeLafarge, B., Freitas, C., Mathivet,
8 T., Bréant, C., Tessier-Lavigne, M., Bikfalvi, A., Eichmann, A., Pardanaud, L.,
9 2008. Netrin-1 inhibits sprouting angiogenesis in developing avian embryos.
10 *Dev Biol.* 318 172-183.
- 11 Bradford, D., Cole, S.J., Cooper, H.M., 2009. Netrin-1: diversity in development.
12 *Int. J. Biochem. Cell. Biol.* 413 487-493.
- 13 Carrello, S., Chamorro, M., Rodriguez-Gago, B., Alvarez, M.J., Molina, J.I.,
14 Rodriguez-Barbosa, A., Sanchez, P., Ramirez, A., Munoz, J., Dominguez, J.,
15 Parrilla, P., Yelamos, J. 2002. Isolation and characterization of immortalized
16 porcine aortic endothelial cell lines. *Vet. Immunol. Immunopathol.* 89, 91-98.
- 17 Castets, M., Coissieux, M.M., Delloye-Bourgeois, C., Bernard, L., Delcros, J.G.,
18 Bernet, A., Laudet, V., Mehlen, P., 2009. Inhibition of endothelial cell apoptosis
19 by netrin-1 during angiogenesis. *Dev. Cell.* 16 614-620.
- 20 Cirulli, V., Yebra, M., 2007. Netrins: beyond the brain. *Nat. Rev. Mol. Cell. Biol.*
21 8, 296-306.
- 22 Dulak, J., Józkowicz, A., Dembinska-Kiec, A., Guevara, I., Zdzenicka, A.,
23 Zmudzinska-Grochot, D., Florek, I., Wójtowicz, A., Szuba, A., Cooke, J.P., 2000.

- 1 Nitric oxide induces the synthesis of vascular endothelial growth factor by rat
- 2 vascular smooth muscle cells. *Arterioscler. Thromb. Vasc. Biol.* 20, 659-666.
- 3 Grasselli, F., Baratta, M., Tamanini, C., 1993. Effects of a GnRH analogue
- 4 (buserelin) infused via osmotic minipumps on pituitary and ovarian activity of
- 5 prepuberal heifers. *Anim. Reprod. Sci.* 32, 153-161.
- 6 Keino-Masu, K., Masu, M., Hinck, L., Leonardo, E.D., Chan, S.S., Culotti, J.G.,
- 7 Tessier-Lavigne, M., 1996. Deleted in Colorectal Cancer (DCC) encodes a netrin
- 8 receptor. *Cell.* 87, 175-185.
- 9 Kimura, H., Esumi, H., 2003. Reciprocal regulation between nitric oxide and
- 10 vascular endothelial growth factor in angiogenesis. *Acta Biochim. Pol.* 50, 49-
- 11 59.
- 12 Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the
- 13 head of bacteriophage T4. *Nature* 227, 680-685.
- 14 Larrivée, B., Freitas, C., Trombe, M., Lv, X., Delafarge, B., Yuan, L., Bouvrée,
- 15 K., Bréant, C., Del Toro, R., Bréchet, N., Germain, S., Bono, F., Dol, F., Claes,
- 16 F., Fischer, C., Autiero, M., Thomas, J.L., Carmeliet, P., Tessier-Lavigne, M.,
- 17 Eichmann, A. 2007. Activation of the UNC5B receptor by Netrin-1 inhibits
- 18 sprouting angiogenesis. *Genes Dev.* 21, 2433-2447.
- 19 Larrivée, B., Freitas, C., Suchting, S., Brunet, I., Eichmann, A., 2009. Guidance
- 20 of vascular development: lessons from the nervous system. *Circ. Res.* 104, 428-
- 21 441.
- 22 Lu, X., Le Noble, F., Yuan, L., Jiang, Q., De Lafarge, B., Sugiyama, D., Bréant,
- 23 C., Claes, F., De Smet, F., Thomas, J.L., Autiero, M., Carmeliet, P., Tessier-

- 1 Lavigne, M., Eichmann, A., 2004. The netrin receptor UNC5B mediates guidance
2 events controlling morphogenesis of the vascular system. *Nature* 432, 179-186.
- 3 Maeda, A., Matsuda, F., Goto, Y., Cheng, Y., Gonda, H., Inoue, N., Nakagawa,
4 S., Manabe, N., 2008. Molecular cloning of a porcine (*Sus scrofa*) apoptosis
5 inhibitory ligand, netrin-1, and its receptor, p53RDL1. *J. Reprod. Dev.* 54, 275-
6 280.
- 7 Masuda, T., Yaginuma, H., Sakuma, C., Ono, K., 2009. Netrin-1 signaling for
8 sensory axons: Involvement in sensory axonal development and regeneration.
9 *Cell. Adh. Migr.* 3, 171-173.
- 10 Nguyen, A., Cai, H., 2006. Netrin-1 induces angiogenesis via a DCC-dependent
11 ERK1/2-eNOS feed-forward mechanism. *Proc. Natl. Acad. Sci. U S A.* 103, 6530-
12 6535.
- 13 Park, K.W., Crouse, D., Lee, M., Karnik, S.K., Sorensen, L.K., Murphy, K.J., Kuo,
14 C.J., Li, D.Y., 2004. The axonal attractant Netrin-1 is an angiogenic factor. *Proc.*
15 *Natl. Acad. Sci. U S A.* 101, 16210-16215.
- 16 Powell, J.A., Mohamed, S.N., Kerr, J.S., Mousa, S.A., 2000. Antiangiogenesis
17 efficacy of nitric oxide donors. *J. Cell. Biochem.* 80, 104-114.
- 18 Sandau, K.B., Fandrey, J., Brüne, B., 2001. Accumulation of HIF-1alpha under
19 the influence of nitric oxide. *Blood* 97, 1009-1015.
- 20 Santini, S.E., Basini, G., Bussolati, S., Grasselli, F., 2009. The phytoestrogen
21 quercetin impairs steroidogenesis and angiogenesis in swine granulosa cells in
22 vitro. *J. Biomed. Biotechnol.* 2009-419891

- 1 Serafini, T., Kennedy, T.E., Galko, M.J., Mirzayan, C., Jessell, T.M., Tessier-
2 Lavigne, M., 1994. The netrins define a family of axon outgrowth-promoting
3 proteins homologous to *C. elegans* UNC-6. *Cell* 78, 409-424.
- 4 Strizzi, L., Mancino, M., Bianco, C., Raafat, A., Gonzales, M., Booth, B.W.,
5 Watanabe, K., Nagaoka, T., Mack, D.L., Howard, B., Callahan, R., Smith, G.H.,
6 Salomon, D.S. 2008. Netrin-1 can affect morphogenesis and differentiation of
7 the mouse mammary gland. *J. Cell. Physiol.* 216, 824-834.
- 8 Towbin, H., Staehelin, T., Gordon, J., 1979. Electrophoretic transfer of proteins
9 from polyacrylamide gels to nitrocellulose sheets: procedure and some
10 applications. *Proc Natl Acad Sci U S A.* 76, 4350-4354.

11

Figure captions

Figure 1. Immunohistochemical localization of Netrin 1 in swine ovary. A: antral follicles. B: vessel. C: follicle walls. D: corpus luteum. A positive signal has been detected only in vessels (B, see arrows).

Figure 2. Netrin 1 protein in swine follicular fluid and spinal marrow revealed by immunoblotting. Protein bands on SDS-PAGE gel were stained with Coomassie blue.

Figure 3. Netrin-1 expression in swine granulosa and theca cells collected from swine follicles. The representative ethidium bromide-stained gel shows ACT and STC 1 amplicons from at least three follicles.

Figure 4. Effect of the 48 h treatment with or without (C) Netrin-1 at the concentrations of 5, 30 or 100 ng/ml on granulosa cell viability evaluated as ATP content. Data, expressed as % vs respective C, represent the mean $\pm$ SEM of six replicates/treatment repeated in four different experiments. Different letters indicate a significant difference ($p < 0.05$).

Figure 5. Effect of the 48 h treatment with or without (C) Netrin-1 at the concentrations of 5, 30 or 100 ng/ml on E2 production in swine granulosa cell culture evaluated by RIA. Data, expressed as ng/ml, represent the mean $\pm$ SEM

of six replicates/treatment repeated in four different experiments. The absence of common letters indicate a significant difference ($p < 0.01$).

Figure 6. Effect of the 48 h treatment with or without (C) Netrin-1 at the concentrations of 5, 30 or 100 ng/ml on P4 production in swine granulosa cell culture evaluated by RIA. Data, expressed as ng/ml, represent the mean $\pm$ SEM of six replicates/treatment repeated in four different experiments. Different letters indicate a significant difference ($p < 0.001$).

Figure 7. Effect of the 48 h treatment with or without (C) Netrin-1 at the concentrations of 5, 30 or 100 ng/ml on AOC growth in fibrin gel. Panel A: phase contrast micrographs showing AOC growth at 48 h in fibrin gel matrix. Panel B: area covered by AOC. Data, expressed as % vs respective C, represent the mean $\pm$ SEM of six replicates/treatment repeated in four different experiments. Different letters indicate a significant difference ($p < 0.05$).

Figure 8. Effect of the 96 h treatment with or without (C) Netrin-1 at the concentrations of 5, 30 or 100 ng/ml on AOC growth in fibrin gel. Panel A: phase contrast micrographs showing AOC growth at 96 h in fibrin gel matrix. Panel B: area covered by AOC. Data, expressed as % vs respective C, represent the mean $\pm$ SEM of six replicates/treatment repeated in four different experiments. Different letters indicate a significant difference ($p < 0.05$).

1 Figure 9. Effect of the 48 h treatment with or without (C) Netrin-1 at the
2 concentrations of 5, 30 or 100 ng/ml on VEGF production, evaluated by ELISA,
3 by AOC culture. Data, expressed as pg/ml, represent the mean $\pm$ SEM of six
4 replicates/treatment repeated in four different experiments. Different letters
5 indicate a significant difference ($p < 0.01$).

6

7 Figure 10. Effect of the 48 h treatment with or without (C) Netrin-1 at the
8 concentrations of 5, 30 or 100 ng/ml on NO production, evaluated by Greiss
9 method, by AOC culture. Data, expressed as μM , represent the mean $\pm$ SEM of
10 six replicates/treatment repeated in four different experiments. The absence of
11 common letters indicate a significant difference ($p < 0.01$).

12

13

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10