A dose escalation study of pemetrexed and docetaxel in non-small cell lung cancer (NSCLC)

Athanasios Kotsakis^{a*}, Sophia Agelaki^a, Nikolaos Vardakis^a, George Stathopoulos^b, Lambros Vamvakas^a, Antonia Kalykaki^a, Nikolaos Kentepozidis^a, Emmanouel Kontopodis^a, Georgia Sfakiotaki^a, Dimitris Mavroudis^a, Vassilis Georgoulas^a

^aDepartment of Medical Oncology, University Hospital of Heraklion, Crete, Greece

^bFirst Department of Medical Oncology, “E. Dynan” General Hospital of Athens, Greece

Running title: Pemetrexed/Docetaxel in NSCLC

Key Words: pemetrexed; docetaxel; NSCLC; phase I study

Acknowledgements: This work was partly supported by a research grant from the Cretan Association for Biomedical Research.

To whom correspondence should be addressed: Athanasios Kotsakis, MD, PhD, Department of Medical Oncology, University General Hospital of Heraklion, PO Box 1352, 711 10 Heraklion, Crete, Greece.

Tel: +30 2810 392783, FAX: +30 2810 392857, E-mail: georgsec@med.uoc.gr

Abstract

A phase I study was conducted to determine the maximum tolerated doses (MTD) and the dose limiting toxicities (DLT) of pemetrexed and docetaxel in patients with advanced unresectable or metastatic non-small cell lung cancer (NSCLC). Patients were treated with escalating doses of pemetrexed ($400\text{-}600\text{mg/m}^2$ as a 10-min intravenous infusion) and docetaxel ($65\text{-}85\text{mg/m}^2$ as a 1 hour intravenous infusion) on day 1, every 3 weeks. An expanded accrual at the level of the recommended dose (RD) had been scheduled. Forty-two patients with metastatic NSCLC were enrolled in the phase I study and 20 additional patients at the RD level. The MTD could not be reached even at the doses of 550mg/m^2 and 85mg/m^2 for pemetrexed and docetaxel respectively, which are higher than the recommended dose for each drug given as a single agent. Therefore, the RD was defined at 500mg/m^2 pemetrexed and 75mg/m^2 docetaxel. Among the 164 administered chemotherapy cycles (phase I part) there were 3 episodes of febrile neutropenia whereas 13 (7.9%) and 11 (6.7%) cycles were complicated with grade III and IV neutropenia, respectively. Three patients developed grade III/IV thrombocytopenia. Non-hematologic toxicity was mild with grade III fatigue occurring in three (6.7%) patients. There was no toxic death. The favorable toxicity profile of the regimen was confirmed in patients treated at the RD level. Overall, one complete (CR) and 13 partial responses (PR) [overall response rate=23%; 95% C.I:12.4% - 33.5%] were documented. The combination of pemetrexed and docetaxel seems to be an effective regimen in NSCLC with acceptable and manageable toxicity which merits further investigation.

Introduction

Lung cancer is the leading cause of cancer death in the United States and Europe and it has been estimated that more patients will die from lung cancer than prostate, breast and colorectal cancer together [1]. In 2006, lung cancer was in incidence the third most common cancer in Europe and the most frequent reason for death caused by a malignant tumor [2]. Non-small cell lung cancer (NSCLC) accounts for approximately 80% of all lung cancers and has become the major cause of cancer-related mortality in both men and women. Despite significant advances in the treatment of NSCLC over the past years, the prognosis of the patients with advanced or metastatic disease is poor with a median survival for the newly diagnosed stage IV patients ranging from 8-11 months and 5-7 months for those with relapsed disease [2].

Platinum-based combination has emerged as the cornerstone first-line treatment for NSCLC. In two large meta-analyses, cisplatin-based combinations revealed superior responses and improved overall survival (OS) in some subgroups of patients [3;4]. Moreover, cisplatin has been shown to be superior to carboplatin when is combined with new chemotherapeutic agents (e.g. paclitaxel, docetaxel and gemcitabine) [3]. Recently, an antibody against vascular endothelial growth factor (bevacizumab) was proved effective in the first-line setting of non-squamous NSCLC, as significantly prolonged the progression-free survival (PFS) [5;6] and OS [5]. Although these agents are active in NSCLC, they add toxicity which, in some cases, is severe. Nausea, vomiting, anemia, neuro-and nephro-toxicity comprise the most common occurred adverse events. Bevacizumab has also been associated with increased toxicity mainly, hypertension, proteinuria, hemorrhage and gastro-intestinal (GI) perforation.

Less toxic and more active regimens are required to be incorporated in the modern oncology armamentarium. Hence, the replacement of platinum compound

with new agents has been emerged as an attractive approach for the treatment of NSCLC. Docetaxel promotes the assembly of microtubules from tubulin dimers, inhibits the depolymerization of tubulin which stabilizes microtubules in the cell and, eventually, inhibits DNA, RNA, and protein synthesis occurring mainly during the M phase of the cell cycle. On the basis of the results of big randomized trials [7-9], docetaxel has been approved as first-line treatment for NSCLC in combination with cisplatin and as monotherapy in the second-line setting. Docetaxel has shown activity in the first-line setting even with non-platinum combinations; we have previously shown that the docetaxel/cisplatin and the docetaxel/gemcitabine regimens had a comparable efficacy in terms of response rate (RR), PFS and OS with a more favorable toxicity profile for the non-platinum regimen [10].

Pemetrexed is a multitargeted antifolate drug that targets the enzymes thymidylate synthase (TS), dihydrofolate reductase (DHFR) and glycinamide ribonucleotide formyl transferase (GARFT) [11]. By inhibition of these enzymes, it affects the synthesis of substrates necessary for cell growth and division and causes cell-cycle arrest in the S phase [12;13]. It has shown activity against NSCLC and has been approved as front-line treatment in combination with cisplatin as well as monotherapy in the second line setting for the non-squamous cell carcinomas [14]. In addition, pemetrexed given as maintenance treatment in NSCLC patients who achieve disease control after platinum-based front-line therapy could significantly improve OS and PFS [15].

The fact that both pemetrexed and docetaxel result to a cell-cycle arrest during different phases, may imply that they could have an additive or synergistic anti-tumor effect. Indeed, preclinical data have shown at least additive effect when pemetrexed was combined with docetaxel [16]. The same combination regimen has also been used in clinical trials in solid tumors with promising efficacy and acceptable toxicity.

Furthermore, the combination of paclitaxel with pemetrexed was recently tested in patients with NSCLC with encouraging results [17]. These data provide us the rationale to conduct a dose-escalation study of the docetaxel/pemetrexed combination in order to determine the dose limiting toxicity (DLT) and the maximum tolerated doses (MTD) of the association in patients with advanced or metastatic NSCLC.

Patients and methods

Patient selection

Patients aged >18 years old with a pathologically or cytologically confirmed, unresectable locally advanced (stage IIIB with pleural infusion) or metastatic NSCLC who had received prior chemotherapy (maximum one prior chemotherapy regimen) as well as chemotherapy-naïve patients were eligible for the study; in addition, patients with CNS disease were also eligible if they had been irradiated and were clinically stable. Prior surgery or radiotherapy (to less than 25% of bone marrow) was allowed. However, a treatment-free interval of at least 4 weeks was required before study enrollment. Other inclusion criteria were: performance status 0-2 (ECOG); adequate blood counts (absolute neutrophil count $>1,500/\mu\text{L}$, hemoglobin $<10/\mu\text{L}$ and platelets $>100,000/\mu\text{L}$); adequate renal function (serum creatinine $<2\text{ mg/dL}$); adequate hepatic function (total bilirubin $<2\text{ mg/dL}$ and transaminases <3 times the upper normal limit); adequate cardiac function without unstable angina or myocardial infarction within the last 6 months; no history of other primary cancer; and life expectancy of at least 3 months. All patients signed a written informed consent before study enrollment. The study had been approved by the Ethics and Scientific Committees of our Institution and is registered in ClinicalTrials.gov (NCT00684099).

Treatment

Treatment schedule and dose escalation are shown in Table 1. Docetaxel (Taxotere; Sanofi-Aventis, Bridgewater, USA) was administered as a 1 hour intravenous infusion (IV) at escalated doses from 60 to 85 mg/m² followed by pemetrexed (Alimta; Eli-Lilly, Indianapolis, USA) given as a 10 minutes IV infusion, at escalated doses from 400 to 550 mg/m²; both drugs were given on day 1 every 3 weeks without prophylactic administration of hematopoietic growth factors. The treatment was continued for up to 6 cycles or occurrence of disease progression or unacceptable toxicity. All patients had standard vitamin supplementation with 1000 mcg vitamin B12 intramuscular every 9 weeks and folic acid 400 mcg/day per os for the administration of pemetrexed. In addition, standard premedication with corticosteroids for docetaxel administration was given. The used antiemetic regimen included ondansetron 16 mg intravenously 30 minutes before chemotherapy administration.

Dose escalation

The following dose-levels for docetaxel/pemetrexed have been evaluated: 65/400; 65/450; 70/450; 70/500; 75/500; 80/500; 80/550; 85/500 (mg/m²) (Table 1). No intra-patient dose escalation was allowed. At least three patients were enrolled at each dose level. If a DLT was observed in one of the first three patients, then three additional patients were enrolled at the same dose level. DLTs were assessed during the first chemotherapy cycle. As DLT was defined the occurrence of any of the following: grade 4 neutropenia or thrombocytopenia; grade 3-4 neutropenia with fever >38.3°C, or a sustained temperature >38°C for more than one hour [18]; grade 3-4 non-hematologic toxicity; and any treatment delay because of toxicity. Dose escalation was discontinued and the DLT level would be reached if at least 50% of the patients

treated at that level developed a DLT (e.g. at least two of three, or three of six patients). The MTD level was defined as the first level below the DLT dose level. At the MTD level, 20 additional patients have to be recruited (a total of 23 patients) in order to further evaluate the tolerance of the regimen at the recommended doses (RD level) for future phase II studies.

Toxicity assessment

Before each chemotherapy cycle, all patients had a detailed toxicity questionnaire and a physical examination in order to define treatment-related adverse events. Hematologic toxicity was assessed weekly with complete blood counts (CBC) with differential and platelet counts; in case of grade III/IV hematologic toxicity, a CBC with differential and platelets count was performed daily until recovery (neutrophils $>1500/\mu\text{l}$ and platelets $>100.000/\mu\text{l}$). A complete biochemical profile was performed at baseline and every three weeks thereafter. Toxicities were graded according to NCI CTC version 3.0. Day 1 treatment was administered on scheduled dates without any delay or dose reduction if the laboratory inclusion criteria were met; otherwise, day 1 treatment was postponed for up to 7 days until resolution of the prohibitive toxicity. Then, treatment was administered according to the predefined schedule and dose and without any missed dose. If the prohibitive toxicity had not resolved after a 7-day treatment delay, the scheduled treatment was missed and upon resolution of the toxicity, treatment was resumed as a new cycle using doses of the previous dose-level. Doses were also reduced to the previous dose-level in case of febrile neutropenia (grade II-IV neutropenia with fever $> 38.0^{\circ}\text{C}$ in two consecutive measurements at least one hour apart) [18] or grade III-IV thrombocytopenia with or without bleeding episode.

Tumor Response

Although patients were not required to have bidimensionally measurable disease to enter the study, response was assessed according to the standard RECIST criteria for those who did have [19]. Tumor response was assessed every three chemotherapy cycles by appropriate imaging studies and all responses have to be confirmed by a repeated imaging evaluation 4 weeks later. The duration of response was defined as the time from the day that objective tumor shrinkage was noted for the first time until the time of documented tumor progression or death due to any cause. The time to progressive (TTP) disease and the overall survival were defined as the time from the date of study enrollment until the date of progressive disease and the date of death due to any cause, respectively.

Results

Patients' demographics

Patients' characteristics are presented in Table 2. Between June 2005 and June 2009 a total of 62 patients were enrolled onto the study; forty-two patients were recruited onto the escalation part of the study and 20 at the RD level of the expanded part of the study. The patients' median age was 63 years and 87% of them had a performance status (ECOG) of 0-1. Fourteen (23%) patients suffering from a squamous cell carcinoma were enrolled in the trial before May 2008 when the published results demonstrated that pemetrexed was mainly active in patients with a non-squamous cell histology (14). From the patients enrolled in the phase I part of the study, 33 (78.6%) had a non-squamous histology whereas in the phase II part of the study, 15 (75%) had a non-squamous histology. Fifty-one (82%) patients had stage IV disease and 21 (34%) were chemotherapy-naïve. Table 2 also indicates that 34 (54.8%) patients had previously received platinum- and 35 (56.5%) taxane-based chemotherapy.

Compliance with the treatment

At the escalation part of the study, a total of 164 chemotherapy cycles were administered in all patients and in all dose-levels with a median of 3 cycles/ patient (range, 1-9). The median interval between cycles was 21 days (range, 21-33). Eleven (7%) cycles were delayed because of hematologic toxicity (n=2 cycles) as well as for reasons unrelated to the disease or treatment (n=9 cycles). Dose reduction was required in 8 (4.9%) cycles because of hematologic (n=6 cycles) and non-hematologic (n=2 cycles) toxicity. The reasons for treatment discontinuation were: completion of treatment as per protocol (n=12 patients), progressive disease (n=25 patients), toxicity (n=3 patients) and patient's consent withdrawn or lost to follow up (n=3 patients). Regarding the expansion at the RD level, the rate and the severity of toxicity was similar to that observed during the escalation and significant delays or dose reductions were not required (data not shown).

Dose-limiting Toxicities

Table 1 shows the dose-escalation levels, the number of patients enrolled at each dose-level and the observed DLTs during the first cycle. The most common DLT was grade III-IV neutropenia or febrile neutropenia observed at dose-levels 2 (n=2 patients), 4 (n=2 patients), 6 (n=1 patient), 7 (n=1 patient) and 8 (n=2 patients). Other DLTs included grade IV thrombocytopenia at dose-level 3 (n=1 patient) and grade III generalized skin rash leading to treatment delay at dose level 6 (n=1 patient). The MTD has not been reached. However, dose escalation has been discontinued when it was ascertained that the regimen was safe, with an acceptable toxicity profile even at dose levels which exceeded the recommended single agent doses and it was observed

that higher doses of the drugs were not associated with better efficacy results (see below). Therefore, the recommended doses for future studies are docetaxel 75mg/m² and pemetrexed 500mg/m² on day 1 every 3 weeks (5th dose-level=RD level).

Hematologic and non-hematologic toxicity

Table 3 indicates the hematologic toxicity observed in all patients and in all chemotherapy cycles irrespectively of the dose-level. A total of 12 (6.7%) and 16 (9.1%) cycles were complicated with grade III and IV neutropenia, respectively; three patients presented grade III (n=2 patients) and IV (n=1 patient) thrombocytopenia; no bleeding episodes were observed and no platelet transfusions were required. There were four episodes of febrile neutropenia occurring at the 1st, the 2nd, the 5th and the 6th dose-levels but none of them occurred during the first chemotherapy cycle. There was no toxic death. One patient developed a non-neutropenic infection requiring hospitalization for intravenous antibiotics and cardio-respiratory support; the patient was uneventfully recovered.

Non-hematologic toxicity was generally mild. Table 4 shows the incidence of moderate to severe non-hematologic toxicity in all patients and all cycles, irrespectively of the dose-levels. The most common toxicity was asthenia which was grade II in 20 (12.2%) cycles corresponding to 11 (26%) patients and grade III in 3 (1.8%) cycles corresponding to three (7.2%) patients. Other toxicities were mild occurring in less than 5% of the administered cycles. Grade III generalized skin rash was observed in one (0.5%) patient two times. No other severe unexpected toxicities were observed. For the patients treated at the higher dose levels (6-8), the observed toxicity was slightly increased, mainly regarding the hematological profile. At the RD level, 20 additional patients were enrolled. The toxicity profile of the regimen as well as the incidence of different adverse events was, practically, similar to those observed

during the dose escalation part of the trial. In particular, grade III and IV neutropenia occurred in three patients each, whereas febrile neutropenia was observed in one patient (Table 5).

Antitumor activity

In total, 61 patients were evaluable for response [41 out of 42 patients who were enrolled in the escalation part of the study and 20 patients enrolled in the RD level]. One patient without a measurable target lesion was not evaluated for response. Complete response (CR) was achieved in one (1.6%) patient and partial response (PR) in 13 (21.0%) [overall response rate [ORR]=23%; 95% C.I:12.4% - 33.5%]; ten (16.1%) patients experienced stable disease (SD) and 37 (60.7%) progressive disease (PD). The disease control rate was 39%. The treatment was 1st-line for five of the responders (ORR=25% in chemotherapy-naïve patients; 95% C.I: 6.0% - 43.9%) and 2nd-line for nine of them. (ORR=22%; 95% CI: 9.3% - 34.6%) (p=0.790). In addition, four (28.5%) and 10 (21.3%) responses occurred in patients with squamous and non-squamous cell carcinoma, respectively. The only CR was observed in the 8th dose-level while the PRs were observed in all dose-levels (2nd: n=1; 3rd: n=2; 4th: n=1; 7th: n=2; 8th: n=1) during the escalation part of the study. Regarding the efficacy of the combination in the expanded cohort of patients (n=20) who were treated with the recommended doses, a PR was achieved in six patients (30.0%), SD in four (20.0%) and PD in 10 (50.0%). The overall median response duration was 6.8 months (range, 0.5-16 months) and the overall median TTP 2.8 months (range, 0.5-17.2 months) [the median TTP was 2.8 (range, 0.7-17.2) and 3.3 (range, 0.4-13.1) months for chemotherapy-naïve and pre-treated patients, respectively (p=0.845)]. After a median follow up period of 18.5 months (0.7-38.2 months), the median overall survival (OS) was 10.0 months (range, 0.7-38.2 months); [the median OS was 10.4 (range, 1.4-38.2) and 9.2 (range, 0.7-31.5) months for patients treated in the 1st- and 2nd-line setting,

respectively ($p=0.356$)). The estimated 1-year survival rate for all patients was 40.5% (40.5% and 40.9% for chemotherapy-naïve and pre-treated patients, respectively).

Discussion

The necessity of more active and less toxic chemotherapy regimens for NSCLC patients is of great interest for the oncologists. In the present study, we combined two active drugs (pemetrexed and docetaxel) against NSCLC [7-9;14;20] in a dose-escalation trial. It is interesting to note that the MTD has not been reached even at doses which exceeded the recommended doses for monotherapy with each one of them. Thus, given that efficacy results were similar in the high and low dose levels, we recommended the dose of 75 mg/m^2 for docetaxel and 500 mg/m^2 for pemetrexed (RD level) for future phase II studies.

The combination of pemetrexed with taxanes (paclitaxel and docetaxel) has been tested in phase I and II studies. The pemetrexed/paclitaxel regimen has been associated with encouraging efficacy results and acceptable toxicity even at doses which approached the recommended dose for each agent in the monotherapy setting [21;22]. In a phase II study, the combination of paclitaxel with pemetrexed, which was evaluated in chemotherapy-naïve patients with NSCLC, showed an encouraging efficacy with an objective response rate of 39.6% associated with a median survival of 14 months and a favorable toxicity profile [17].

There are few clinical trials evaluating the pemetrexed/docetaxel combination in patients with solid tumors [23-26]. In a phase I study, 17 patients with solid tumors were treated with a fixed dose of docetaxel (60 mg/m^2 on day 8) combined with escalated doses of pemetrexed ($300\text{-}600 \text{ mg/m}^2$ on day1); the first level of 300 mg/m^2 for pemetrexed and 60 mg/m^2 for docetaxel was the MTD. Then, the investigators amended the treatment schedule and the same doses of pemetrexed were administered

on day 1 and docetaxel on day 15. This sequential modified administration was feasible and well tolerated; at the time of study presentation, the MTD had not been reached (pemetrexed 600mg/m² and docetaxel 60mg/m²). Hematologic toxicity was the main adverse event of the combination [26]. In another phase I study, both agents were administered on days 1 and 15. At the dose level of pemetrexed 400mg/m² and docetaxel 40 mg/m² the MTD had not been reached [23]. Similar to our study, another phase I trial investigated the administration of pemetrexed and docetaxel on day 1 every 3 weeks, in patients with metastatic disease who had received ≥ 2 prior regimens of chemotherapy. This study was suspended and the planned phase II study was abandoned due to the limited tolerability and efficacy of this regimen [24]. This is in overt contrast with the favorable toxicity profile which was observed in the current study; this discrepancy could be attributed to the fact that patients suffering from a wide range of tumors, which have a variable sensitivity to both agents, were enrolled in this trial. Based on the tolerability of the docetaxel/pemetrexed regimen it seems that both the biweekly as well as the 3-weekly schedule should be preferred for further evaluation.

In the present study, the docetaxel/pemetrexed regimen was well tolerated and neutropenia and fatigue were the most frequently observed adverse events. In the initial clinical trials, pemetrexed was associated with severe toxicity including hematologic toxicity, skin rash, lethargy, nausea and vomiting [27-29]. The incidence of these adverse events was significantly reduced with the administration of vitamin supplementation [30]. Consistently, the administration of vitamin B12 and folates was standard practice in the current study explaining, thus, the low incidence of severe toxicity.

Although efficacy was not the primary endpoint of the study, the pemetrexed/docetaxel regimen resulted in an ORR of almost 23%, a disease control

rate (CR+PR+SD) of 39% and a median overall survival of 10.0 months. These efficacy results should be considered encouraging, given that 66% of the enrolled patients were already pre-treated. Moreover, it should be taken into account that the study was initiated in April 2005 and 23% of the enrolled patients had the diagnosis of squamous cell carcinoma, for which pemetrexed has been proved recently ineffective [14]; therefore, it could be argued that these particular patients could achieve the same clinical response with docetaxel alone.

In conclusion, this dose-escalation study demonstrated that the combination of docetaxel and pemetrexed is well-tolerated in patients with NSCLC. The drugs can be safely combined at the recommended, for each agent, monotherapy doses. The antitumor activity of this regimen was promising and its combination with anti-angiogenic agents in non-squamous lung carcinoma could be of great interest in future studies. In addition, we consider that it merits further evaluation in second line treatment compared to monotherapy with docetaxel, pemetrexed or erlotinib.

Conflict of interest

None declare

References

1. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ (2009) Cancer Statistics, 2009. *CA Cancer J Clin*
2. Longo-Sorbello GS, Chen B, Budak-Alpdogan T, Bertino JR (2007) Role of pemetrexed in non-small cell lung cancer. *Cancer Invest* 25: 59-66
3. Hotta K, Matsuo K, Ueoka H, Kiura K, Tabata M, Tanimoto M (2004) Meta-analysis of randomized clinical trials comparing Cisplatin to Carboplatin in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 22: 3852-3859
4. Ardizzoni A, Boni L, Tiseo M, Fossella FV, Schiller JH, Paesmans M, Radosavljevic D, Paccagnella A, Zatloukal P, Mazzanti P, Bisset D, Rosell R (2007) Cisplatin- versus carboplatin-based chemotherapy in first-line treatment of advanced non-small-cell lung cancer: an individual patient data meta-analysis. *J Natl Cancer Inst* 99: 847-857
5. Sandler A, Gray R, Perry MC, Brahmer J, Schiller JH, Dowlati A, Lilenbaum R, Johnson DH (2006) Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 355: 2542-2550
6. Reck M, von PJ, Zatloukal P, Ramlau R, Gorbounova V, Hirsh V, Leigh N, Mezger J, Archer V, Moore N, Manegold C (2009) Phase III trial of cisplatin plus gemcitabine with either placebo or bevacizumab as first-line therapy for nonsquamous non-small-cell lung cancer: AVAIL. *J Clin Oncol* 27: 1227-1234
7. Fossella F, Pereira JR, von PJ, Pluzanska A, Gorbounova V, Kaukel E, Mattson KV, Ramlau R, Szczesna A, Fidias P, Millward M, Belani CP (2003) Randomized, multinational, phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for advanced non-small-cell lung cancer: the TAX 326 study group. *J Clin Oncol* 21: 3016-3024

8. Shepherd FA, Dancey J, Ramlau R, Mattson K, Gralla R, O'Rourke M, Levitan N, Gressot L, Vincent M, Burkes R, Coughlin S, Kim Y, Berille J (2000) Prospective randomized trial of docetaxel versus best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy. *J Clin Oncol* 18: 2095-2103
9. Fossella FV, DeVore R, Kerr RN, Crawford J, Natale RR, Dunphy F, Kalman L, Miller V, Lee JS, Moore M, Gandara D, Karp D, Vokes E, Kris M, Kim Y, Gamza F, Hammershaimb L (2000) Randomized phase III trial of docetaxel versus vinorelbine or ifosfamide in patients with advanced non-small-cell lung cancer previously treated with platinum-containing chemotherapy regimens. The TAX 320 Non-Small Cell Lung Cancer Study Group. *J Clin Oncol* 18: 2354-2362
10. Georgoulas V, Papadakis E, Alexopoulos A, Tsiafaki X, Rapti A, Veslemes M, Palamidas P, Vlachonikolis I (2001) Platinum-based and non-platinum-based chemotherapy in advanced non-small-cell lung cancer: a randomised multicentre trial. *Lancet* 357: 1478-1484
11. Adjei AA (2000) Pemetrexed: a multitargeted antifolate agent with promising activity in solid tumors. *Ann Oncol* 11: 1335-1341
12. Tonkinson JL, Marder P, Andis SL, Schultz RM, Gossett LS, Shih C, Mendelsohn LG (1997) Cell cycle effects of antifolate antimetabolites: implications for cytotoxicity and cytostasis. *Cancer Chemother Pharmacol* 39: 521-531
13. Dubey S, Schiller JH (2005) Three emerging new drugs for NSCLC: pemetrexed, bortezomib, and cetuximab. *Oncologist* 10: 282-291
14. Scagliotti GV, Parikh P, von PJ, Biesma B, Vansteenkiste J, Manegold C, Serwatowski P, Gatzemeier U, Digumarti R, Zukin M, Lee JS, Mellemaard A,

- Park K, Patil S, Rolski J, Goksel T, de MF, Simms L, Sugarman KP, Gandara D (2008) Phase III study comparing cisplatin plus gemcitabine with cisplatin plus pemetrexed in chemotherapy-naïve patients with advanced-stage non-small-cell lung cancer. *J Clin Oncol* 26: 3543-3551
15. Ciuleanu T, Brodowicz T, Zielinski C, Kim JH, Krzakowski M, Laack E, Wu YL, Bover I, Begbie S, Tzekova V, Cucevic B, Pereira JR, Yang SH, Madhavan J, Sugarman KP, Peterson P, John WJ, Krejcy K, Belani CP (2009) Maintenance pemetrexed plus best supportive care versus placebo plus best supportive care for non-small-cell lung cancer: a randomised, double-blind, phase 3 study. *Lancet* 374: 1432-1440
 16. Teicher BA, Alvarez E, Liu P, Lu K, Menon K, Dempsey J, Schultz RM (1999) MTA (LY231514) in combination treatment regimens using human tumor xenografts and the EMT-6 murine mammary carcinoma. *Semin Oncol* 26: 55-62
 17. Stathopoulos GP, Dimitroulis J, Toubis M, Katis C, Karaindros D, Stathopoulos J, Koutandos J (2007) Pemetrexed combined with paclitaxel in patients with advanced or metastatic non-small-cell lung cancer: a phase I-II trial. *Lung Cancer* 57: 66-71
 18. Hughes WT, Armstrong D, Bodey GP, Bow EJ, Brown AE, Calandra T, Feld R, Pizzo PA, Rolston KV, Shenep JL, Young LS (2002) 2002 guidelines for the use of antimicrobial agents in neutropenic patients with cancer. *Clin Infect Dis* 34: 730-751
 19. Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van GM, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National

Cancer Institute of the United States, National Cancer Institute of Canada. J Natl Cancer Inst 92: 205-216

20. Hanna N, Shepherd FA, Fossella FV, Pereira JR, De MF, von PJ, Gatzemeier U, Tsao TC, Pless M, Muller T, Lim HL, Desch C, Szondy K, Gervais R, Shaharyar, Manegold C, Paul S, Paoletti P, Einhorn L, Bunn PA, Jr. (2004) Randomized phase III trial of pemetrexed versus docetaxel in patients with non-small-cell lung cancer previously treated with chemotherapy. J Clin Oncol 22: 1589-1597
21. Hanauske AR, Dumez H, Piccart M, Yilmaz E, Graefe T, Gil T, Simms L, Musib L, Awada A (2009) Pemetrexed combined with paclitaxel: a dose-finding study evaluating three schedules in solid tumors. Invest New Drugs 27: 356-365
22. Graefe T, Lubbing C, Bolling C, Muller-Hagen S, Leisner B, Fleet J, Ludtke FE, Blatter J, Suri A, Hanauske AR (2004) Phase I study of pemetrexed plus paclitaxel in patients with solid tumor. J Clin Oncol ASCO Annual Meeting Proceedings 22: No 14S (abstr 2103)
23. Garland LL, Cranmer L, Gerstner G, Walker R, Obregon Y, Ebbinghaus S (2007) Phase I dose escalation trial of biweekly pemetrexed in combination with docetaxel in advanced solid tumors. J Clin Oncol ASCO Annual Meeting Proceedings Part I 25: No 18S (abstr 13019)
24. Petrylak DP, Hood K, Xu R, Gill JF, Nicol SJ (2008) Phase I trial of pemetrexed (P) plus docetaxel (D) in advanced malignancies. J Clin Oncol ASCO Annual Meeting Proceedings 26: abstr 13517
25. Mackay HJ, Carmichael J, Roberts PJ, Glasspool R, Molife R, Macham M, Readett D, Vasey P (2002) Phase I study of pemetrexed in combination with

- docetaxel in patients with advanced solid malignancies. J Clin Oncol ASCO Annual Meeting Proceedings 21: abstr 2120
26. Lopez-Martin A, Cubedo R, Garcia-Rivas I, Castellano D, Perez-Romero A, Savulsky C, Rodriguez R, Espana P, Cortes-Funes H, Paz-Ares L (2005) Phase I trial of the sequential administration of pemetrexed (P) and docetaxel (D) in patients (pts) with advanced solid tumours. J Clin Oncol ASCO Annual Meeting Proceedings Part I 23: No. 16S (abstr 2108)
 27. Rusthoven JJ, Eisenhauer E, Butts C, Gregg R, Dancey J, Fisher B, Iglesias J (1999) Multitargeted antifolate LY231514 as first-line chemotherapy for patients with advanced non-small-cell lung cancer: A phase II study. National Cancer Institute of Canada Clinical Trials Group. J Clin Oncol 17: 1194-1199
 28. Clarke SJ, Abratt R, Goedhals L, Boyer MJ, Millward MJ, Ackland SP (2002) Phase II trial of pemetrexed disodium (ALIMTA, LY231514) in chemotherapy-naïve patients with advanced non-small-cell lung cancer. Ann Oncol 13: 737-741
 29. Smit EF, Mattson K, von PJ, Manegold C, Clarke S, Postmus PE (2003) ALIMTA (pemetrexed disodium) as second-line treatment of non-small-cell lung cancer: a phase II study. Ann Oncol 14: 455-460
 30. Bajetta E, Celio L, Buzzoni R, Ferrari L, Marchiano A, Martinetti A, Longarini R, Becerra C, Ilardi C, John W (2003) Phase II study of pemetrexed disodium (Alimta) administered with oral folic acid in patients with advanced gastric cancer. Ann Oncol 14: 1543-1548

Table 1 Dose levels and DLTs

Level (no of Patients)	Doses (mg/m ²)		No of DLTs	DLT events
	D	P		
1 st (n=3pts)	65	400	-	None
2 nd (n=6pts)	65	450	2	Neutropenia grIV (n=1), Febrile neutropenia grIII (n=1)
3 rd (n=6pts)	70	450	1	Thrombocytopenia grIV (n=1)
4 th (n=6pts)	70	500	2	Neutropenia grIV (n=2)
5 th (n=3pts)	75	500	-	None
6 th (n=6pts)	80	500	2	Neutropenia grIV (n=1), rash grIII (n=1)
7 th (n=6pts)	80	550	1	Neutropenia grIV (n=1)
8 th (n=6pts)	85	500	2	Neutropenia grIV (n=2)

Table 2 Patient characteristics

No of enrolled patients	62
Age, median (range)	63 (37-80)
Sex (no)	
Male	50 (81%)
Female	12 (19%)
Stage (no)	
IIIB	11 (18%)
IV	51 (82%)
Performance status (no)	
0-1	54 (87%)
2	8 (13%)
Histology (no)	
Squamous	14 (23%)
Adenocarcinoma	36 (58%)
Large cell	2 (3%)
Mixed	2 (3%)
Non otherwise specified	8 (13%)
No of prior regimens	
No prior therapy	21 (34%)
1 regimen	41 (66%)
Prior chemotherapy regimens	
Platinum-based	34 (55%)
Taxane-based	35 (57%)

Table 3 Hematologic toxicity in all cycles and dose levels

Level/ No of cy	Docetaxel	Pemetrexed	<u>Neutropenia</u> no of patients (%)			<u>Anemia</u> no of patients (%)			<u>Thrombocytopenia</u> no of patients (%)			<u>Febrile neutropenia</u>
			Grade			Grade			Grade			
			II	III	IV	II	III	IV	II	III	IV	
1 ^o / 7	65	400	-	2 (28.6)	-	1 (14.3)	-	-	-	-	-	GrIII=1
2 ^o / 25	65	450	1 (4.0)	2 (8.0)	1 (4.0)	-	-	-	-	-	-	GrIII=1
3 ^o / 20	70	450	2 (10.0)	2 (10.0)	1 (5.0)	-	1 (5.0)	-	-	-	1 (5.0)	-
4 ^o / 28	70	500	3 (10.7)	1 (3.6)	2 (7.1)	6 (21.4)	-	-	-	-	-	-
5 ^o / 11	75	500	-	-	-	-	-	-	-	-	-	GrIII=1
6 ^o / 22	80	500	4 (18,2)	1 (4.5)	1 (4.5)	-	-	-	-	1 (4.5)	-	GrIII=1
7 ^o / 30	80	550	1 (3.3)	2 (6.7)	3 (10.0)	1 (3.3)	-	-	-	-	-	-
8 ^o / 23	85	500	3 (13.0)	3 (13.0)	3 (13.0)	1 (4.3)	-	-	-	1 (4.3)	-	-

Table 4 Non- hematologic toxicity in all cycles and dose levels

			<u>Nausea/ Vomiting</u> no of patients (%)		<u>Diarrhea</u> no of patients (%)			<u>Asthenia</u> no of patients (%)			<u>Others</u>
n=42			Grade		Grade			Grade			
Level/ No of cy	Docetaxel	Pemetrexed	II	III	II	III	IV	II	III	IV	
1 ^o / 7	65	400	2 (28.6)	-	-	-		-	-	-	-
2 ^o / 25	65	450	-	-	-	-	-	3 (12.0)	-	-	Non neutropenic infection (n=1)
3 ^o / 20	70	450	1 (5.0)	-	1 (5.0)	-	1 (5.0)	1 (5.0)	2 (10.0)	-	-
4 ^o / 28	70	500	-	-	2 (7.1)	-	-	4 (14.3)	-	-	-
5 ^o / 11	75	500	2 (18.2)	-	-	-	-	1 (9.1)	-	-	-
6 ^o / 22	80	500	1 (4.5)	-	1 (4.5)	-	-	3 (13.6)	1 (4.5)	-	Generalized Skin rash GrIII (n=2)
7 ^o / 30	80	550	2 (6.7)	-	1 (3.3)	-	-	6 (20.0)	-	-	-
8 ^o / 23	85	500	-	-	-	3 (13.0)	-	2 (8.7)	-	-	

Table 5 Toxicity at the RD (docetaxel 75mg/m² plus pemetrexed 500mg/m²) level

Grade	II	III	IV
Toxicity#	<u>no of patients (%)</u>	<u>no of patients (%)</u>	<u>no of patients (%)</u>
Neutropenia	1 (4.3)	3 (13)	3 (13)
Febrile neutropenia	-	1 (4.3)	-
Anemia	7 (30.4)	1 (4.3)	-
Thrombocytopenia	1 (4.3)	-	-
Nausea	1 (4.3)	-	-
Vomiting	1 (4.3)	1 (4.3)	-
Diarrhea	1 (4.3)	-	-
Asthenia	2 (8.7)	-	-

Worst toxicity in all cycles and all patients enrolled in the RD level