Lithospheric folding and sedimentary basin evolution: a review and analysis of formation mechanisms

Sierd Cloething, Evgenii E.B. Burov

► To cite this version:

Sierd Cloething, Evgenii E.B. Burov. Lithospheric folding and sedimentary basin evolution: a review and analysis of formation mechanisms. *Basin Research*, 2011, 23 (3), pp.257-290. 10.1111/j.1365-2117.2010.00490.x . hal-00640964

HAL Id: hal-00640964

<https://hal.science/hal-00640964>

Submitted on 24 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lithospheric folding and sedimentary basin evolution: a review and analysis of formation mechanisms

Journal:	<i>Basin Research</i>
Manuscript ID:	BRE-070-2009.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	
Complete List of Authors:	Cloetingh, Sierd; VU University, Earth & Life Sciences Burov, Evgenii; Université Pierre et Marie Curie, Laboratoire de Tectonique
Keywords:	tectonics and sedimentation, geodynamics, rift basins, foreland basins, Intracratonic basins, strike-slip basins, tectonic geomorphology

Lithospheric folding and sedimentary basin evolution:
a review and analysis of formation mechanisms

Sierd Cloetingh^{a,*} and Evgenii Burov^b

^aNetherlands Research Centre for Integrated Solid Earth Sciences, Faculty of Earth and Life Sciences, VU University Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands.

^bUniversité Pierre et Marie Curie, Laboratoire de Tectonique, 4 Place Jussieu, 75252 Paris Cedex 05, France.

Basin Research

Submitted: August 17, 2009

Revised: March 18, 2010

* Corresponding author: Fax: +31-20- 598 99 43
E-mail addresses: sierd.cloetingh@falw.vu.nl (Sierd Cloetingh), evgenii.burov@upmc.fr (Evgenii Burov)

ABSTRACT

Lithospheric folding is an important mode of basin formation in compressional intraplate settings. Basins formed by lithospheric folding are characterized by distinct features in subsidence history. A comparison with extensional basins, foreland basins, intracratonic basins and pull-apart basins provides criteria for the discrimination between these modes of basin formation. These findings are important in deciphering the feed-backs between tectonics and surface processes. In addition, inferences on accommodation space and thermal regime have important consequences for hydrocarbon maturity. Lithospheric folding is coupled to compressional basin and fault reactivation and, therefore, strongly affects reservoir characteristics of sedimentary basins.

INTRODUCTION

Over the last few years much progress has been made in understanding the mechanisms of the formation of sedimentary basins (Cloetingh & Ziegler, 2007; Roure *et al.*, 2010). Quantitative models for several basin types have significantly advanced the understanding of extensional (McKenzie, 1978; Van Wees *et al.*, 2009) and foreland basins (Beaumont, 1981; Naylor & Sinclair, 2008). The mechanisms of basin formation as a result of lithospheric folding have received considerably less attention (Cloetingh & Ziegler, 2007) except for some clear-cut cases, in particular in central Asia (Burov & Molnar, 1998; Thomas *et al.*, 1999a,b). Many data exist on the geometry of sedimentary basins affected by large-scale compressional intraplate deformation (Cobbold *et al.*, 1993; Lefort & Agarwal, 1996, 2000, 2002; Burov & Molnar, 1998). These observations and results of analytical, numerical (Martinod & Davy, 1992; Burov *et al.*, 1993; Gerbault *et al.*, 1998; Cloetingh *et al.*, 1999) and analogue modelling (Martinod & Davy, 1994; Sokoutis *et al.*, 2005) demonstrate that the thermo-mechanical age of the lithosphere exerts a prime control on the wavelength of lithospheric folds. These studies focused on the role of tectonic stress in lithospheric folding

1
2
3 50 (Stephenson *et al.*, 1990; Burov & Molnar, 1998; Stephenson & Cloetingh, 1991; Perez-Gussinye &
4
5 51 Watts, 2005). However, surface processes also play a significant role in the mechanics of
6
7 52 lithospheric folding. Erosion enhances the development of folding and has a pronounced effect on
8
9
10 53 its wavelengths (Cloetingh *et al.*, 1999; Burov & Toussaint, 2007), specifically in the short-
11
12 54 wavelength domain. In addition, both sedimentation and erosion are likely to significantly prolong
13
14 55 the lifetime of folding. Sedimentation decreases the effect of gravity by filling the downward flexed
15
16 56 basins and thus reducing the isostatic restoring force, whereas erosion of the upward flexed
17
18 57 basement unloads the lithosphere on the basins' uplifted flanks. Lithospheric folding has significant
19
20 58 effects for the geometry of sedimentary sequences deposited on folded lithosphere. However, the
21
22 59 effect has been studied little for several reasons (Cobbold *et al.*, 1993). The hanging walls of
23
24 60 emerging low-angle thrust faults tend to collapse rapidly, blurring the structural relationships at
25
26 61 basin edges. Further, ongoing sedimentation may onlap and bury the hanging walls, whereas scarp
27
28 62 erosion may destroy them. Seismic images of basin edges may not be well resolved where footwall
29
30 63 conglomerates contain few reflecting horizons or when low-angle thrusts reflect and refract seismic
31
32 64 waves, thus hampering the imaging of deeper parts of the basin. Downwarping of folded basins
33
34 65 renders surface geological mapping difficult. At the same time, high plateaux are not at first sight
35
36 66 linked with basins of high hydrocarbon potential, leaving them in many places without well control or
37
38 67 seismic constraints on basin architecture.
39
40
41
42 68 In this paper we review evidence from a number of basins in Eurasia, where folding has been
43
44 69 documented and alternative basin formation mechanisms such as lithospheric extension or foreland
45
46 70 flexure cannot fully explain key features of basin geometry and evolution. We restrict ourselves to
47
48 71 cases where independent evidence points to a major role for lithospheric folding in the basin
49
50 72 development. We demonstrate that an interpretation in terms of folding can sometimes resolve
51
52 73 inconsistencies resulting from interpreting basins origin via other mechanisms. For example
53
54 74 (Ritzmann & Faleide, 2009), the formation mechanism of intracratonic basins is still a matter of
55
56 75 debate. A striking feature of these basins is the prolonged intervals of low rate subsidence
57
58
59
60

alternating with episodic accelerations in subsidence rates, often coeval with orogenic activity at plate boundaries. Due to the frequent absence of evidence for thinned lithosphere below these basins, alternative models have been proposed. Stel *et al.* (1993) propose that non-extensional crustal thinning is induced by basaltic underplating, which causes thermal uplift and erosional thinning. Tectonic loading at plate boundaries is also proposed as a cause of or contributor to large-scale sag-type subsidence in plate interiors (Cloetingh, 1988; Leighton & Kolata, 1990; Quinlan, 1987) as well as phase transformations (Artyushkov, 2007). Ritzmann & Faleide (2009) point out that intraplate stresses and crustal inhomogeneities coupled with loading scenarios provide the best explanation for the Barents Sea Basin and other intracratonic basin evolution. The basins reviewed here, however, do not include cases such as the Barents Sea where the effect of stresses on topography and accommodation space has been considerably complicated by other basin formation processes.

We present inferences from thermo-mechanical models of lithospheric folding and a comparison with the results of analogue modelling of folding. We concentrate on the overall geometry of basins formed by lithospheric folding, the typical timescales intrinsic to their development and the brittle deformation patterns in the underlying lithosphere. We then discuss subsidence patterns and heat flow characteristics for such basins. We conclude by comparing key features of basin accommodation shape, vertical motions, thermal history and brittle deformation patterns of basins formed on folded lithosphere, foreland basins, intracratonic basins, extensional basins and pull-apart basins.

LITHOSPHERIC FOLDING: AN IMPORTANT MODE OF INTRAPLATE BASIN FORMATION AND DEFORMATION

Folding of the lithosphere, involving positive and negative deflections (**Fig. 1**), appears to be of more importance in the large-scale deformation of intraplate domains than hitherto realized

(Cloetingh *et al.*, 1999). As has been shown (Burov *et al.*, 1993; Nikishin *et al.*, 1993; Cloetingh *et al.*, 1999; Gerbault *et al.*, 1998; Schmalholtz & Podladchikov, 2000), visco-elastic folding starts to develop from the onset of compression and in contrast to elastic folding, does not require specifically large intraplate stresses. Folding may continue until the back-ground strain-rate drops, for example due to the localization of deformation in a specific area or to reduction of far-field forces. Such intraplate deformation results from transmission of intraplate stress fields away from plate boundaries into continental forelands (Ziegler *et al.*, 1998; Van der Pluijm *et al.*, 1997; Tesauro *et al.*, 2005).

In continental lithosphere, several wavelengths of folding are expected to develop as result of the presence of several rheologically competent layers decoupled from each other by weak layers (**Fig. 1, Fig. 3**). Crustal folding can be traced from the surface topography, neotectonic movements and free air gravity. The crust-mantle boundary (Moho) topography reflects mantle folding, because the mantle strength is highest at this interface and thus the geometry of Moho deflection is most likely to correspond to that of mantle folding. This wavelength can be traced from Bouguer gravity anomalies because the crust-mantle boundary represents also the main density contrast in the lithosphere. It is to be expected that the base of the Mechanical Lithosphere (MLB) has roughly the same geometry as the Moho deflection. Likely, the lithosphere-asthenosphere boundary (LAB) has little to do with mantle lithosphere folding except some specific cases, as the MLB (roughly corresponding to 700°C depth) is found well above the LAB (1330°C). Consequently, the low viscosity zone between MLB and LAB may damp the deformation in a way that the LAB remains even flat. Due to the low viscosity of gravitating mantle between the LAB and MLB, the LAB is not necessarily advected or down-warped with deflection of the MLB, or at least the relations between the deformation of the MLB and LAB are not straightforward. In particular, gravity instabilities or convective movements not related to folding can perturb the LAB without impact on the MLB.

Folding has important implications for vertical motions, sedimentary basin architecture and the evolution of hydrocarbon systems (Ziegler *et al.*, 1995; 1998). The large wavelength of vertical motions associated with lithospheric folding necessitates the integration of data from large areas (Elfrink, 2001; Allen & Davies, 2007), often beyond the scope of regional structural and geophysical studies that target specific structural provinces. Recent studies on the North German Basin show neotectonic reactivation by lithospheric folding (Marotta *et al.*, 2000). Similarly, acceleration of the Plio-Pleistocene subsidence in the North Sea Basin is attributed to stress-induced buckling of its lithosphere (Van Wees & Cloetingh, 1996). Folding of the Variscan lithosphere has been documented for Brittany (Bonnet *et al.*, 2000; Lagarde *et al.*, 2000), the adjacent Paris Basin (Lefort & Agarwal, 1996) and the Vosges-Black Forest arch (Dèzes *et al.*, 2004; Ziegler & Dèzes, 2007; Bourgeois *et al.*, 2007). Lithospheric folding, therefore, appears to be an effective mechanism for the propagation of tectonic deformation from active plate boundaries far into intraplate domains (e.g. Stephenson & Cloetingh, 1991; Burov *et al.*, 1993; Ziegler *et al.*, 1995, 1998).

Folding can be observed at different spatial scales. At the scale of a micro-continent that was affected by a succession of collisional events, Iberia illustrates lithospheric folding and interplay between neotectonics and surface processes (Cloetingh *et al.*, 2002). An important factor favouring a lithosphere-folding scenario for Iberia is the compatibility of the thermo-tectonic age of its lithosphere and the wavelength of observed deformations.

Other well-documented examples of continental lithospheric folding occur in other cratons (**Fig. 2**). Prominent examples occur in the Western Gobi area, in the Ferghana basin, but also probably in the Tadjik basin of Central Asia, involving a lithosphere with thermo-tectonic ages from 150 Ma (Kazakh shield, Ferghana, Tadjik basins and likely Dzhungaria basin) to 400 Ma (Tarim basin). In this area, mantle and crustal wavelengths are 360 km and 50 km, respectively, with a shortening rate of ~10-20 mm/yr and 200-250 km of shortening during 10-15 Myr (Burov *et al.*, 1993; Burov & Molnar, 1998). Three-dimensional fold structures in the Tibetan Plateau have been inferred from

GRACE satellite gravity data, demonstrating prevailing fold wavelengths of 300 to 420 km (Shin *et al.*, 2009).

The inferred wavelength of these neotectonic lithosphere folds is consistent with the general relationship between the wavelength of lithospheric folds and the thermo-tectonic age of the lithosphere shown by a global inventory of lithospheric folds (**Fig. 3**) (Cloetingh & Burov, 1996; Cloetingh *et al.*, 2005). In some other areas of continental lithosphere folding, smaller wavelength crustal folds have also been detected (Burov *et al.*, 1993; Nikishin *et al.*, 1993). Thermal thinning of the mantle lithosphere, often associated with volcanism and doming, enhances lithospheric folding and appears to affect the wavelengths of folds (Burov & Cloetingh, 2009).

OBSERVATIONS

Basins developed on folded continental lithosphere have characteristic features. We illustrate this by largely concentrating on the basins of Central Asia, the folded lithosphere of Iberia, the NW European platform, the Pannonian Basin and the South Caspian Basin.

Compressional basins of Central Asia (Tianshan, Ferghana)

The young Ferghana and Tadjik basins are compressional basins northwest and north of the Pamir and south of the Tien Shan ranges (Burov & Molnar, 1998). In this area lithosphere underwent Jurassic reactivation and is characterized by relatively young thermo-mechanical ages (175 Ma) (Burg *et al.*, 1994; Burov & Molnar, 1998). In addition to having undergone thermal weakening, the lithosphere underlying these basins also probably has a weak, quartz-dominated lower crustal rheology. This has resulted in low values for the effective elastic plate thickness (EET) of the order of 15 km (Burov & Molnar, 1998).

The Central Tien Shan consists of an alternation of ranges and basins separated by reverse faults.

The Tien Shan terminates west of the right-lateral Talas-Ferghana fault by splaying into two narrow

mountain chains that surround the Ferghana Valley. A narrow belt of mountains parallel to the central section of the fault slopes downward to the Ferghana Valley, underlain by a deep basin with as much as 8 km of Cretaceous-Cenozoic sediments (Cobbold *et al.*, 1993). Gravity data suggest that the Ferghana and Tadjik basins are gravitationally overcompensated. The negative Bouguer gravity anomalies indicate a Moho several km deeper than predicted by models of local Airy isostasy (Burov & Molnar, 1998). This points to Moho down-warping due to non-isostatic processes and most probably, in this compressional context, to lithospheric folding. As demonstrated by numerical-thermomechanical experiments (Burov & Molnar, 1998), the approximately north-south shortening of the relatively thin lithosphere could have created central down-warping and anticline mountains north and south of the basin. According to this model, this shortening has produced a folding instability in the mantle lithosphere that has warped the basement immediately surrounding the basin upward, and has forced the basin floor down beneath the Ferghana Valley (with an estimate for the folding wavelength of the order of 200-250 km), possibly also producing deep mantle faulting (seen as Moho offsets in seismic reflection data).

Pre-existing thermal structure and variations in crustal thickness have played a major control on the styles and distribution of faulting in this region (**Fig. 4**), (see also Burov & Molnar 1998; Cobbold *et al.*, 1993). Primary faults appear before the folding develops with a spacing that is proportional to brittle layer thickness. Subsequently the two processes, faulting and folding, co-exist in such a way that faults localize at the inflection points of folds. At this stage, the appearance of faults does not significantly influence the wavelength of folding. Because of the weakness of the lower crust, the upper crust should be completely decoupled from the mantle and interact with it only by flow in the lower crust. Lower crustal flow changes the wavelength and amplitude of the surface folding, which is terminated by the development of a single down-warped megafold (Cloetingh *et al.*, 1999). Liu *et al.* (2007) reported evidence for lithospheric folding in the Central/ North China Tarim Basin region, initiated during Pliocene times, with good time constraints provided by Ar/Ar dating. Two wavelengths are found of 30 and 400 km. The thermo-mechanical age of the lithosphere is

estimated to be 250 Ma, as the main thermal perturbation was associated with a major orogenic phase at 250 Ma ago. These findings are consistent with models in which decoupled lithosphere folds with 30 km and 400 km wavelength correspond to, respectively, crustal and mantle folding. Deep seismic data for basins north of and within the Tibetan plateau (Li *et al.*, 2006; Liu *et al.*, 2006; Zhao *et al.*, 2006) show that folding is not limited to the Tibetan plateau (see also Shin *et al.*, 2009).

Folded lithosphere of Iberia

As reviewed by Cloetingh *et al.* (2002), Tertiary lithospheric folding of Iberia occurs in Variscan lithosphere with wavelengths of 300 km, leading to the development of a system of parallel trending basins and highs (**Fig. 5a**). This folding generated the distribution of basins and mountain chains, bounded by folds and faults (Cloetingh *et al.*, 2002; De Vicente *et al.*, 2007). The regularity of the fluvial network pattern of the central-western part of Iberia also indicates large-scale lithospheric folding. Alpine (mainly Cantabrian-Pyrenean related) compressional tectonics was the principal factor for closing internally drained sedimentary basins, including the Ebro, Duero and Tagus basins (Casas-Sainz & De Vicente, 2009).

A large body of geophysical and geological observations is available on the crustal geometries and stress regimes of Iberia (De Vicente *et al.*, 2007, 2008). This is supplemented by thermogeochronology data (De Bruijne & Andriessen, 2002; Ter Voorde *et al.*, 2004), demonstrating accelerated uplift of the highs during Late Miocene-Pliocene times. In spite of the existence of recent localized volcanism in the inner part of the Iberian plate, an important contribution from thermal uplift is questionable due to the distribution of crustal thickness (Casas-Sainz & De Vicente, 2009). Fernandez-Lozano *et al.* (2010) have constructed a three-layer analogue model consisting of brittle upper crust, ductile lower crust and ductile upper mantle for Iberia. These models show that folding is associated with deformation of narrow mountain ranges separated by basins (see **Fig. 5b**). The narrow mountain ranges representing upper crustal pop-ups form the main topographic reliefs. Shortening is accommodated within the viscous crust beneath the pop-ups leading to lateral

thickness variations of the ductile crust. These results are consistent with seismic and gravity data collected in the Cantabrian mountains, the Spanish Central System, and the Toledo Mountains (see also **Fig. 5a**).

The earliest sediments in the terrestrial basins (Calvo et al., 1993) were deposited during the Eocene, with very well developed continuous successions, dominated by continental facies with high sedimentation rates. The Duero and Tagus basins began to be individualized at the onset of uplift and the formation of a topographic range in the Central System (Casas-Sainz & De Vicente, 2009). In contrast with this early more homogeneous stage, the Oligocene-Lower Miocene stratigraphy of the basins is characterized by several tectono-stratigraphic units, separated by unconformities representing short periods of time. The number and specific ages of these units varies between basins, indicating variable activity at basin borders. The fact that the largest number of breaks in the sedimentary succession is concentrated during this time interval points to a tectonic control (De Vicente & Vegas, 2009).

The final stages in the evolution of these onshore basins are characterized by widespread Miocene lacustrine sedimentation in the endorheic basins and during Pliocene and Quaternary times, by the beginning of river capture and exorheism in the Duero, Ebro and Tagus basins (Garcia-Castellanos *et al.*, 2003). Alluvial deposits were deposited by the fluvial network developed on the ancient, internally drained basins (Casas-Sainz & De Vicente, 2009). During the endorheic-exorheic transition, paleo-karstification processes and erosion occurred in the Late Miocene lacustrine limestones. Two main types of high plains can be distinguished on the Iberian Peninsula. The coupled origin of both types of surfaces can be related to uplift/sedimentation histories along the Tertiary compressional basin margins in the Iberian Peninsula (Casas-Sainz & De Vicente, 2009). The first are high plains with Tertiary sedimentary basins, containing well preserved Late Miocene to Pliocene strata with subhorizontal bedding. Typical examples are the Duero and Tagus basins. The second consists of eroded sediments of Mesozoic and Paleozoic rocks with planation surfaces developed on the Paleogene-Lower Miocene ranges created by fold and thrust systems and

1
2
3 255 basement uplifts. Different patterns in evolution and stratigraphy between the eastern and western
4
5 256 Tagus basin may be related to the mechanical decoupling between Iberia and Africa during the
6
7 257 Miocene emplacement of the Alboran Domain in the southern part of the plate. This generated
8
9 258 extensional stresses in the eastern Iberian plate, coeval with compression and lithospheric folding in
10
11 259 the western part of the plate (De Vicente *et al.*, 2008; De Vicente & Vegas, 2009).
12
13
14 260 In general the dimensions of peneplains linked to extensional basins and passive margins are
15
16 261 larger, and their evolution involves time scales much longer than those characteristic for high plains
17
18 262 of the Iberian plate (Casas-Sainz & De Vicente, 2009). Peneplains associated with compressional
19
20 263 or convergent regimes (e.g. Anatolia, Central Iran and Morocco) are characterized by spatial
21
22 264 dimensions comparable with Iberia. In addition, their tectonic setting in the hanging wall of the
23
24 265 African-Eurasian convergence zone is also similar. Thus the Iberian plate may be an analogue for
25
26 266 other present-day intra-mountain basins.
27
28
29
30

31 267
32
33 268 **The European Alpine foreland**

34 269 Geophysical and geomorphological studies (e.g. Lefort & Agarwal, 1996; 2000; 2002; Marotta et al.,
35
36 270 2000) indicate large-scale compressional deformation of the European Alpine foreland lithosphere.
37
38 271 Although different in structural grain and rheological structure (Tesauro et al., 2007; Cloetingh et al.,
39
40 272 2005) inherited from differences in their Paleozoic-Mesozoic history, different segments of the
41
42 273 lithosphere in the northern Alpine European foreland share many similarities in their Cenozoic
43
44 274 intraplate deformation. Seismic and geological evidence shows that the North German Basin cannot
45
46 275 be explained by classical models of basin formation (Marotta et al., 2000). The basin underwent a
47
48 276 polyphase history (Scheck & Bayer, 1999; Littke et al., 2008) of extension in the late Triassic
49
50 277 followed by lithospheric scale folding in the late Cretaceous-Early Cenozoic (Mazur *et al.*, 2005;
51
52 278 Marotta *et al.*, 2000). 3D volumetric analysis shows that the compressional event was followed by
53
54 279 faster subsidence during the Cenozoic (Scheck & Bayer, 1999). A similar observation was made by
55
56 280 Van Wees & Cloetingh (1996) for the North Sea Basin. Lithospheric folding modified a basin formed
57
58
59
60

initially by deep seated thermal perturbations (Scheck & Bayer, 1999) with a thinned crust and shallow Moho. The Moho updoming with a high of 3 km for the height of the bulge in the southern part of the North German basin appears to be the consequence of flexural buckling of the previously thinned lithosphere by a compressive stress perpendicular to the strike of the basin.

From earliest Triassic to Late Jurassic, the Paris Basin (**Fig. 6a**) subsided in an extensional framework and was larger than the present basin (Guillocheau *et al.*, 2000). The western margin of the Paris Basin and the rifted Atlantic margin of France were subject to thermal rejuvenation during Mesozoic extension related to North Atlantic rifting (Ziegler & Dèzes, 2006; Robin *et al.*, 2003). Subsequent compressional intraplate deformation (Ziegler *et al.*, 1995) also affected the Paris Basin (Lefort & Agarwal, 1996). Numerical modelling indicates that the amplitude of the compression induced vertical deflection in the Paris Basin is relatively small, requiring additional mechanisms such as flexure due to the load by the Alpine system (Bourgeois *et al.*, 2007; Ziegler & Dèzes, 2007). However, periodic deformation at the same wavelength occurs northwest of the Paris Basin (**Fig. 6 a**, Bourgeois *et al.*, 2007), which would not be the case if the north-eastern part of the basin was flexed down by the load of the Alps. Bourgeois *et al.* (2007) carefully analysed basement geometry and timing of vertical motions in the NW European platform. They separate the contributions from the European Cenozoic Rift System (ECRIS) and the contributions from long-wavelength folding striking NE and located between the Alpine front and the North Sea. According to their analysis, the ECRIS developed mostly between 37 and 17 Ma, whereas lithospheric folds developed between 17 Ma and present, with a wavelength of 270 km and amplitude of 1500 m (see **Fig. 6b**).

Quaternary folding of the Variscan lithosphere in the area of the Armorican Massif (Bonnet *et al.*, 2000) developed folds with a wavelength of 250 km, pointing to a mantle-lithospheric control on deformation. The timing and spatial pattern of uplift inferred from river incision studies in Brittany is incompatible with a glacio-eustatic origin. Therefore, Bonnet *et al.* (2000) related the vertical motions to deflection of the lithosphere by the present-day NW-SE directed compressional intraplate

1
2
3 307 stress field. Stress-induced uplift of the area appears to control fluvial incision rates and the position
4
5 308 of the main drainage divides. Leveling studies (LeNotre *et al.*, 1999) indicate its ongoing
6
7 309 deformation.
8
9 310
10
11 311 **Pannonian Basin System**
12
13 312 Recent studies of the Pannonian Basin system - formed as a Miocene back-arc basin behind the
14
15 313 Carpathian arc - show that active crustal deformation resulted in significant differential vertical
16
17 314 motions during Late Neogene times (Horvath & Cloetingh, 1996; Cloetingh *et al.*, 2005, 2006;
18
19 315 Horvath *et al.*, 2006; Matenco *et al.*, 2007; **Fig. 7a**). The river network in the area is affected by
20
21 316 differential neotectonic crustal motions (Necea *et al.*, 2005; Dombradi *et al.*, 2010).
22
23 317 The Late Neogene evolution of the Pannonian Basin is largely controlled by its interaction with the
24
25 318 Carpathian arc and the Adriatic indenter impinging the Pannonian basin lithosphere (e.g. Bada *et al.*,
26
27 319 2001). As a result of the consumption of subductable lithosphere of the European foreland, the
28
29 320 Pannonian basin became locked. At the same time, the continuous N-NE directed indentation of the
30
31 321 Adriatic microplate built up a high level of compressional stresses in the Pannonian Basin. The
32
33 322 structural inversion of the basin is a direct consequence of the temporal changes in the stress field
34
35 323 from extension to compression. Recent studies of the stress field together with an extensive array of
36
37 324 geological and geophysical data have shown that horizontal stresses are transmitted from the plate
38
39 325 boundaries into the interior of the Pannonian Basin (Bada *et al.*, 2007).
40
41 326 The Pannonian Basin is underlain by what is probably the hottest lithosphere of continental Europe,
42
43 327 characterized by a very low lithospheric strength (Tesauro *et al.*, 2007). Miocene extension resulted
44
45 328 in a high level of mantle stretching, leading to a strength distribution with practically zero strength in
46
47 329 the subcrustal lithosphere and lower crust (Cloetingh *et al.*, 2005), making it prone to tectonic
48
49 330 reactivation. Therefore, in this area relatively low compressional stresses induced mainly by plate
50
51 331 boundary interactions, and further amplified by topography-induced stresses (Bada *et al.*, 2001),
52
53 332 could initiate significant intraplate deformation.
54
55
56
57
58
59
60

It should be noted that in the Pannonian basin system, seismic tomography (Wortel & Spakman, 2000) supports slab detachment under the adjacent Romanian Carpathians. At the same time, the seismic tomography shows the presence of a hot upper mantle below the folded lithosphere in the Pannonian basin. In this area, therefore, the upper mantle structure reflects more the back-arc extension prior to the folding than the consequences of folding on the density stratification of the mantle. The reverse is true for the crustal configuration. The geometries imaged by the crustal scale deep seismic reflection profiling are clearly dominated by compressional structures resulting from folding (**Fig. 7b**).

Quaternary patterns of differential uplift and subsidence with a timing coeval with an increased level of compressional stresses are interpreted as caused by lithospheric folding (Horvath & Cloetingh, 1996; Cloetingh *et al.*, 2006; Horvath *et al.*, 2006; Bada *et al.*, 2007). Late-stage compressional deformation of the Pannonian basin is demonstrated to be of key importance for the geometry of basin fill (Sacchi *et al.*, 1999, see **Fig. 7b**) and for its reservoir structures, both crucial in assessment of its hydrocarbon potential (Horváth, 1995). Analogue tectonic experiments by Dombradi *et al.* (2010) further examine accumulation and transmission of stresses in the hot and weak back-arc lithosphere of the Pannonian basin and the interplay between lithospheric processes and the intrabasinal topography development. Based on available rheological and seismotectonic constraints, the basin system is represented by a two-layer model with a 10-15 km thick brittle upper crust. The lower crust and the mantle are represented by a low-strength layer of 45 km thickness (Dombradi *et al.*, 2010). The strain rates applied were constrained by GPS velocities (Grenerczy *et al.*, 2005). In the experiments lithospheric folds develop with wavelengths equivalent to 200-250 km (**Fig. 7c**). The experiments also show that crustal thickness variations largely determine the localisation of stresses. Topographic profiles extracted by laser scans of the model show that surface topography reflects long wavelength buckling of the lithosphere. These results are compatible with Pliocene-Quaternary subsidence, which occurs in the centre of the Pannonian basin, simultaneously with accelerated uplift at the basin margins. The latter has lead to elevated

1
2
3 359 topography of up to 1 km in areas at the rim of the Pannonian Basin such as the Styrian, Zala and
4
5 360 Transylvanian basins (Cloetingh *et al.*, 2006).
6
7 361
8
9
10 362 **The South Caspian Basin**
11
12 363 The South Caspian Basin with an estimated sedimentary thickness up to 20 km (Brunet *et al.*, 2003;
13
14 364 Egan *et al.*, 2009), is probably one the deepest basins in the world. It is flanked by the Alborz
15
16 365 Mountains to the south, with an elevation of 2-4 km (see **Fig. 8**). The transition between basin and
17
18 366 flanking high is abrupt and coincides with a coastline bounded by a major fault system. The
19
20 367 formation of the South Caspian Basin is enigmatic. Attempts to explain it as an extensional basin or
21
22 368 as a foreland basin have been unsuccessful. There is a lack of evidence for basin extension and
23
24 369 both stretching models and topographic loads and slab pull operating on down flexed lithosphere in
25
26 370 foreland basin systems cannot explain differential vertical motions of this order of magnitude. As a
27
28 371 result, phase changes in continental crust have been proposed as a mechanism for the formation of
29
30 372 this superdeep basin (Artyushkov, 2007). A limitation in conclusively resolving the basin formation
31
32 373 mechanism is the lack of multi-channel seismic reflection data capable of imaging crustal structure
33
34 374 below the thick sedimentary sequences. Over the last few years, however, other constraints on the
35
36 375 area have become available (e.g. Guest *et al.*, 2007).
37
38 376 **Figure 8** presents a cross section of northern Iran and the South Caspian Basin (Guest *et al.*,
39
40 377 2007). As shown, wavelengths for compressional folding of the lithosphere are typically in the range
41
42 378 of 400 km. Although the actual basin formation mechanism for the Caspian Basin is not well
43
44 379 resolved (e.g. Artyushkov, 2007), late Neogene folding has been affecting a lithosphere probably
45
46 380 thermally reset by middle Late Jurassic marginal basin formation (Guest *et al.*, 2007) with a thermo-
47
48 381 mechanical age at the onset of collision of 130-150 Ma. Satellite data demonstrate an exceptionally
49
50 382 large gravity anomaly over the basin (Kaban *et al.*, 2002). Seismic tomography shows slab
51
52 383 detachment under the Iranian plateau (Hafkenscheid *et al.*, 2006; Alinaghi *et al.*, 2007) beginning at
53
54 384 about 10-15 Ma.
55
56
57
58
59
60

Basin analysis has shown that basement inversion has occurred 20 Ma ago, simultaneously with the slowing of Arabian/Eurasian plate convergence and the onset of accumulation of Neogene clastics in foreland basins (e.g. Fakhari *et al.*, 2008). By 10 Ma, contraction occurred by underplating of the Arabian crustal units beneath the Iranian plate. Estimated shortening rates from current geodetic surveys (Vernant *et al.*, 2004) are 7 mm/yr. Results from geothermochronology demonstrate rapid uplift of the Alborz mountains, with rates of 0.7 km/my exhumation between 6 and 4 Ma (Axen *et al.*, 2001), implying approximately 10 km of uplift. This uplift was nearly synchronous with rapid South Caspian subsidence (Nadirov *et al.*, 1997) and subsequent folding (Devlin *et al.*, 1999). South Caspian sedimentation rates locally increased more than tenfold at ca 6 Ma, with more than 10 km of sediments deposited since then. Axen *et al.* (2001) argue that if approximately 10 km of post 6 Ma sediments are present in this basin, then as much as 20 km (equivalent to 80 %) of the structural relief of about 25 km between the high Alborz and the southernmost Caspian basement may be younger than 6 Ma.

Folds in the southernmost Caspian Basin (Devlin *et al.*, 1999) and in the Neogene of the Northern Alborz foothills imply contraction. The lack of a crustal root under the Alborz Mountains also points to a flexural support by the South Caspian basement (Axen *et al.*, 2001). The simultaneous reversal of Zagros strike slip and extrusion of central Iran, coarse Zagros molasse deposition, Dead Sea transform reorganisation, Red Sea oceanic spreading (Chu & Gordon, 1998) all initiated around 5 ± 2 Ma (e.g. Axen *et al.*, 2001; Smit *et al.*, 2010), was interpreted by these authors in terms of a widespread tectonic event. Slab detachment could explain part of the recent uplift of Iran, but does not directly explain the simultaneous dramatic subsidence in the adjacent Caspian Basin. However, slab detachment could have an indirect effect if reducing slab pull forces acting on the downgoing lithosphere promote the development of compressional stresses in both the downgoing and overriding plates (Cloetingh *et al.*, 2004; Matenco *et al.*, 2007).

Guest *et al.* (2007) argue in favour of compressional deformation of the South Caspian Basin/Alborz mountains. This interpretation was largely based on available industrial seismic reflection data and

1
2
3 411 gravity data for the basin and followed a similar proposition made for the late-stage deformation of
4
5 412 the adjacent Black Sea (Nikishin *et al.*, 2003; Cloetingh *et al.*, 2008). Guest *et al.* (2007) present
6
7 413 kinematic models, without addressing the reason for the difference in the order of magnitude of the
8
9 414 basin depression and the adjacent high. Analogue experiments (Sokoutis *et al.* 2005) show that
10
11 415 when compression acts on two blocks with contrasting thickness (and rigidity), a major syncline will
12
13 416 develop on top of the suture zone separating these blocks, flanked by an anticline of much lower
14
15 417 amplitude. These results are consistent with predictions from folding theory that as a result of the
16
17 418 acting gravity field, down-warping is mechanically more effective than up-warping against the action
18
19 419 of gravity forces. The presented numerical experiments illustrate this (see for example **Fig. 9**).
20
21
22
23
24

25 421 **Basins on folded lithosphere in Eurasia: differences and similarities in lithospheric**
26
27 422 **deformation and basin habitat**
28

29 423 Comparison of lithospheric deformation in Central Asia, Iberia, the NW European Alpine foreland,
30
31 424 the Pannonian Basin and the South Caspian Basin demonstrates common features compatible with
32
33 425 an interpretation in terms of basins developed on folded lithosphere. Amongst these are their
34
35 426 symmetrical shape, dimensions and the temporal association of their basin formation history with
36
37 427 the build up of intraplate stresses. Important differences can be observed between the Central
38
39 428 Asian and Iberian basins and the South Caspian Basin on one side and the European Alpine
40
41 429 foreland and the Pannonian Basin on the other side.
42
43

44 430 The Central Asian Basins and the basins of the Iberian plate were initiated by Tertiary folding,
45
46 431 whereas the European Alpine foreland underwent a polyphase history, following Carboniferous
47
48 432 and Triassic stretching (Scheck & Bayer, 1999; Le Solleuz *et al.*, 2004), prior to Late Cretaceous-
49
50 433 Early Cenozoic folding. Similarly, the Pannonian Basin, formed by back-arc extension in Miocene
51
52 434 times, was subsequently compressionally reactivated (Cloetingh *et al.*, 2008). The origin of the
53
54 435 South Caspian Basin is not well resolved, but km scale vertical motions are compatible with a
55
56 436 megafold as a prime control on present-day basin geometry. These differences in evolution have a
57
58
59
60

pronounced impact on basin dimensions and on the nature of basin fill. The Central Asian basins, the folded basins of Iberia, and the South Caspian Basin have a wavelength compatible with theoretical predictions. In contrast, the fold wavelengths of the European Alpine foreland and the Pannonian Basin appear to be primarily controlled by the pre-existing geometry of the sediment-basement interface and pre-existing Moho topography inherited from the pre-folding stage. Differences occur also in the development of the ranges flanking the basins. In Central Asia and Iberia, these highs have amplitudes and widths comparable to the prediction for periodic folding, whereas in the Paris Basin and the North German Basin these highs are less well developed. The ratio of the depth of the South Caspian Basin to the height of the flanking Alborz Mountains is compatible with a megafold under this area of dramatic differential topography. The large amplitudes of folding at the flanks of the Pannonian Basin system are compatible with its extremely weak rheology. Differences in the amplitude of the highs might also partly result from the differences in neotectonic shortening. The highs bounding the basins of Iberia, Pannonian Basin, South Caspian Basin and Central Asia, located relatively close to the site of ongoing convergence, are still experiencing active uplift. This is much less the case for the low strain deforming areas of the NW European platform and the North German basin where the peak compression occurred during Late Cretaceous-Early Cenozoic. Another difference occurs in the depositional regime prevailing in these basins. The closed basin systems of Central Asia, Iberia and the intra-Carpathian Pannonian Basin are dominated by continental and lacustrine deposits, whereas capture is usually a late-stage feature of these basins. The Paris basin of the NW European platform has a basin fill largely dominated by marine sediments deposited during the pre-folding stage of their evolution, followed by marine Tertiary successions in the basin centre with truncated basin margins. The creation of basin highs in this case was in general not capable of interrupting open connections to marine environments.

A general observation for all these basin systems is the flatness of sedimentary sequences in the depocentres. Faulting plays a minor role in the basin development. Faulting appears to be

1
2
3 463 concentrated at the basin margins, with a highly variable depth extent, varying from upper crustal
4
5 464 scale faults bounding pop-up structures in the Central System of Iberia and the Pannonian Basin to
6
7 465 minor faulting in the basement of the North German Basin. The thermal regimes of the basins are
8
9
10 466 similar with thermal histories strongly affected by subsidence and sediment deposition on down
11
12 467 flexed lithosphere, pre-conditioned by its thermo-mechanical age.
13

14 468
15
16 469 **THE ROLE OF LITHOSPHERIC RHEOLOGY**
17

18 470 Folding is commonly associated with periodic deformation of layered structures with contrasting
19
20 471 mechanical properties. As a consequence, the harmonic spatial periodicity is often believed to be its
21
22 472 primary, almost synonymous, identifying factor. However, from a mechanical point of view, constant
23
24 473 wavelength is not a necessary requirement. Folding is a compressional instability developing in stiff
25
26 474 layers embedded in a weaker material. The harmonic solution of the equilibrium and conservation
27
28 475 equations is a possible solution (Muehlhaus *et al.*, 1998; Burov & Molnar, 1998; Cloetingh *et al.*,
29
30 476 1999). The physical mechanism of folding is well understood. In a continuous layered medium the
31
32 477 stresses and strains must be continued across the interfaces between the layers. In multi-layer
33
34 478 systems with contrasting mechanical properties (e.g. strong and weak layers) this requirement may
35
36 479 be difficult to satisfy, because the same amount of shortening in a stiffer layer would require much
37
38 480 larger stress than in the neighbouring weaker layer. The system becomes unstable and, in
39
40 481 attempting to reduce the stress and strain unconformities at the interfaces between the layers, starts
41
42 482 to fold (or buckle) in response to very small perturbations. In non-elastic media with strain-
43
44 483 dependent properties, locally increased flexural strain at the fold limbs can create weakened zones
45
46 484 that significantly facilitate further deformation. These weak or softened plastic or viscous zones are
47
48 485 often referred to as inelastic hinges, since the system easily folds at such weakened zones.
49
50 486 Homogeneous shortening of the lithosphere under horizontal compression requires more work than
51
52 487 required by an equivalent amount of shortening by folding. However, shortening by
53
54 488 underthrusting/subduction of the lithosphere may be more mechanically efficient than folding but
55
56
57
58
59
60

may be blocked or unable to start immediately after the onset of shortening. For this reason, folding is likely to be a primary and common response to tectonic compression. Folding probably may continue, in an attenuated form, even after the beginning of subduction (Cloetingh *et al.*, 1999), or re-appear when the subduction channel is locked or during the aftermath of collision (Cloetingh *et al.*, 2004; Matenco *et al.*, 2007).

The rheological structure of the lithosphere (Watts & Burov, 2003) is a key factor in lithospheric folding. Compressional stresses must build up to a level comparable with the integrated strength of the lithosphere in order to induce folding. Prior to the advent of rheological models based on extrapolation of laboratory experiments (Goetze & Evans, 1979; Carter & Tsenn, 1987) lithospheric stresses were thought to be incapable of reaching the failure levels required for folding (Turcotte & Schubert, 2002).

The simplest expression of the folding process is in oceanic lithosphere, characterized by an absence of a rheological stratification, and with thermo-mechanical ages spanning a limited age window with a maximum of 200 Myr (Geller *et al.*, 1993; Zitellini *et al.*, 2010). Because weak lithosphere is easier to fold than strong lithosphere, one would expect the first observations of lithospheric folding from the rheologically weak continental plate interiors. However, the observation of folding of the relatively strong lithosphere of the central Indian Ocean with a thermo-mechanical age of 80 Ma and with wavelengths of 200-250 km (Geller *et al.*, 1983; McAdoo & Sandwell, 1985; Stein *et al.*, 1989; Bull & Scrutton, 1992) triggered the interest in this mode of deformation. There may be two reasons for this. First, the recognition of folding, by mapping of basin reflectors and gravity anomalies requires spatial scales of several hundreds of kilometres. This is feasible with marine geophysical surveys, but generally in excess of the spatial scales covered by land surveys. An exception has been the systematic mapping of basins in Central Australia, located on strong lithosphere with a thermo-mechanical age of approximately 700 Ma (Lambeck, 1983; Stephenson & Lambeck, 1985). The Indo-Australian plate is under an exceptionally high level of compression (Stein *et al.*, 1989), as a result of its collision with the Eurasian plate. Under these conditions high

1
2
3 515 stresses might have been more important than the rheology of the plate interior (Beekman et al.,
4
5 516 1996; Gerbault *et al.*, 1998; Gerbault, 2000).
6
7 517 Secondly, folding in oceanic lithosphere is not affected by surface erosion, making it relatively easy
8
9 518 to recognize. The presence of only one dominant folding wavelength characteristic of non-stratified
10
11 519 oceanic lithosphere also facilitated a quantitative interpretation. Following the early studies in the
12
13 520 Indian Ocean and in Central Australia in the rear of the Indian-Eurasian collision, attention shifted to
14
15 521 the other side of the plate contact in Central Asia. Large scale folding patterns were recognized
16
17 522 through analysis of geophysical data, including gravity and topography (Burov *et al.*, 1993; Nikishin
18
19 523 *et al.*, 1993; Burov & Molnar, 1998) as well as detailed geological studies (Cobbald *et al.*, 1993).
20
21 524 On the basis of an inventory of currently available results of studies of folded continental
22
23 525 lithosphere, it appears that in general the observed wavelengths of folding follow theoretical
24
25 526 predictions for wavelengths as a function of thermo-mechanical age remarkably well (see **Fig. 3**).
26
27
28
29
30
31
32

33 529 **NUMERICAL EXPERIMENTS FOR SELECTED THERMO-MECHANICAL AGES OF**
34
35 **LITHOSPHERE**
36
37
38
39

40 532 Folding is an unstable deformation that may develop in stratified media with significant competence
41
42 533 contrasts, as occur in the continental lithosphere, in response to horizontal loading (Biot, 1961;
43
44 534 Ramberg, 1961). These instabilities may develop under compression or basal shear as a result of
45
46 535 strain/stress incompatibility on the interfaces between layers of different competence. In the initial
47
48 536 stages of deformation, compressional instabilities cause periodic folds with an exponential growth
49
50 537 rate and a dominant wavelength λ roughly proportional to 5 -10 times the thickness(es) of the
51
52 538 competent layer(s). In the lithosphere, the layer thickness is that of the competent core h . So, $h =$
53
54 539 10-100 km, gives rise to $\lambda = 50-1000$ km.
55
56
57
58
59
60

"Biot's", or linear, folding encompasses the cases where compression of the lithosphere takes place from the beginning, leading to the formation of alternating (synclinal) basins and (anticlinal) highs, and where the conditions of the linear theory are more or less satisfied (e.g. thin layer approximation, no strain-softening, plane layers). For linear folding in a Newtonian medium, an asymptotic relationship derived from the thin layer equilibrium equation is (Biot, 1961; Ramberg, 1961):

$$\lambda_1 = 2\pi h (\mu_{11}/6\mu_{12})^{1/3} \quad (1)$$

Here λ_1 is the "Laplacian" dominant wavelength of folding in the absence of gravity, h is the thickness of the competent layer (crustal or mantle), and μ_{11} and μ_{12} are the effective viscosities (or competencies) of the strong layer and weak surrounding material, respectively. The above equation, derived assuming zero gravity, provides estimates of $\lambda_1/h = 20-40$ for typical competence contrasts. This does not hold for most lithospheric-scale cases, where $\lambda/h = 4 - 6$ is more common due to the influence of the gravity-dependent terms. A reduction of the normalized characteristic wavelength to values around 4 is also predicted for power-law rheology or plastic layers (Smith, 1979). For a single stiff viscous layer on top of an inviscid medium, the dominant gravity-dependent wavelength is (Burov & Molnar, 1998):

$$\lambda_g \sim 2\pi(2h \dot{\epsilon} \mu_{eff} / (\Delta\rho g))^{1/2} \quad (2)$$

where $\Delta\rho$ is the density contrast, $\dot{\epsilon}$ is the strain rate, μ_{eff} is the layers' effective ductile viscosity (see Appendix A), n is the power-law exponent, and h is the thickness of the competent layer.

At large amounts of shortening this deformation may become a-periodic (Hunt *et al.*, 1996) so that the wavelength and amplitude of folding vary along the deformed transect. Plume-induced periodic deformation (Burov & Cloetingh, 2009) also includes folding or boudinage caused by shear on the base of the lithosphere. However, the prime cause for folding of continental lithosphere is tectonic loading by horizontal far-field forces. For characteristic parameters of various layers within the

continental lithosphere ($h = 10 - 120$ km, $\mu = 10^{19} - 10^{26}$ Pa s; $\Delta\rho = 10 - 600$ kg m⁻³), equations (1-2) predict folding wavelengths that may vary from 30 km (upper crustal folding) to as high as 600-800 km. The findings of numerical models (Cloetingh *et al.*, 1999) are in excellent agreement with analogue tectonic experiments (Sokoutis *et al.*, 2005). Numerical experiments show that the wavelength and amplitude of folding is largely controlled by the thermo-mechanical age and less by the rate of shortening (Burov & Cloetingh, 2009). Lithospheres older than 500 Ma develop nearly the same wavelength of folding as 1000 Ma old lithosphere because of the similar nearly stationary thermal state. Here, therefore, we restrict ourselves to a set of experiments for ages of 150 and 300 Ma and convergence rates of 1.5 and 3 cm/yr.

Surface topography and sedimentary deposition is described through diffusion-like erosion laws and fluvial transport laws (Gossman, 1976; Kirkby, 1986; Leeder, 1991; Beaumont *et al.*, 1992; Kooi and Beaumont, 1994):

$$dh/dt = \text{div}(k^*(x, h, \nabla h) \nabla h) \quad (\text{slope erosion})$$

where h is topography, t is time, x is the horizontal coordinate, k^* is an experimentally established scale-dependent coefficient of erosion (500 – 1000 m²/y). For simplicity, we used zero order linear diffusion (k^* constant) for the short-range erosion and flat deposition for long-range fluvial transport (Avouac & Burov, 1996). The hill-slope erosion law ensures conservation of matter but holds only for small-scales. At large scales it is no longer applicable, and conservation of matter is not observed, because some sediments are lost due to transport out of the system by long-range processes.

Figure 9 shows the results of the experiments for relatively young (150 Ma) lithosphere compressed at rates of 1.5 and 3 cm/yr. Strongly decoupled low-amplitude short wavelength (mantle wavelength $\lambda = 250$ km) folding developed leading to crustal thickening above a synclinal mantle fold. At later stages, harmonic folding was followed by mega-folding (Cloetingh *et al.*, 1999; Burg & Podladchikov, 1999). For higher rates of shortening these mantle synclines developed in a mode

resembling symmetric subduction. For lower convergence rates (**Fig. 9a**) folding does not develop due to the low Peclet number of the system, leading to heat diffusion and weakening of folds.

Folding is well developed for medium (300 Ma) age lithosphere (**Fig. 10**). This case is characterized by long mantle wavelengths ($\lambda=360$ km) and high surface amplitudes (2000 m after 10 Myr). These compare well with observations from Iberia (Fernandez-Lozano *et al.*, 2010), Western Goby and the Ferghana basin in Central Asia (Burov *et al.*, 1993; Burov & Molnar, 1998). At late stages (10 - 26 Myr since the onset of shortening for 3 cm/y or 20- 50% of shortening), folding becomes aperiodic, leading to mega-folding (Cloetingh *et al.*, 1999; Burg & Podladchikov, 1999) and subsequent formation of high-amplitude crustal down-warps. The amplitude of vertical movements may reach 20 km (± 10 km) or even more. As pointed out earlier, such high amplitudes of vertical motions are observed in the South Caspian Basin (Guest *et al.*, 2007) and the Barents Sea (Ritzmann & Faleide, 2009). However, it may be relatively rare for folding to continue for periods in excess of 10 Myr. More typically is that at a certain stage deformation localizes along single major fault zones (Cloetingh *et al.*, 1999; Gerbault *et al.*, 1998).

As was discussed by Bird (1991), Avouac & Burov (1996) and Cloetingh *et al.* (1999), large-scale undulations of the lithosphere cannot be preserved for a long time (longer than 10 Myr) in the absence of sufficient compression, except for plates with very strong (especially lower crustal) rheology. Otherwise, the folds either will be flattened by gravity-driven crustal flow associated with the large crust-mantle density contrast at the Moho, or deformation will localize along some of the faults created at the inflection points of folds.

The fact that folding with wavelengths of 400-500 km and 500-600 km persists long after cessation of the compression in the presumably high-rigidity Australian craton and Russian platform points to a strong rheology yielding a high value for the effective elastic thickness EET (in excess of 60 km), as expected from its thermo-mechanical age (Beekman *et al.*, 1997). The long wavelengths of 400-600 km of crust-mantle down-warping in the eastern Barents Sea, where Mid-Cenozoic

1
2
3 611 compression has affected a basin system formed during Late Permian-Triassic time (Ritzmann &
4
5 612 Faleide, 2009), with a total infill of sediments of the order of 15-20 km is another impressive
6
7 613 example of compressional deformation on relatively cold lithosphere.
8
9
10 614 Most of the areas with present-day long-wavelength continental lithosphere folding formed during or
11
12 615 short after the Alpine collision 60 Myr ago. It thus seems that 60 Ma is a characteristic timescale of
13
14 616 gravity collapse of the large-scale folds in such intermediate-age lithospheres. In the case of very
15
16 617 weak quartz-dominated lower crust, folds may disappear rapidly within 8-15 Myr. Experiments
17
18 618 (Cloetingh *et al.*, 1999) for 1000 Ma old lithosphere, with a diabase lower crust, demonstrate that
19
20 619 folding-induced deformation persists for 10-50 Myr following the cessation of compression. During
21
22 620 this period the amplitude of the folds decreased by less than 10 percent. At this stage, crustal faults
23
24 621 may accelerate the gravity collapse of folds, creating inverted basins.
25
26
27 622
28
29 623

30
31 624 **BASIN GEOMETRY AND ACCOMMODATION SPACE**
32
33 625

34
35
36 626 **Basin shape**

37
38 627 Folding of the lithosphere leads to a symmetrical pattern of down-warped areas (synclines) flanked
39
40 628 by highs (anticlines) of similar amplitude and wavelength, like for instance the Ferghana basin in
41
42 629 Central Asia (Burov & Molnar, 1998).
43
44 630 This symmetry is in marked contrast to the asymmetrical shape of foreland basins, formed by
45
46 631 flexure in front of an orogenic wedge, which are flanked by a flexural bulge of an amplitude that is
47
48 632 only up to 10 percent of the maximum depth of the foreland depression (Royden, 1988; Zoetemeijer
49
50 633 *et al.*, 1999). Both types of compressional basins show linear nature alignment of parallel
51
52 634 depocentres and flanking highs. An important difference is the presence of parallel trending
53
54 635 depocentres for lithospheric folding, whereas foreland flexural basins have a single depositional
55
56 636 system. In addition, basins developed on folded lithosphere have a static location of the axis of their
57
58
59
60

depocentres, in contrast to foreland basins, where the axis of depocentres can migrate with time (e.g. Zoetemeijer *et al.*, 1993).

In both cases, the integrated strength of the lithosphere (Watts and Burov, 2003; Tesauro *et al.*, 2007) defines the characteristic width of the basins. For folding, basin width roughly corresponds to one half of the fold wavelength (λ), which equals 5 to 10 times the thickness of the strong core of the lithospheric plate, h ($h = 10$ to 100 km, i.e. $\lambda = 50 - 1000$ km). In the case of crust-mantle decoupling, which occurs in relatively young plates, two dominant folding wavelengths develop (typically $250 - 400$ km and $50-100$ km). Thus two basin populations may be observed, one imbricated within the other. In the case of flexural foreland deformation, the width of the basin is controlled by the flexural parameter and is thus of the same order as the largest fold wavelength.

Subsidence patterns

Lithospheric folding results from an instability in the lithosphere due to stress/strain incompatibilities that develop in rheologically stratified layers under compressive strain. This process typically operates with timescales of the order of 1-10 Myr (Cloetingh *et al.*, 1999). The preservation of lithospheric folds appears to depend strongly on the thermo-mechanical age of the underlying lithosphere (Cloetingh *et al.*, 1999). Folds in young lithosphere will not be preserved upon relaxation of the stress field, whereas ones in cratonic lithosphere will be well preserved (Cloetingh *et al.*, 1999).

Figure 11a displays characteristic basin subsidence patterns predicted by folding of 150 and 300 Ma old lithosphere due to a 3 cm/year shortening rate. As illustrated (**Figure 11b**), the differential motions occur in three distinct phases:

Stage 1 is the basin formation phase, coinciding with the initiation and development of folding, which is marked by an acceleration of subsidence in the basin and uplift in the flanking highs. As a result of the time lag between sediment supply and the creation of accommodation space, during the few Myr needed to form the basin, deposition cannot keep up with subsidence causing

1
2
3 663 sediment-starved basins. A similar pattern occurs in pull-apart basins (Pitman & Golovchenko,
4
5 664 1983) which are also associated with ultra rapid subsidence in their formation stage also lasting only
6
7 665 a few Myr.
8
9
10 666 **Stage 2** is the basin preservation stage, where equilibrium develops between sediment supply and
11
12 667 sediment deposition. As a result the basin will be rapidly filled to overfilled.
13
14 668 **Stage 3** is the basin destruction phase, characterized by basin capture and removal of sediments to
15
16 669 areas outside the folding system. During this phase, the size of the accommodation space is
17
18 670 reduced and erosion occurs on both flanks and depocentre. Thus, the net effect of these three
19
20 671 stages is that lithospheric folding will lead to the development of distinct depositional-erosional
21
22 672 cycles.
23
24 673 Characteristic patterns for vertical motions for folding of continental lithosphere of 150 and 300 Ma
25
26 674 old (**Fig. 11**) demonstrate a remarkably short time scale in which substantial amounts of tectonic
27
28 675 subsidence are induced by the process of lithospheric folding. As shown by **Figure 11**, within 1 Myr
29
30 676 after the initiation of folding induced by shortening the lithosphere at a rate of 3 cm/yr, subsidence
31
32 677 rates are in the order of 5-15 km/Myr, depending on the thermo-mechanical age of the lithosphere.
33
34 678 Thus, folding appears to be more effective in middle aged lithosphere of 300 Ma than in lithosphere
35
36 679 of younger ages (Cloetingh *et al.*, 1999; Burov & Cloetingh, 2009). After 2 Myr, a slow uplift phase
37
38 680 starts of relatively minor magnitude of the order of a few hundreds of meters to a km. The predicted
39
40 681 subsidence in **Figure 11** is for the centre of the syncline. As pointed out above, erosion of the
41
42 682 uplifted flanks is taken into account, adopting a diffusive equation approach to erosion (see Burov &
43
44 683 Cloetingh, 1997; Cloetingh *et al.*, 1999, for further details).
45
46
47
48 684 The alignment of parallel highs and basins of similar dimensions has an important consequence for
49
50 685 the areal extent of the source areas for sediments available for deposition in the folded depressions.
51
52 686 In comparison with foreland basins the distribution of sources is more symmetrical, superseding the
53
54 687 volume of sediments that can be eroded from, for example, the flexural foreland bulge. As noted
55
56 688 earlier, erosion reduces the contribution of gravity-dependent terms and accelerates local
57
58
59
60

deformation. Erosion, therefore, has an important feed-back with the geometry of the accommodation space in changing the spectrum of wavelengths (Cloetingh *et al.*, 1999). Erosion acts as a filter, suppressing the shorter wavelengths in folded basin topography. Strong erosion, insufficiently compensated by tectonic deformation, can even wipe out most of the topography. However, if the erosion is tuned to the average elevation rates (**Fig. 12**), it may dramatically accelerate folding (Cloetingh *et al.*, 1999).

There has been a rapid development and application of stratigraphic modelling packages to sequence stratigraphy over the last two decades (e.g. Cross, 1989; Alzaga-Ruiz *et al.*, 2008). Among the different types of modelling programs, the forward stratigraphic modelling approach has been applied extensively by basin modellers from academia and petroleum industry (e.g. Lawrence *et al.*, 1990; Granjeon, 2009). Forward stratigraphic modelling commonly starts with an estimate of the initial conditions related to an adopted basin formation mechanism and controlling parameters, followed by a forward prediction in time. Most of these software packages are based on sequence stratigraphic concepts (Wilgus *et al.*, 1988). Assumptions for initial subsidence are either ad-hoc or related to stretching (e.g. Lawrence *et al.*, 1990) or foreland flexure (Garcia-Castellanos *et al.*, 1997), but to date do not include lithospheric folding as a basin formation mechanism.

Marine versus continental deposits and closed basins versus capture of drainage

Erosion of uplifted areas and sedimentation in the depressions created by folded lithosphere is a self reinforcing process promoting continuing uplift of the highs and subsidence in the depocentres. Thus, such basins will remain closed during much of their evolution. Folded basins will hence be predominantly characterized by continental deposits with only minor deposition of marine sediments during marine incursions or during basin capture. This seems to be the case for folded basins in Iberia (e.g. Duero Basin) and Central Asia (Ferghana, Tarim Basin, Lake Issyk-Kul) (Cobbold *et al.*, 1993). As has been pointed out for flexural foreland basins (Garcia-Castellanos *et al.*, 2003), the capture time for opening basins decreases with increasing flexural rigidity. The flexural upwarp to

1
2
3 715 restoring the topography removed by erosion will be almost instantaneous for very weak
4
5 716 lithosphere, but requires up to 50-100 Myr for high lithospheric rigidities (Garcia-Castellanos *et al.*,
6
7 717 2003). Thus, basins created by folding of cratonic lithosphere probably have a shorter capture time
8
9 718 than ones formed in younger lithosphere.
10
11
12 719

13
14 720 **THERMAL REGIME, STRESS REGIME AND STYLE OF FAULTING**
15

16 721
17
18 722 **Thermal regime**
19

20 723 Lithospheric folding is controlled by the interplay of lithospheric stresses and inherited strength of
21
22 724 the lithosphere. The thermal regime controls the rheological profile and differs between folded
23
24 725 basins developed in young lithosphere and basins in cratonic lithosphere. The latter are associated
25
26 726 with much lower thermal gradients than basins developed on young continental lithosphere. As for
27
28 727 foreland basins, the initiation of folding is not associated with a thermal instability, unless folding
29
30 728 occurs in conjunction with plume activity (Burov & Cloetingh, 2009). Following the deposition of
31
32 729 radiogenic sediments in the folded depression, sediment blanketing will affect the heat flow
33
34 730 (Stephenson *et al.*, 1990; Lavier & Steckler, 1997; Van Wees *et al.*, 2009), modifying the surface
35
36 731 heat flow in the basin centre. As pointed out by Lavier & Steckler (1997) and Ziegler *et al.* (1998),
37
38 732 the effect of sediment fill is to weaken the lithosphere. The low thermal conductivities of the
39
40 733 sediments lead to high temperatures in the upper lithosphere and consequently low local yield
41
42 734 strength. During basin subsidence sediments deposited in the central parts of the basin might be
43
44 735 exposed to temperature windows corresponding to hydrocarbon generation. At the same time,
45
46 736 sediments pre-dating the folding may have undergone extra burial due to syn-folding sedimentation.
47
48 737 An example is the Paris basin, where source rocks of Toarcian age were folded in Cenozoic time
49
50 738 (Guillocheau *et al.*, 2000; Le Solleuz *et al.*, 2004).
51
52
53
54
55 739 During basin capture overall cooling takes place. This effect is illustrated by characteristic thermal
56
57 740 evolution for different thermo-tectonic ages of 150 and 300 Ma of the folded continental lithosphere,
58
59
60

incorporating radioactive heat production in the sediments filling in the synclinal depression, shown in **Figure 13**. Subsidence induced by folding is calculated for the centre of the basin. Basins developing on folded lithosphere are characterized by their relatively low heat flow at the onset of folding, followed by a steady increase with time, doubling the heat flow over a time interval of the order of 5 Myr following the cessation of shortening. This increase is primarily due to the contribution of radioactive heat production in the very substantial pile of sediments accumulating in a relatively short time interval. The patterns of heat flow are similar for different thermo-mechanical ages of the lithosphere, with the oldest lithosphere having the lower heat flow. Older ages enhance the accommodation space and the contribution of the sediments to heat flow but are compensated by a larger decrease in heat flow with aging lithosphere. Another important factor is the mode of folding. Crustal scale folding, which is characteristic for Lake Issyk-Kul, will be associated with shorter wavelengths, shallower basin depths and thinner sequences of heat producing sediments than for mantle lithospheric folding. This effect might explain the relatively low heat flow observed in Lake Issyk-Kul (Vermeesch *et al.*, 2004).

Modes of brittle deformation and faulting

The role of pre-existing faults on the development of folding has been investigated through numerical experiments on folding of brittle-elasto-viscous lithosphere (Beekman *et al.*, 1996; Gerbault *et al.*, 1998; Cloetingh *et al.*, 1999) and through analogue experiments on plastic-elastic lithosphere (e.g. Martinod & Davy, 1994; Sokoutis *et al.*, 2005). These studies show that lithospheric folding and faulting can develop simultaneously and that pre-existing crustal scale faults do not prevent, but instead promote the development of folding instabilities. The instabilities, in turn, can initiate the formation of new faults at the inflection points of the folds. Although the usual expectation is that faulting prevents folding, this view neglects the role of the gravity and friction on faults in the behaviour of the large-scale fault-and-fold systems. As also observed in analogue experiments (Martinod & Davy, 1994), this continuous behaviour of faulted lithosphere can be explained by fault

1
2
3 767 locking due to gravity and friction. After some sliding (uplift) on a fault, the potential gravitational
4
5 768 energy working against friction generated by horizontal shortening forces becomes too high. As a
6
7 769 result, the fault locks and the rock mass transmits horizontal stresses as a continuous medium. As
8
9
10 770 the compression continues, one of the folds finally starts to grow faster than others, resulting in a
11
12 771 loss of the periodicity and the formation of a mega-fold (see also the discussion in the context of the
13
14 772 Ferghana and Tadjik basins of Central Asia).
15
16 773 Following the initiation of folding, and the further build up of compressional stresses, folding will,
17
18 774 therefore, usually be accompanied by brittle deformation and faulting in the crust and upper
19
20 775 lithosphere. The surface expression of folding in uniform lithosphere is frequently in the form of
21
22 776 fault-controlled pop-up and pop-down structures and inverted basins (e.g. Cobbold *et al.*, 1993),
23
24 777 accompanied by development of imbrications of bivergent thrust wedges at the areas where
25
26 778 shortening is applied (Sokoutis *et al.*, 2005). These findings (see **Fig. 14**) are corroborated by
27
28 779 observations in the Central System of Iberia, where folding of Variscan lithosphere with a thermo-
29
30 780 mechanical age of 280 Ma occurs (Cloetingh *et al.*, 2002). Here the brittle deformation associated
31
32 781 with crustal scale folding expressed by a folded Moho is mechanically decoupled from folding of the
33
34 782 mantle lithosphere (Fernandez-Lozano *et al.*, 2010; see **Fig. 5b**).
35
36 783 Crustal and mantle faults may develop as a result of folding (Burov & Molnar, 1998) (**Fig. 15**).
37
38 784 Folding can continue after these faults develop and folding and faulting can co-exist for times of
39
40 785 several Myr (**Fig. 5b**). For crust-mantle decoupling, a complex distribution of crustal faults and
41
42 786 surface undulations can be expected, with crustal faults not necessarily connected to mantle faults.
43
44 787 Numerical experiments suggest that mantle faults are unlikely to be longer than a few km in depth
45
46 788 (Burov & Molnar, 1998). Such faults may also not develop at longer wavelengths of folding
47
48 789 connected with stronger lithosphere. However, deep seated faults in mantle lithosphere can develop
49
50 790 when cratonic lithosphere is folded (Cloetingh *et al.*, 1999). The mode of faulting, depth extent and
51
52 791 spacing of faults depend on the thermo-mechanical age of the lithosphere. Once the development
53
54 792 of folding begins in a basin, the intensity and depth penetration of faulting will accelerate. The mode
55
56
57
58
59
60

of decoupling between crustal and mantle folding is crucial. Discontinuities within the folded lithosphere will localize the development of deep synclines, flanked by more modest anticlines (Sokoutis *et al.*, 2005) and affect the distribution of faulting (**Fig. 14**).

During the next phase of basin evolution, when a steady state is reached between the creation of accommodation space and sediment supply, only minor faulting will occur. This is in marked contrast with the later phase of basin capture, which is in general characterized by erosion and narrowing of the basin. Due to erosional unloading, stress relaxation will occur, manifested in upward propagating faults.

Interaction of folding with other tectonic processes

Folding, in addition to being a basin forming mechanism, frequently interacts with other tectonic processes. As pointed out by Cloetingh (1988), intraplate compression can modify pre-existing basins. As discussed above, the Pannonian Basin of Central Europe created by Miocene back-arc extension appears to be an example of such a configuration, characterized by a-typical a-periodic folding. Other examples are cratonic sag basins where intraplate compression (Cloetingh, 1988) is thought to occur in interaction with phase changes in the lithosphere (Artyushkov, 2007). Examples of these might be the Barents Sea (Ritzmann & Faleide, 2009) and possibly the South Caspian Basin, although in the latter case pre-orogenic extension cannot be ruled out (Guest *et al.*, 2007). Basins exposed to changes in tectonic regime will have a polyphase record (Cloetingh & Ziegler, 2007), characterized by a superposition of more than two regimes, such as pre-orogenic extension, foreland flexure and late-stage folding. This sequence appears to be characteristic for some very deep foreland basins, such as the Focsani depression of the Romanian Carpathians, with more than 16 km of sediments, which was affected by extension due to the opening of the Black Sea Basin, followed by foreland flexure and overprinted by Late Miocene compression (Tarapoanca *et al.*, 2003). As pointed out above (Ziegler & Dèzes, 2007), the Northwestern European foreland indicates folding, overprinting, rifting and foreland flexure. Quantification of the topographies created

1
2
3 819 by rifting and foreland flexure and correcting for them is essential to reconstruct the shape of the
4
5 820 additional accommodation shape created by subsequent lithospheric folding (Bourgeois *et al.*,
6
7 821 2007).
8
9
10 822 The impingement of plumes on the base of continental lithosphere can induce differential
11
12 823 topography similar to the surface deflections characteristic for lithosphere folds (Guillou-Frottier *et*
13
14 824 *al.*, 2007; Burov *et al.*, 2007). Plumes and lithospheric folds frequently interact in space and time in
15
16 825 the geological record (Ziegler & Dèzes, 2007; Burov & Cloetingh, 2009). Therefore, the evolution of
17
18 826 folded lithosphere basins can be overprinted by the signatures of a plume or vice-versa. This will
19
20 827 lead to an amplification of the induced vertical motions, particularly significant for young and
21
22 828 intermediate age lithosphere. Emplacement of hot upper mantle material will raise temperatures in
23
24 829 the lithosphere and increase heat flow. Due to the lag time in the propagation of heat, the effect at
25
26 830 the surface might become manifest only after several tens of Myr. Plume emplacement might also
27
28 831 weaken the lithosphere, making it more prone to compressional fault reactivation after cessation of
29
30 832 the folding.
31
32
33 833 Interplay of lithosphere folding and plume impingement on the continental lithosphere occurred
34
35 834 probably almost simultaneously in late Neogene times in the Northwestern European foreland. Two
36
37 835 examples are the Eifel and Massif Central areas of the Alpine foreland of NW Europe (Cloetingh &
38
39 836 van Wees, 2005; Ziegler & Dèzes, 2007). Both areas are sites of main Late Neogene volcanic
40
41 837 activity in the European Cenozoic rift system (ECRIS). Seismic tomography (Ritter *et al.*, 2001)
42
43 838 shows finger shaped baby plumes with a characteristic spatial diameter of 100 km, extending
44
45 839 downward to 400 km. Ziegler & Dèzes (2007) propose that plume activity occurs simultaneously
46
47 840 with recent compressional deformation in the Massif Central area (Guillou-Frottier *et al.*, 2007) and
48
49 841 the Ardennes/Eifel area. Geomorphological studies constrain the recent uplift of the Ardennes and
50
51 842 the Eifel area. An order in magnitude difference occurs between the uplift of the Eifel area, underlain
52
53 843 by a plume and the adjacent Ardennes area where evidence for a plume is lacking. The patterns of
54
55 844 uplift appear to be radial, superimposed by a linear NE-SW trend perpendicular to the main axis of
56
57
58
59
60

compression. Baby plumes primarily develop in the anticlines of lithospheric folds (Burov & Cloetingh, 2009). The plume activity presumably accelerated the rate of uplift by a factor of 3-5 (Cloetingh & Ziegler, 2007). According to presently available data, the plumes in this segment of the European Cenozoic Rift System arrived about 1 Myr ago. This activity was preceded and followed by lithospheric folding that continues to present-day since 17 Myr ago (Bourgeois *et al.*, 2007). Thermo-mechanical modelling illustrates the relative effectiveness of amplification of lithosphere deformation and topographic effects induced by plumes through folding and vice-versa. Burov & Cloetingh (2009) examined the response times and time-lags involved and whether these baby plumes were more efficient in localizing deformation than large plumes. Plume-affected folding appears to accelerate surface uplift, whereas folding goes into saturation and stagnates when plastic hinges form. For the discrimination of plumes and folding it is critical to access constraints on the presence or absence of radial versus linear symmetry, heat flow anomalies, gravity and geoid data. Plume activity facilitates folding, by dramatically lowering the stress levels required (Burov & Cloetingh, 2009). Plume impact also reduces the fold wavelength and localizes folding above the plume impact area. A general outcome of the modelling and observations is that lithospheric folding as a mechanism for producing thermal perturbations in the lithosphere/upper mantle system is a less feasible scenario.

FOLDING AND OTHER MODES OF BASIN FORMATION: DIFFERENCES AND SIMILARITIES IN STRUCTURE AND EVOLUTION

Sedimentary basins formed through lithospheric folding have characteristic features including vertical motions, basin architecture, thermal regime and fault activity, distinguishing them from other basin types (Cloetingh & Ziegler, 2007; Xie & Heller, 2009). **Figure 1b** and **Table 2** illustrate differences and similarities between basins developed on folded lithosphere (FLB), foreland basins (FB), intracratonic basins (ICB), extensional basins (EB), and pull-apart basins (PAB).

1
2
3 870 Sedimentary basin systems are, by their nature, prone to tectonic reactivation and, therefore,
4
5 871 frequently characterised by a poly-phase evolution (Cloetingh & Ziegler, 2007). Extensional basins
6
7 872 are, for example, often formed in areas previously thickened by tectonic compression and
8
9
10 873 subsequently often subject to late-stage compression (Lundin & Dorè, 1997). Examples include the
11
12 874 Pannonian basin of Hungary (Horváth & Cloetingh, 1996) and the North Atlantic rifted margins
13
14 875 (Cloetingh *et al.*, 2008). Similarly, foreland basins are frequently characterised by pre-orogenic
15
16 876 extension. Examples include the Carpathian foredeep (e.g. Tarapoinca *et al.*, 2003) and the
17
18 877 Aquitaine basin of Southern France, which was the retro-arc foreland basin of the Pyrenees
19
20 878 (Desegaulx *et al.*, 1991). The actual subsidence patterns of these poly-phase systems are often
21
22 879 more complex than predicted by the end-member models discussed below that only consider the
23
24 880 basin formation mechanism. Retro-arc foreland basins are characterised by a prolonged subsidence
25
26 881 history with a distinct thermal signature including the effects of thermal cooling inherited from pre-
27
28 882 orogenic extension. This is sometimes neglected in comparing pro-arc and retro-arc foreland basins
29
30 883 (Naylor & Sinclair, 2007).
31
32
33 884 Basin infill records allow main features, including geometry, vertical motions and faulting
34
35 885 characteristics to be unravelled (Zoetemeijer *et al.*, 1993). However, heat flow histories need to be
36
37 886 determined through kinematic reconstruction and forward modelling. Results of heat flow history
38
39 887 modelling for FB and EB settings (Van Wees *et al.*, 2009) can be subsequently compared to FLB
40
41 888 results presented in this paper (**Fig. 13**). For ICB and PAB we have refrained from a quantitative
42
43 889 approach for heat flow predictions as vertical motions driving subsidence cannot be linked in a
44
45 890 straightforward way to a single or uniformly distributed lithosphere process. As a consequence, for
46
47 891 these settings tentative heat flow patterns have been drawn.
48
49
50 892 *Folded lithosphere basins* (FLB) that develop as a result of periodic folding are highly symmetrical,
51
52 893 with dimensions that can vary from 50 km to 600 km, depending on lithospheric age and shortening
53
54 894 rate. These basins can accumulate thick sedimentary sequences up to the order of 20 km. The time
55
56 895 scales associated with this process of basin formation are very short, typically a few Myr. The
57
58
59
60

subsidence is so fast that erosion and sediment supply at the basin formation stage typically lags behind, leading to the development of starved basins, followed by shallowing upward sequences after stress relaxation. Subsidence patterns are characteristically convex upward with time. A noticeable feature is the development of significant topography of the order of several km flanking the synclinal depression. No initial heating is predicted, with an increase in heat flow with time following the basin formation. Significant brittle deformation is expected to occur in the folded basement, accelerating during the folding phase.

Foreland basins (FB) are strongly asymmetrical, with depths usually less than 10 km, and have modest flexural bulges of a few hundred meters (Royden, 1993; DeCelles & Gilles, 1996; Zoetemeijer *et al.*, 1999). Important differences occur in the nature of the subsidence of pro- and retro-foreland basins (Naylor & Sinclair, 2007). The first are typically associated with short life spans and convex up subsidence patterns with time, whereas the latter show more prolonged subsidence histories and concave up subsidence patterns. Subsidence patterns will be interrupted during thrusting phases, also leading to uplift of the flexural marginal bulge (Quinlan & Beaumont, 1984; Zoetemeijer *et al.*, 1993). The predicted heat flow pattern of a pro-foreland setting shown for the Ebro foreland basin (well Jabali from Verges, 1999), marked by rapid basement subsidence and sedimentary infill of ca 4 km in ca 10 My period (other parameters as in Van Wees *et al.*, 2009). Flexure and rapid sediment infill is marked by a reduction of heat flow, followed by a progressive increase related to radiogenic heat production of foreland sediments. Basement fracturing is concentrated at the points of maximum curvature in the flexed lithosphere, at the maximum points of upward and downward deflection, respectively, with intensity higher than inferred for FLB.

Intracratonic basins (ICB) share their symmetrical shape and great thickness of their sedimentary basin fill with FLB. An important difference is their evolution in time, characteristically associated with very long time spans (typically of the order of 500-800 Myr) with long phases of low subsidence rate punctuated by phases of subsidence acceleration. As pointed out above, this is why ICB are probably affected by short-term phases of enhanced intraplate compression superimposed on other

1
2
3 922 mechanisms operating on longer time scales. The large dimensions of these basins point to
4
5 923 formation in high rigidity lithosphere, that freezes in basin deformation over long time scales. The
6
7 924 long time scales involved also imply that the highs induced by compressional upwarping will be
8
9 925 commonly eroded. A major difference between ICB and FLB is that current hypotheses for ICB
10
11 926 formation are casted only in terms of the tectonic processes operation at their earlier stage such as
12
13 927 phase changes in the lithosphere. It appears, therefore, that lithospheric stress fields and the
14
15 928 lithosphere rheology (inferred from gravity and flexure) are far more diagnostic than the lithosphere-
16
17 929 asthenosphere boundary, which is only giving information on the thermal state of the lithosphere.
18
19 930 Heat flow patterns of ICB will be dominated by sediment infill. ICB are in general characterized by a
20
21 931 low intensity of faulting. However, in the absence of a rigorous understanding of the formation
22
23 932 mechanisms of ICB, predictions for heat flow are often not available. As a consequence, their heat
24
25 933 flow is most often described primarily in terms of the thermal consequences of their burial history.
26
27 934 *Extensional basins* (EB) have received major attention from basin modellers, both because of their
28
29 935 relative abundance and importance and because of the availability of a relatively simple quantitative
30
31 936 model for their formation (McKenzie, 1978). The latter can be relatively easily be incorporated in
32
33 937 forward modelling of their subsidence and heat flow histories (Van Wees *et al.*, 2009). The basin
34
35 938 shape is symmetrical or asymmetrical depending on a pure shear or simple shear mode of
36
37 939 extension. Rifts can be narrow or wide, depending on the mode of rifting and the rheology of the
38
39 940 extending lithosphere (Huisman and Beaumont, 2002). Dependent on the rigidity of the extending
40
41 941 lithosphere, rift shoulders with asymmetrical shape can be formed and be prone to erosion (Van
42
43 942 Balen *et al.*, 1995, Van der Beek *et al.*, 1995). The development of rift shoulders and their erosion
44
45 943 can lead to feed back with lower crustal flow and faulting in the extending lithosphere (Burov &
46
47 944 Cloetingh, 1997). During the post-rift phase, flexural widening of the basin commonly occurs. A
48
49 945 characteristic feature of EB predicted by simple stretching models (McKenzie, 1978) is the
50
51 946 succession of rapid initial subsidence in the syn-rift phase and exponentially decaying subsidence in
52
53 947 their post-rift phase with a typical thermally controlled time constants of the order of 70 Myr. Feed
54
55
56
57
58
59
60

backs between rift shoulder erosion and lower crustal flow (Burov & Cloetingh, 1997; Burov & Poliakov, 2001), have pronounced effects on subsidence in the final phases of rifting and in the early post-rift phase. Both can be influenced by changes in the mechanical properties of the lithosphere and its strengthening upon rifting. Multiple rifting phases is another common feature of EB. The duration of the syn-rift phase can vary from quasi-instantaneous (shorter than 20 Myr) to several hundreds of Myr (Cloetingh and Ziegler, 2007). In the latter case basin migration is a common feature (Corti *et al.*, 2003). Extensional faulting occurs during the rift phase, with absence of faulting in the post-rift phase. Heat flow reaches a peak during the syn-rift phase, followed by a rapid decrease in heat flow (Van Wees *et al.*, 2009). The syn-rift peak in heat flow is low or almost absent in continental extensional settings as a consequence of removal of heat producing elements in the crust and cooling by sediment infill. **Fig. 16** shows such a setting, which is based on the benchmark model presented in Van Wees *et al.* (2009). As a consequence of the loss of radiogenic heat production in the crust at the end of the post-rift stage values, heat flow evolves to values considerably lower than the initial ones. The EB heat flow pattern is, therefore, the reverse of that in FB where syntectonic heat flow is first lowered due to sediment infilling (Kombrink *et al.*, 2008), but eventually increases due to thermal relaxation and additional radiogenic heat production in the sediments.

Pull-apart basins (PAB) are characterized by a narrow fault bounded geometry (Smit *et al.*, 2008), developing over a very short time span. Models have been developed to calculate subsidence histories and heat flow histories for these basins (Pitman & Golovchenko, 1983), incorporating lateral heat transport in stretching models. These models predict rapid subsidence of several km's over time intervals of the order of 0.1 – 1 Myr, followed by a decay in the subsidence with a concave shape upward. Erosion and sediment flux can not keep pace with the subsidence at the basin formation phase, leading initially to a starved basin, followed by a shallowing upward sequence (Pitman & Golovchenko, 1983). These basins are intensively faulted, also due to their strike-slip origin, whereas heat flow decays on very short time scales of the order of a few Myr. Although this

1
2
3 974 scenario is a good first order description, many PAB show a more punctuated fault controlled
4
5 975 subsidence history, often terminating by late-stage uplift, as observed for Late Neogene pull-apart
6
7 976 basins in the internal zone of the Betics in SE Spain (Cloetingh *et al.*, 1992) and for the flanks of
8
9 977 pull-apart basins in the Dead Sea Fault System (Smit *et al.*, 2008, 2010).
10
11 978 From the above overview, it is apparent that FLB have in common with FB a compressional origin
12
13 979 and thermal histories controlled to a large extent by burial histories. At the same time, pronounced
14
15 980 differences occur in the duration of subsidence, varying from ultra short (FLB 1-10's Myr), medium
16
17 981 (FB, tens of Myr) and ultra long (ICB, hundreds of Myr). An important difference occurs in the spatial
18
19 982 scales of the topography generated by folding and foreland fold-and-thrust belt deformation. In the
20
21 983 latter case a much steeper differential topography is generated over typical horizontal differences of
22
23 984 the order of 10 km between peak and trough. In contrast, in the case of lithospheric folding
24
25 985 horizontal distances between the topographic highs and the axes of the depocentres are typically of
26
27 986 the order of hundreds of kilometres. As a result, diffusive transfer of mass will occur at a much
28
29 987 slower rate of sediment supply in the case of folded lithosphere than rates of sediment supply
30
31 988 locally generated by thrusting. This together with a very rapid initial subsidence of folded lithosphere
32
33 989 basins has important consequences for the paleobathymetry of these systems.
34
35 990 FLB contrast with extensional and pull-apart basins in the shape of the subsidence curves, being
36
37 991 convex up for FLB and concave for EB and PAB. The heat flow histories differ also. EB and PAB
38
39 992 are characterized by a phase of initial heating, whereas FLB and FB are characterized by an
40
41 993 absence of initial heating. A common feature of FLB and PAB formation is their capability to cause
42
43 994 dramatic differential vertical motions in a short time interval, predicting shallowing upward
44
45 995 sequences. FLB are to some extent unique in terms of the magnitude of the uplifted topography
46
47 996 surrounding the down warped depocentres. Similar magnitudes are only reached at rift shoulders of
48
49 997 basins developing in intracratonic lithosphere (Van der Beek *et al.*, 1995). FLB can be associated
50
51 998 with localized brittle deformation and pop-up structures in the upper crust and widespread brittle
52
53 999 deformation at deeper crustal and upper mantle levels.
54
55
56
57
58
59
60

CONCLUSIONS

Sedimentary basins controlled by lithospheric folding are characterized by a number of features in their basin architecture, subsidence history, thermal evolution and faulting patterns making them distinctly different from other basin systems. Of particular importance are the relatively short temporal scales involved in their evolution, whereas their spatial scales vary from tens to several hundreds of km, depending on the rheological stratification of the lithosphere and its thermo-mechanical age. Subsidence patterns are characterized by an acceleration of subsidence in the depocentres during compressional basin formation, occurring simultaneously with accelerated uplift of and erosion from flanking basin highs. In contrast with asymmetrical foreland basins, basins on folded lithosphere are symmetrical with equal dimensions for linearly trending topographic highs and lows.

Basins on folded lithosphere have a strong tendency to remain closed systems during a major part of their evolution, with capture times decreasing with thermo-mechanical age from several tens of Myr to less than 10 Myr. The thermal history of these basins is not characterized by an initial thermal perturbation, unless folding occurs in interaction with plume activity. In the absence of an interplay with lithospheric scale thermal perturbations, their thermal evolution will be primarily controlled by the burial history of the sediments deposited in the accommodation space created by the folding process.

Intrabasinal faulting and the formation of pop-up structures and inverted basins at upper crustal levels are important in the structural evolution of these basin systems. The intensity of the brittle deformation peaks with ongoing shortening of the lithosphere during folding but faulting activity is dramatically reduced after the cessation of folding. At this stage, with an equilibrium between sedimentation and erosion, minor faulting occurs as a result of stress relaxation due to erosion of

1
2
3 1026 folded highs. During capture and overall uplift of basins on folded lithosphere, their accommodation
4
5 1027 space is further reduced by unflexing and associated stress release through faulting.
6
7 1028
8
9 1029
10
11
12 1030 **ACKNOWLEDGEMENTS**
13
14 1031 This work was initiated during S. Cloetingh's visit to Paris VI as Visiting Professor and finalized
15
16 1032 during a subsequent Visiting Professorship at IFP. Comments and stimulating suggestions by Fred
17
18 1033 Beekman, Jan-Diederik van Wees, Francois Roure, Seth Stein, Jeroen Smit and Didier Granjeon
19
20 1034 are gratefully acknowledged. Thomas Kruijer and Elishevah van Kooten provided valuable technical
21
22 1035 assistance. We benefitted from constructive criticism from Dr. Mary Ford and two anonymous
23
24 1036 reviewers.
25
26
27 1037
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1038 REFERENCES

- 1039
- 1040 ALINAGHI, A., KOULAKOV, I. & THYBO, H. (2007) Seismic tomographic imaging of *P*- and *S*-waves velocity
1041 perturbations in the upper mantle beneath Iran. *Geophys. J. Int.*, 169, 1089-1102.
- 1042
- 1043 ALZAGA-RUIZ, H., GRANJEON, D., LOPEZ, M., SERANNE, M., ROURE, F. (2008) Gravitational collapse
1044 and Neogene sediment transfer across the western margin of the Gulf of Mexico: Insights from numerical
1045 models. *Tectonophysics*, 470, 21-41.
- 1046
- 1047 ALLEN, M.B. & DAVIES, C.E. (2007) Unstable Asia: active deformation of Siberia revealed by drainage shifts.
1048 *Basin Res.*, 19, 379–392.
- 1049
- 1050 ARTYUSHKOV, E. V. (2007) Formation of the superdeep Caspian Basin: subsidence driven by phase change
1051 in continental crust. *Russ. Geol. Geophys.*, 48, 1102-1014.
- 1052
- 1053 AVOUAC, J., & BUROV, E. (1996) Erosion as a driving mechanism of intracontinental mountain growth. *J.*
1054 *Geophys. Res.*, 101, 17747-17769.
- 1055
- 1056 AXEN, G.J., LAM, P.J., GROVE, M., STOCKLI, D.F. & HASSANZADEH, J. (2001) Exhumation of the west-
1057 central Alborz Mountains, Iran, Caspian subsidence, and collision-related tectonics. *Geology*, 29, 559-562.
- 1058
- 1059 BADA, G., HORVÁTH, F., CLOETINGH, S., COBLENTZ, D.D. & TÓTH, T. (2001) The role of topography
1060 induced gravitational stresses in basin inversion: the case study of the Pannonian basin. *Tectonics*, 20, 343–
1061 363.
- 1062
- 1063 BADA G., HORVÁTH F., DÖVÉNYI P., SZAFIÁN P., WINDHOFFER G. & CLOETINGH S. (2007) Present-
1064 day stress field and tectonic inversion in the Pannonian basin. *Global Planet. Change*, 58, 165-180.
- 1065
- 1066 BEAUMONT, C. (1981) Foreland basins. *Geophys. J. R. Astron. Soc.*, 65, 291–329.
- 1067
- 1068 BEAUMONT, C., FULLSACK, P., & J. HAMILTON (1992) Erosional control of active compressional orogens. In:
1069 *Thrust Tectonics* (Ed. by K.R. McClay), Chapman & Hall, London, 1-18.
- 1070
- 1071 BEEKMAN, F., BULL, J., CLOETINGH, S., SCRUTTON, R., (1996) Crustal fault reactivation facilitating
1072 lithospheric folding/buckling in the Central Indian Ocean. In: *Modern developments in structural interpretation*
1073 *validation and modelling* (Ed. by P. Buchanan and D.A. Nieuwland). Geol. Soc. Lond. Spec. Publ., 99, 251-
1074 263.

- 1075
- 1076 BEEKMAN, F., STEPHENSON, R.A. & KORSCH, R.J. (1997) Mechanical stability of the Redbank Thrust
- 1077 Zone, Central Australia: dynamic and rheological implications. *Australian J. Earth Sci.*, 44, 215-226.
- 1078
- 1079 BERCOVICI, D., RICARD, Y. & SCHUBERT, G. (2001) A two-phase model for compaction and damage 3.
- 1080 Applications to shear localization and plate boundary formation. *J. Geophys. Res.*, 106, 8925-8939.
- 1081
- 1082 BIOT, M.A. (1961) Theory of Folding of Stratified viscoelastic media and its implications in tectonics and
- 1083 orogenesis. *Geol. Soc. Am. Bull.*, 72, 1595-1620.
- 1084
- 1085 BIRD, P. (1991) Lateral extrusion of lower crust from under high topography in the isostatic limit. *J. Geophys.*
- 1086 *Res.*, 96, 10275-10286.
- 1087
- 1088 BONNET, S., GUILLOCHEAU, F., BRUN, J-P. & VAN DEN DRIESSCHE, J. (2000) Large-scale relief
- 1089 development related to Quaternary tectonic uplift of a Proterozoic-Paleozoic basement: The Armorican
- 1090 Massif, NW France. *J. Geophys. Res.*, 105, 19273-19288.
- 1091
- 1092 BOURGEOIS, O., FORD, M., DIRAISON, M., LE CARLIER DE VESLUD, C., GERBAULT, M., PIK, R., RUBY,
- 1093 N. & BONNET, S. (2007) Separation of rifting and lithospheric folding signatures in the NW-Alpine foreland.
- 1094 *Int. J. Earth Sci.*, 96, 1003-1031.
- 1095
- 1096 BRUNET, M.-F., KOROTAEV, M.V., ERSHOV, A.V. & NIKISHIN, A.M. (2003) The South Caspian Basin: a
- 1097 review of its evolution from subsidence modelling. *Sediment. Geol.*, 156, 119-148.
- 1098
- 1099 BULL, J.M. & SCRUTTON, R.A. (1992) Seismic reflection images of intraplate deformation, central Indian
- 1100 Ocean, and their tectonic significance. *J. Geol. Soc. London*, 149, 955-966.
- 1101
- 1102 BURG, J.P., DAVY, P. & MARTINOD, J. (1994) Shortening of analogue models of the continental lithosphere:
- 1103 New hypothesis for the formation of the Tibetan plateau. *Tectonics*, 13, 475-483.
- 1104
- 1105 BURG, J.P. & PODLADCHIKOV, Y. (1999) Lithosphere scale folding: numerical modelling and application to
- 1106 the Himalayan syntaxes. *Int. J. Earth. Sci.*, 88, 190-200.
- 1107
- 1108 BUROV, E.B. & POLIAKOV, A. (2001) Erosion and rheology controls on syn- and post-rift evolution: verifying
- 1109 old and new ideas using a fully coupled numerical model. *J. Geophys. Res.*, 106, 16461-16481.
- 1110
- 1111 BUROV, E.B. & CLOETINGH, S. (1997) Erosion and rift dynamics: new thermo-mechanical aspects of post-
- 1112 rift evolution of extensional basins. *Earth Planet. Sci. Lett.*, 150, 7-26.

- 1113
- 1114 BUROV, E.B. & CLOETINGH, S. (2009) Controls of mantle plumes and lithospheric folding on modes of intra-
1115 plate continental tectonics: differences and similarities. *Geophys. J. Int.*, 178, 1691-1722
- 1116
- 1117 BUROV, E.B., NIKISHIN, A.M., CLOETINGH, S. & LOBKOVSKY, L.I. (1993) Continental lithosphere folding in
1118 central Asia (Part II): Constraints from gravity and tectonic modelling. *Tectonophysics*, 226, 73-87.
- 1119
- 1120 BUROV, E. B. & MOLNAR, P. (1998) Gravity anomalies over the Ferghana Valley (Central Asia) and
1121 intracontinental deformation. *J. Geophys. Res.*, 103, 18137-18152.
- 1122
- 1123 BUROV, E.B., JOLIVET, L., LE POURHIET, L., POLIAKOV, A. (2001) A thermomechanical model of
1124 exhumation of HP and UHP metamorphic rocks in Alpine mountain belts. *Tectonophysics*, 342, 113-136.
- 1125
- 1126 BUROV, E., GUILLOU-FROTTIER, L., D'ACREMONT, E., LE POURHIET, L. & CLOETINGH, S. (2007)
1127 Plume head–lithosphere interactions near intra-continental plate boundaries. *Tectonophysics*, 434, 15-38.
- 1128
- 1129 BUROV E. & TOUSSAINT, G. (2007) Surface processes and tectonics: forcing of continental subduction and
1130 deep processes. *Global Planet. Change*, 58, 141-164.
- 1131
- 1132 BYERLEE, J. D. (1978) Friction of rocks. *Pure Appl. Geophys.* 116, 615-626.
- 1133
- 1134 CALVO, J.P., DAAMS, R., MORALES, J., LÓPEZ MARTÍNEZ, N., AGUSTÍ, J., ANADÓN, P.,
1135 ARMENTEROS, I., CABRERA, L., CIVIS, J., CORROCHANO, A., DÍAZ MOLINA, M., ELIZAGA, E., HOIOS,
1136 M., MARTÍN, E., MARTÍNEZ, J., MOISSENET, E., MUÑOZ, A., PEREZ GARCÍA, A., PÉREZ GONZÁLEZ, A.,
1137 PORTERO, J.M., ROBLES, F., SANTISTEBAN, C., TORRES, T., VAN DER MEULEN, A.J., VERA, J.A. &
1138 MEIN, P. (1993) Up-to-date Spanish continental Neogene synthesis and paleoclimatic interpretation. *Rev.*
1139 *Soc. Geol. España*, 6, 29-40.
- 1140
- 1141 CARTER, N.L. & TSENN, M.C. (1987) Flow properties of continental lithosphere. *Tectonophysics*, 136, 27–
1142 63.
- 1143
- 1144 CASAS-SAINZ, A.M & DE VICENTE, G. (2009) On the tectonic origin of Iberian topography. *Tectonophysics*,
1145 474, 214-235.
- 1146
- 1147 CHU, D. & GORDON, R.G. (1998) Current plate motions across the Red Sea. *Geophys. J. Int.*, 135, 313-328.
- 1148

1
2
3 1149 CLOETINGH, S. (1988) Intraplate stresses: a new element in basin analysis. In: *New perspectives in Basin*
4 1150 *Analysis* (Ed. by K.L. Kleinspehn and C. Poala), Springer-Verlag, New York, 205-230.
5
6 1151
7
8 1152 CLOETINGH, S. & BUROV, E. (1996) Thermomechanical structure of European continental lithosphere:
9 1153 constraints from rheological profiles and EET estimates. *Geophys. J. Int.*, 124, 695–723.
10
11 1154
12
13 1155 CLOETINGH, S. & WEES, VAN J-D. (2005) Strength reversal in Europe’s intraplate lithosphere: transition of
14 1156 basin inversion to lithospheric folding. *Geology*, 33, 285-288.
15
16 1157
17
18 1158 CLOETINGH, S. & ZIEGLER P.A. (2007). Tectonic Models for the Evolution of Sedimentary Basins. In:
19 1159 *Treatise on Geophysics*, 6 (Ed. by A.B. Watts), Elsevier, Amsterdam, 486-611.
20
21 1160
22 1161 CLOETINGH, S., BEEK VAN DER, P.A., REES VAN, D., ROEP, TH.B., BIERMANN, C. & STEPHENSON,
23 1162 R.A. (1992) Flexural interaction and the dynamics of Neogene extensional basin formation in the Alboran-
24 1163 Betic region. *Geo-Mar. Lett.*, 12, 66-75.
25
26 1164
27
28 1165 CLOETINGH, S., BUROV, E. & POLIAKOV, A. (1999) Lithosphere folding: primary response to compression?
29 1166 (from Central Asia to Paris Basin). *Tectonics*, 18, 1064-1083.
30
31 1167
32 1168 CLOETINGH, S., BUROV, E., BEEKMAN, F., ANDEWEG, B., ANDRIESSEN, P., GARCIA-CASTELLANOS,
33 1169 D., DE VICENTE, G. & VEGAS, R. (2002) Lithospheric folding in Iberia. *Tectonics*, 21, 5,
34
35 1170 doi:10.1029/2001TC901031.
36
37 1171
38 1172 CLOETINGH, S., BUROV, E., MATENCO, L., TOUSSAINT, G., BERTOTTI, G., ANDRIESSEN, P.A.M.,
39 1173 WORTEL, M.J.R. & SPAKMAN, W. (2004) Thermo-mechanical controls on the mode of continental collision in
40 1174 the SE Carpathians (Romania). *Earth Planet. Sci. Lett.*, 218, 57-76.
41
42 1175
43
44 1176 CLOETINGH, S., ZIEGLER, P.A., BEEKMAN, F., ANDRIESSEN, P.A.M., MATENCO, L., BADA, G., GARCIA-
45 1177 CASTELLANOS, D., HARDEBOL, N., DÉZES, P. & SOKOUTIS, D. (2005) Lithospheric memory, state of
46 1178 stress and rheology: neotectonic controls on Europe’s intraplate continental topography. *Quat. Sci. Rev.*, 24,
47 1179 241–304.
48
49
50 1180
51
52 1181 CLOETINGH, S., BEEKMAN, F., VAN WEES, J.D., ZIEGLER, P.A. & SOKOUTIS, D. (2008) Post-rift
53 1182 compressional reactivation potential of passive margins and extensional basins. In: *Compressional*
54 1183 *Deformation within Passive Margins: Nature, Causes and Effects* (Ed. by H. Johnson et al.), Geol. Soc.
55
56 1184 London Spec. Pub., 306, 27-70.
57
58 1185
59
60

- COBBOLD, P.R., DAVY, P., GAPAIS, E.A., ROSSELLO, E.A., SADYBASAKOV, E., THOMAS, J.C., TONDJI BIYO, J.J. & DE URREIZTIETA, M. (1993) Sedimentary basins and crustal thickening. *Sediment. Geol.*, 86, 77–89.
- CORTI, G., VAN WIJK, J., BONINI, M., SOKOUTIS, D., CLOETINGH, S., INNOCENTI, F., MANNETI, P. (2003) Transition from continental break-up to seafloor spreading: how fast, symmetric and magmatic. *Geophys. Res. Lett.*, 30, 12, doi:10.1029/2003GL017374.
- CUNDALL, P.A. (1989) Numerical experiments on localization in frictional materials, *Ing. Arch.* 59, 148-159.
- DECELLES, P.G., & GILES, K.A. (1996) Foreland basin systems. *Basin Res.*, 8, 105-123.
- DESEGAULX, P., KOOI, H. & CLOETINGH, S., (1991) Consequences of foreland basin development on thinned continental lithosphere: application to the Aquitaine basin (SW France). *Earth. Planet. Sci. Lett.*, 106, 116-132.
- DE VICENTE, G. & VEGAS, R. (2009) Large-scale distributed deformation controlled topography along the western Africa–Eurasia limit: Tectonic constraints. *Tectonophysics*, 474, 124-143.
- DE VICENTE, G., VEGAS, R., MUÑOZ MARTÍN, A., SILVA, P.G., ANDRIESSEN, P., CLOETINGH, S., GONZÁLES CASADO, J.M., WEES VAN, J-D., ALVAREZ, J., CARBÓ, A. & OLAIZ, A. (2007) Cenozoic thick-skinned deformation and topography evolution of the Spanish Central System. *Global Planet. Change*, 58, 335-381.
- DE VICENTE G., CLOETINGH, S., MUÑOZ-MARTÍN, A., OLAIZ, A., STICH, D., VEGAS, R., GALINDO-ZALDÍVAR, J., FERNÁNDEZ-LOZANO, J. (2008) Inversion of moment tensor focal mechanisms for active stresses around the Microcontinent Iberia: Tectonic implications. *Tectonics*, 27, TC1009.
- DEVLIN, W.J., COGSWELL, J.M., GASKINS, G.M., ISAKSEN, G.H., PITCHER, D.M., PIUS, D.P., STANLEY, K.O. & WALL, J.R.T. (1999) South Caspian basin: young, cool, and full of promise. *GSA Today*, 9, 1-9.
- DÉZES, P., SCHMID, S. M. & ZIEGLER, P. A. (2004) Evolution of the European Cenozoic Rift System: Interaction of the Alpine and Pyrenean orogens with their foreland lithosphere. *Tectonophysics*, 389, 1-33.

1
2
3 1221 DOMBRADI, E., SOKOUTIS, D., BADA, G., CLOETINGH, S., & HORVATH, F. (2010) Modelling of recent
4 1222 deformation of the Pannoanian lithosphere: lithospheric folding and tectonic topography. *Tectonophysics*, 484,
5 1223 103-118.
6 1224
7
8 1225 EGAN, S.S., MOSAR, J., BRUNET, M.-F. & KANGARLI, T. (2009) Subsidence and uplift mechanisms within
9 1226 the South Caspian Basin: insights from the onshore and offshore Azerbaijan region. *Geol. Soc. London Spec.*
10 1227 *Pub.*, 312, 219-240.
11 1228
12
13 1229 ELFRINK, N.M. (2001) Quaternary groundwater avulsions: Evidence for large-scale midcontinent folding?
14 1230 *Assoc. Env. Eng. Geol. News*, 44, 60.
15 1231
16
17 1232 FAKHARI, M.D., AXEN, G.J., HORTON, B.K., HASSANZADEH, J. & AMINI, A. (2008) Revised age of
18 1233 proximal deposits in the Zagros foreland basin and implications for Cenozoic evolution of the High Zagros.
19 1234 *Tectonophysics*, 451, 170-185.
20 1235
21
22 1236 FERNANDEZ-LOZANO, J., SOKOUTIS, D., WILLINGSHOFER, E., DE VICENTE, G. & CLOETINGH, S.
23 1237 (2010) Lithospheric-Scale Folding in Iberia from the perspective of analogue modelling. *Trabajos de Geología.*
24 1238 *Univ. Oviedo. Volumen Especial*, 0474-9588. Coden TBGLA9, in press.
25 1239
26
27 1240 GARCIA-CASTELLANOS, D., FERNANDEZ, M., & TORNE, M. (1997) Numerical modelling of foreland basin
28 1241 formation: a program relating thrusting, flexure, sediment geometry and lithosphere rheology. *Comp. &*
29 1242 *Geosc.*, 23, 993-1003.
30 1243
31
32 1244 GARCIA-CASTELLANOS, D., VERGES, J., GASPAR-ESCRIBANO, J. & CLOETINGH, S. (2003) Interplay
33 1245 between the tectonics, climate, and fluvial transport during the Cenozoic evolution of the Ebro Basin (NE
34 1246 Iberia). *J. Geophys. Res.*, 108, 2347.
35 1247
36
37 1248 GELLER, C. A., WEISSEL, J. K. & ANDERSON, R. N. (1983) Heat transfer and intraplate deformation in the
38 1249 central Indian Ocean. *J. Geophys. Res.*, 88, 1018-1032.
39 1250 .
40
41 1251 GERBAULT, M., BUROV, E., POLIAKOV, A.N.B. & DAIGNIERES, M. (1998) Do faults trigger folding in the
42 1252 lithosphere? *Geophys. Res. Lett.*, 26, 271-274.
43 1253
44
45 1254 GERBAULT, M. (2000) At what stress level is the central Indian Ocean lithosphere buckling? *Earth. Planet.*
46 1255 *Sci. Lett.*, 178, 165-181.
47 1256
48
49 1257 GOETZE, C. & EVANS, B. (1979) Stress and temperature in the bending lithosphere as constrained by
50 1258 experimental rock mechanics. *Geophys. J. Roy. Astron. Soc.*, 59, 463-478.
51
52
53
54
55
56
57
58
59
60

- 1259
- 1260 GOSSMAN, H. (1976) Slope modelling with changing boundary conditions - effects of climate and lithology, *Z.*
1261 *Geomorphol. Suppl.*, 25, 72-88.
- 1262
- 1263 GRANJEON, D. (2009) 3D stratigraphic modelling of sedimentary basins, from source to sink. *AAPG Annual*
1264 *Convention and Exhibition*, June 7-10, 2009, Denver Colorado.
- 1265
- 1266 GRENERCZY, Y., SELLA, G.F., STEIN, S. & KENYERES, A. (2005) Tectonic implications of the GPS velocity
1267 field in the northern Adriatic region. *Geophys. Res. Lett.*, 32, L16311.
- 1268
- 1269 GUEST, B., GUEST, A. & AXEN, G. (2007). Late Tertiary tectonic evolution of northern Iran: a case for simple
1270 crustal folding. *Global. Planet. Change*, 58, 435-453.
- 1271
- 1272 GUILLOCHEAU *et al.* (2000) Meso-Cenozoic geodynamic evolution of the Paris Basin: 3D stratigraphic
1273 constraints. *Geodin. Acta*, 13, 189-246.
- 1274
- 1275 GUILLOU-FROTTIER, L., BUROV, E., NEHLIG, P. & WYNS, R. (2007) Deciphering plume–lithosphere
1276 interactions beneath Europe from topographic signatures. *Global Planet. Change*, 58, 119-140.
- 1277
- 1278 HAFKENSCHIED, E., WORTEL, M.J.R. & SPAKMAN, W. (2006) Subduction history of the Tethyan region
1279 derived from seismic tomography and tectonic reconstructions. *J. Geophys. Res.*, 111, B08401.
- 1280
- 1281 HORVÁTH, F. (1995) Phases of compression during the evolution of the Pannonian Basin and its bearing on
1282 hydrocarbon exploration. *Mar. Petrol. Geol.*, 12, 837-844.
- 1283
- 1284 HORVÁTH, F. & CLOETINGH, S. (1996) Stress-induced late-stage subsidence anomalies in the Pannonian
1285 basin. *Tectonophysics*, 266, 287–300.
- 1286
- 1287 HORVÁTH, F., BADA, G., SZAFIÁN, P., TARI, G., ÁDÁM, A. & CLOETINGH, S. (2006) Formation and
1288 deformation of the Pannonian basin: constraints from observational data. *Geol. Soc. London Mem.*, 32, 191-
1289 206.
- 1290
- 1291 HUISMANS, R.S. & BEAUMONT, C.B. (2002) Asymmetric lithospheric extension: the role of frictional plastic
1292 strain softening inferred from numerical experiments. *Geology*, 30, 211-214.
- 1293
- 1294 HUNT, G., MUHLHAUS, H., HOBBS, B., & ORD, A. (1996) Localized folding of viscoelastic layers. *Geol.*
1295 *Rundsch.*, 85, 58-64.
- 1296

1
2
3 1297 KABAN, M.K. (2002) A gravity model of the north Eurasia crust and upper mantle: 2. The Alpine-
4 1298 Mediterranean fold belt and adjacent structures of the southern former USSR. *Russ. J. Earth. Sci.*, 4, 19-33.
5
6 1299
7
8 1300 KIRBY, S.H. & KRONENBERG, A.K. (1987) Rheology of the lithosphere: selected topics. *Rev. Geophys.* 25,
9 1301 1219-1244.
10 1302
11
12 1303 KIRKBY, M.J. (1986) A two-dimensional model for slope and stream evolution. In: *Hillslope Processes* (Ed. by
13 1304 Abrahams, A.D.) Allen and Unwin, Boston, 203-224.
14
15 1305
16
17 1306 KOHLSTEDT, D.L., EVANS, B. & MACKWELL, S.J. (1995) Strength of the lithosphere: constraints imposed
18 1307 by laboratory experiments. *J. Geophys. Res.*, 100, 17587-17602.
19
20 1308
21 1309 KOMBRINK, H., LEEVER, K.A., VAN WEES, J-D., VAN BERGEN, F., DAVID, P., & WONG, T.E. (2008) Late
22 1310 Carboniferous foreland basin formation and Early Carboniferous stretching in Northwestern Europe:
23 1311 inferences from quantitative subsidence analyses in the Netherlands. *Basin Res.*, 20, 377-395.
24
25 1312
26
27 1313 KOOI, H. & BEAUMONT, C. (1994) Escarpment evolution on high-elevation rifted margins: insights derived
28 1314 from a surface processes model that combines diffusion, advection and reaction, *J. Geophys. Res.*, 99,
29 1315 12191-12210.
30
31 1316
32
33 1317 LAGARDE, J.L. BAIZE, S., AMORESE, D., DELCCAILLAU, B., FORT, M. & VOLANT, V. (2000) Active
34 1318 tectonics, seismicity and geomorphology with special reference to Normandy (France). *J. Quat. Sci.*, 15, 745-
35 1319 758.
36
37 1320
38
39 1321 LAMBECK, K. (1983) The role of compressive forces in intracratonic basin formation and mid-plate orogenies.
40 1322 *Geophys. Res. Lett.*, 10, 845-848.
41
42 1323
43
44 1324 LAVIER, L.L. & STECKLER, M.S. (1997) The effect of sedimentary cover on the flexural strength of the
45 1325 continental lithosphere. *Nature*, 389, 476-479.
46 1326
47 1327 LAWRENCE D.T., DOYLE. M., & AIGNER, T. (1990) Stratigraphic simulation of sedimentary basins:
48 1328 Concepts and calibration. *AAPG Bull.*, 74, 273-295.
49
50 1329
51 1330 LEEDER, M. R. (1991) Denudation, vertical crustal movements and sedimentary basin infill. *Geol. Rundsch.*,
52 1331 80, 441-458.
53
54 1332
55
56 1333 LEFORT, J-P. & AGARWAL, B.N. (1996) Gravity evidence for an Alpine buckling of the crust beneath the
57 1334 Paris Basin. *Tectonophysics*, 258, 1-14.
58
59
60

- 1335
- 1336 LEFORT, J.P. & AGARWAL, B.N. (2000) Gravity and geomorphological evidence for a large crustal bulge
1337 cutting across Brittany (France): a tectonic response to the closure of the Bay of Biscay. *Tectonophysics*, 323,
1338 149-162.
- 1339
- 1340 LEFORT, J.P. & AGARWAL, B.N. (2002) Topography of the Moho undulations in France from gravity data:
1341 their age and origin. *Tectonophysics*, 350, 193-213.
- 1342
- 1343 LEIGHTON, M.W. & KOLATA, D.R. (1990) Selected interior cratonic basins and their place in the scheme of
1344 global tectonics: A synthesis. In: *AAPG Memoir* (Ed. by M.W. Leighton, D.R. Kolata, D.F. Oltz and J.J. Eidel),
1345 51, 729-797.
- 1346
- 1347 LENOTRE, N., THIERRY, P., BLANCHIN, R. & BROCHARD, G. (1999) Current vertical movement
1348 demonstrated by comparative levelling in Brittany (France). *Tectonophysics*, 301, 333-344.
- 1349
- 1350 LE SOLLEUZ, A., DOIN, M.-P. DOIN, ROBIN, C., & GUILLOCHEAU, F. (2004) From a mountain belt collapse
1351 to a sedimentary basin development: 2-D thermal model based on inversion of stratigraphic data in the Paris
1352 Basin. *Tectonophysics*, 386, 1-27.
- 1353
- 1354 LI, S., MOONEY, W.D. & FAN, J. (2006) Crustal structure of mainland China from deep seismic sounding
1355 data. *Tectonophysics*, 420, 239-252.
- 1356
- 1357 LITCKE, R., BAYER, U., GAJEWSKI, D., & NELSKAMP, S. (2008) Dynamics of Complex Intracontinental
1358 Basins, Berlin. Heidelberg, Springer, 520 p.
- 1359
- 1360 LIU, M., MOONEY, W.D., LI, S., OKAYA, N., & DETWEILER, S. (2006) Crustal structure of the northeast
1361 margin of the Tibetan plateau from the Songpan-Ganzi terrane to the Ordos basin. *Tectonophysics*, 420, 253-
1362 266.
- 1363
- 1364 LIU Y., NEUBAUER, F., GENSER, J., GE, X., TAKASU, A., YUAN, S., CHANG, L., & LI, W. (2007)
1365 Geochronology of the initiation and displacement of the Altyn strike-slip fault, western China. *J. Asian Earth*
1366 *Sci.*, 29, 243-252.
- 1367
- 1368 LUNDIN, E.R. & DORÉ, A.G. (1997) A tectonic model for the Norwegian passive margin with implications for
1369 the NE Atlantic: Early Cretaceous to break-up. *J. Geol. Soc.*, 154, 545-550.
- 1370
- 1371 MAROTTA, A. M., BAYER, U. & THYBO, H. (2000) The legacy of the NE German Basin – Reactivation by
1372 compressional buckling. *Terra Nova*, 12, 132-14

1
2
3 1373
4
5 1374 MARTINOD, J. & DAVY, P. (1992) Periodic instabilities during compression or extension of the lithosphere: 1.
6 1375 Deformation modes from an analytical perturbation method. *J. Geophys. Res.*, 97, 1999-2014.
7
8 1376
9 1377 MARTINOD, J. & DAVY, P. (1994) Periodic instabilities during compression of the lithosphere: 2. Analogue
10 1378 experiments. *J. Geophys. Res.*, 99, 12057-12069.
11
12 1379
13 1380 MATENCO, L., BERTOTTI, G., LEEVER, K., CLOETINGH, S., SCHMID, S., TĂRĂPOANCĂ, M. & DINU, C.,
14 1381 2007. Large-scale deformation in a locked collisional boundary: Interplay between subsidence and uplift,
15 1382 intraplate stress, and inherited lithospheric structure in the late stage of the SE Carpathians evolution.
16 1383 *Tectonics*, 26, TC4011
17
18 1384
19
20 1385 MAZUR, S., SCHECK-WENDEROTH M. & KRZYWIEC, P. (2005) Different modes of the Late Cretaceous –
21 1386 Early Tertiary inversion in the North German and Polish basins. *Int. J. Earth Sci.*, 94, 782-798.
22
23 1387
24 1388 MCADOO, D.C. & SANDWELL, D.T. (1985) Folding of the oceanic lithosphere. *J. Geophys. Res.*, 90, 8563–
25 1389 8569.
26
27 1390
28
29 1391 MCKENZIE, D.P. (1978) Some remarks on the development of sedimentary basins. *Earth. Planet. Sci. Lett.*,
30 1392 95, 53-72.
31
32 1393
33 1394 MUEHLHAUS, H.B., SAHAGUCHI, H., & HOBBS, B.E. (1998) Evolution of three-dimensional folds for a non
34 1395 Newtonian plate in a viscous medium. *Proc. R. Soc. London, series A, Math. Phys. NG. Sci.*, 454, 3121-3143.
35 1396
36 1397 NADIROV, R.S., BAGIROV, E., TAGIYEV, M., & LERCHE, I. (1997) Flexural plate subsidence, sedimentation
37 1398 rates and structural development of the super-deep south Caspian Basin. *Mar. Petrol. Geol.*, 14, 383-400.
38 1399
39 1400 NAYLOR, M. & SINCLAIR, H.D. (2007) Punctuated thrust deformation in the context of doubly vergent thrust
40 1401 wedges: implications for the localization of uplift and exhumation. *Geology*, 35, 559–562.
41 1402
42 1403 NAYLOR, M. & SINCLAIR, H.D. (2008) Pro- vs. Retro foreland Basins. *Basin Res.*, 20, 1365-2117.
43 1404
44 1405 NECEA, D., FIELITZ, W., & MATENCO, L. (2005) Late Pliocene-Quaternary tectonics in the frontal part of the
45 1406 SE Carpathians: Insights from tectonic geomorphology. *Tectonophysics*, 410, 137-156.
46 1407
47 1408 NIKISHIN, A.M., CLOETINGH, S., LOBKOVSKY, L., & BUROV, E.B. (1993) Continental lithosphere folding in
48 1409 Central Asia (Part I): constraints from geological observations. *Tectonophysics*, 226, 59-72.
49 1410
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 1411 NIKISHIN, A.M., BRUNET, M.F., CLOETINGH, S., & ERSHOV, A.V. (1997) Northern Peri-Tethyan Cenozoic
4 1412 intraplate deformations: Influence of the Tethyan collision belt on the Eurasian continent from Paris to Tian-
5 1413 Shan. *C.R. Acad. Sci., Ser. Ila: Sci. Terre Planets*, 234, 49-57.
6
7 1414
8
9 1415 NIKISHIN, A.M., KOROTAEV, M.V., ERSHOV, A.V., & BRUNET, M.F. (2003) The Black Sea Basin: tectonic
10 1416 history and Neogene-Quaternary rapid subsidence modelling. *Sediment. Geol.*, 156, 149-168.
11 1417
12 1418 PEREZ-GUSSINYE, M. & WATTS, A.B. (2005) The long-term strength of Europe and its implications for
13 1419 plate-forming processes. *Nature*, 436, 381-384.
14
15 1420
16
17 1421 PITMAN, W.C. & GOLOVCHENKO, X. (1983) The effect of sea-level change on the shelf edge and slope of
18 1422 passive margin. In: *The shelf break: critical interface on continental margins* (Ed. by D.J. Stanley and G.T.
19 1423 Moore), SEPM, Spec. Publ., 33, 41-58.
20
21 1424
22
23 1425 QUINLAN, G. & BEAUMONT, C. (1984) Appalachian thrusting, lithospheric flexure and the Paleozoic
24 1426 stratigraphy of the eastern interior of North America. *Can. J. Earth Sci.*, 21, 973-996.
25
26 1427
27
28 1428 QUINLAN, G. (1987) Models of subsidence mechanisms in intracratonic basins and their applicability to North
29 1429 American examples. *Can. Soc. Pet. Geol. Mem.*, 12, 463-481.
30
31 1430
32
33 1431 RAMBERG, H. (1961) Contact strain and folding instability of multilayered body under compression. *Geol.*
34 1432 *Rundsch.*, 51, 405-439.
35
36 1433
37 1434 RANALLI, G. (1995) *Rheology of the Earth*, 2nd edn., Chapman and Hall, London, 413 pp.
38
39 1435
40 1436 RITTER, J.R.R., JORDAN, M., CHRISTENSEN, U.R. & ACHAUER, U. (2001) A mantle plume under the Eifel
41 1437 volcanics fields, Germany. *Earth Planet. Sci. Lett.*, 186, 7-14.
42
43 1438
44 1439 RITZMANN, O. & FALEIDE, J.I. (2009) The crust and mantle lithosphere in the Barents Sea/Kara Sea region.
45 1440 *Tectonophysics*, 470, 89-104.
46
47 1441
48
49 1442 ROBIN, C., ALLEMAND, P., BUROV, E., DOIN, M.P., GUILLOCHEAU, F., DROMART, G. & GARCIA, J.P.
50 1443 (2003) Vertical movements of the Paris Basin (Triassic - Pleistocene): from 3D stratigraphic database to
51 1444 numerical models. In: *New insights in structural interpretation and modelling* (Ed. by P. Buchanan and D.A.
52 1445 Nieuwland) Geol. Soc. London Spec. Publ., 212, 225-250.
53
54 1446
55
56 1447 ROYDEN, L. (1988) Flexural behaviour of the continental lithosphere in Italy: constraints imposed by gravity
57 1448 and deflection data. *J. Geophys. Res.*, 93, 7747-7766.
58
59
60

1
2
3 1449
4
5 1450 ROYDEN, L.H. (1993) The tectonic expression of the slab pull at continental convergent boundaries.
6 1451 *Tectonics*, 12, 303-325.
7
8 1452
9 1453 ROURE, F., CLOETINGH, S., SCHECK-WENDEROTH, M., & ZIEGLER, P.A., (2010) Achievements and
10 1454 challenges in sedimentary basin dynamics: a review. In: *New Frontiers in Integrated Solid Earth Sciences* (Ed.
11 1455 by S. Cloetingh and J. Negendank), Springer-Verlag, 145-233.
12
13 1456
14
15 1457 SACCHI, M., HORVATH, F. & MAGYARI, O. (1999) Role of unconformity bounded units in the stratigraphy of
16 1458 the continental record: a casestudy from the Late Miocene of the western Pannonian basin, Hungary. *Geol.*
17 1459 *Soc. London Spec. Publ.*, 156, 357-390.
18
19 1460
20
21 1461 SCHECK, M. & BAYER, U., (1999) Evolution of the Northeast German Basin – inferences of a 3D structural
22 1462 model and subsidence analysis. *Tectonophysics*, 373, 55-73.
23
24 1463
25 1464 SCHMALHOLTZ, S.M. & PODLADCHIKOV, Y.Y. (2000) Finite amplitude folding: transition from exponential
26 1465 to taper length controlled growth. *Earth Planet. Sci. Lett.*, 181, 619 -633.
27
28 1466
29
30 1467 SCHUBERT, G., TURCOTTE, D.L. & OLSON, P. (2001) Mantle convection in the Earth and Planets.
31 1468 *Cambridge University Press*, 956 pp.
32
33 1469
34 1470 SHIN, Y.H., SHUM, C.-K., BRAITENBERG, C., LEE, S.M., XU, H., CHOI, K.S., BAEK, J.H., & PARK, J.U.
35 1471 (2009) Three-dimensional fold structure of the Tibetan Moho from GRACE gravity data. *Geophys. Res. Lett.*,
36 1472 36, L01302.
37
38 1473
39
40 1474 SMIT, J., BRUN, J.P., CLOETINGH, S., & BEN-AVRAHAM, Z. (2008) Pull-apart basin formation and
41 1475 development in narrow transform zones with application to the Dead Sea Basin. *Tectonics*, 27, TC6018.
42
43 1476
44 1477 SMIT, J., BRUN, J.P., CLOETINGH, S. & BEN-AVRAHAM, Z. (2010) The rift-like structure and asymmetry of
45 1478 the Dead Sea Fault. *Earth Planet. Sci. Lett.*, 290, 74-82.
46
47 1479
48
49 1480 SMITH, R.B. (1979) The folding of a strongly non-Newtonian layer. *Am. J. Sci.*, 79, 272-287.
50 1481
51
52 1482 SOKOUTIS, D., BURG, J.P., BONINI, M., CORTI, G. & CLOETINGH, S. (2005) Lithospheric-scale structures
53 1483 from the perspective of analogue continental collision. *Tectonophysics*, 406, 1-15.
54
55 1484
56 1485 SOLOMATOV, V.S. & MORESI, L.N. (2000) Scaling of time-dependent stagnant lid convection; application to
57 1486 small-scale convection on Earth and other terrestrial planets. *J. Geophys. Res.*, 105, 21795-21817.
58
59
60

- 1487
- 1488 STEIN, C.A., CLOETINGH, S. & WORTEL, R. (1989) Seasat-derived gravity constraints on stress and
- 1489 deformation in the northeastern Indian Ocean. *Geophys. Res. Lett.*, 16, 823-826.
- 1490
- 1491 STEL, H., CLOETINGH, S., HEEREMANS, M., & VAN DER BEEK, P.A. (1993) The kinematics of anorogenic
- 1492 basement tectonics and the origin of sedimentary basins. *Tectonophysics*, 226, 285-299.
- 1493
- 1494 STEPHENSON, R.S. & CLOETINGH, S. (1991) Some examples and mechanical aspects of continental
- 1495 lithospheric folding. *Tectonophysics*, 188, 27-37.
- 1496
- 1497 STEPHENSON, R., & LAMBECK, K. (1985) Isostatic response of the lithosphere with in-plane stress:
- 1498 application to Central Australia. *J. Geophys. Res.*, 90, 8581-8588.
- 1499
- 1500 STEPHENSON, R.S., RICKETTS, B.D., CLOETINGH, S., & BEEKMAN, F. (1990) Lithosphere folds in the
- 1501 Eureka orogen, Arctic Canada? *Geology*, 18, 603-606.
- 1502
- 1503 TARAPOANCA, M., BERTOTTI, G., MATENCO, L., DINU, C. & CLOETINGH, S. (2003) Architecture of the
- 1504 Focsani Depression: A 13km deep basin in the Carpathians bend zone (Romania). *Tectonics*, 22, 1074.
- 1505
- 1506 TER VOORDE, M., DE BRUIJNE, C.H., CLOETINGH, S.A.P.L. & ANDRIESSEN, P.A.M. (2004) Thermal
- 1507 consequences of thrust faulting: simultaneous versus successive fault activation and exhumation. *Earth*.
- 1508 *Planet. Sci. Lett.*, 223, 395-413.
- 1509
- 1510 TESAURO, M., HOLLENSTEIN, C., EGLI, R., GEIGER, A. & KAHLE, H.-G. (2005) Continuous GPS and
- 1511 broad-scale deformation across the Rhine Graben and the Alps. *Int. J. Earth. Sci.*, 94, 525-537.
- 1512
- 1513 TESAURO, M., KABAN, M., CLOETINGH, S., HARDEBOL, N. & BEEKMAN, F. (2007) 3D strength and
- 1514 gravity anomalies of the European lithosphere. *Earth Planet. Sci. Lett.*, 263, 56-73.
- 1515
- 1516 THOMAS, J.C., COBBOLD, P.R., SHEIN, V.S. & LE DOUARAN, S. (1999a) Sedimentary record of late
- 1517 Paleozoic to Recent tectonism in central Asia - analysis of subsurface data from the Turan and south Kazak
- 1518 domains. *Tectonophysics*, 313, 243-263.
- 1519
- 1520 THOMAS, J.C., GRASSO, J.R., BOSSU, R., MARTINOD, J., & NURTAEV, B. (1999b) Recent deformation in
- 1521 the Turan and South Kazakh platforms, western central Asia, and its relation to Arabia-Asia and India-Asia
- 1522 collisions. *Tectonics*, 18, 201-214.
- 1523

1
2
3 1524 TIKOFF, B. & MAXSON, J. (2001) Lithospheric buckling of the Laramide foreland during Late Cretaceous and
4 1525 Paleogene, western United States. *Rocky Mt Geol.*, 36, 13-35.
5
6 1526
7
8 1527 TURCOTTE, D.L. & SCHUBERT, G. (2002) *Geodynamics, applications of continuum physics to geological*
9 1528 *problems*, 2nd edn., *Cambridge University Press*, 456 pp.
10 1529
11
12 1530 VAN BALEN, R.T., VAN DER BEEK, P.A. & CLOETINGH, S. (1995). The effect of rift shoulder erosion on
13 1531 stratal patterns at passive margins: implications for sequence stratigraphy. *Earth Planet. Sci. Lett.*, 134, 527-
14 1532 544.
15
16 1533
17
18 1534 VAN DER BEEK., P., ANDRIESSEN, P.A.M., & CLOETINGH, S. (1995). Morphotectonic evolution of rifted
19 1535 continental margins: inferences from a coupled tectonic-surface processes model and fission track
20 1536 thermochronology. *Tectonics*, 14, 406-421.
21 1537
22
23 1538 VAN DER PLUIJM, B., CRADDOCK, J.P., GRAHAM, B.R. & HARRIS, J.H. (1997) Paleostress in cratonic
24 1539 North America: implications for deformation of continental interiors. *Science*, 277, 794-796.
25
26 1540
27
28 1541 VAN WEES, J-D. & CLOETINGH, S. (1996) 3D flexure and intraplate compression in the North Sea basin.
29 1542 *Tectonophysics*, 266, 343-359.
30 1543
31
32 1544 VAN WEES, J.D., VAN BERGEN, F., DAVID, P., NEPVEU, M., BEEKMAN, F. & CLOETINGH, S. (2009)
33 1545 Probabilistic Tectonic heat flow modelling for basin maturation: method and applications. *Mar. Petrol. Geol.*,
34 1546 26, 536-551.
35
36 1547
37
38 1548 VERGES, J, (1999) Estudi Geologic Del Vessant Sud Del Pirineu Oriental I Central. In: *Evolucio Cinematica*,
39 1549 3rd edn. (Ed. by Col·leccio Monografies Techniques), Institut Cartografic de Catalunya, Barcelona, 7, 194.
40
41 1550
42
43 1551 VERNANT, P., NILFOROUSHAN, F., HATZFELD, D., ABBASSI, M.R., VIGNY, C., MASSON, F., NANKALI,
44 1552 MARTINOD, J., ASHTIANI, A., BAYER, R., TAVAKOLI, F., CHERY, J. (2004) Present-day crustal
45 1553 deformation and plate kinematics in the Middle East constrained by GPS measurements in Iran and northern
46 1554 Oman. *Geophys. J. Int.*, 157, 381-398.
47
48 1555
49
50 1556 VERMEESCH, P., POORT, J., DUCHKOV, A.D., KLERKX, J. & DE BATIST, M. (2004) Lake Issyk-Kul (Tien
51 1557 Shan): Unusually low heat flow in an active intramontane basin. *Russ. Geol. Geophys.*, 45, 616-625.
52
53 1558
54
55 1559 WATTS, A.B. & BUROV E.B. (2003) Lithospheric strength and its relationship to the elastic and seismogenic
56 1560 layer thickness. *Earth. Planet. Sci. Lett.*, 213, 113-131.
57
58 1561
59
60

- WEINBERG, R.F. & PODLADCHIKOV, Y. (1994) Diapiric ascent of magmas through power law crust and mantle. *J. Geophys. Res.*, 99, 9543-9559.
- WILGUS, C.K., HASTINGS, B.S., KENDALL, C.G.St.C., PASAMENTIER, H.W., ROSS, C.A., VAN WAGONER, J.C. (1988) Sea-level changes – an integrated approach. *Spec. Pub. Soc. Econ. Pal. Min.*, 42, 402.
- WORTEL, M.J.R. & SPAKMAN, W. (2000) Subduction and slab detachment in the Mediterranean Carpathian region. *Science*, 290, 1910-1917.
- XIE, X. & HELLER, P.L. (2009) Plate tectonics and basin subsidence history. *Geol. Soc. Am. Bull.*, 121, 55-64.
- ZHAO, J., MOONEY, W.D, ZHANG, X., LI, Z., JIN, Z., & OKAYA, N. (2006) Crustal structure across the Atityn Tagh Range at the northern margin of the Tibetan plateau and tectonic implications. *Earth Planet. Sci. Lett.*, 241, 804-841.
- ZIEGLER, P.A., CLOETINGH, S. & VAN WEES, J-D. (1995) Dynamics of intraplate compressional deformation: the Alpine foreland and other examples. *Tectonophysics*, 252, 7–59.
- ZIEGLER, P.A., VAN WEES, J-D. & CLOETINGH, S. (1998) Mechanical controls on collision-related compressional intraplate deformation. *Tectonophysics*, 300, 103-129.
- ZIEGLER, P.A. & DÈZES, P. (2006) Crustal evolution of Western and Central Europe. In: *European Lithosphere Dynamics* (Ed. by D.G. Gee and R.A. Stephenson), *Geol. Soc. London Mem.*, 32, 43-56.
- ZIEGLER, P.A. & DÈZES, P. (2007) Cenozoic uplift of Variscan Massifs in the Alpine foreland: Timing and controlling mechanisms. *Global Planet. Change*, 58, 237-269.
- ZITELLINI, N., CLOETINGH, S., D'ORIANO, F., & BUROV, E.B., (2010) Synclinal deformation as prime expression of compressional deformation in the lithosphere: the Atlantic segment of the Iberia-Africa plate boundary. *Geology*, submitted.
- ZOETEMEIJER, R., CLOETINGH, S., SASSI, W. & ROURE, F. (1993) Modelling of piggy-back basin stratigraphy: record of tectonic evolution. *Tectonophysics*, 226, 253-269.
- ZOETEMEIJER, R., TOMEK, C. & CLOETINGH, S. (1999) Flexural expression of European continental lithosphere under the western outer Carpathians. *Tectonics*, 18, 843-861.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

APPENDIX A: MODEL DESCRIPTION

A1. Equations. The Flamar-Para(o)voz is a “2.5 D” FLAC-like code (Cundall, 1989). It has a mixed finite-difference/finite element numerical scheme, in which the coordinate frame is Cartesian 2D, but stress/strain relations are computed in a 3D formulation. The Lagrangian mesh of Paravoz is composed of quadrilateral elements subdivided onto 2 couples of triangular sub-elements with triangular shape functions. Para(o)voz is a large strain fully explicit time-marching algorithm. It locally solves full Newtonian equations of motion in continuum mechanics approximation:

$$\langle \rho \ddot{\mathbf{u}} \rangle - \text{div} \boldsymbol{\sigma} - \rho \mathbf{g} = 0 \quad (\text{A1})$$

coupled with constitutive equations:

$$\frac{D\boldsymbol{\sigma}}{Dt} = F(\boldsymbol{\sigma}, \mathbf{u}, \nabla \dot{\mathbf{u}}, \dots, T \dots) \quad (\text{A2})$$

and with equations of heat transfer (heat advection $\mathbf{u} \nabla T$ in the equation below is solved separately):

$$\rho C_p D T / D t + \dot{\mathbf{u}} \nabla T - k \text{div}(\nabla T) - H_r = 0 \quad (\text{A3})$$

$$\rho = \rho_0(1 - \alpha T)$$

Here $\mathbf{u}$, $\boldsymbol{\sigma}$, $\mathbf{g}$, k are the respective terms for the displacement, stress, acceleration due to body forces and thermal conductivity. The triangular brackets in (A1) specify conditional use of the related term (in quasi-static mode inertial terms are dumped using inertial mass scaling (Cundall, 1989)). The terms t , ρ , C_p , T and H_r designate respectively time, density, specific heat, temperature and internal heat production. The terms D/Dt , $D\boldsymbol{\sigma}/Dt$, F are a time derivative, an objective (Jaumann) stress time derivative and a functional, respectively. In the Lagrangian method, incremental displacements are added to the grid coordinates allowing the mesh to move and deform with material. This enables solution of large-strain problems locally using a small-strain formulation. On

1
2
3 1619 each time step the solution is obtained in local coordinates, which are then updated in a large strain
4
5 1620 mode.
6
7
8 1621 Solution of (A1) provides velocities at mesh points used for computation of element strains and of
9
10 1622 heat advection $\mathbf{u} \nabla T$. These strains are used in (A2) to calculate element stresses and equivalent
11
12 1623 forces as input for the computation of the velocities for the next time step. Due to the explicit
13
14 1624 approach, there are no convergence issues, which is rather a common problem of implicit methods
15
16 1625 in case of non-linear rheologies. The algorithm automatically checks and adopts the internal time
17
18 1626 step using 0.1 – 0.5 of Courant’s criterion for propagation of information, which warrants a stable
19
20 1627 solution.
21
22
23

24 1628
25
26 1629 **A2. Explicit EVP rheology.**
27
28

29 1630 We use a serial (Maxwell type) body (eq. 1), in which total strain increment in each numeric element
30
31 1631 is defined by a sum of elastic, viscous and brittle strain increments. Consequently, in contrast to
32
33 1632 fluid dynamic approaches, where non-viscous rheological terms are simulated using pseudo-plastic
34
35 1633 and pseudo-elastic viscous terms (e.g. Bercovici *et al.*, 2001; Solomatov & Moresi, 2000), our
36
37 1634 method explicitly treats all rheological terms. The parameters of elastic-ductile-plastic rheology laws
38
39 1635 for crust and mantle come from rock mechanics data (**Tables A1 and A2**; Kirby & Kronenberg,
40
41 1636 1987; Kohlstedt *et al.*, 1995).
42
43
44

45 1637
46
47 1638 **Plastic (brittle) behavior.**
48
49

50 1639 The brittle behavior of rocks is described by Byerlee’s law (Byerlee, 1978; Ranalli, 1995), which
51
52 1640 corresponds to Mohr-Coulomb material with a friction angle $\phi = 30^\circ$ and cohesion $|C_0| < 20$ MPa
53
54 1641 (e.g., Gerbault *et al.*, 1998):
55
56
57
58
59
60

$$\tau = C_0 - \sigma_n \tan \phi \quad (A4)$$

where σ_n is normal stress $\sigma_n = \frac{1}{3}\sigma_1 + \sigma_{II}^{\text{dev}} \sin \phi$, $\frac{1}{3}\sigma_1 = P$ is the effective pressure, σ_{II}^{dev} is the second invariant of deviatoric stress, or effective shear stress. The condition of transition to brittle deformation (function of rupture f) reads as: $f = \sigma_{II}^{\text{dev}} + P \sin \phi - C_0 \cos \phi = 0$ and $\partial f / \partial t = 0$. In terms of principal stresses, the equivalent of the yield criterion (5) reads as:

$$\sigma_1 - \sigma_3 = -\sin \phi (\sigma_1 + \sigma_3 - 2C_0 / \tan \phi) \quad (A5)$$

Elastic behavior.

The elastic behavior is described by the linear Hooke's law:

$$\sigma_{ij} = \lambda \varepsilon_{ij} \delta_{ij} + 2G \varepsilon_{ij} \quad (A6)$$

where λ and G are Lamé's constants. Repeating indexes mean summation and δ is Kronecker's operator.

Viscous (ductile) behavior.

Mantle convection and a part of lithospheric deformation is controlled by thermally activated creep (Ranalli, 1995; Kirby & Kronenberg, 1987). Within deep lithosphere and underlying mantle regions, the creeping flow is non-Newtonian since the effective viscosity can vary within 10 orders of magnitude as a function of differential stress:

$$\dot{\varepsilon}_{ij}^d = A (\sigma_1 - \sigma_3)^n \exp(-Q / RT) \quad (A7)$$

Where $\dot{\varepsilon}_{ij}^d = \dot{\varepsilon}_{ij}^d$ is shear strain rate, A is material constant, n is the power law exponent, Q is the activation enthalpy, R is Boltzman's gas constant, and T is temperature in K°, σ_1 and σ_3 are the principal stresses. The effective viscosity μ_{eff} for this law is defined as:

$$\tau_{ij} \equiv \mu_{eff} e_{ij}^d$$

1660 which yields:

$$\mu_{eff} = e_{ij}^{d(1-n)/n} A^{-1/n} \exp(Q(nRT)^{-1}) \quad (A8)$$

1661 For non-uniaxial deformation, the law (A8) is converted to a triaxial form, using invariant of strain
1662 rate and geometrical proportionality factors:

$$\mu_{eff} = e_{ij}^{d(1-n)/n} A^{-1/n} \exp(Q(nRT)^{-1})$$

1663 where

$$e_{II}^d = (\ln v_{II}(\mathbf{e}_{ij}))^{1/2} \text{ and } A^* = \frac{1}{2} A \cdot 3^{(n+1)/2} \quad (A9)$$

1664 parameters A, n, Q are experimentally determined material constants (**Table A1**). Using olivine
1665 parameters (**Table A1**), one can verify that the predicted effective viscosity at the base of the
1666 lithosphere is $10^{19} - 5 \times 10^{19}$ Pa s matching post-glacial rebound data (Turcotte & Schubert, 2002).
1667 In the depth interval of 200 km - 0 km the effective viscosity grows from 10^{19} to $10^{25} - 10^{27}$ Pa s with
1668 decreasing temperature. Within the adiabatic temperature interval at depth, the dislocation flow law
1669 (A8) is replaced by nearly Newtonian diffusion creep, which results in a quasi-constant mantle
1670 viscosity of $10^{19} - 10^{21}$ Pa s (e.g., Turcotte & Schubert, 2002). Weinberg & Podladchikov (1994)
1671 have also shown that the effective viscosity in close vicinity of an ascending diapir is influenced by
1672 the local strain rate field and partly by heat exchanges between the diapir and surrounding rock,
1673 which suggests possible changes.

1674

FIGURE CAPTIONS

Figure 1. Concepts of folding in rheologically stratified lithosphere and feedback with sedimentation in downfolded areas and erosion of adjacent highs. Different simultaneously occurring wavelengths of crustal and mantle folding are a consequence of the rheological stratification of the lithosphere. Surface wavelengths can be affected also by feedback with surface processes.

Figure 2. Map with examples of well-documented areas affected by lithospheric folding. Areas: Central Indian Ocean Basin (CIOB) (Geller *et al.*, 1983 ; Stein *et al.*, 1989) ; NW European platform (Bourgeois *et al.*, 2007 ; Lefort & Agarwal, 1996; Marotta *et al.*, 2000); Pannonian Basin (Dombradi *et al.*, 2010); Iberia (Cloetingh *et al.*, 2002; De Vicente *et al.*, 2007; Fernandez-Lozano *et al.*, 2010); Central Asia (Burov *et al.*, 1993; Nikishin *et al.*, 1993; Burov & Molnar, 1998); Tibet/ Himalayan syntaxis belt (Burg & Podlachikov, 1999; Shin *et al.*, 2009); Central Australia (Stephenson & Lambeck, 1985; Lambeck, 1983); Arctic Canada (Stephenson *et al.*, 1990); Transcontinental Arch of North America (Ziegler *et al.*, 1995); South Caspian Basin (Guest *et al.*, 2007); Laramide Foreland (USA) (Tikoff & Maxson, 2001); Barents Sea (Ritzmann & Faleide, 2009) (map credit: NASA).

Figure 3. Relationship between wavelengths of lithospheric folding and thermo-tectonic age of lithosphere. The grey zones correspond to theoretical folding wavelengths derived for the upper crust, mantle and coupled whole mantle folding, based on an analytical model that accounts for strength variations as a function of thermo-mechanical age (Cloetingh *et al.*, 1999). Black squares are estimates for wavelengths of differential topography and thermo-tectonic ages inferred from observational studies. Numbers refer to sites listed in Table 1.

Figure 4. (a) Map showing major tectonic features of Tien Shan area of Central Asia , with location of folding profiles discussed in Burov *et al.* (1993), Burov & Molnar (1998) and Cloetingh *et al.* (1999). (b) Neotectonic movements along the profiles AD and CD (Burov *et al.*, 1993). Modified after Cloetingh *et al.* (1999). (c) Measured and calculated Bouguer gravity anomalies for lithosphere under horizontal compression along profile F across the Ferghana basin with corresponding topography profile (d) (after Burov & Molnar, 1998). Predicted topography and Moho geometry for Airy and best fitting folding scenario. During compression the amplitude of plate deflection changed significantly while the wavelength of folding stayed constant (Burov & Molnar, 1998). Red lines are Bouguer anomaly predicted by folding model and Moho depression, respectively. Solid grey line is Airy isostasy. Solid black line is Airy Moho. Triangles indicate observed Bouguer anomaly. Green is topography above sea level. FB: Ferghana Basin.

Figure 5. First order features of basin geometry of folded lithosphere of Iberia. (a) Topography displaying cylindrical patterns of alternating parallel trending highs and lows and corresponding gravity anomalies (red line) (see Cloetingh *et al.*, 2002). (b) Results from analogue modelling experiments, displaying pop-up structures underlying topographic highs, accompanied by folding of the Moho (Fernandez-Lozano *et al.*, 2010). Top panel shows final top view of the analogue model and location of the cross-section shown below. Arrow indicates the direction of the moving wall. λ is the wavelength of folding.

Figure 6. (a) Map of elevation of basement in the NW Alpine foreland with rift signature removed from the map of actual basement (from Bourgeois *et al.*, 2007). Dashed lines show axes of Late Neogene lithospheric folds, with a succession of anticlines and synclines: the Normandy-Vogelsberg anticline (NVA), Sologne-Franconian Basin Syncline (SFS) and the Burgundy-Swabian Jura Anticline (BSJA). LBR: London Brabant Ridge with thick crust. BF: Black Forest; FB Franconian Basin; E: Eifel; H: Hunsruck; S: Sauerland; T: Taunus; V: Vosges. (b) Cross sections of

top basement in the NW Alpine foreland with rift signatures removed (Bourgeois *et al.*, 2007; see for location Fig. 6a). Thin line: actual topography surface. Bold line: elevation of top basement as predicted in the absence of rifting. Fold signatures are similar with a wavelength λ of 270 km and amplitude A of 1500 m in the Franconian Basin (Cross section G-G') and in the upper Rhine Graben area (Cross section H-H'). In the Paris basin, development of folds with assumed same wavelength and amplitude (stippled bold line in cross section C-C'), implies subsidence of Sologne and uplift of Normandy.

Figure 7. (a) Schematic E-W cross section through the Pannonian-Carpathian basin system (Cloetingh *et al.*, 2008), displaying anomalous Late Neogene acceleration of the subsidence between isochrons 2 and 0 Ma in the centre of the Pannonian basin and the Focsani depression of the Southern Carpathian foredeep, flanked by uplifting highs. Wavelengths of the differential vertical motions are typically of the order of 250 km. **(b)** Interpreted seismic section in the central part of the Pannonian Basin (Sacchi *et al.*, 1999) showing evidence for late Neogene crustal scale folding at scales of 20-40 km. Note syn-rift grabens formed during the Miocene back-arc extension of the basin. **(c)** Results of analogue tectonic experiments carried out to examine the patterns of differential vertical motions in the northern part of the Pannonian Basin and surrounding areas (Dombradi *et al.*, 2010), demonstrating a-typical folding of variable amplitude with a wavelength equivalent to 200-250 km.

Figure 8. Schematic north-south trending cross-section illustrating the basic tectonic setting for northern Iran and the South Caspian Basin (Guest *et al.*, 2007), illustrating basin inversion of the central Iranian basin, pronounced differential topography at the transition of the Alborz mountains, with a topographic high of several km's and the South Caspian Basin with about 20 km of sedimentary infill.

Figure 9. Characteristic patterns of continental lithosphere deformation induced by folding of the lithosphere for with time increasing amounts of horizontal shortening for a 150 Ma old lithosphere with convergence rates of (a) 1.5 cm/yr and (b) 3 cm/yr. From left to right: material phase distribution (with blue for the crust, orange for the mantle lithosphere and red for the upper mantle below the lithosphere), temperature, effective viscosity, and topography.

Figure 10. Characteristic patterns of continental lithosphere deformation induced by folding of lithosphere with time. Increasing amounts of horizontal shortening for a 300 Ma old lithosphere with convergence rates of (a) 1.5 cm/yr and (b) 3 cm/yr. From left to right: phase distribution, temperature, effective viscosity, and topography.

Figure 11. (a) Characteristic subsidence patterns in centre of synclinal depression for thermo-mechanical age of 150 and 300 Ma, respectively. Shortening rate is 3 cm/year. **(b)** Characteristic stages in the evolution of a basin formed by lithospheric folding. Stage 1: Basin formation stage, acceleration of subsidence and uplift during folding. Stage 2: Steady-state: equilibrium between tectonic subsidence and sediment supply from eroding highs. Stage 3: Capture of folded basin; overall uplift and erosion. Wiggled waved blue line marks position of base level.

Figure 12. Illustration of the effect of erosion, which acts as a filter suppressing the short wavelengths. In this case, the wavelength and amplitude vary along the cross section at different stages of deformation due to a partial crust-mantle coupling and strain localization for a 400 Ma old lithosphere with weak quartz -dominated rheology. After 5 percent shortening (top), after 25 percent shortening (middle), after 25 percent shortening, strong zero order diffusional erosion (Avouac & Burov, 1996) tuned to keep mean elevation at the level of 3000 m. Erosion reduces the contribution of gravity-dependent terms (middle wavelength) and accelerates local deformation. Strong erosion,

insufficiently compensated by the tectonic deformation wipes out most of the topography (after Cloetingh *et al.*, 1999).

Figure 13. Thermal consequences of lithosphere folding, and associated basin evolution depicted in Fig. 11. The heat flow response is marked by an initial decrease in heat flow as a consequence of rapid sedimentation, followed by a progressive long-term heat flow increase after termination of shortening. Thermal conductivity of sediments is $2.5 \text{ W m}^{-1} \text{ C}^{-1}$. Surface radiogenic heat production H_s is $9.5 \times 10^{-10} \text{ W kg}^{-1}$. Concentration of radiogenic heat sources within sedimentary basin fill is homogeneous. Exponential decay of heat production is assumed for underlying crustal rocks. Note that the figure shows only 7 Ma of thermal evolution. The heat flow does not increase anymore after some tens of Ma and subsequently decreases at later stages.

Figure 14. Results of analogue tectonic experiments for lithospheric folding (after Sokoutis *et al.*, 2005) (Top): folding of uniform lithosphere. Cross section demonstrating pop-up structures in the upper crust above highs induced by lithospheric folding. λ_1 and λ_2 indicate first-order and second-order wavelengths, respectively. Imbrication of bivergent thrust wedges is responsible for the relatively high topography belt (Σ -belt) at the right-hand of the model and closure of prismatic basins with a rather undisturbed bottom. The progressive sinking of the prismatic basin allows transferring of the crustal material (i.e. cover sediments) that was initially situated at the surface of the model down to significant depth. Left-side (a) final top view of model after 34% of bulk shortening. Right side (b) cross-section located on the top view. Arrows indicate the direction of the moving wall. Note the similarity to Central Iberia (Fig. 5). (Bottom): Geometries predicted by analogue experiment, shortening a lithosphere with two contrasting blocks, bounded by suture. A deep synclinal depocentre develops over the suture zone, flanked by an anticline of smaller amplitude. Left-side (c) final top view of model after 26% of bulk shortening. Right side (d) cross-section located on the top

1
2
3 1802 view. λ_1 indicates a first-order wavelength of the folding. Note the striking similarity to the
4
5 1803 configuration of the South Caspian Basin/ Alborz Mountain system displayed in **Fig. 8**.
6
7 1804

9
10 1805 **Figure 15.** Numerical experiment on faulting developed in folded continental lithosphere of 150 Ma
11
12 1806 (Burov & Molnar, 1998). Note different wavelengths of the crustal and mantle part of the lithosphere
13
14 1807 and differences in spacing and character of faults that are conditioned by folding. In each layer, two
15
16 1808 fault spacings occur: a shorter one proportional to layer thickness (also controlled by the strength
17
18 1809 contrast between the stiff layer and embeddings), and the longer wavelength controlled by the
19
20 1810 wavelength of folding. The insert shows plastic (brittle) strain localization characterizing initialization
21
22 1811 of mantle faulting.
23
24 1812

25
26
27 1813 **Figure 16.** Comparison of main characteristics of basins developed on folded lithosphere (FLB with
28
29 1814 a shortening rate of 3 cm/yr) with other basin types, including foreland basins (FBr = retro-arc
30
31 1815 foreland basin; FBp = pro-arc foreland basin), intracratonic basins (ICB), extensional basins (EB)
32
33 1816 and pull-apart basins (PAB). Different panels give theoretical predictions. (Top): Basin subsidence
34
35 1817 history in basin centre. (Bottom): Thermal history. See text for explanation.
36
37 1818

TABLE 1

Wavelengths and ages of folded lithosphere displayed in Fig. 3 (see Fig. 2 for location)

	Area	Thermo-tectonic age (Ma)	Folding wavelength λ (km)	References
1	Tien Shan	175	200-250	Burov & Molnar, 1998; Burg <i>et al.</i> , 1994
2	Western Goby	175-400	300-360	Nikishin <i>et al.</i> , 1993; Burov <i>et al.</i> , 1993
3	Central Asia	370-430	50-70 (crust) 300-400 (litho-mantle)	Nikishin <i>et al.</i> , 1993; Burov <i>et al.</i> , 1993
4	Himalayan syntaxis belt	8-10	150	Burg & Podlachikov, 1999
5	Central Australia	500-900	550-650	Lambeck, 1983; Stephenson & Lambeck, 1985; Beekman <i>et al.</i> , 1997
6	Russian platform	400-600	500-600	Nikishin <i>et al.</i> , 1997
7	South Caspian Basin	125-155	350-450	Guest <i>et al.</i> , 2007
8	Eastern Black Sea	40-80	50-100 (crust) 100-150 (litho-mantle)	Cloetingh <i>et al.</i> , 2008
9	Western Black Sea	75-125	50-100 (crust) 100-200 (litho-mantle)	Cloetingh <i>et al.</i> , 2008
10	Pannonian Basin System	20	200-250	Horvath & Cloetingh, 1996; Matenco <i>et al.</i> , 2007 ; Dombradi <i>et al.</i> , 2010
11	NW European platform	180-230	270	Bourgeois <i>et al.</i> , 2007
12	Brittany	210-290	225-275	Bonnet <i>et al.</i> , 2000
13	Iberia	330-370	40-80 (crust) 125-275 (litho-mantle)	Cloetingh <i>et al.</i> , 2002
14	Barents Sea	215-245	550-650	Ritzmann & Faleide, 2009
15	Canadian Arctic	150-250	170-230	Stephenson <i>et al.</i> , 1990
16	Transcontinental Arch of North America	1000-1400	500-700	Ziegler <i>et al.</i> , 1995
17	Laramide foreland (USA)	175-225	190	Tikoff & Maxson, 2001

TABLE 2. Key characteristic features of different classes of sedimentary basins

Basin type	Basin shape	Dimensions	Subsidence characteristics	Thermal evolution	Faulting history
FLB	Symmetrical	Width: 50-600 km; depth: up to 20 km	Accelerated subsidence, time scales 1-10 Myr; simultaneously uplift at highs	No initial heating; heat flow increases in first 10's Myr with sediment deposition and burial, and subsequently decreases	Intensive deformation, accelerating trough basin formation phase
FB	Asymmetrical	Width: 50-250 km; depth: up to 10 km	Linear subsidence, interrupted by short-lived faster subsidence during thrusting phases; timescales 10-100 Myr; simultaneously uplift from thrust front and foreland bulge	No initial heating; heat flow increases with time with sediment deposition and burial	Faulting and thrusting limited to orogenic wedge; minor deformation through faulting in foreland
EB	Symmetrical (pure shear); asymmetrical (simple shear)	Width: 30-500 km; depth: up to 10 km	Rapid initial subsidence, followed by post-rift decay in subsidence; in case of multiple rifting repeated accelerated subsidence; time scales for post-rift subsidence of the order of 70 Myr; At rifting stage development takes place of asymmetrical rift shoulder topography, flanking the rift	Initial heating event, followed by decaying heat flow	Extensional faulting limited to basin formation phase; during post-rift absence of faulting
PAB	Symmetrical	Width: 20-50 km; depth: up to 10 km	Very rapid initial subsidence, followed by post-rift decay in subsidence, time scales: 1-10 Myr In early stage, sedimentation cannot keep up with subsidence	Initial heating event, followed by very rapid decay in heat flow	Extensional faulting limited to basin formation phase; during post-rift absence of faulting
ICB	Symmetrical	Width: 500-1000 km; depth: up to 10 km	Slow subsidence, punctuated by periods of faster subsidence, possibly related to large-scale tectonic events; Characteristic time scales: 500-800 Myr	In absence of understanding of basin formation mechanism no prediction for initial heat flow; Heat flow at later stage dominated by burial history	Very low level of faulting throughout tectonic history

Table A1

Notations and physical values common for all experiments (Turcotte and Schubert, 2002; Schubert *et al.*, 2001)

Parameter	Values and units	Definition
σ, τ	Pa, MPa	stress (complete, deviatoric)
P	Pa, MPa	pressure
u	m, mm	displacement vector
v	m s^{-1} , mm an^{-1}	velocity vector
$\dot{\epsilon}$	s^{-1}	strain rate
μ	10^{19} - 10^{25} Pa s	effective viscosity ($\tau/\dot{\epsilon}$)
T	$^{\circ}\text{C}$, K	temperature
h_c	40 km	Moho depth
h_l	100 - 250 km	thickness of lithosphere
D	100 - 200 km	plume diameter
ρ	3330 kg m^{-3}	density of mantle lithosphere
ρ_m	3340 kg m^{-3}	reference deep mantle density
ρ_p	$\rho_m + \Delta\rho_{ch} + \alpha\rho_m\Delta T$	plume density
$\Delta\rho_{ch}$	$0\text{-}25 \text{ kg m}^{-3}$	chemical density contrast
g	9.8 m s^{-2}	acceleration due to gravity
C_p	$10^3 \text{ J Kg}^{-1} \text{ }^{\circ}\text{C}^{-1}$	specific heat
ΔT	250°C	initial temperature contrast plume - background
α	$3 \times 10^{-5} \text{ }^{\circ}\text{C}^{-1}$	thermal expansion coefficient at 0 km depth
β	$8 \times 10^{-12} \text{ Pa}^{-1}$	isothermal compressibility

Table A2. Specific rheology and related thermal parameters. Compilation by Burov *et al.* (2001). ρ is density; Q , n , A are material-dependent parameters of ductile flow laws (Kirby & Kronenberg, 1987; Kohlstedt *et al.*, 1995). Other parameters from Turcotte & Schubert (2002).

Parameter	Value
All rocks	
λ , G Lamé elastic constants ($\lambda = G$)	30 GPa (above 250 km depth)
λ , G Lamé elastic constants ($\lambda = G$)	60 GPa (below 250 km depth)
ϕ friction angle (Mohr-Coulomb criterion)	30°
C_0 cohesion (Mohr-Coulomb criterion)	20 MPa
<u>Specific upper or weak (quartz) lower-crust properties</u>	
ρ (upper crust)	2700 kg m ⁻³
ρ (lower crust)	2900 kg m ⁻³
n	2.4
A	6.7 × 10 ⁻⁶ MPa ⁻ⁿ ·s ⁻¹
Q	1.56 × 10 ⁵ kJ·mol ⁻¹
<u>Specific strong lower crust properties (diabase or basalt)</u>	
ρ	2980 kg m ⁻³
n	3.4
A	2 × 10 ⁻⁴ MPa ⁻ⁿ ·s ⁻¹
Q	2.6 × 10 ⁵ kJ·mol ⁻¹
<u>Specific mantle properties (olivine)</u>	
ρ (lithosphere)	3330 kg m ⁻³
n	3
A	1 × 10 ⁴ MPa ⁻ⁿ ·s ⁻¹
Q	5.2 × 10 ⁵ kJ·mol ⁻¹
<u>Thermal model</u>	
Surface temperature (0 km depth)	0 °C
Temperature at the bottom of thermal lithosphere	1330 °C
Temperature at 660 km depth	2000 °C
Thermal conductivity of crust k	2.5 Wm ⁻¹ °C ⁻¹
Thermal conductivity of mantle k	3.5 Wm ⁻¹ °C ⁻¹
Thermal diffusivity of mantle χ	10 ⁻⁶ m ² ·s ⁻¹
Surface radiogenic heat production H_s	9.5×10 ⁻¹⁰ W kg ⁻¹
Radiogenic heat production decay depth h_r	10 km
Thermo-tectonic age of the lithosphere a	60 (young) to 1000 Ma (old)

Figure 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 2

Figure 3

Figure 4

Figure 5a

Figure 5b

Figure 6a

Figure 6b

Figure 7a

Figure 7b

Figure 8

Figure 9

a) $t=150$ Ma, $v=1.5$ cm/yr

b) $t=150$ Ma, $v=3.0$ cm/yr

Figure 10

a) $t=300\text{ Ma}$, $v=1.5\text{ cm/yr}$

b) $t=300\text{ Ma}$, $v=3.0\text{ cm/yr}$

Figure 11

Figure 12

Figure 13

Figure 14

Figure 15

Figure 16