Either taking it easy or feeling too tired: old Cory's Shearwaters display reduced activity levels while at sea

Paulo Catry, José Pedro Granadeiro, Jaime Ramos, Richard A. Phillips, Paulo Oliveira

► To cite this version:

Paulo Catry, José Pedro Granadeiro, Jaime Ramos, Richard A. Phillips, Paulo Oliveira. Either taking it easy or feeling too tired: old Cory's Shearwaters display reduced activity levels while at sea. *Journal für Ornithologie = Journal of Ornithology*, 2010, 152 (3), pp.549-555. 10.1007/s10336-010-0616-7 . hal-00646464

HAL Id: hal-00646464

<https://hal.science/hal-00646464>

Submitted on 30 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Either taking it easy or feeling too tired: old Cory's Shearwaters display reduced activity levels while at sea.

Paulo Catry^{1,2*}, José Pedro Granadeiro², Jaime Ramos³, Richard A. Phillips⁴, Paulo Oliveira⁵

¹ Eco-Ethology Research Unit & Centro de Biociências, ISPA, Rua Jardim do Tabaco 34, 1149-041 Lisboa, Portugal.

² – Centro de Biologia Ambiental, Museu Nacional de História Natural, Rua da Escola Politécnica, Lisboa, Portugal

³ - IMAR-Institute of Marine Research, Department of Zoology, University of Coimbra, Portugal

⁴ - British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge CB3 0ET, UK

⁵ – Serviço do Parque Natural da Madeira, Secretaria Regional do Ambiente e dos Recursos Naturais, Funchal, Portugal.

* Correspondence: Paulo Catry. Tel: (351) 96 2445005. Email: paulo.catry@gmail.com

Running title: Activity levels in old Shearwaters

Abstract

It has long been known that birds change their behaviour, reproductive performance and survival as they mature, including in the first few years after recruitment into the breeding population. However, and contrasting with the description of patterns of actuarial and reproductive senescence in latter years, there are surprisingly few studies documenting changes in behaviour in old individuals. Such studies are important, as birds provide particularly interesting models for studying the biology of senescence. It has been suggested that unlike mammals, birds may remain physically fit until an advanced age, yet this has limited empirical support. In this paper, we used activity (immersion) loggers to show that old (> 26 years) Cory's Shearwaters *Calonectris diomedea* are less active when foraging at sea, spend more time resting on the water and have a smaller number of take-offs and landings during darkness, when compared to experienced mid-aged individuals (13-20 years old). Old individuals also tended to have reduced immune response against an experimental challenge using phytohaemagglutinin. These results are in line with observed reductions in activity levels with age in a wide range of non-avian taxa, and may suggest that old seabirds are physically less fit than younger individuals. Alternatively, old birds might simply be more experienced and their reduction in activity might reflect a strategic regulation of investment in different activities. Our study illustrates the potential for gaining insights into avian aging patterns and processes by looking into the behaviour of model organisms. We therefore encourage more research focusing on behavioural parameters that may reflect variations in physical condition or strategic choices, during both the breeding and non-breeding seasons.

Zusammenfassung

Nimms' leicht oder Ermüdung: Alte Gelbschnabel-Sturmtaucher zeigen reduzierte Aktivität auf See.

Es ist bereits seit längerem bekannt, dass sich das Verhalten, der Bruterfolg und die Überlebensrate von Vögeln mit zunehmendem Alter verändert. Trotz der Beschreibung solcher Muster von Sterblichkeits- und reproduktiver Vergreisung in späteren Jahren gibt es erstaunlich wenige Studien, die eine derartige Veränderung im Verhalten alter Individuen dokumentieren. Solche Studien sind wichtig, weil Vögel ein besonders interessantes Modellsystem für die Untersuchung der Biologie von Vergreisung darstellen. Anders als bei Säugetieren wurde vorgeschlagen, dass Vögel bis ins hohe Alter physisch gesund bleiben. Dies wird jedoch nur im begrenzten Maße von empirischen Daten unterstützt. In der vorliegenden Studie verwendeten wir Aktivitätslogger, um zu zeigen, dass 26-jährige Gelbschnabel-Sturmtaucher *Calonectris diomedea* weniger aktiv sind, während sie auf See furagieren. Sie haben kürzere Ruhepausen auf dem Wasser und eine geringere Anzahl Starts und Landungen während der Dunkelheit, im Vergleich mit erfahrenen mittel alten Individuen (13-20 Jahre alt). Alte Individuen neigten auch zu einer reduzierten Immunreaktion auf eine experimentelle Injektion von Phytohaemagglutinin. Diese Ergebnisse stimmen im Wesentlichen überein mit der beobachteten Abnahme von Aktivität mit zunehmendem Alter in einer großen Anzahl anderer Tiergruppen. Es kann sein, dass alte Seevögel physisch weniger fit sind als jüngere. Alternativ könnten ältere Vögel auch erfahrener sein, und die Aktivitätsreduktion könnte eine strategische Regulierung der Allokation von Investment in unterschiedliche Aktivitäten sein. Unsere Untersuchung zeigt, dass detaillierte Studien des Verhaltens von Modellorganismen das Potential haben, Muster und Alterungsprozesse bei Vögeln zu erklären. Wir regen deshalb mehr Verhaltensstudien an, die die Variation in der physischer Kondition oder verschiedene strategischen Entscheidungen reflektieren, und dies sowohl in wie außerhalb der Brutsaison.

81 **Introduction**

82 In recent decades, many studies have focused on age-related changes in the behaviour
83 and fitness of birds. Some general trends are now relatively well-established. For
84 example, in the first months or years of life, birds apparently improve their foraging
85 efficiency (Wunderle 1991), dominance-mediated competitive ability (Sergio et al.
86 2009), migratory performance (Thorup et al. 2003), and breeding success and survival
87 (Newton 1989). Such changes are apparent in cross-sectional studies and result not only
88 from the selective appearance and disappearance of phenotypes, but also, as
89 longitudinal analyses indicate, from the acquisition of relevant experience (Forslund and
90 Pärt 1995).

91 Changes in behaviour may also occur at a more advanced age and could result
92 from strategic choices, as predicted by classic life-history theory. This includes, for
93 example, that in their last breeding attempt, individuals should raise their reproductive
94 effort in order not to waste unused resources (Stearns 1992). However, recent
95 theoretical research suggests that increased parental investment may not be adaptive
96 before the terminal breeding attempt, and that instead, a reduction in reproductive effort
97 should be typical response to an advancing age (McNamara et al. 2009). Furthermore, in
98 the last breeding attempt, it is possible that birds are already somewhat debilitated by a
99 rapidly advancing terminal disease (Ricklefs 2000, 2008, Coulson and Fairweather
100 2001), which would hinder the detection of an increased terminal effort. In fact, only a
101 handful of experimental studies have found evidence for the terminal investment
102 hypothesis (e.g. Hanssen 2006, Velando et al. 2006).

103 Studies to date have either documented actuarial senescence (Ricklefs 2000), or,
104 if they dealt with loss of function in old individuals, have mostly focused on
105 reproduction (e.g. Bennett and Owens 2002, Jones et al. 2008). This reflects the desire

to understand age-specific reproductive performance in relation to life-history theory, and also the relative ease with which relevant parameters can be measured. However, reproductive performance may be affected by a number of external factors, such as the quality of the nesting partner or changes in the environment, which confound interpretation (Nisbet 2001). Therefore, measuring other aspects of individual state and performance may contribute to a better understanding of the mechanisms of aging, and hence of the biology of senescence. Although there has been recent research focusing on hormonal, physiological and cellular parameters (Hausmann et al. 2005, Vleck et al. 2007), surprisingly few studies have explored the effects of aging of wild animals on traits such as foraging behaviour and efficiency (but see Catry et al. 2006, MacNulty et al. 2009, Lecomte et al. 2010). Such traits are highly relevant because they influence the rate of resource acquisition, and hence have major implications for fitness correlates that include reproductive output and survival.

Declines in activity levels of old individuals have been described for taxa ranging from nematodes and insects, to rodents, dogs and primates (e.g. Emborg et al. 1998, Fernández et al. 1999, Siwak et al. 2002, see Sallis 2000 and Ingram 2000 for reviews). Such declines are more obvious for demanding actions that require higher energetic expenditure or greater motor coordination, and, at least in mammals, seem to have a biological basis, resulting from impaired motor and neurobiological function, rather than a simple adaptive reduction in motivation or as an effect of social influences (Emborg et al. 1998, Ingram 2000, Carter et al. 2002, Némoy-Bertholet and Aujard 2003). Few studies of wild birds have attempted to measure activity levels in relation to aging, but very old male Grey-headed Albatrosses *Thalassarche chrysostoma* were reported to display reduced foraging efficiency when compared to mid-aged individuals, which suggests a deterioration in physical performance (Catry et al. 2006). Furthermore,

131 in Wandering Albatrosses *Diomedea exulans*, differences were found in the at-sea
132 activity budgets and distribution of mid-age and old male individuals, and old birds had
133 higher levels of the stress hormone corticosterone after a foraging trip (Lecomte et al.
134 2010). These results contrasts with recent suggestions that birds remain physically fit
135 into old age (Ricklefs 2000, 2008). Clearly this is an interesting question that deserves
136 further attention.

137 New data logger systems are revolutionising our ability to continuously monitor
138 bird behaviour, including in species such as seabirds whose pelagic lifestyles were
139 formerly largely unknown (Ropert-Coudert and Wilson 2005). Seabirds are particularly
140 useful models for studying age-related changes in ecology and behaviour, given their
141 unusual longevity and the ease with which they can be marked (ringed), such that many
142 individuals are of known age in populations subject to long-term study (Wooller et al.
143 1992). Cory's Shearwaters *Calonectris diomedea* are medium-sized seabirds that
144 typically forage for fish and squid in the upper levels of the pelagic domain. While one
145 bird incubates, the partner engages in long foraging trips that normally last for several
146 days and can take the off-duty individual to hundreds of kilometres from the nesting
147 site. Here we report a study on the activity levels and water mass selection of foraging
148 Cory's Shearwaters of known age.

149 We were particularly interested in exploring the possibility that old individuals
150 might suffer from some form of physical senescence effects that could affect their at-sea
151 distribution or foraging behaviour. If so, we would make the following predictions: (1)
152 old individuals should spend more time resting on the water during a foraging trip
153 (Passos et al. 2010), and (2) old birds should perform a smaller number of take-offs and
154 landings, as these are known to be energy-demanding activities in medium to large
155 sized petrels (Weimerskirch et al. 2000, Shaffer et al 2001). If the first two predictions

are confirmed, we would also predict as a corollary that (3) old individuals should take longer to recover body mass lost in incubation and hence make longer foraging trips. It is important to note from the outset that similar patterns (predictions 1 and 2) could be observed if old birds were simply more efficient, allowing them more discretionary resting time than mid-aged individuals. Hence, in this respect our study is exploratory, as to attribute any detected changes to senescence requires more detailed, longitudinal analyses. Our aim is therefore primarily to identify variables that deserve scrutiny in future research.

Methods

This study took place at Selvagem Grande, Savage islands (30°09'N, 15°52'W), where a large colony (currently ca 30,000 pairs) of Cory's Shearwaters has been the object of long-term research (Mougin et al. 2000, Granadeiro et al. 2006). Cory's Shearwaters nest in burrows and incubate their single egg for a period of ca 54 days (Thibault et al. 1997). In some parts of the colony, chicks and breeding adults have been marked with monel rings on the leg since 1977. In June 2007, during early incubation, we searched those sectors for individuals ringed as chicks, and therefore of known age. To increase sample sizes, we also selected birds that had been ringed as adults in 1977-87 and hence were of considerably advanced age. On average, Cory's Shearwaters start nesting at 8.9 ± 1.7 years and fewer than 1% of the individuals do so before the age of 6 (Mougin et al. 2000), hence these birds would have been at least 26 years old by the time of our study. Based on previous studies of senescence in this population that reported declines in breeding success for birds older than 25 years (Mougin 2002), we defined two groups of ringed birds, as follows: "Old" individuals were those ringed as chicks (40%) or adults (60%) and, respectively, known or assumed to be at least 26 years old, and "Mid-

aged” individuals were those aged 13-20 years (the majority were 13-16 years old).

These groups were defined *a priori* (i.e. before subsequent study).

In the last week of June 2007, we deployed activity and temperature loggers on incubating birds. The loggers (Mk 7; British Antarctic Survey, Cambridge) weigh only 3.5g and were attached to the monel ring using a plastic, weather-resistant cable-tie. These loggers record a single sea-surface temperature (SST) value after 20 min. of continuous immersion, and will only record a new value after a subsequent dry period, followed by immersion for a further 20 min. They also record the timings of all changes of state (from wet to dry, and *vice versa*) of 6 s or more, allowing the reconstruction of detailed activity patterns. These loggers are not known to have any significant deleterious effects (Igual et al. 2005). Loggers were recovered 3 to 4 weeks later, by which time all individuals had performed at least one foraging trip. The recovery rate was 96%.

The approximate start and end date of each foraging trip (which typically lasted several days, see results) could be estimated from the first and last wet event between incubation stints (Edwards et al. 2007). However, after leaving its nest, a bird could potentially fly for several hours before landing on the water, which would be indistinguishable from time spent at the nest, as would the last flight at the end of the foraging trip. Hence, when estimating number of hours flying or resting on the water per day, we excluded information on the first and the last day of the foraging trip. During a foraging trip, birds spent most of the night on the water. On rare occasions, loggers recorded a long continuous dry period during the night, which could result from the bird attending the colony, rather than represent time in flight. To avoid the disproportionate influence of these rare events, we used median values, rather than means, when calculating summary statistics for individuals.

206 Shearwater at-sea nocturnal activity may be influenced by variation in ambient
207 light levels related to the moon phase (Yamamoto et al. 2008). To assess if this factor
208 might confound our results, we compared the likely light levels experienced by Mid-
209 aged and Old individuals by the use of a moon score. A night with full moon was given
210 a score 15, while the new moon had a score 0, with intermediate dates having
211 intermediate scores, between 1 and 14. We then calculated the median score
212 experienced by each individual during the nights it was out at sea and finally compared
213 the two age-groups. This analysis is unlikely to be further confounded by cloud-cover,
214 as only thin clouds were occasionally present during the study. Furthermore, departure
215 date had no influence on the behavioural variables measured and when we repeated the
216 analyses statistically controlling for departure date, the patterns reported in this paper
217 remained unchanged.

218 To assess potential differences in selection of foraging areas, we calculated the
219 mean logger SST value for each bird during darkness (to avoid direct effects of sunlight
220 on the loggers) on each day, took the mean of those values during each foraging trip,
221 and then compared this between between Old and Mid-aged groups.

222 In our analyses, we deliberately excluded birds that might be affected by an
223 advanced terminal illness (Catry et al. 2006). Survival rates of Cory's Shearwaters are
224 typically high and only a few individuals were not resighted in the following year.
225 These birds may have died in the intervening period, or be deferring breeding. This
226 small sample was insufficient for exploring patterns relating to terminal investment, but
227 could introduce considerable noise in the data sets. Hence, analyses presented here
228 focus exclusively on birds that were resighted in the season following our field
229 measurements.

Phytohaemagglutinin skin-test. Mougin (2002) has shown that Cory's

Shearwaters aged above 25 years old displayed a decline in reproductive function consistent with senescence. To further assess whether our group of Old birds showed physiological evidence of senescence, we evaluated their immune status by means of the phytohaemagglutinin (PHA) test (Smits et al. 1999, Haussmann et al. 2005, Kennedy and Nager 2006). This test consists of injecting a mitogen (PHA) and measuring the swelling at the injection site 24 hours later. The swelling results from a perivascular accumulation of T lymphocytes followed by macrophage infiltration (Smits et al. 1999), and its measurement is thought to provide an indication of the T-cell mediated immune response.

We attempted to use the same procedure as Forero et al. (2006), which involved the injection of a solution of PHA into the foot-web and measurement of subsequent swelling. However, we were unsuccessful at web injections and instead injected 0.05 ml of 2 mg ml⁻¹ PHA (Sigma) in phosphate-buffered saline (PBS) at a marked point on the tarsus. Three measurements of tarsus thickness over the injection point were made before and 24 hours after the injection using an electronic Mitutoyo micrometer (± 0.01 mm). From these, we calculated three different PHA response measures to assess the repeatability of our estimates, which was high ($r = 0.87$, $F_{64,194} = 20.1$, $P < 0.001$). We then averaged the three measurements before and the three measurements after and calculated a final PHA response value based on this difference.

Study birds were sexed using a discriminant function based on bill measurements (Granadeiro 1993). Individuals with a score > 0.1 were classified as males and those with a score < -0.1 as females. Birds with intermediate values were sexed if paired with an individual of known sex, or if their distinctive vocalizations

(Thibault et al. 1997) were heard during handling. By using these criteria, and cross-validation with nesting partners, sex classification accuracy was >99% (unpubl. data).

Variables related to the number of take-offs and landings were log-transformed to meet the requirements of normality and homogeneity of variances. Means are presented $\pm$ SD.

Results

Sex had no influence on foraging trip variables reported below and hence this factor was excluded from all models. During foraging trips, Mid-aged individuals took off and landed more often than did Old individuals (Fig. 1). This difference was only significant during darkness (Table 1). The proportion of time spent resting on the water was higher in Old individuals, again with the most pronounced differences observed during darkness (Table 1). There were no differences between age groups in foraging trip duration, mean daily SST values or absolute minimum SST values attained during the trip (Table 1). Trip duration correlated negatively with minimum SST values ($r = -0.53$, $n = 33$, $P = 0.001$), indicating that only on longer trips did birds reach the cold waters of the Canaries upwelling, near the African coast (own unpubl. data). This illustrates the usefulness of SST as an indicator of foraging region.

Sex had no influence on PHA response and hence was excluded from the model. Cory's Shearwaters from the Old group had a weaker PHA response (0.663 ± 0.423 mm, $N = 33$) than Mid-aged individuals (0.853 ± 0.451 mm, $N = 32$; ANOVA $F_{1,64} = 3.06$, $P = 0.085$). Note that the difference is significant if we adopt a one-tailed test, in accordance with our directional prediction that senescent animals should show a depressed immune response.

Discussion

The concept that, unlike mammals such as humans, birds may remain physically fit until old age (Ricklefs 2000, 2008) has seldom been tested (but see Coulson and Fairweather 2001, Catry et al. 2006), despite its importance in the context of the biology of senescence. The results from the present work, and the ones from a recent study with albatrosses (Lecomte et al. 2010), suggest that old, free-living seabirds display a reduction in activity levels that parallels patterns reported for a diversity of other taxa (e.g. Emborg et al. 1998, Fernández et al. 1999, Ingram 2000, Siwak et al. 2002, Némoy-Bertholet and Aujard 2003). The causes for such reduction in activity levels are so far unknown, but circumstantial evidence suggests that senescence due to progressive wear and tear (somatic deterioration) of aging bodies is a strong candidate explanation.

For medium to large petrels that rely on wind-assisted dynamic soaring for travel, flight is often considered to be a relatively economical activity, only slightly more costly than resting: instead, most of the variation in energy expenditure during foraging trips of albatrosses is explained by the number of take-offs, which are energetically expensive (Bevan et al. 1995, Weimerskirch et al. 2000, Shaffer et al 2001). However, those studies involved birds that proceed by flap-gliding in the windy southern ocean and it is likely that flight is much more costly for Cory's Shearwaters, which are frequently seen using flapping flight.

The discovery that Old Cory's Shearwaters differed from Mid-aged individuals mostly in the number of take-offs and landings fits well with evidence from other taxa that declines in spontaneous activity are more pronounced in the frequency of demanding physical tasks. For example, in Grey Mouse Lemurs *Microcebus murinus*, differences between young and old adults were clearest in activities such as difficult

jumps, where an intense physical effort was required (Némoz-Bertholet and Aujard 2003).

Petrels are known to both travel and forage during darkness as well as daylight, although they tend to be more active during the latter (Catry et al. 2004, Phalan et al. 2007, this study). The disproportionate reduction of activity among old Shearwaters during darkness might reflect a strategy to avoid demanding activities when they are less likely to yield a benefit, given that foraging at night seems to be less profitable for visual predators such as seabirds (Catry et al. 2004). An experimental study with Cory's Shearwaters showed that handicapped birds (carrying extra mass) also reduced their activity levels (less time spent flying) during darkness (Passos et al. 2010). It seems less likely that Old birds sleep more during darkness, as sleep is more often interrupted in aged animals (e.g. Hornung et al. 2005), including birds (Paredes et al. 2006). That sea surface temperatures in feeding areas and foraging trip duration were similar in Mid-aged and Old individuals, suggests that differences in activity levels between age groups cannot be explained by disparate habitat preferences.

The decline in the PHA response by aged Cory's Shearwaters is consistent with results from recent studies indicating that immunosenescence may be common in both short and long-lived bird species (Hausmann et al. 2005, Vleck et al. 2007). It also suggests that our Old birds in our study had reached an age where senescence effects might be apparent (see also Mougin 2002).

There are alternative explanations to deterioration in physical function for the observed declines in activity. Old individuals were not forced to prolong their foraging trips as a result of their lower activity levels, and it is therefore possible that they were more efficient at foraging as a result of accumulated experience. In this context, it should be mentioned that the handicapped (carrying extra mass) Cory's shearwaters

studied by Passos et al. (2010), and alluded to above, did make longer foraging trips, which may suggest that this is a relevant variable when studying performance during foraging movements.

Both the activity and the immune response trends presented in this paper result from cross-sectional analyses. Cross-sectional studies have limitations, because patterns may result from cohort heterogeneity or even individual heterogeneity in longevity and other parameters of interest (Nisbet 2001). However, as far as we are aware, no study has demonstrated that differential mortality of phenotypes is responsible for apparent senescence in breeding performance in long-lived birds. Hence, despite this being a possibility, it is relatively unlikely to have influenced our results. Nevertheless, the hypotheses tested in this study should be further evaluated using longitudinal data sets. Neither can we discard the idea that old individuals might be strategically reducing their parental effort (McNamara *et al.* 2009). This explanation could be ruled out in our case by determining if differences in individual activity patterns persist during the nonbreeding season.

In conclusion, our study demonstrates the potential for gathering insights into avian senescence patterns and processes by looking beyond the classical variables of adult mortality and breeding output. We would therefore encourage more research focusing on parameters that reflect physical performance and behaviour, both in the breeding and non-breeding seasons.

Acknowledgements

Parque Natural da Madeira, and particularly Dília Menezes provided permissions to carry out the work and, together with the wardens at the Nature Reserve where this study took place, gave important logistical support. Hany Alonso, Rafael Matias, Maria

Dias, Miguel Lecoq, Rui Rebelo, Filipe Moniz, Ana Leal, Ricardo Martins and others helped with fieldwork. The manuscript benefited from the comments of two anonymous referees. This study is an output from a project on the ecology and senescence in Cory's Shearwaters (PDCT/MAR/58778/2004) supported by Fundação para a Ciência e a Tecnologia (FCT - Portugal) and FEDER. P. Catry benefited from postdoctoral fellowships from FCT (BPD/11631/02 and SFRH/BPD/30031/2006) and further support was received through Programa Plurianual (UI&D 331/94).

References

- Bennett PM, Owens IPF (2002) Evolutionary Ecology of Birds. Oxford University Press, Oxford
- Bevan RM, Butler PJ, Woakes AJ, Prince PA (1995) The energy expenditure of free-ranging black-browed albatrosses. *Phil Trans R Soc Lond B* 350:119-131
- Carter CS, Sonntag WE, Onder WE, Pahor M (2002) Physical performance and longevity in aged rats. *J Gerontol* 57:B193-B197
- Catry P, Phillips RA, Phalan B, Silk JRD, Croxall JP (2004) Foraging strategies of grey-headed albatrosses *Thalassarche chrysostoma*: integration of movements, activity and feeding events. *Mar Ecol Progr Ser* 280:261-273
- Catry P, Phillips RA, Phalan B, Croxall JP (2006) Senescence effects in an extremely long-lived bird: the grey-headed albatross *Thalassarche chrysostoma*. *Proc R Soc B* 273:1625-1630
- Coulson JC, Fairweather JA (2001) Reduced reproductive performance prior to death in the Black-legged Kittiwake: senescence or terminal illness? *J Avian Biol* 32:146-152

377 Edwards AM, Phillips RA, Watkins NW, Freeman MP, Murphy EJ, Afanasyev V,
 378 Buldyrev SV, da Luz MGE, Raposo EP, Stanley HE, Viswanathan GM (2007)
 379 Overturning evidence of Levy flight searches by wandering albatrosses,
 380 bumblebees and deer. *Nature* 449:1044-1049
 381 Emborg ME, Ma SY, Mufson EJ, Levey AI, Taylor MD, Brown WD, Holden JE
 382 Kordower JH (1998) Age-related declines in nigral neuronal function correlate
 383 with motor impairments in rhesus monkeys. *J Comp Neurol* 401:253-265
 384 Fernández JR, Grant MD, Tulli NM, Karkowski LM, McClearn GE (1999) Differences
 385 in locomotor activity across the lifespan of *Drosophila melanogaster*. *Exp*
 386 *Gerontol* 34:621-631
 387 Forero MG, González-Solis J, Igual JM, Hobson KA, Ruíz X, Viscor G (2006)
 388 Ecological and physiological variance in T-cell mediated immune response in
 389 Cory's Shearwaters. *Condor* 108:865-876
 390 Forslund P, Pärt T (1995) Age and reproduction in birds – hypotheses and tests. *Trends*
 391 *Ecol Evol* 10:374-378
 392 Granadeiro JP (1993) Variation in measurements of Cory's Shearwater between
 393 populations and sexing by discriminant analysis. *Ring Migr* 14:103–112
 394 Granadeiro JP, Dias MP, Rebelo R, Santos CD, Catry P (2006) Numbers and population
 395 trends of Cory's Shearwater *Calonectris diomedea* at Selvagem Grande,
 396 Northeast Atlantic. *Waterbirds* 29:56–60
 397 Hanssen SA (2006) Cost of an immune challenge and terminal investment in a long-
 398 lived bird. *Ecology* 87:2440-2446
 399 Haussmann MF, Winkler DW, Huntington CE, Vleck D, Sanneman CE, Hanley D,
 400 Vleck CM (2005) Cell-mediated immunosenescence in birds. *Oecologia*
 401 145:270-275

402 Hornung OP, Danker-Hopfe H, Heuser I (2005) Age-related changes in sleep and
403 memory: commonalities and interrelationships. *Exp Gerontol* 40:279-285

404 Igual JM, Forero MG, Tavecchia G, González-Solis J, Martínez-Abraín A, Hobson KA,
405 Ruiz X, Oro D (2005) Short-term effects of data-loggers on Cory's Shearwaters
406 *Calonectris diomedea*. *Mar Biol* 146:619-624

407 Ingram DK (2000) Age-related decline in physical activity: generalization to
408 nonhumans. *Med Sci Sports Exerc* 32:1623-1629

409 Jones OR and 34 co-authors (2008) Senescence rates are determined by ranking on the
410 fast-slow life-history continuum. *Ecol Lett* 11:664-673

411 Kennedy MW, Nager MG (2006) The perils and prospects of using
412 phytohaemagglutinin in evolutionary ecology. *Trends Ecol Evol* 21:653-655

413 Lecomte VJ, Sorci G, Cornet S, Jaeger A, Faivre B, Arnoux E, Gaillard M, Trouvé C,
414 Besson D, Chastel O, Weimerskirch H (2010) Patterns of aging in the long-lived
415 wandering albatross. *Proc Nat Acad Sci* 107:6370-6375

416 MacNulty DR, Smith DW, Vucetich JA, Mech LD, Stahler DR, Packer C (2009)
417 Predatory senescence in ageing wolves. *Ecol Lett* 12:1347-1356.

418 McNamara JM, Houston AI, Barta Z, Scheuerlein A, Fromhage L (2009) Deterioration,
419 death and the evolution of reproductive restraint in late life. *Proc R Soc B*
420 276:4061-4066

421 Mougin J-L (2002) Influence de la sénilité sur la réussite de la reproduction chez le
422 puffin cendré *Calonectris diomedea*. *Bol Mus Mun Funchal* 53 :27-34

423 Mougin J-L, Jouanin C, Roux F (2000) Démographie du puffin cendré *Calonectris*
424 *diomedea* de Selvagem Grande. *Rev Ecol (Terre Vie)* 55:275-290

425 Némoy-Bertholet F, Aujard F (2003) Physical activity and balance performance as a
 426 function of age in a prosimian primate (*Microcebus murinus*). *Exp Gerontol*
 427 38:407-414
 428 Newton I (ed.) (1989) Lifetime reproduction in birds. London, Academic Press.
 429 Nisbet ICT (2001) Detecting and measuring senescence in wild birds: experience with
 430 long-lived seabirds. *Exp Gerontol* 36:833-843
 431 Paredes SD, Terrón MP, Cubero J, Valero V, Barriga C, Reiter RJ, Rodríguez AB
 432 (2006) Comparative study of the activity/rest rhythms in young and old ringdove
 433 (*Streptopelia risoria*): correlation with serum levels of melatonin and serotonin.
 434 *Chronobiol Internat* 23:779-793
 435 Passos C, Navarro J, Giudici A, González-Solís J (2010) Effects of extra mass on the
 436 pelagic behavior of a seabird. *Auk* 127:100-107
 437 Phalan B, Phillips RA, Silk JRD, Afanasyev V, Fukuda A, Fox J, Catry P, Higuchi H,
 438 Croxall JP (2007) Foraging behaviour of four albatross species by night and day.
 439 *Mar Ecol Progr Ser* 340:271-286
 440 Ricklefs RE (2000) Intrinsic aging-related mortality in birds. *J Avian Biol* 31:103-111
 441 Ricklefs RE (2008) The evolution of senescence from a comparative perspective. *Funct*
 442 *Ecol* 22: 379-392
 443 Ropert-Coudert Y, Wilson RP (2005) Trends and perspectives in animal-attached
 444 remote sensing. *Front Ecol Environ*. 3:437-444
 445 Sallis JF (2000) Age-related decline in physical activity: a synthesis of human and
 446 animal studies. *Med Sci Sports Exerc* 32:1598-1600
 447 Sergio F, Blas J, Hiraldo F (2009) Predictors of floater status in a long-lived bird: a
 448 cross-sectional and longitudinal test of hypotheses. *J Anim Ecol* 78:109-118

449 Shaffer SA, Costa DP, Weimerskirch H (2001) Behavioural factors affecting foraging
 450 effort of breeding wandering albatross. *J Anim Ecol* 70:864-874
 451 Siwak CT, Murphey HL, Muggenburg BA, Milgram NW (2002) Age-dependent decline
 452 in locomotor activity in dogs is environment specific. *Physiol Behav* 75:65-70
 453 Smits JE, Bortolotti GR, Tella JL (1999) Simplifying the phytohaemagglutinin skin-
 454 testing technique in studies of avian immunocompetence. *Funct Ecol* 13:567-
 455 572
 456 Stearns SC (1992) The evolution of life histories. Oxford University Press, Oxford.
 457 Thibault J-C, Bretagnolle V, Rabouam C (1997) *Calonectris diomedea* Cory's
 458 Shearwater. *BWP Update* 1:75-98
 459 Thorup K, Alerstam T, Hake M, Kjellén M (2003) Bird orientation: compensation for
 460 wind drift in migrating raptors is age dependent. *Proc R Soc Lond B (Suppl.)*
 461 270:S8-S11
 462 Velando A, Drummond H, Torres R (2006) Senescent birds redouble reproductive effort
 463 when ill: confirmation of the terminal investment hypothesis. *Proc R Soc B*
 464 273:1443-1448
 465 Vleck CM, Haussmann ME, Vleck D (2007) Avian senescence: underlying
 466 mechanisms. *J Orn* 148: S611-S624
 467 Weimerskirch H, Guionnet T, Martin J, Shaffer SA, Costa DP (2000) Fast and fuel
 468 efficient? Optimal use of wind by flying albatrosses. *Proc R Soc Lond B*
 469 267:1869-1874
 470 Wooller RD, Bradley JS, Croxall JP (1992) Long-term population studies of seabirds.
 471 *Trends Ecol Evol* 7:111-114
 472 Wunderle JM (1991) Age-specific foraging proficiency in birds. *Curr Ornithol* 8:273-
 473 324

474 Yamamoto T, Takahashi A, Yoda K, Katsumata N, Watanabe S, Sato K, Trathan PN
475 (2008) The lunar cycle affects at-sea behaviour in a pelagic seabird, the Streaked
476 Shearwater *Calonectris leucomelas*. Anim Behav 76:1647–1652.
477
478
479

480 Table 1. Characteristics of foraging trips carried out by Mid-aged and Old Cory's
 481 Shearwaters during incubation. Number of take-offs and landings and mean sea surface
 482 temperature (SST) are means of mean daily values per individual. Proportions of time
 483 resting are means of medians of daily values per individual. See text for details.
 484

	Mid-aged birds (N = 17)	Old birds (N = 18)	Statistical comparison
Wet-dry changes during darkness	30.8 ± 24.0	13.4 ± 7.2	F _{1,34} = 12.6 P < 0.001
Wet-dry changes in daylight	100.8 ± 32.7	84.7 ± 24.5	F _{1,34} = 0.84 P = 0.34
Proportion time resting in water	0.46 ± 0.13	0.55 ± 0.13	F _{1,34} = 3.67 P = 0.06
Resting in water during darkness	0.72 ± 0.18	0.83 ± 0.20	F _{1,34} = 3.96 P = 0.05
Resting in water during daylight	0.38 ± 0.14	0.44 ± 0.12	F _{1,34} = 1.89 P = 0.18
Mean SST (°C)	18.65 ± 1.10	18.59 ± 0.62	F _{1,34} = 0.45 P = 0.83
Minimum SST (°C)	17.12 ± 1.25	17.26 ± 1.16	F _{1,34} = 0.14 P = 0.71
Foraging trip duration (days)	9.23 ± 2.69	8.52 ± 1.93	F _{1,34} = 0.76 P = 0.39
Light levels (median moon score)	7.5	9.0	M-W U = 100.5 P = 0.20

485

Fig. 1. Box-plots depicting the number of take-offs and landings (per day or night) of Old (N = 18) and Mid-aged (N=17) Cory's shearwaters. Differences were tested on log-transformed data (see Table 1 for statistics).