Differential Sensitivity To Adrenergic Stimulation Underlies The Sexual Dimorphism In The Development Of Diabetes Caused By Irs-2 Deficiency

Maria Jose Garcia-Barrado, Maria Carmen Iglesias-Osma, Veronica Moreno-Viedma, Maria Francisca Pastor Mansilla, Silvia Sanz Gonzalez, Jose Carretero, Julio Moratinos, Deborah J. Burks

► To cite this version:

Maria Jose Garcia-Barrado, Maria Carmen Iglesias-Osma, Veronica Moreno-Viedma, Maria Francisca Pastor Mansilla, Silvia Sanz Gonzalez, et al.. Differential Sensitivity To Adrenergic Stimulation Underlies The Sexual Dimorphism In The Development Of Diabetes Caused By Irs-2 Deficiency. *Biochemical Pharmacology*, 2010, 81 (2), pp.279. 10.1016/j.bcp.2010.10.008 . hal-00649886

HAL Id: hal-00649886

<https://hal.science/hal-00649886>

Submitted on 9 Dec 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Differential Sensitivity To Adrenergic Stimulation Underlies The Sexual Dimorphism In The Development Of Diabetes Caused By Irs-2 Deficiency

Authors: Maria Jose Garcia-Barrado, Maria Carmen Iglesias-Osma, Veronica Moreno-Viedma, Maria Francisca Pastor Mansilla, Silvia Sanz Gonzalez, Jose Carretero, Julio Moratinos, Deborah J. Burks

PII: S0006-2952(10)00764-1
DOI: doi:10.1016/j.bcp.2010.10.008
Reference: BCP 10740

To appear in: *BCP*

Received date: 30-6-2010
Revised date: 7-10-2010
Accepted date: 11-10-2010

Please cite this article as: Garcia-Barrado MJ, Iglesias-Osma MC, Moreno-Viedma V, Mansilla MFP, Gonzalez SS, Carretero J, Moratinos J, Burks DJ, Differential Sensitivity To Adrenergic Stimulation Underlies The Sexual Dimorphism In The Development Of Diabetes Caused By Irs-2 Deficiency, *Biochemical Pharmacology* (2010), doi:10.1016/j.bcp.2010.10.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**DIFFERENTIAL SENSITIVITY TO ADRENERGIC STIMULATION
UNDERLIES THE SEXUAL DIMORPHISM IN THE DEVELOPMENT
OF DIABETES CAUSED BY IRS-2 DEFICIENCY**

Abbreviated Title: Female *Irs2*^{-/-} Mice Display Catecholamine Resistance.

Maria Jose Garcia-Barrado¹, Maria Carmen Iglesias-Osma¹, Veronica Moreno-Viedma⁴,
Maria Francisca Pastor Mansilla², Silvia Sanz Gonzalez⁴, Jose Carretero³, Julio Moratinos¹,
Deborah J. Burks⁴

¹ Department of Physiology and Pharmacology, Faculty of Medicine, University of
Salamanca, Salamanca, Spain

² University of Valladolid, Valladolid, Spain

³ Department of Anatomy, Faculty of Medicine, University of Salamanca, Salamanca, Spain

⁴ Regenerative Medicine Program, CIBER de Diabetes y Enfermedades Metabólicas
(CIBERDEM), Centro de Investigación Príncipe Felipe, Valencia, Spain

Key Words: Insulin, Diabetes, Adrenergic Receptors, Lipolysis, cAMP

Address Correspondence to:
Deborah Burks, Ph.D.
Centro de Investigacion Principe Felipe
Avenida del Autopista del Saler, 16
46013 Valencia, Spain
TEL: (34) 963289680
FAX: (34) 963289701
Email: dburks@cipf.es

Disclosure Statement: The authors have no conflicts of interest to disclose.

ABSTRACT

The diabetic phenotype caused by the deletion of insulin receptor substrate-2 (Irs-2) in mice displays a sexual dimorphism. Whereas the majority of male Irs-2^{-/-} mice are overtly diabetic by 12 weeks of age, female Irs-2^{-/-} animals develop mild obesity and progress less rapidly to diabetes. Here we investigated β -cell function and lipolysis as potential explanations for the gender-related differences in this model. Glucose-stimulated insulin secretion was enhanced in islets from male null mice as compared to male WT whereas this response in female Irs-2^{-/-} islets was identical to that of female controls. The ability of α_2 -adrenoceptor (α_2 -AR) agonists to inhibit insulin secretion was attenuated in male Irs2 null mice. Consistent with this, the expression of the α_{2A} -AR was reduced in male Irs-2^{-/-} islets. The response of male Irs-2^{-/-} islets to forskolin was enhanced, owing to increased production of cAMP. Basal lipolysis was increased in male Irs-2^{-/-} but decreased in female Irs-2^{-/-} mice, concordant with the observation that adipose tissue is sparse in males whereas female Irs2 null mice are mildly obese. Adipocytes from both male and female Irs-2^{-/-} were resistant to the anti-lipolytic effects of insulin but female Irs-2^{-/-} fat cells were additionally resistant to the catabolic effects of beta-adrenergic agonists. This catecholamine resistance was associated with impaired generation of cAMP. Consequently, targets of cAMP-dependent protein kinase (PKA) which mediate lipolysis were not phosphorylated in adipose tissue of female Irs-2^{-/-} mice. Our findings suggest that IRS-2 deficiency in mice alters the expression and/or sensitivity of components of adrenergic signaling.

1. INTRODUCTION

The presence of a sexual dimorphism is observed in almost all mouse models of diabetes, including those generated by altering insulin signaling, with males displaying a predominance of the diabetic phenotype [1-4]. However, with the non-obese diabetes (NOD) mouse model of type 1 diabetes, there is a strong tendency for females to develop diabetes, even though both genders present lymphocytic infiltration of the islets [5, 6]. The specific factors which mediate sexual dimorphisms remain largely unexplained in these experimental models of diabetes. Thus, the molecular mechanisms underlying the gender-specific differences which modulate the development and progression of diabetes may provide important clues for the design of anti-diabetic drugs or therapies in human patients.

Various lines of evidence suggest that catecholamines and adrenergic receptors are involved in the etiology and pathogenesis of type 2 diabetes mellitus [7, 8], which is characterized by an insufficient secretion of insulin to compensate for resistance to insulin action in peripheral tissues [9]. Indeed, the plasma concentrations of catecholamines are higher in diabetic than in healthy subjects [10, 11] and glucose intolerance is frequently observed in patients with endocrine disorders such as pheochromocytoma [12, 13]. The anti-secretagogue effect of catecholamines is mediated primarily by the stimulation of α_2 -adrenoceptors (α_2 -AR) in pancreatic beta cells, suggesting that increased sympathetic innervation or a reinforced α_2 -adrenergic response in pancreatic islets may account for the impaired secretory response to glucose observed in certain forms of type 2 diabetes [14, 15]. Adrenaline and other α_2 -adrenoceptor agonists inhibit insulin secretion by a number of mechanisms coupled to the Gi/Go signaling system, including inhibition of adenylate cyclase (AC) and cAMP production [16], activation of K^+ channels [17, 18], and inhibition of L-type voltage-dependent Ca^{+2} channels [19]. Generation of mice deficient in α_2 -AR subtypes has demonstrated that insulin secretion is modulated principally by the α_{2A} AR; the absence of

1 inhibitory pancreatic beta-cell α_{2A} -AR function causes hyperinsulinaemia, reduced blood
2 glucose levels and improved glucose tolerance in knockout mice [20]. Conversely,
3
4 overexpression of α_{2A} -AR specifically in beta cells causes hyperglycemia in response to α_2 -
5 AR agonists due to reduced insulin secretion [21]. A recent study has demonstrated that a
6
7 single-nucleotide polymorphism in the human *ADRA2A* gene is associated with
8
9 overexpression of α_{2A} -AR, reduced insulin secretion, and increased risk for type 2 diabetes
10
11 [22]. Consistent with these findings, a genome-wide association (GWA) project has recently
12
13 linked the *ADRA2A* locus with beta-cell dysfunction in humans with Type 2 diabetes [23].
14
15
16
17
18

19 Given that obesity is closely associated with the development of diabetes, the effects
20
21 of catecholamines on the metabolic parameters of white and brown adipose tissue represent
22
23 another target for intervention in this disease ([24]. The three β -adrenergic receptor (β -AR)
24
25 subtypes (β -AR1, β -AR2, and β -AR3) are coupled to $G_{\alpha s}$ for increasing intracellular cAMP
26
27 levels [25]. In white adipose tissue, lipolysis is regulated by activation of adenylyl cyclase and
28
29 cAMP-dependent protein kinase (PKA) which stimulates lipase, the enzyme that catalyzes the
30
31 breakdown of triacylglycerol into glycerol and free fatty acids (FFA) [26]. By contrast,
32
33 activation of α_2 - adrenoceptors inhibits lipolysis [27]. In most models of obesity, the β -AR
34
35 system is dysfunctional [28] leading to impairments of lipolysis and thermogenesis. However,
36
37 whereas activation of the β -AR subtypes can stimulate lipolysis, the involvement of each
38
39 subtype varies according to fat location, species, gender, age, and degree of obesity [29].
40
41 Interestingly, selective agonists for the β -AR3, the subtype expressed predominantly in
42
43 adipocytes [30], prevent or reverse obesity and accompanying insulin resistance in animal
44
45 models. Whether these agonists represent a viable therapeutic option for human obesity is
46
47 much debated. Nevertheless, the physiological changes in adrenoceptor function associated
48
49 with obesity in rodent models have yielded beneficial insights into β -AR signaling and
50
51 adipocyte physiology.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Members of the insulin receptor substrate (IRS) protein family mediate the physiological effects of insulin and IGF-I [31]. Deletion of IRS-2 produces diabetes in mice owing to peripheral insulin resistance and a reduction in pancreatic β -cell mass [32]. However, this diabetic phenotype displays a sexual dimorphism; male *Irs-2*^{-/-} mice often die of diabetic complications by 12 weeks of age but female *Irs-2*^{-/-} develop a milder form of diabetes and many live up to 6 months [33, 34]. Additionally, female *Irs-2* deficient mice display hyperleptinemia and develop moderate obesity [35, 36], in contrast to male *Irs-2*^{-/-} which are often leaner than control mice. In the present study, we used a pharmacological approach to 1) characterize in greater detail the sexual dimorphism of the *Irs-2*^{-/-} model with respect to beta cell function and lipolysis and 2) to explore the contributions of adrenergic signaling to the diabetic phenotype of *Irs-2* deficient animals.

2. MATERIAL AND METHODS

2.1. Animal Experimentation. The generation and genotyping of mice deficient for *Irs-2*^{-/-} have been described previously [3]. The mice used in the present study were maintained on a C57Bl6 background and were allowed free access to food (irradiated chow, Harlan 20/14) and water during controlled light-dark cycles of 12 hours. All mice were studied at 8-10 weeks of age and any which were overtly diabetic (>220 fed glucose mg/dl or 120 fasted glucose mg/dl) were excluded in order to examine the consequences of *Irs-2*^{-/-} deficiency under conditions which minimize the complications of diabetic metabolism. For routine measurements of fasting glucose and insulin, mice were fasted for 16 hours and a small quantity of tail blood was extracted for immediate analysis by a glucometer (Bayer Elite model, Bayer Healthcare, Barcelona, Spain). Circulating insulin was measured by a mouse ultra-sensitive ELISA (Mercodia, Uppsala, Sweden, <http://www.mercodia.se>). When fed values for insulin and glucose were assessed by these same methods, tail blood was collected consistently between 10 and 11 AM. All experimental procedures were approved by the institutional committee for animal experimentation.

2.2. Adipocyte isolation and lipolysis assay. The details of this experimental procedure have been described previously [37]. Briefly, intra-abdominal white adipose tissue (WAT) of perirenal and epididymal origin was removed from fed animals. Adipose tissue was digested for 35 to 45 min at 37 °C under shaking with 1.5 mg/ml collagenase A (Roche Diagnostics, Mannheim, Germany) in Krebs-Ringer containing 15 mM sodium bicarbonate, 6 mM glucose, 10 mM HEPES, and bovine serum albumin (35 mg/ml) adjusted to pH 7.4 (KRBA buffer). Isolated fat cells were filtered and washed three times in KRBA buffer to eliminate collagenase. The packed cells were diluted in around 10-fold their volume of KRBA, and 500 µl of cell suspension was distributed into plastic incubation vials. Therefore, after 90-min

incubation under gentle shaking at 37 °C with the indicated final concentration of tested drugs, the glycerol released into the medium was assayed and lipolytic activity was expressed as μ moles of glycerol released/100 mg of cellular lipid/90 min. Changes in glycerol release are expressed as percentages of mean control. A one-way analysis of variance (ANOVA) and the Newman–Keuls test were used for statistical analysis. Results were considered significant if $P < 0.05$.

2.3. Islet isolation and measurement of insulin secretion/content. Islets were isolated by collagenase digestion (Immunogenetics, Madrid, Spain) of the pancreas of fed female and male WT and IRS-2^{-/-} mice, followed by manual selection using a dissecting microscope [38]. They were free of visible exocrine contamination. The medium used for islet isolation was a bicarbonate-buffered solution (HCO₃ medium) containing 120 mM NaCl, 4.8 mM KCl, 2.5 mM CaCl₂, 1.2 mM MgCl₂, 5 mM HEPES and 24 mM NaHCO₃. It was equilibrated with O₂-CO₂ (94:6) to maintain a pH of 7.4 and was supplemented with 1 mg ml⁻¹ BSA and 10 mM glucose. After isolation, the islets were pre-incubated for 60 min in HCOS medium containing 15 mM glucose before being distributed into batches of three. Each batch of islets was then incubated for 60 min in 1 ml at 37°C of medium containing glucose and test substances. A portion of the medium was withdrawn at the end of the incubation to measure the insulin concentration. Additionally, the islets were recovered following the incubation and total insulin content was determined after extraction in acid-ethanol [39]. Insulin was measured by a double-antibody RIA (Schering Laboratories, Madrid, Spain). Calculations and statistics were performed with the Instat software (GraphPad, San Diego, CA, USA). Results are provided as mean $\pm$ S.E.M. Statistical significance was assessed by the Student's t-test or the

one-way analysis of variance (ANOVA) corrected by the Newman–Keuls test. Results were considered significant if $P < 0.05$.

2.4. Immunohistochemistry and quantification of beta cell area. Pancreata were removed at the time of sacrifice and fixed for 16 hours in Bouin's solution (Sigma, Steinheim, Germany, <http://www.sigmaaldrich.com>). Subsequently, pancreatic tissue was embedded in paraffin and sections of 5 μ m were prepared. Following re-hydration and permeabilization with 1% Triton X-100, sections were incubated with anti-insulin (Sigma, <http://www.sigmaaldrich.com>) and anti-glucagon (Sigma, <http://www.sigmaaldrich.com>) antibodies overnight at 4°C. Detection was performed with rhodamine and fluorescein conjugated secondary antibodies (Jackson ImmunoResearch, USA, www.jacksonimmuno.com). For quantification of β -cell area, sections were viewed using a Zeiss Axiovert S100 TV microscope at a magnification of 10x. The islet cross-sectional area and total pancreatic area were measured using Openlab Image analysis software (Improvision Imaging). At least 3 sections, separated by 200 μ m were measured per animal. For quantification of the number of islets per area, only islets with more than 5 cells were scored.

2.5. Western Blotting. Abdominal fat depots were collected and frozen immediately in liquid nitrogen. Tissue was lysed in RIPA buffer (NaCl 150mM, Tris 50mM, EDTA 1mM, EGTA 1mM, SDS 0.1%, Sodium deoxycholate 0.5%, NP-40 1%, Na_3VO_4 1mM, NaF 1mM) by polytron and homogenates were clarified by centrifugation at 12,000 $\times$ g for 10 min. Protein determination was by the Biorad assay. 50 μ g of total protein was separated by SDS-PAGE. Gels were transferred to Immun-BlotTM PVDF Membrane (Bio-Rad Laboratories, Hercules CA, www.bio-rad.com) and incubated with one of the following antibodies: rabbit anti-phospho PKA substrate (Cell Signaling, <http://www.cellsignal.com>), anti-perilipin A/B

(Sigma, <http://www.sigmaaldrich.com>), rabbit anti-alpha2A adrenergic receptor (Acris Antibodies, Hiddenhausen, Germany, www.acris-antibodies.com), or anti-hormone sensitive lipase (Santa Cruz, <http://www.scbt.com/>), or rabbit anti-beta tubulin (Sigma, <http://www.sigmaaldrich.com>). Westerns were developed by ECL (Pierce, Thermo Scientific, Rockford IL, <http://www.piercenet.com/>).

2.6. Measurement of Intracellular cAMP Content. Islets were isolated as described above in batches of 100 size-matched islets. Subsequently, the medium was changed to adjust glucose concentrations to 3mM or 15mM, with or without forskolin, in KRBH containing 1mM isobutylmethylxanthine (IBMX) which prevents cAMP degradation by inhibiting cyclic nucleotide phosphodiesterases. Following a 30 min incubation at 37°C, islets were collected by centrifugation and the pellet was frozen immediately in liquid N₂. Adipocytes were isolated and incubated with pharmacological agents as described above. Intracellular cAMP levels were assessed using the AlphaScreen cAMP kit (Perkin-Elmer, Massachusetts, USA <http://las.perkinelmer.com>). Statistical significance was evaluated by the Student's t-test. Results are expressed as mean $\pm$ S.E.M.

3. RESULTS

3.1. Metabolic differences between male and female IRS2-deficient mice cannot be explained by beta cell number or insulin content.

To explore the physiological basis of the sexual dimorphism in the *Irs2* knockout model, we first measured fed blood glucose and insulin levels in mice of 8-10 weeks of age. Fed plasma glucose values in both male and female *Irs2*^{-/-} mice were higher than in their WT controls (Fig. 1A), although the differences were less pronounced in female mice. Mice which were overtly diabetic (>220 fed glucose mg/dl or 120 fasted glucose mg/dl) were excluded from study in order to examine the contributions of *Irs2*^{-/-} deficiency under conditions which

minimize the complications of diabetic metabolism. Both male and female *Irs-2^{-/-}* displayed higher fed insulin values in comparison to their WT controls, although this was statistically significant only in males (Fig. 1B). Consistent with previous reports, the female *Irs2^{-/-}* mice in our study weighed more than their WT controls (*Irs2^{-/-}*: 19.3 grams $\pm$ 2.40, WT: 15.9 g $\pm$ 2.23; $p < 0.05$, $n=10$ mice of each genotype) whereas no differences were observed between the body weights of male transgenic and control mice (*Irs2^{-/-}*: 20.6 grams $\pm$ 2.19, WT: 20.3 g $\pm$ 2.05; $n=10$ mice of each genotype). To determine whether the metabolic differences between male and female IRS2-deficient mice could be explained by differences in the beta cell population, histological analysis was performed. Beta cell mass was reduced in both males and females, although to a slightly lesser extent in females (male *Irs-2^{-/-}*: 40.2% vs. female *Irs-2^{-/-}*: 28.5%) (Fig. 1C). When total pancreas insulin content was evaluated (Figure 1D), it was reduced similarly in male and female *Irs-2^{-/-}* mice (males: WT 1,719 $\pm$ 145 vs. *Irs-2^{-/-}*: 1.287 $\pm$ 44 mIU/islet; females: WT 1,202 $\pm$ 105 vs. *Irs-2^{-/-}* 928 $\pm$ 76 mIU/islet). The integrity of the insulin secretory response to glucose was evaluated by measuring glucose-stimulated insulin release from isolated islets. In response to various concentrations of glucose, islets from *Irs-2^{-/-}* males secreted more insulin than WT controls whereas the response of female *Irs-2^{-/-}* was identical to WT control females (Fig. 1E, F). Thus, the rapid progression of diabetes in males *Irs2* null mice could not be explained by gender differences in beta cell area and insulin content but was associated with defective insulin secretion.

3.2. Secretion of insulin by male *Irs-2^{-/-}* islets displays altered sensitivity to adrenergic agonists and forskolin

Since it is well established that the sympathetic nervous system can modulate insulin secretion, we tested the effect of various types of pharmacological agonists on insulin release under conditions where glucose was maintained at 15 mM. The muscarinic agonist carbachol increased insulin secretion equivalently in islets from all genotypes (Fig. 2A). Brimonidine

(UK 14,304), an agonist for α_2 adrenoceptors, at a dose of 1 μ M reduced insulin secretion as expected in control islets (figure 2A). However, inhibition of insulin secretion by this α_2 agonist in male *Irs2*^{-/-} was significantly less efficient than its effect on WT islets. We reasoned that this decreased sensitivity to brimonidine might reflect down-regulation of α_2 -AR or modifications of cAMP signalling components at the post-receptor level. To test these possibilities, we performed Western blot analysis and, interestingly, observed that expression of α_{2A} -AR was reduced in islets of male *Irs-2*^{-/-} mice (Figure 2B).

Based on these observations, we tested the response of *Irs-2*^{-/-} islets to forskolin, a direct activator of adenylate cyclase which is known to promote insulin secretion by increasing intracellular cAMP. The response of male *Irs-2*^{-/-} islets to forskolin was enhanced in comparison to WT controls, whereas female *Irs-2*^{-/-} islets displayed a slight decrease in the response to this cAMP agonist (Figure 2C). To further explore this altered response, we measured cAMP production in islets stimulated with either glucose or forskolin. The generation of cAMP in islets of male *Irs-2*^{-/-} displayed enhanced sensitivity to both glucose and forskolin (Figure 2D), consistent with the insulin secretion results. In contrast, islets of female *IRS2*-deficient mice produced less cAMP in response to forskolin than female controls. These results suggest that the increased sensitivity of cAMP-generation in islets of male *Irs2*^{-/-} males, combined with reduced expression of α_{2A} -AR, contributes to the dysregulated insulin secretion and hyperinsulinemia observed in these animals.

3.3. Basal and Insulin-inhibited Lipolysis in Males vs. Females

As mentioned previously, female *Irs-2*^{-/-} develop moderate obesity whereas body weight of males is comparable to their WT controls, at least during the pre-diabetic phase. Although *Irs2*^{-/-} is known to have an important role in hypothalamic regulation of appetite and obesity [36, 40], we considered the possibility that adipose metabolism might also contribute

to differences between male and female *Irs-2^{-/-}*. Therefore, we measured glycerol release in adipocytes isolated from male and female mice of both genotypes. Basal lipolysis in male *Irs-2^{-/-}* was increased by 21% whereas it was reduced in females *Irs-2^{-/-}* by 19%, in comparison to their respective WT controls (Figure 3A). When we tested the effects of insulin on lipolysis in isolated adipocytes, this hormone suppressed glycerol release effectively at both concentrations (10^{-8} and 10^{-7} M) in WT animals (Figure 3B, C). In contrast, lipolysis was not inhibited significantly by insulin in male (Figure 3B) or female *Irs-2^{-/-}* adipocytes (Figure 3C), suggesting that these animals are resistant to insulin action. Previous studies have demonstrated that IRS2-deficiency causes peripheral insulin resistance, particularly in liver [41].

3.4. Adipocytes of female *Irs-2^{-/-}* mice display resistance to β -adrenergic-mediated lipolysis

One of the major counter-regulatory mechanisms for controlling lipolysis is the adrenergic system. Catecholamines modulate lipolysis through lipolytic β -adrenoceptor and anti-lipolytic α_2 -adrenoceptor [24]. Given the altered basal lipolysis in male and female *Irs-2^{-/-}*, we examined the effects of adrenergic stimulation on isolated adipocytes. Isoproterenol, a classic β -agonist, increased lipolysis in adipocytes from male WT and *Irs-2^{-/-}* in a dose-dependent manner (Figure 4A). In sharp contrast, adipocytes isolated from female *Irs-2^{-/-}* displayed a blunted response to isoproterenol, particularly at high concentrations (10^{-7} and 10^{-6} M) of this agonist. (Figure 4B). This defective response to β -adrenergic agonists in female *Irs-2^{-/-}* fat cells may reflect an inability to elevate intracellular cAMP, owing either to impaired cAMP generation or an increase to its degradation. We next tested the effects of the α_2 -adrenergic agonist brimonidine on isoproterenol-stimulated lipolysis which would be expected to inhibit given that α_{2A} -AR decreases adenylate cyclase activity. At the highest dose

(10⁻⁶ M) isoproterenol, brimonidine attenuated significantly glycerol release in male and female WT adipocytes (Figure 4C and D). However, in male *Irs-2^{-/-}* fat cells, brimonidine had no effect on isoproterenol-induced lipolysis, consistent with the reduced sensitivity to brimonidine observed in islets of male null mice. The inhibitory effects of brimonidine on lipolysis were also not significant in female *Irs-2^{-/-}* adipocytes but this is most likely explained by the apparent resistance of these cells to isoproterenol.

3.5. Female *Irs-2^{-/-}* adipocytes are less sensitive to forskolin

To investigate whether the resistance to β -adrenergic agonists in female *Irs-2^{-/-}* adipocytes reflects alterations at the receptor or post-receptor level, the effects of forskolin on lipolysis were tested. In male *Irs-2^{-/-}* fat cells, forskolin (10⁻⁵ M) increased basal lipolysis to a greater extent than in WT (40% of control, $p < 0.001$, Figure 5A). However, forskolin was less effective at stimulating lipolysis in adipocytes of female transgenics as compared to female WT controls (20% less glycerol was released when compared with WT, $p < 0.001$, figure 5B).

Given the dampened response to both β -adrenergic agonists and forskolin in female *Irs-2^{-/-}* adipocytes, we next evaluated cAMP generation in response to these agents. Consistent with the lipolysis data, generation of cAMP in adipocytes of male *Irs-2^{-/-}* mice displayed increased sensitivity to forskolin. In contrast, the ability of isoproterenol as well as forskolin to promote cAMP accumulation was attenuated in adipocytes of female *Irs-2^{-/-}* as compared with female controls. Thus, these results reveal a potential explanation for the failure of forskolin and isoproterenol to stimulate lipolysis in adipocytes derived from female *IRS2*-deficient mice.

3.6. cAMP signalling and expression of hormone sensitive lipase are reduced in adipose tissue of female *Irs-2^{-/-}* mice

Perilipin A (Plin) is a major lipid droplet protein that regulates basal and PKA-stimulated lipolysis [42, 43]. Plin is required for the translocation of hormone-sensitive lipase (HSL) from the cytosol to lipid droplets upon stimulation [44]. When catecholamines bind to their receptors and initiate signals that increase cAMP, PKA phosphorylates Plin A which then promotes the translocation of HSL to facilitate maximal lipolysis. Thus, to determine whether the reduced ability to elevate cAMP agents in female *Irs-2*^{-/-} adipocytes has consequences for downstream targets of the lipolytic pathway, we examined the phosphorylation of Plin A/B in adipose tissue lysates by Western blotting with an antibody specific for substrates of PKA. Although total expression levels were similar between *Irs-2* males and WT males as well as between *Irs-2* females and WT females, basal PKA-mediated phosphorylation of Plin A/B was notably reduced in the adipose tissue of *Irs-2* deficient females (Fig. 6A). These results suggest that reduced sensitivity to cAMP-elevating agents impairs the PKA pathway in adipose tissue of female *Irs2* null mice. Additionally, the expression of HSL was notably reduced in adipocytes of female *Irs-2*^{-/-} mice (Fig. 6B). Collectively, these alterations may explain the reduced lipolysis and increased body weight of female *Irs2*-deficient mice.

4. DISCUSSION

The development of diabetes in *Irs-2*^{-/-} null mice has been well-characterized and is attributed to beta cell insufficiency paired with severe insulin resistance [32]. However, the sexual dimorphism associated with this diabetic phenotype is poorly understood. In our study, circulating insulin levels were not significantly different between WT and *Irs2*^{-/-} females. However, male *Irs-2*^{-/-} animals were hyperinsulinemic when compared with their WT controls. Consistent with this, glucose-stimulated insulin secretion was enhanced in islets from male *Irs-2*^{-/-} whereas no difference was observed between female WT and *Irs-2*^{-/-}

animals. The metabolic differences between male and female IRS2-deficient mice could not be explained on the basis of differential defects in the β -cell population and/or insulin content. β -cell mass was reduced in *Irs2* null mice of both genders, though to a slightly lesser extent in females and insulin content was diminished to a similar extent in both male and female transgenic mice.

This compensatory hyperinsulinemia could derive from an adaptive response of the autonomic nervous system to the pathology of pre-diabetes since differences were observed when we analysed the inhibitory effect of the adrenergic agents on insulin secretion. Brimonidine, an α_2 -AR agonist, inhibited insulin secretion less efficiently in male *Irs-2^{-/-}* mice than in their WT controls. Additionally, the expression of this α_{2A} -AR was down-regulated in male but not in female IRS2-deficient islets, consistent with the reduced response to the inhibitory effects of α_2 -AR on insulin release. Since α_{2A} -AR appears to be the main receptor in beta cells and mediates the inhibition of adenylate cyclase, reduced expression of this receptor would be expected to favor increased intracellular levels of cAMP [45, 46]. Therefore, dysregulation of cAMP could explain the enhanced insulin secretory response to both glucose and forskolin observed in islets of male *Irs-2^{-/-}* mice. Indeed, the cAMP-dependent guanine nucleotide exchange factor EPAC potentiates exocytosis by interacting with K^+ ATP and Ca^{+2} voltage channels and by promoting granule fusion events. [47]. Chronic, elevated insulin levels have been reported in α_2 -AR^{-/-} mice and animals deprived of noradrenergic tonic inhibition [20, 48]. Recent studies in humans have suggested that the α_2 -AR may also play an important role in development of diabetes. A single-nucleotide polymorphism in the human *ADRA2A* gene has been correlated with overexpression of α_{2A} -AR, reduced insulin secretion, and increased risk for type 2 diabetes [22]. Consistent with these findings, a GWA study has recently linked the *ADRA2A* locus with beta-cell dysfunction in humans with Type 2 diabetes [23].

Basal lipolysis was significantly increased in male *Irs-2^{-/-}* adipocytes whereas it was markedly reduced in female *Irs-2^{-/-}* in comparison to their WT controls. When we tested the anti-lipolytic effect of insulin on isolated fat cells, glycerol release was suppressed in WT samples but insulin had no significant inhibitory effect on lipolysis in adipocytes of male or female *Irs-2^{-/-}* mice. This results suggest that IRS2 may play a role in regulating insulin sensitivity in adipose tissue. Thus, our lipolysis data confirm the presence of insulin resistance and hyperinsulinemia in male *Irs-2^{-/-}* mice. In female *Irs-2^{-/-}* animals, not only was the basal rate of lipolysis reduced in comparison to WT females but the expected lipolytic response to either isoproterenol or forskolin was clearly attenuated; even at a concentration of 10^{-6} M, the glycerol release evoked by isoproterenol was reduced by approximately 50% in adipocytes of *Irs-2^{-/-}* females. Consistent with this altered lipolytic response, cAMP generation in response to isoprenaline and forskolin was reduced in adipocytes of *Irs-2^{-/-}* females as compared with female controls, suggesting that post-receptor defects may contribute to the catecholamine resistance in these animals. The enhanced lipolytic response to forskolin in adipocytes of male *Irs-2^{-/-}* adipocytes was associated with increased production of cAMP. Given that we detected reduced expression of α_{2A} -AR in islets of male *Irs-2^{-/-}*, it is possible that the absence of an α_2 AR inhibitory effect also increases cAMP levels in male *Irs-2^{-/-}* adipocytes, thereby predisposing them to an increased basal rate of lipolysis.

The observation that PKA-mediated phosphorylation of perilipin is significantly reduced in adipose tissue is consistent with the defective accumulation of cAMP in fat cells of female *Irs-2^{-/-}* mice. Moreover, female *Irs-2^{-/-}* adipocytes display another defect not present in their male counterparts which is a significant reduction in the expression of HSL. In *Irs1^{-/-}* mice, which are lean and resistant to the effects of high-fat diet, HSL expression in adipocytes is enhanced more than 4-fold [49,50], suggesting that IRS1 and IRS2 may differentially regulate expression of lipolytic enzymes. These data provide a molecular explanation for the

development of moderate obesity in female *Irs2*-deficient animals; since phosphorylation of perilipin by PKA is required for the translocation of HSL to lipid droplets, the observed defects in adipocytes of female *Irs-2*^{-/-} would be expected to impair lipolysis and facilitate fat storage. Interestingly, diminished expression of HSL has been observed in patients with obesity and Type 2 diabetes [51, 52]. Studies in humans have also demonstrated that the expression of β -AR2 and the regulatory II β -subunit of PKA are reduced in adipocytes of female patients with polycystic ovary disease (PCOS), a common endocrine disorder characterized by obesity and insulin resistance [53, 54]. These defects are associated with catecholamine resistance and decreased lipolytic activity *in vivo* which promotes obesity in PCOS patients. In addition to the metabolic abnormalities characterized by beta cell failure and insulin resistance, *Irs-2*^{-/-} females are infertile as IRS-2 signals are required for ovarian function and proper regulation of the reproductive axis [36, 55]. Thus, the presence of catecholamine resistance and perturbed cAMP signaling in adipose tissue of *Irs-2*^{-/-} females suggest that these mice may represent a valid model for unravelling the molecular mechanisms underlying the pathology of PCOS.

In summary, the sexual dimorphism described for the diabetic phenotype of *Irs-2*^{-/-} mice can be explained, at least partially, by differential defects in pancreatic islets and adipose tissue. In female *Irs-2*^{-/-} mice at 8-10 weeks of age, reduced basal lipolysis and catecholamine resistance in adipocytes are paired with a normal secretory response to glucose in pancreatic beta cells. Conversely, in male IRS2-deficient mice of the same age, basal lipolysis is enhanced as adipocytes are resistant to both insulin and to the inhibitory effects of α_2 AR agonists, a situation that requires increased secretion of insulin and favours a more rapid progression of diabetes. These metabolic differences between male and female *Irs-2*^{-/-} mice must also reflect some influence of sex steroid hormones. In addition to its role in the physiology of reproduction, estrogen modulates metabolic parameters. For example, estrogen

plays an essential role in adaption to pregnancy by enhancing insulin biosynthesis and glucose-stimulated insulin secretion [56]. However, as both gonadotropins and steroid hormones are reduced in the infertile *Irs2^{-/-}* females [36, 55], it is unlikely that the influence of estrogen and progesterone on gene expression has a major role in delaying the progression of diabetes in females versus males of this experimental model. The metabolic alterations which underlie this sexual dimorphism appear to be associated with differential defects in the cAMP system, including reduced expression of α_2 AR in β -cells of male *Irs-2^{-/-}* and impaired generation of cAMP in adipose tissue of female *Irs-2^{-/-}* mice.

ACKNOWLEDGEMENTS: This work was supported by Instituto de Salud Carlos III-FIS grant PI03/0652, Fundación Mutua Madrileña, CIBER de Diabetes y Enfermedades Metabólicas del Instituto de Salud ISCIII, EMER007 (ISCIII), Ministerio de Ciencia y Innovación SAF2008-00011, Programa de Juan de la Cierva del MICINN (Silvia Sanz), and the Regenerative Medicine Program of the Valencian Community.

REFERENCES

- [1] Kadowaki T. Insights into insulin resistance and type 2 diabetes from knockout mouse models. *J Clin Invest* 2000;106:459-465.
- [2] Oriel RC, Wiley CD, Dewey MJ, Vrana PB. Adaptive genetic variation, stress and glucose regulation. *Dis Model Mech* 2008;1:255-263.
- [3] Withers DJ, Gutierrez JS, Towery H, Burks DJ, Ren JM, Previs S, et al. Disruption of *IRS-2* causes type 2 diabetes in mice. *Nature* 1998;391:900-904.
- [4] Wells JC. Sexual dimorphism of body composition. *Best Pract Res Clin Endocrinol Metab* 2007;21:415-430.
- [5] Amrani A, Durant S, Throsby M, Coulaud J, Dardenne M, Homo-Delarche F. Glucose homeostasis in the nonobese diabetic mouse at the prediabetic stage. *Endocrinology* 1998;139:1115-1124.
- [6] Kikutani H, Makino S. The murine autoimmune diabetes model: NOD and related strains. *Adv Immunol* 1992;51:285-322.
- [7] Ahren B. Evidence that autonomic mechanisms contribute to the adaptive increase in insulin secretion during dexamethasone-induced insulin resistance in humans. *Diabetologia* 2008;51:1018-1024.
- [8] Ahren B. Autonomic regulation of islet hormone secretion--implications for health and disease. *Diabetologia* 2000;43:393-410.

- [9] Redmon JB, Towle HC, Robertson RP. Regulation of human insulin gene transcription by glucose, epinephrine, and somatostatin. *Diabetes* 1994;43:546-551.
- [10] Robertson RP, Halter JB, Porte D, Jr. A role for alpha-adrenergic receptors in abnormal insulin secretion in diabetes mellitus. *J Clin Invest* 1976;57:791-795.
- [11] Christensen NJ. Catecholamines and diabetes mellitus. *Diabetologia* 1979;16:211-224.
- [12] Isles CG, Johnson JK. Pheochromocytoma and diabetes mellitus: further evidence that alpha 2 receptors inhibit insulin release in man. *Clin Endocrinol (Oxf)* 1983;18:37-41.
- [13] Stenstrom G, Sjostrom L, Smith U. Diabetes mellitus in pheochromocytoma. Fasting blood glucose levels before and after surgery in 60 patients with pheochromocytoma. *Acta Endocrinol (Copenh)* 1984;106:511-515.
- [14] Ostenson CG, Cattaneo AG, Doxey JC, Efendic S. Alpha-adrenoceptors and insulin release from pancreatic islets of normal and diabetic rats. *Am J Physiol* 1989;257:E439-443.
- [15] Guillot E, Coste A, Angel I. Alpha 2-adrenoceptors and the regulation of glucose, insulin and amylin levels in diabetic rats. *Life Sci* 1995;57:2081-2090.
- [16] Nakaki T, Nakadate T, Ishii K, Kato R. Postsynaptic alpha-2 adrenergic receptors in isolated rat islets of Langerhans: inhibition of insulin release and cyclic 3':5'-adenosine monophosphate accumulation. *J Pharmacol Exp Ther* 1981;216:607-612.
- [17] Drews G, Debuyser A, Nenquin M, Henquin JC. Galanin and epinephrine act on distinct receptors to inhibit insulin release by the same mechanisms including an increase in K⁺ permeability of the B-cell membrane. *Endocrinology* 1990;126:1646-1653.
- [18] Rorsman P, Bokvist K, Ammala C, Arkhammar P, Berggren PO, Larsson O, et al. Activation by adrenaline of a low-conductance G protein-dependent K⁺ channel in mouse pancreatic B cells. *Nature* 1991;349:77-79.
- [19] Nilsson T, Arkhammar P, Rorsman P, Berggren PO. Inhibition of glucose-stimulated insulin release by alpha 2-adrenoceptor activation is paralleled by both a repolarization and a reduction cytoplasmic free Ca²⁺ concentration. *J Biol Chem* 1988;263:1855-1860.
- [20] Savontaus E, Fagerholm V, Rahkonen O, Scheinin M. Reduced blood glucose levels, increased insulin levels and improved glucose tolerance in alpha2A-adrenoceptor knockout mice. *Eur J Pharmacol* 2008;578:359-364.
- [21] Devedjian JC, Pujol A, Cayla C, George M, Casellas A, Paris H, Bosch F. Transgenic mice overexpressing alpha2A-adrenoceptors in pancreatic beta-cells show altered regulation of glucose homeostasis. *Diabetologia* 2000;43:899-906.
- [22] Rosengren AH, Jokubka R, Tojjar D, Granhall C, Hansson O, Li DQ, et al. Overexpression of alpha2A-adrenergic receptors contributes to type 2 diabetes. *Science* 2010;327:217-220.
- [23] Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, et al. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* 2010;42:105-116.
- [24] Langin D. Adipose tissue lipolysis as a metabolic pathway to define pharmacological strategies against obesity and the metabolic syndrome. *Pharmacol Res* 2006;53:482-491.
- [25] Alexander SP, Mathie A, Peters JA. Guide to Receptors and Channels (GRAC), 3rd edition. *Br J Pharmacol* 2008;153:S1-209.
- [26] Fu L, Isobe K, Zeng Q, Suzukawa K, Takekoshi K, Kawakami Y. The effects of beta(3)-adrenoceptor agonist CL-316,243 on adiponectin, adiponectin receptors and

- tumor necrosis factor- α expressions in adipose tissues of obese diabetic KKAY mice. *Eur J Pharmacol* 2008;584:202-206.
- [27] Fain JN, Garcija-Sainz JA. Adrenergic regulation of adipocyte metabolism. *J Lipid Res* 1983;24:945-966.
- [28] Butler AA, Cone RD. Knockout models resulting in the development of obesity. *Trends Genet* 2001;17:S50-54.
- [29] Arner P. Human fat cell lipolysis: biochemistry, regulation and clinical role. *Best Pract Res Clin Endocrinol Metab* 2005;19:471-482.
- [30] Emorine LJ, Marullo S, Briend-Sutren MM, Patey G, Tate K, Delavie-Klutchko C, Strosberg AD. Molecular characterization of the human beta 3-adrenergic receptor. *Science* 1989;245:1118-1121.
- [31] Kitamura T, Kahn CR, Accili D. Insulin receptor knockout mice. *Annu Rev Physiol* 2003;65:313-332.
- [32] Burks DJ, White MF. IRS proteins and beta-cell function. *Diabetes* 2001;50:S140-145.
- [33] Kubota N, Tobe K, Terauchi Y, Eto K, Yamauchi T, Suzuki R, et al. Disruption of insulin receptor substrate 2 causes type 2 diabetes because of liver insulin resistance and lack of compensatory beta-cell hyperplasia. *Diabetes* 2000;49:1880-1889.
- [34] Hashimoto H, Arai T, Ohnishi Y, Eto T, Ito M, Hioki K, et al. Phenotypes of IRS-2 deficient mice produced by reproductive technology are stable. *Exp Anim* 2007;56:149-154.
- [35] Tobe K, Suzuki R, Aoyama M, Yamauchi T, Kamon J, Kubota N, et al. Increased expression of the sterol regulatory element-binding protein-1 gene in insulin receptor substrate-2(-/-) mouse liver. *J Biol Chem* 2001;276:38337-38340.
- [36] Burks DJ, Font de Mora J, Schubert M, Withers DJ, Myers MG, Towery HH, et al. IRS-2 pathways integrate female reproduction and energy homeostasis. *Nature* 2000;407:377-382.
- [37] Iglesias-Osma MC, Garcia-Barrado MJ, Visentin V, Pastor-Mansilla MF, Bour S, Prevot D, Valet P, Moratinos J, Carpenne C. Benzylamine exhibits insulin-like effects on glucose disposal, glucose transport, and fat cell lipolysis in rabbits and diabetic mice. *J Pharmacol Exp Ther* 2004;309:1020-1028.
- [38] Garcia-Barrado MJ, Ravier MA, Rolland JF, Gilon P, Nenquin M, Henquin JC. Inhibition of protein synthesis sequentially impairs distinct steps of stimulus-secretion coupling in pancreatic beta cells. *Endocrinology* 2001;142:299-307.
- [39] Detimary P, Jonas JC, Henquin JC. Possible links between glucose-induced changes in the energy state of pancreatic B cells and insulin release. Unmasking by decreasing a stable pool of adenine nucleotides in mouse islets. *J Clin Invest* 1995;96:1738-1745.
- [40] Kubota N, Terauchi Y, Tobe K, Yano W, Suzuki R, Ueki K, et al. Insulin receptor substrate 2 plays a crucial role in beta cells and the hypothalamus. *J Clin Invest* 2004;114:917-927.
- [41] Valverde AM, Burks DJ, Fabregat I, Fisher TL, Carretero J, White MF, Benito M. Molecular mechanisms of insulin resistance in IRS-2-deficient hepatocytes. *Diabetes* 2003;52:2239-2248.
- [42] Ducharme NA, Bickel PE. Lipid droplets in lipogenesis and lipolysis. *Endocrinology* 2008;149:942-949.
- [43] Granneman JG. Delivery of DNA into adipocytes within adipose tissue. *Methods Mol Biol* 2008;423:191-195.
- [44] Shen WJ, Patel S, Miyoshi H, Greenberg AS, Kraemer FB. Functional interaction of hormone-sensitive lipase and perilipin in lipolysis. *J Lipid Res* 2009.

- [45] Grill V, Cerasi E. Activation by glucose of adenylyl cyclase in pancreatic islets of the rat. *FEBS Lett* 1973;33:311-314.
- [46] Delmeire D, Flamez D, Hinke SA, Cali JJ, Pipeleers D, Schuit F. Type VIII adenylyl cyclase in rat beta cells: coincidence signal detector/generator for glucose and GLP-1. *Diabetologia* 2003;46:1383-1393.
- [47] Tengholm A, Gylfe E. Oscillatory control of insulin secretion. *Mol Cell Endocrinol* 2009;297:58-72.
- [48] Ste Marie L, Palmiter RD. Norepinephrine and epinephrine-deficient mice are hyperinsulinemic and have lower blood glucose. *Endocrinology* 2003;144:4427-4432.
- [49] Tsuji Y, Kaburagi Y, Terauchi Y, Satoh S, Kubota N, Tamemoto H, et al. Subcellular localization of insulin receptor substrate family proteins associated with phosphatidylinositol 3-kinase activity and alterations in lipolysis in primary mouse adipocytes from IRS-1 null mice. *Diabetes* 2001;50:1455-1463.
- [50] Selman C, Lingard S, Choudhury AI, Batterham RL, Claret M, Clements M, et al. Evidence for lifespan extension and delayed age-related biomarkers in insulin receptor substrate 1 null mice. *FASEB J* 2008;22:807-818.
- [51] Large V, Arner P. Regulation of lipolysis in humans. Pathophysiological modulation in obesity, diabetes, and hyperlipidaemia. *Diabetes Metab* 1998;24:409-418.
- [52] Watt MJ, Carey AL, Wolsk-Petersen E, Kraemer FB, Pedersen BK, Febbraio MA. Hormone-sensitive lipase is reduced in the adipose tissue of patients with type 2 diabetes mellitus: influence of IL-6 infusion. *Diabetologia* 2005;48:105-112.
- [53] Faulds G, Ryden M, Ek I, Wahrenberg H, Arner P. Mechanisms behind lipolytic catecholamine resistance of subcutaneous fat cells in the polycystic ovarian syndrome. *J Clin Endocrinol Metab* 2003;88:2269-2273.
- [54] Ek I, Arner P, Bergqvist A, Carlstrom K, Wahrenberg H. Impaired adipocyte lipolysis in nonobese women with the polycystic ovary syndrome: a possible link to insulin resistance?. *J Clin Endocrinol Metab* 1997;82:1147-1153.
- [55] Neganova I, Al-Qassab H, Heffron H, Selman C, Choudhury AI, Lingard SJ, et al. Role of central nervous system and ovarian insulin receptor substrate 2 signaling in female reproductive function in the mouse. *Biol Reprod* 2007;76:1045-1053.
- [56] Nadal A, Alonso-Magdalena P, Soriano S, Ropero AB, Quesada I. The role of oestrogens in the adaptation of islets to insulin resistance. *J Physiol* 2009;587:5031-5037.

FIGURE LEGENDS

Figure 1. Characterization of beta cell function in male and female mice.

(A) Fed blood glucose levels were measured in animals of 8-10 weeks of age. n= 14 male WT, 14 male *Irs-2^{-/-}*, 16 females WT and 18 females *Irs-2^{-/-}*. *** p< 0.001, ** p< 0.01. (B) Circulating insulin levels from fed animals were measured by ELISA. n= 14 male WT, 14 male *Irs-2^{-/-}*, 16 female WT and 18 female *Irs-2^{-/-}*. **p< 0.01 (C) Quantification of beta cell area. Animals were sacrificed at 8-10 weeks of age. Pancreas sections were stained with anti-

insulin antibodies and beta cell area was determined using serial sections. $n=4$ male WT, 4 male $Irs-2^{-/-}$, 3 female WT and 4 female $Irs-2^{-/-}$. $**p < 0.01$. (D) Total insulin content was determined from isolated islets by radioimmunoassay and plotted as μ units of total insulin content per islet. For the experiment, 25-30 batches of three islets from the experimental groups were matched carefully by size. Data represent the average $\pm$ SEM of 6 WT and 6 $Irs-2^{-/-}$ animals. $**p < 0.01$. (E) Glucose-stimulated insulin release was measured in the presence of 3, 15 and 30 mM of glucose in islets isolated from male or (F) female animals (8-10 weeks of age). Insulin levels were determined by radioimmunoassay and plotted as μ units of insulin per islet/h. $n=9$ males of each genotype and $n=10$ females of each genotype. $***p < 0.001$, $**p < 0.01$. Note: The results of the female $Irs-2^{-/-}$ were identical to female WT control and thus, the lines of the graph in F are superimposed.

Figure 2. Analysis of insulin release in male and female mice. (A) The effects of carbachol (100 μ M) and brimonidine (UK14,304, 1 μ M) on insulin release were tested in islets of male and female mice (8-10 weeks of age). Pancreatic islets were incubated for 60 min in 1 ml of bicarbonate-buffered medium (HCOS medium supplemented with 1 mg/ml of BSA and 5 mM HEPES, see Methods) containing the indicated drug concentrations and 15 mM glucose. Values are expressed as percentages of mean control (100%, 15mM glucose of each animal) $\pm$ SEM of 10-15 batches of three islets from 8-10 animals. $*p < 0.05$. (B) Expression of the α_2A -AR in isolated islets of male mice was analysed by Western blotting. Anti-beta tubulin was used to confirm equal loading of samples. Islets were pooled from three animals of each genotype. (C) The effects of forskolin (FK, 1 μ M) on insulin release were tested in isolated islets as indicated in A. Values are expressed as percentages of mean control (100%, glucose 15mM) $\pm$ SEM of 10-15 batches of three islets from 8-10 animals. $**p < 0.05$. (D) Islets were isolated as described and incubated in HCOS medium (see Methods) containing 3 mM

glucose. Subsequently, batches of 100 size-matched islets from each genotype were selected. Islets were then stimulated with either forskolin or 15 mM glucose for 10 minutes as described in C. The stimulation with forskolin was performed in triplicate and samples were lysed to measure intracellular cAMP. $**p < 0.01$, $*p < 0.05$.

Figure 3. Basal glycerol release in isolated adipocytes and inhibition by insulin. (A)

Following isolation of adipocytes from abdominal fat, basal lipolytic activity in wild-type vs. *Irs2*^{-/-} mice (8-10 weeks of age) was measured. Values are expressed as means $\pm$ SE obtained from the analysis of the following groups of mice: male WT (n = 29), male *Irs2*^{-/-} (n = 28), female WT (n = 46) and female *Irs2*^{-/-} (n = 41). Data are expressed relative to basal lipolysis, which was set as 100% in WT animals of each gender (absolute values for basal glycerol released in WT mice were 0.34 ± 0.05 and 0.42 ± 0.02 $\mu\text{mol}/100$ mg lipid/90 min for male and female, respectively). $**p < 0.01$ and $*p < 0.05$. (B and C) Anti-lipolytic effect of insulin on basal glycerol release. Increasing concentrations of insulin (10^{-8} and 10^{-7} M) were added to the incubation media and the percentage of inhibition of glycerol release from adipocytes of male (B) and female (C) mice was measured. $**p < 0.01$ and $*p < 0.05$.

Figure 4. Analysis of adrenergic-regulated lipolysis in adipocytes.

(A and B) Analysis of isoproterenol-induced glycerol release. Fat cells from WT and *Irs2*^{-/-} male (A) and female (B) mice of 8-10 weeks of age were incubated 90 min with increasing concentrations (10^{-9} – 10^{-6} M) of the β -adrenergic agonist, isoproterenol (Iso). Basal lipolysis was considered as 100% in WT animals of each gender. Data are presented as means $\pm$ SE (see Fig. 3A for absolute values in WT mice). $*p < 0.05$ and $**p < 0.01$. n= 6 mice of each experimental group. (C and D) Effects of the $\alpha 2$ -adrenergic agonist brimonidine (UK 14,304) on isoproterenol-induced glycerol release from mouse adipocytes. Values are expressed as

mean $\pm$ SE relative to the percentage of WT basal values (100% of WT of each gender). * p <0.05 and ** p <0.01. n = 4 mice of each experimental group.

Figure 5. cAMP-mediated glycerol liberation in isolated mouse adipocytes.

(A and B) Effects of 10 μ M forskolin (FK) on basal lipolysis. Basal lipolysis was established as 100% in WT animals of each gender. *** p <0.001. n = 4 animals of each experimental group. (C) Triplicates of isolated adipocytes from each experimental group were stimulated with the indicated concentrations of either isoproterenol or forskolin for 10 min. Samples were then lysed and assayed for intracellular cAMP. n = 6 mice of each experimental group. * P <0.05 and ** P <0.01.

Figure 6. Analysis of cAMP-dependent signaling in adipose tissue. (A) Western analysis of perilipin A/B expression and phosphorylation. Visceral adipose tissue was removed and protein lysates were prepared from four animals of each genotype. 500 μ g of total protein were immunoprecipitated (from pools of two animals of each experimental group) using anti-perilipin A/B and these immunocomplexes were then separated by SDS-PAGE. The membrane was first incubated with anti-phospho PKA substrate to detect phosphorylated perilipin A/B. Subsequently, the blot was stripped and re-probed to reveal total levels of perilipin A/B. Blots were scanned and phospho-perilipin A (approximately 62 kDa) was quantified by densitometry. Total perilipin A/B was used to normalize the levels of phospho-perilipin A/B. A representative blot from two independent experiments is presented. Total n = 8 animals of each group. **(B)** Western analysis of hormone-sensitive lipase expression in adipose tissue. 50 μ g of the adipose tissue lysates prepared in A were probed with anti-HSL antibodies. Anti-beta actin was used to confirm equal protein loads. Molecular weight is indicated at the left of each membrane.

Accepted Manuscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

A **B**

Gender Differences in IRS2 Model of Diabetes