Characterisation of hepatitis B virus in Turkish blood donors, and the prevalence of the SP1 splice variant.

Laura Elizabeth Cox, Onder Arslan, Jean-Pierre Allain

► To cite this version:

Laura Elizabeth Cox, Onder Arslan, Jean-Pierre Allain. Characterisation of hepatitis B virus in Turkish blood donors, and the prevalence of the SP1 splice variant.. *Journal of Medical Virology*, 2011, 83 (8), pp.1321. 10.1002/jmv.22118 . hal-00652148

HAL Id: hal-00652148

<https://hal.science/hal-00652148>

Submitted on 15 Dec 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Characterisation of hepatitis B virus in Turkish blood donors, and the prevalence of the SP1 splice variant.

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-10-2247.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	01-Apr-2011
Complete List of Authors:	Cox, Laura; University of Cambridge, Haematology Arslan, Onder; Ankara University, Haematology Allain, Jean-Pierre; University of Cambridge, Haematology
Keywords:	Turkey, HBV, sequencing, splicing

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Characterisation of hepatitis B virus in Turkish blood donors, and the prevalence of the SP1 splice variant.

Laura E. Cox¹
Önder Arslan²
Jean-Pierre Allain¹

¹ Division of Transfusion Medicine, Department of Haematology, University of Cambridge, Cambridge Blood Centre, Long Road, Cambridge CB2 0PT, UK. (Institution where work was performed)

² Ankara University, School of Medicine, Department of Haematology, Blood Bank & Apheresis Unit, Cebeci, 06590, Ankara, Turkey.

Correspondence to:

Laura Cox: Email: lec48@cam.ac.uk

Jean-Pierre Allain: Email: jpa1000@cam.ac.uk

Division of Transfusion Medicine, Department of Haematology, University of Cambridge, Cambridge Blood Centre, Long Road, Cambridge CB2 0PT, UK.

Running title: HBV characterisation in Turkish blood donors

Key words: Turkey, HBV, sequencing, splicing

Abstract

Hepatitis B is a disease of the liver that can manifest acutely, or persist chronically as a result of infection with the hepatitis B virus (HBV). Turkey has a moderate endemicity level of HBV infection, and all data published to date has shown this to be of genotype D, predominantly of subgenotype D1. However the sequences of very few full genomes have been published. The aim of this study was to characterise the molecular profile of hepatitis B virus in asymptomatic, first-time Turkish blood donors. These results confirm that genotype D, subgenotype D1 is the most prevalent HBV strain in Turkey, accounting for 94% of cases. Subgenotypes D2 and D3 were present as minority strains (4% and 2%, respectively). A singly-spliced HBV variant that is capable of forming defective HBV particles and has been associated with apoptosis and activation of T-cell responses was also detected in 52.5% of samples screened, co-circulating with wild type genomes.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Hepatitis B virus (HBV) carrier prevalence in Turkey is reported to be around 4% [Degertekin and Gunes, 2008], which is in line with other Mediterranean and Middle Eastern countries and meets the criteria for intermediate endemicity level. HBV belongs to the family Hepadnaviridae, and has a 3.2kb genome that, when encapsidated within the virion, is a relaxed circular DNA molecule that is only partially double-stranded. It was first reported in 1989 that HBV could undergo splicing by the host's machinery in human hepatoma cells transfected with full-length HBV [Su et al., 1989]. This 2kb variant is commonly termed SP1, and contains a 1223 nucleotide deletion beginning at the last codon of the core gene and spanning through to the middle of the S gene. To date, this splice variant has been identified in chronic hepatitis B patients infected with HBV genotypes A and D, and an additional 10 splice variants have been identified in patients infected with genotypes A, B and D [Gunther et al., 1997].

HBV is divided into eight well-documented genotypes (A-H), some of which are further divided into two or more sub-genotypes [Kay and Zoulim, 2007]. Recently, two putative new genotypes, I and J, have been reported [Tatematsu et al., 2009; Tran et al., 2008]. Genotype D is predominant in the Mediterranean basin, and is also frequently found in populations from Europe, Africa and Asia. Of the HBV genotypes, this is the least defined geographically, which may be a result of diverging earlier than the other genotypes [Norder et al., 2004]. The majority of the current data on HBV in Turkey has come from patients with clinical Hepatitis B, with few full genome sequences published. Sequencing of full genomes and the S gene from patients with chronic HBV infection identified only genotype D in Turkish samples, and where subgenotyping was documented, D1 was the most prevalent subgenotype, with a minority of subgenotype D2 strains present [Bozdayi et al., 2005; Sayiner et al., 2008; Serin et al., 2005; Sertoz et al., 2008]. The aim of this study was to characterise HBV strains obtained from HBsAg-positive plasma

samples from asymptomatic, first-time, replacement blood donors collected in Ankara, Turkey, using serological and molecular techniques.

Materials & Methods

HBsAg-positive plasma samples were collected during 3 separate periods in 2004, 2005 and 2006, with a total collection period of 13 months. Samples were heat-inactivated at 60°C for 30 min prior to shipping on dry ice, and upon arrival in the UK were stored at -80°C until ready to use. Viral DNA was extracted and quantified as described previously [Allain et al., 2003; Garmiri et al., 2009]. To generate sequence for the entire HBV genome from 50 samples, the Expand High Fidelity PCR system (Roche) was used for two nested PCR assays. Firstly, a nearly full-length fragment of approximately 3000bp, minus a 50bp precore section was amplified as previously described [Zahn et al., 2008]. A second 300bp fragment including the basic core promoter/precore (BCP/PC) 50bp gap was produced to complete the full genome [Candotti et al., 2006]. PCR products were purified using the EZNA cycle pure kit (Omega; Crawley, UK).

A semi-nested PCR was used to screen for the SP1 variant, using newly designed primers located in the core and S genes. SP1_core (5'-GTCGCAGAAGATCTCAATCTCGGG-3'; position 2421–2444) was used as the sense primer in both first and second round PCR; SP1_R1 (5'-ATACAAAGGCATTAATGCAGGGTA-3'; position 1065–1042) and SP1_R2 (5'-TGTGTAAATGGAGCGGCAAAGCC-3'; position 1012–1034) were used as the anti-sense primers in first and second round PCR, respectively. Using the Roche Expand High Fidelity system, a 50µL reaction mixture contained: 1X PCR buffer II (including MgCl₂; final concentration 2.5mM), 0.8 mM dNTPs, 0.3 µM each primer, and 5.25 U HiFi. For samples with a viral load of $\times 10^8$ IU ml⁻¹ or above, 3µl of viral DNA was used as first-round template; 5µl was used for samples with viral loads of $\times 10^4$ – $\times 10^7$ IU ml⁻¹ and 10µL for samples $\times 10^3$ IU ml⁻¹ or below. Six microlitres of the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

84 first-round reaction was loaded into the second-round PCR. An identical touchdown cycling
85 programme was used for both rounds, and consisted of: an initial denaturation step at 94°C for
86 5 minutes, followed by 10 cycles of 94°C/40s; 65°C/45s (reduced by 1°C each cycle); 72°C/1
87 min, and a following 30 cycles of 94°C/40s; 55°C/45s; 72°C/1 min. A final extension step was
88 performed at 72°C for 10 minutes. In the presence of full-length HBV DNA (3.2kb) a 1.8kb
89 product is amplified, in comparison to a 580bp product for the SP1 variant, visualised using
90 agarose gel electrophoresis.

91 Sequences with interesting deletions or insertions were examined further by fresh full
92 genome or BCP/PC PCR reactions. Following addition of a 3' single adenosine overhang to the
93 PCR product, amplicons were ligated into either the pCR2.1 TOPO® TA vector (BCP/PC) or the
94 TOPO® XL vector (Invitrogen; Paisley, UK) as per the manufacturer's instructions. Following
95 transformation of chemically competent *E. coli* cells, plasmid DNA was isolated using a QIAprep
96 spin miniprep kit (Qiagen), and a minimum of 10 clones were sequenced.

97 Sequencing and phylogenetics analysis was performed using Seqman, SeqBuilder
98 (DNASTAR) and MacVector software. PAUP* version 4.0 beta 10 was used to calculate genomic
99 distances. Microsoft Excel and Prism (version 4) were used for analysis of HBV DNA and HBsAg
100 load.

101 Antibodies against HBcAg (anti-HBc) and HBsAg (anti-HBs) were tested by ELISA with a
102 Monolisa® anti-HBc PLUS kit (Bio-Rad; Hemel Hempstead, UK) and a Monolisa® anti-HBs PLUS kit
103 (Bio-Rad), respectively.

104
105 **Results**

106 Over the collection period samples were taken from 20,960 first-time, replacement
107 blood donors (18,898 male: 2,062 female). Routine screening identified 200 samples as positive

108 for HBsAg (187 male: 13 female), giving a prevalence of 1% for males and 0.6% for females.
109 Plasma was obtained for 199 HBsAg positive samples, and quantification of HBV DNA load by Q-
110 PCR was successful in 176 (88.4%) of these samples. The median viral load was 1.19×10^3 IU ml⁻¹
111 (range 1.62×10^1 - 8.95×10^8 IU ml⁻¹). In the remaining 23 samples (11.6%), HBV DNA was
112 undetected by Q-PCR. However amplification of the BCP/PC region was achieved by nested-PCR
113 in one of the 23 samples, meaning detection of HBV DNA was possible for a total of 177/199
114 samples (89%).

115 Complete genomic sequences were obtained for 50 samples (GenBank Accession
116 numbers: JF754586-JF754635), from 57 randomly selected samples in which this was attempted.
117 Phylogenetic analysis identified all strains as belonging to HBV genotype D; 47 clustered as
118 subgenotype D1 (94%), 2 as subgenotype D2 (4%), and 1 as subgenotype D3 (2%). Sequence
119 analysis of the S gene revealed the predominant serotype to be *ayw2* (*n*=41), 5 strains specified
120 *ayw3* and 1 strain was *ayw4*. An additional strain belonged to serotype *adw*, but could not be
121 classified further, and serotype determination was not possible in one strain due to a deletion in
122 this region (Table I).

123 124 *PreC/C*

125 One sample contained a deletion of 37 nucleotides, resulting in a frameshift mutation
126 and premature protein truncation. In another strain, insertion of two adenine nucleotides in the
127 precore region caused a frameshift mutation, which is predicted to result in the inclusion of 31
128 amino acids in the precore protein, whilst leaving the core protein unaffected. Cloning of the
129 BCP/PC for this strain identified the double adenine insertion in all 10 clones sequenced. Serum
130 testing of these two samples identified the presence of anti-HBc. Another sample contained a
131 single adenine insertion within the core region, introducing a premature stop codon and protein

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

132 truncation. Serum was not available for anti-HBc testing of this sample. Seven samples (14%)
133 contained mutations that abolished the precore initiation codon. The G1896A mutation, which
134 introduces a premature stop codon and abolishes precore protein and HBeAg synthesis, was
135 present in 33 strains (66%).

136

137 *Pre-S/S*

138 Mutations disrupting the initiation codon of Pre-S2 were identified in 3 samples (6%).
139 Five samples contained deletions in the Pre-S/S transcript, ranging in size from 6 to 51
140 nucleotides. These were present throughout the three ORFs. Three samples contained nonsense
141 mutations that are predicted to result in premature truncation of the protein. One further
142 sample contained a 66 nucleotide deletion spanning the Pre-S1/Pre-S2 boundary, in addition to
143 a nonsense mutation leading to premature truncation of the S protein at amino acid residue 69.
144 Testing for the presence of anti-HBs in the serum from which this strain was obtained was
145 negative. Cloning of this sample identified the presence of wild-type species, in addition to a
146 much smaller product with a 1223 nucleotide deletion (positions 2449-489), predicted to result
147 from splicing of viral mRNA. This deletion started at the final codon of the core gene and
148 extended into the middle of the S gene.

149 The mean overall amino acid divergence within the Pre-S/S region was 2.7% (range 0-
150 6.9%). The major hydrophilic region (positions 100-169) was more variable, with a mean amino
151 acid substitution rate of 5.3% (range 0-20%). No substitutions associated with vaccine-escape
152 mutants were identified in this cohort.

153

154 *X gene*

155 No deletions were present within the X transcript in any samples. However, one strain
156 contained a 6 nucleotide insertion, leading to the inclusion of two additional amino acid residues
157 (glycine and methionine) in the X protein, inserted between wild-type residues 79 and 80. The
158 basal core promoter (BCP) variants, A1762T and G1764A were present as a dual mutation in 10
159 strains, with an additional 6 samples carrying G1764A as a single change.

160

161 *Pol*

162 Six strains carrying deletions in the polymerase gene ranging between 6-66 nucleotides
163 were identified; these were all located in the spacer domain of the protein. There were no
164 disruptions to the highly conserved YMDD motif in any of the 50 strains sequenced. One further
165 sample had a 6 nucleotide insertion within the RNase H domain of the protein.

166

167 *SP1 screening*

168 Screening for the presence of the SP1 variant was successful in 40/43 samples in which
169 this was attempted (Figure 1). A full genome sequence was available for 27 of the 40
170 successfully amplified samples. The SP1 variant was present in 10/10 (100%) samples with viral
171 loads $\geq 10^6$, in 8/18 (44%) samples with viral loads of $\geq 10^4 - < 10^6$, and in 3/12 (25%) of samples
172 with viral loads $< 10^4$.

173

174 **Discussion**

175 Analysis of 50 full HBV genotypes extracted from asymptomatic Turkish blood donors
176 selected at random supports previous findings that the most prevalent HBV strain in this country
177 is subgenotype D1 and serotype *ayw2* [Bozdayi et al., 2005; Sertoz et al., 2008].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In previously published complete HBV genomes from chronic Hepatitis B patients from Turkey, the G1896A variant was identified in 18% of samples, and these patients were HBeAg negative [Bozdayi et al., 2005]. In this cohort from healthy volunteer donors, the G1896A precore stop variant was identified in 66% of samples. This is in keeping with data that shows a high prevalence of the G1896A mutation in samples from Mediterranean populations [Funk et al., 2002]. G1896A has previously been documented to be present in 15% of HBeAg-positive Turkish patients, compared to 85% of HBeAg-negative patients [Bozdayi et al., 1999; Bozdayi et al., 2005]. This mutation is of interest as it results in the prevention of HBeAg synthesis, and seroconversion of HBeAg to anti-HBe is correlated with a decline in viral load [Tedder et al., 2002]. The core promoter dual mutations (A1762T, G1764A) were identified previously in 3/11 chronic-HBV infected patients, one of whom was HBeAg positive [Bozdayi et al., 2005], and was identified in 20% of the strains sequenced in this study. The double mutation has been implicated with a reduction in the generation of precore mRNA and HBeAg production, and is more prevalent in patients with advanced liver disease [Poustchi et al., 2008]. Given the relatively high frequency of these mutations in this study population, and their clinical implications, it may be of use to sequence the viral genome in subjects testing positive for HBV-infection in Turkey.

This is the first time that screening of the SP1 variant has been conducted in asymptomatic blood donors. The data demonstrated that the SP1 variant was present in more than half of the samples screened (21/40), and was ubiquitous in samples with viral loads $\geq 10^6$. However, that SP1 is detected at a much lower frequency in samples with low viral loads (3/10 VL $< 10^4$ IU/mL) may not be directly linked to viral load, but may be a matter of the ratio at which the transcripts are present and the ability of the PCR to detect very low levels of the spliced variant. Previous studies have examined the prevalence in symptomatic HBV-infected

patients, and have correlated its presence with chronic infection and hepatocellular carcinoma [Lin et al., 2002]. The association between prevalence of the HBV splice variant and severity of liver disease is not yet elucidated; one study reported an association between a high proportion of spliced HBV: wild-type HBV in patients with more severe liver necroinflammation [Soussan et al., 2008], whereas a different study found no association between spliced HBV prevalence and disease status [Preiss et al., 2008]. Decreased levels of SP1 have been observed in patients with a lamivudine-resistant HBV strain, suggesting that antiviral therapy disrupts the balance between circulating HBV splice variants and full-length HBV genomes [Preiss et al., 2008].

In addition, a novel 10.4kDa protein (HBV splice-generated protein; HBSP) arose from the fusion of a section of the polymerase and a new ORF created downstream of the splicing event [Soussan et al., 2000]. Detection by western blot analysis has revealed the presence of this protein in 46% of serum samples from chronic HBV carriers, as opposed to <1% of healthy HBV-negative controls [Soussan et al., 2003]. Functionally, this protein has been demonstrated *in vitro* to increase hepatocyte apoptosis [Soussan et al., 2000], and in humans and transgenic mice to induce a T-cell response [Mancini-Bourgine et al., 2007].

Acknowledgements

The authors would like to thank Mrs Yeşim Özer and the staff of the Blood Bank & Apheresis Unit, Ankara, Turkey who participated in the collection and screening of blood donor samples, in addition to Dr B.H.M. Meldal and Dr I.H.A. Barnes for their contribution to DNA extraction and quantification. This study was supported by grant BS05/2 from the National Health Service Blood and Transplant.

References:

Allain JP, Candotti D, Soldan K, Sarkodie F, Phelps B, Giachetti C, Shyamala V, Yeboah F, Anokwa M, Owusu-Ofori S, Opare-Sem O. 2003. The risk of hepatitis B virus infection by transfusion in Kumasi, Ghana. *Blood* 101(6):2419-2425.

Bozdayi AM, Bozkaya H, Turkyilmaz A, Aslan N, Verdi H, Kence A, Uzunlimoglu O. 1999. Polymorphism of precore region of hepatitis B virus DNA among patients with chronic HBV infection in Turkey. *Infection* 27(6):357-360.

Bozdayi G, Turkyilmaz AR, Idilman R, Karatayli E, Rota S, Yurdaydin C, Bozdayi AM. 2005. Complete genome sequence and phylogenetic analysis of hepatitis B virus isolated from Turkish patients with chronic HBV infection. *J Med Virol* 76(4):476-481.

Candotti D, Opare-Sem O, Rezvan H, Sarkodie F, Allain JP. 2006. Molecular and serological characterization of hepatitis B virus in deferred Ghanaian blood donors with and without elevated alanine aminotransferase. *J Viral Hepat* 13(11):715-724.

Degertekin H, Gunes G. 2008. Horizontal transmission of hepatitis B virus in Turkey. *Public Health* 122(12):1315-1317.

Funk ML, Rosenberg DM, Lok AS. 2002. World-wide epidemiology of HBeAg-negative chronic hepatitis B and associated precore and core promoter variants. *J Viral Hepat* 9(1):52-61.

Garmiri P, Loua A, Haba N, Candotti D, Allain JP. 2009. Deletions and recombinations in the core region of hepatitis B virus genotype E strains from asymptomatic blood donors in Guinea, west Africa. *J Gen Virol* 90(Pt 10):2442-2451.

Gunther S, Sommer G, Iwanska A, Will H. 1997. Heterogeneity and common features of defective hepatitis B virus genomes derived from spliced pregenomic RNA. *Virology* 238(2):363-371.

Kay A, Zoulim F. 2007. Hepatitis B virus genetic variability and evolution. *Virus Res* 127(2):164-176.

Lin X, Wen Y, Wan D, Qian G, Gu J. 2002. Structural and functional analysis of 2.2 kb spliced variant of hepatitis B virus genomes isolated from liver tissues from hepatocellular carcinoma patients. *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 16(1):11-15.

Mancini-Bourguine M, Bayard F, Soussan P, Deng Q, Lone YC, Kremsdorf D, Michel ML. 2007. Hepatitis B virus splice-generated protein induces T-cell responses in HLA-transgenic mice and hepatitis B virus-infected patients. *J Virol* 81(10):4963-4972.

Norder H, Courouce AM, Coursaget P, Echevarria JM, Lee SD, Mushahwar IK, Robertson BH, Locarnini S, Magnus LO. 2004. Genetic diversity of hepatitis B virus strains derived worldwide: genotypes, subgenotypes, and HBsAg subtypes. *Intervirology* 47(6):289-309.

Poustchi H, Mohamadkhani A, Bowden S, Montazeri G, Ayres A, Revill P, Farrell GC, Locarnini S, George J, Malekzadeh R. 2008. Clinical significance of precore and

- core promoter mutations in genotype D hepatitis B-related chronic liver disease. *J Viral Hepat* 15(10):753-760.
- Preiss S, Littlejohn M, Angus P, Thompson A, Desmond P, Lewin SR, Sasadeusz J, Matthews G, Dore GJ, Shaw T, Sozzi V, Yuen L, Lau G, Ayres A, Thio C, Avihingsanon A, Ruxrungtham K, Locarnini S, Revill PA. 2008. Defective hepatitis B virus DNA is not associated with disease status but is reduced by polymerase mutations associated with drug resistance. *Hepatology* 48(3):741-749.
- Sayiner AA, Ozcan A, Sengonul A. 2008. Naturally occurring MHR variants in Turkish patients infected with hepatitis B virus. *J Med Virol* 80(3):405-410.
- Serin MS, Akkiz H, Abayli B, Oksuz M, Aslan G, Emekdas G. 2005. Genotyping of hepatitis B virus isolated from chronic hepatitis B patients in the south of Turkey by DNA cycle-sequencing method. *Diagn Microbiol Infect Dis* 53(1):57-60.
- Sertoz RY, Erensoy S, Pas S, Ozacar T, Niesters H. 2008. Restriction fragment length polymorphism analysis and direct sequencing for determination of HBV genotypes in a Turkish population. *New Microbiol* 31(2):189-194.
- Soussan P, Garreau F, Zylberberg H, Ferray C, Brechot C, Kremsdorf D. 2000. In vivo expression of a new hepatitis B virus protein encoded by a spliced RNA. *J Clin Invest* 105(1):55-60.
- Soussan P, Pol J, Garreau F, Schneider V, Le Pendeven C, Nalpas B, Lacombe K, Bonnard P, Pol S, Kremsdorf D. 2008. Expression of defective hepatitis B virus particles derived from singly spliced RNA is related to liver disease. *J Infect Dis* 198(2):218-225.
- Soussan P, Tuveri R, Nalpas B, Garreau F, Zavala F, Masson A, Pol S, Brechot C, Kremsdorf D. 2003. The expression of hepatitis B spliced protein (HBSP) encoded by a spliced hepatitis B virus RNA is associated with viral replication and liver fibrosis. *J Hepatol* 38(3):343-348.
- Su TS, Lai CJ, Huang JL, Lin LH, Yauk YK, Chang CM, Lo SJ, Han SH. 1989. Hepatitis B virus transcript produced by RNA splicing. *J Virol* 63(9):4011-4018.
- Tatematsu K, Tanaka Y, Kurbanov F, Sugauchi F, Mano S, Maeshiro T, Nakayoshi T, Wakuta M, Miyakawa Y, Mizokami M. 2009. A genetic variant of hepatitis B virus divergent from known human and ape genotypes isolated from a Japanese patient and provisionally assigned to new genotype J. *J Virol* 83(20):10538-10547.
- Tedder RS, Ijaz S, Gilbert N, Barbara JA, Corden SA, Gilson RJ, Boxall EH. 2002. Evidence for a dynamic host-parasite relationship in e-negative hepatitis B carriers. *J Med Virol* 68(4):505-512.
- Tran TT, Trinh TN, Abe K. 2008. New complex recombinant genotype of hepatitis B virus identified in Vietnam. *J Virol* 82(11):5657-5663.
- Zahn A, Li C, Danso K, Candotti D, Owusu-Ofori S, Temple J, Allain JP. 2008. Molecular characterization of occult hepatitis B virus in genotype E-infected subjects. *J Gen Virol* 89(Pt 2):409-418.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legends:

Table I: Molecular and serological characteristics of HBV strains from Turkish blood donors.

Summary of the features identified in the HBV strains sequenced, including deletions, truncations and insertions in the four ORFs, and viral load and SP1 distribution. (HBV: hepatitis B virus)

Figure 1: Representative gel of SP1 screening. Amplification of full-length sequence results in a 1802bp product, whereas spliced HBV DNA, which contains a 1223 nucleotide deletion, generates a 579bp product. Depending on the ratios present, both products may be amplified or alternatively only one amplicon is generated. Sample 1: TK89; Sample 2: TK88 (JF754601); Sample 3: TK129 (JF754618); Sample 4: TK32; Sample 5: TK28 (JF754624); Sample 6: TK74; Sample 7: TK171 (JF754623); Sample 8: TK178; Sample 9: TK63 (JF754608); Sample 10: TK38 (JF754591); Sample 11: TK70). (Marker: Bioline, Hyperladder I; HBV: hepatitis B virus)

325 Table I

Feature	Results				
	(No. of strains in which feature present/No. of strains screened)				
Genotype	D1 (47/50)	D2 (2/50)	D3 (1/50)		
Serotype	adw (1/50)	ayw2 (41/50)	ayw3 (5/50)	ayw4 (1/50)	Unknown (2/50)
X ORF	Deletions (0/50)	Insertions (1/50) (6bp)	Truncations (0/50)		
Pol ORF	Deletions (6/50) (6-66bp)	Insertions (1/50) (6bp)	Truncations (0/50)		
Pre-S/S ORF	Deletions (6/50) (6-66bp)	Insertions (0/50)	Truncations (4/50) (12-474bp)		
Pre-C/C ORF	Deletions (1/50) (37bp)	Insertions (2/50) (1-2bp)	Truncations (2/50) (369- 417bp)	G1896A (33/50)	
Viral load (VL) (IU/mL)	Undetectable (23/199)	<100 (26/199)	$1 \times 10^2 - 9 \times 10^3$ (120/199)	$1 \times 10^4 - 9 \times 10^6$ (22/199)	$>1 \times 10^7$ (8/199)
Splice variant (SP1) prevalence	$VL < \times 10^4$ (3/12)	$VL \geq \times 10^4 -$ $< \times 10^6$ (8/18)	$VL > \times 10^6$ (10/10)		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

330 **Figure 1**

331

For Peer Review